

ORAL PRESENTATION

Open Access

Phototoxicity—from molecular descriptors to classification models

Friedemann Schmidt^{1*}, Andreas Evers¹, Alexander Amberg¹, Gerhard Hessler¹, Catherine Robles², Karl-Heinz Baringhaus¹

From 6th German Conference on Chemoinformatics, GCC 2010
Goslar, Germany. 7-9 November 2010

Potential photoactivation of certain pharmaceuticals, cosmetic ingredients and natural products by sunlight (e.g., phenothiazines, arylsulfonamides, or coumarins) has to be considered early on in development in order to avoid serious adverse effects (for example phototoxic or photoallergic reactions). Current clinical trial registration guidelines (FDA May 2003 [1], EMEA Dec. 2002 [2]) recommend photosafety testing of molecules if they exhibit strong absorption bands between 290-700nm and if they are significantly partitioned in human skin or eyes.

The UV absorption coefficients and the tissue partitioning of a compound are considered as important factors for phototoxic effects. However, the rationalization and prediction of phototoxicity by (quantitative) structure-property relationships ((Q)SPR) offers a valuable strategy to reduce experimental testing if an appropriate precision level of the underlying model is guaranteed.

A diverse data set of known phototoxicants and non-phototoxicants including various molecular chemotypes (90 % of them are pharmaceuticals) was compiled. After geometry optimization the maximum absorption wavelength of each compound was calculated by semi-empirical methods followed by subsequent computation of molecular descriptors. Our *insilico* analysis (e.g., PLS and recursive partitioning) of quantum chemical as well as classical molecular descriptors (e.g., LUMO, HOMO/LUMO gap, electron affinity, ionization energy, molecular fragments, physicochemical descriptors such as logD, pK_a and logP_{eff}) has led to predictive photosafety classifiers. Model validation was performed with a proprietary external test set of an *in vitro* photosafety assay (3T3 neutral red assay).

Our photosafety models are currently applied in a prospective manner in the prioritization, classification and labeling of newly designed molecules.

Author details

¹Sanofi-Aventis, R&D Frankfurt, Germany. ²Sanofi-Aventis, R&D Montpellier, France.

Published: 19 April 2011

References

1. Food and Drug Administration: 5600 Fishers Lane, Rockville, MD 20857, <http://www.fda.gov/cder/guidance/index.htm>.
2. The European Agency for the Evaluation of Medicinal Products: 7 Westferry Circus, Canary Wharf, London E14 4HB, <http://www.fda.gov/cder/guidance/>.

doi:10.1186/1758-2946-3-S1-O10

Cite this article as: Schmidt et al.: Phototoxicity—from molecular descriptors to classification models. *Journal of Cheminformatics* 2011 **3** (Suppl 1):O10.

Publish with **ChemistryCentral** and every scientist can read your work free of charge

“Open access provides opportunities to our colleagues in other parts of the globe, by allowing anyone to view the content free of charge.”

W. Jeffery Hurst, The Hershey Company.

- available free of charge to the entire scientific community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
<http://www.chemistrycentral.com/manuscript/>


ChemistryCentral

* Correspondence: Friedemann.Schmidt@sanofi-aventis.com

¹Sanofi-Aventis, R&D Frankfurt, Germany

Full list of author information is available at the end of the article