

Interaction of *Medicago truncatula* Lysin Motif Receptor-Like Kinases, NFP and LYK3, Produced in *Nicotiana benthamiana* Induces Defence-Like Responses

Anna Pietraszewska-Bogiel¹, Benoit Lefebvre^{2,3}, Maria A. Koini¹, Dörte Klaus-Heisen^{2,3}, Frank L. W. Takken⁴, René Geurts⁵, Julie V. Cullimore^{2,3}, Theodorus W.J. Gadella^{1*}

1 Section of Molecular Cytology, Swammerdam Institute for Life Sciences, University of Amsterdam, Amsterdam, The Netherlands, **2**INRA, Laboratoire des Interactions Plantes-Microorganismes (LIPM), UMR441, F-31326 Castanet-Tolosan, France, **3**CNRS, Laboratoire des Interactions Plantes-Microorganismes (LIPM), UMR2594, F-31326 Castanet-Tolosan, France, **4** Section of Plant Pathology, Swammerdam Institute for Life Sciences, University of Amsterdam, Amsterdam, The Netherlands, **5** Department of Plant Science, Laboratory of Molecular Biology, Wageningen University, Wageningen, The Netherlands

Abstract

Receptor(-like) kinases with Lysin Motif (LysM) domains in their extracellular region play crucial roles during plant interactions with microorganisms; e.g. *Arabidopsis thaliana* CERK1 activates innate immunity upon perception of fungal chitin/chitooligosaccharides, whereas *Medicago truncatula* NFP and LYK3 mediate signalling upon perception of bacterial lipo-chitooligosaccharides, termed Nod factors, during the establishment of mutualism with nitrogen-fixing rhizobia. However, little is still known about the exact activation and signalling mechanisms of MtNFP and MtLYK3. We aimed at investigating putative molecular interactions of MtNFP and MtLYK3 produced in *Nicotiana benthamiana*. Surprisingly, heterologous co-production of these proteins resulted in an induction of defence-like responses, which included defence-related gene expression, accumulation of phenolic compounds, and cell death. Similar defence-like responses were observed upon production of AtCERK1 in *N. benthamiana* leaves. Production of either MtNFP or MtLYK3 alone or their co-production with other unrelated receptor(-like) kinases did not induce cell death in *N. benthamiana*, indicating that a functional interaction between these LysM receptor-like kinases is required for triggering this response. Importantly, structure-function studies revealed that the MtNFP intracellular region, specific features of the MtLYK3 intracellular region (including several putative phosphorylation sites), and MtLYK3 and AtCERK1 kinase activity were indispensable for cell death induction, thereby mimicking the structural requirements of nodulation or chitin-induced signalling. The observed similarity of *N. benthamiana* response to MtNFP and MtLYK3 co-production and AtCERK1 production suggests the existence of parallels between Nod factor-induced and chitin-induced signalling mediated by the respective LysM receptor(-like) kinases. Notably, the conserved structural requirements for MtNFP and MtLYK3 biological activity in *M. truncatula* (nodulation) and in *N. benthamiana* (cell death induction) indicates the relevance of the latter system for studies on these, and potentially other symbiotic LysM receptor-like kinases.

Citation: Pietraszewska-Bogiel A, Lefebvre B, Koini MA, Klaus-Heisen D, Takken FLW, et al. (2013) Interaction of *Medicago truncatula* Lysin Motif Receptor-Like Kinases, NFP and LYK3, Produced in *Nicotiana benthamiana* Induces Defence-Like Responses. PLoS ONE 8(6): e65055. doi:10.1371/journal.pone.0065055

Editor: James Porter, University of North Dakota, United States of America

Received: January 24, 2013; **Accepted:** April 20, 2013; **Published:** June 4, 2013

Copyright: © 2013 Pietraszewska-Bogiel et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by the European Community Marie Curie Research Training Network Programme through contract MRTN-CT-2006-035546 "NODPERCEPTION". Work in Toulouse was supported by the French National Research Agency (ANR) through contracts "NodBindsLysM", "SYMPASIGNAL" and "LCDinNONLEGUMES", and has been done as part of the Laboratoire d'Excellence (LABEX) entitled TULIP (ANR-10-LABX-41). FLWT acknowledges the Centre for BioSystems Genomics and Netherlands Genomics Initiative program. RG acknowledges the Dutch Science Organisation NWO (VIDI 864.06.007). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: Th.W.J.Gadella@uva.nl

Introduction

Legumes can establish a mutualism with compatible rhizobia ultimately leading to nodulation, i.e. a formation of specialized symbiotic organs (nodules) in which atmospheric dinitrogen is converted into ammonia by the bacteria in exchange for plant carbohydrates. Nod factors (NFs) play a central role during most *Rhizobium*-legume (RL) symbioses [1]. They are secreted rhizobial signals whose perception by host legume roots is required for root nodule organogenesis, invasion of rhizobia toward a nodule primordium, and accommodation of bacteria inside nodule cells [2–4]. In two model legumes, NF-induced responses during the pre-infection step of RL interaction require *M. truncatula* (*Medicago*)

Nod Factor Perception (*MtNFP*), and *Lotus japonicus* (*Lotus*) *Nod Factor Receptor 1* and *5* (*LjNFR1* and *LjNFR5*) [5–11]. At a later step, *LjNFR1*, *LjNFR5*, *MtNFP*, and an additional *Medicago* gene, *LysM domain-containing Receptor-Like Kinase/Root Hair Curling* (*MtLYK3/MtHCL*), are required for rhizobial infection via so-called infection threads through which the bacteria penetrate nodule primordia [9,12–15]. Additionally, *MtNFP* and *MtLYK3* might co-function during nodule development and/or accommodation of rhizobia inside nodule cells [9,16–18]. Recently demonstrated binding of NF derivatives to *LjNFR1* and *LjNFR5* confirmed their role as NF receptors [19], whereas the exact mechanism of *MtNFP* and *MtLYK3* activation by compatible NFs remains to be shown.

All four genes encode receptor-like kinases (RLKs) with an extracellular region (ExR) predicted to contain three LysM domains, a transmembrane helix, and a protein kinase domain (KD) within the intracellular region (InR) [5,20–22]. Remarkably, in contrast to MtLYK3 and LjNFR1, which both display kinase activity, MtNFP and LjNFR5 seem to function as pseudokinases that neither show nor rely on the intrinsic kinase activity to signal [9,20–22]. LjNFR5 is hypothesized to form a receptor complex with LjNFR1: a notion consistent with their demonstrated co-functioning during the determination of RL specificity [23]. Similarly, a receptor complex composed of MtNFP and a yet-unidentified LysM-RLK or MtLYK3 is predicted to initiate the pre-infection responses and the infection process, respectively [12,15]. Since mutagenesis studies in *Medicago* have not identified alterations in genes other than *MtNFP* that lead to complete lack of responsiveness to NFs, a function of this additional LysM-RLK in the pre-infection stage is most likely redundant. In addition, MtNFP has been implicated in *Medicago* interactions with pathogens (*Aphanomyces euteiches* and *Colletotrichum trifolii*), and with beneficial arbuscular mycorrhiza (AM) fungi [24–27]. However, it remains to be shown whether MtNFP functions in these processes alone or in co-operation with (an)other RLK(s).

LysM-RLKs in non-legume species also govern plant-microbe interactions. An *MtNFP/LjNFR5* homolog in *Parasponia andersonii*, *PaNFP*, is involved in interactions with *Sinorhizobium* sp. NGR234 and *Rhizophagus irregularis* (formerly *Glomus intraradices*), resulting in nitrogen-fixing and arbuscular mycorrhiza symbiosis, respectively [28]. *Arabidopsis thaliana* (*Arabidopsis*) *LysM-RLK1/CERK1* (*Chitin Elicitor Receptor Kinase 1*) and its ortholog from rice (*Oryza sativa*), OsCERK1, are essential for microbe-associated molecular pattern (MAMP)-triggered immunity. MAMPs are specific molecules conserved in various classes of microorganisms that activate receptor-mediated defence signalling [29–30]. CERK1-mediated innate immunity to fungal and bacterial pathogens is activated upon perception of chitin/chitooligosaccharides (COs), or peptidoglycan (PGN), respectively [31–35]. In the latter case, both in rice and in *Arabidopsis* PGN binds not to OsCERK1/AtCERK1 but to extracellular LysM domain-containing proteins, termed LYPs or LYMs [35–36]. This in turn is postulated to induce a formation of AtCERK1/AtLYMs receptor complexes, and subsequent signal transduction via the kinase activity of AtCERK1. A similar mechanism operates during COs-induced signalling in rice, involving OsCERK1 and (a) LYP protein(s) [34,36], whereas in *Arabidopsis* COs bind directly to AtCERK1 [37–38]. Therefore, modes of CERK1 activation, even upon perception of the same MAMP, can differ between plant species.

We are interested in NF-induced signalling mediated by MtNFP and MtLYK3, focusing on their postulated interaction *in situ*. However, our attempts to visualize these proteins in *Medicago* root have been unsuccessful, presumably due to stringent regulation of their accumulation (even in the situation of an attempted overproduction). *Nicotiana benthamiana* (*Nicotiana*) has proved to be a useful model for heterologous production and structure-function studies on multiple proteins, providing invaluable insights that guided their subsequent analyses in the respective homologous systems [39]. Therefore, we employed an *Agrobacterium tumefaciens* (*Agrobacterium*)-mediated transient transformation of *Nicotiana* leaves [40], which allowed us to produce both proteins to levels suitable for fluorescence microscopy. Remarkably, we found that heterologous co-production of MtNFP and MtLYK3 resulted in the induction of defence-like responses that are typically observed upon treatment with pathogen-derived molecules. As the apparent (functional) interaction of these LysM-RLKs in *Nicotiana* activated defence-like responses, similar to these mediated by AtCERK1,

our results indicate the existence of parallels between NF-induced and COs-induced signalling.

Materials and Methods

Constructs for Plant Expression

The cDNA of *MtNFP*, *MtLYK3*, and *MtDMI2* (in the latter case the first intron was included in the cDNA), and the genomic sequence of *AtCERK1* were PCR amplified and cloned in a pMON999 vector containing a CaMV 35Sp:: (*sYFP2*, *mCherry*, *3xFLAG* or empty) 35S terminator cassette. The stop codon was removed from the coding sequences during cloning (except when generating untagged *MtNFP* and *MtLYK3* constructs) to allow a translational fusion. Sequences of the primers and linkers are given in Table S1. All point mutations were introduced as described [20]. All constructs were sequenced to verify the correct insert sequence. Constructs generated in pMON999 vector were subsequently re-cloned into a pBin+ (all *MtNFP* constructs) or pCambia1390 (all *MtLYK3*, *AtCERK1*, and *MtDMI2* constructs) vector using HindIII and SmaI sites. *MtNFP*ΔInR, *MtNFP-YFP_N*, *MtLYK3-YFP_N*, *MtLRR1.1-YFP_C*, *AtBRI1-YFP_C* (where *YFP_N* or *YFP_C* encode, respectively, the N- and C-terminal part of split YFP used in BiFluorescence Complementation assay) constructs are described [22,41].

Nicotiana Transformations

Agrobacterium tumefaciens GV3101::pMP90 and LBA4404 strains were transformed with the respective constructs via electroporation. The LBA4404 strain was used only in the experiments that compared the effect of different *Agrobacterium* strains on cell death (CD) induction upon MtNFP and MtLYK3 co-production. All results presented in Figures 1–5, Figure S1 and S3, and Tables 1–3 were obtained with *Agrobacterium* GV3101::pMP90 strain. *Agrobacterium*-mediated transformation of *Nicotiana* was performed essentially as described [42], except that *Agrobacterium* cultures were grown in LB medium supplemented with 25 µg/mL of rifampicin and 50 µg/mL of kanamycin. Resuspended cells were incubated at room temperature for at least 1 h before being infiltrated into fully expanded leaves of green house-grown plants using needleless syringes. *Agrobacterium* transformants carrying the respective construct were resuspended in the infiltration medium to desired OD₆₀₀: all *MtNFP* and *MtNFP*ΔInR-*sYFP2* constructs - OD₆₀₀ = 0.4; all *MtLYK3* and *AtCERK1* constructs - OD₆₀₀ = 0.7; *MtDMI2-sYFP2* - OD₆₀₀ = 1.0. Then, they were mixed 1:1 with GV3101::pMP90 transformants carrying: pCambia1390 vector with an empty CaMV 35Sp::35S terminator cassette (for separate expression), a desired *MtNFP* construct or a desired *MtLYK3* construct before being infiltrated into *Nicotiana* leaves. All experiments included mock infiltration with GV3101::pMP90 transformants carrying pCambia1390 vector with an empty CaMV 35Sp::35S terminator cassette, and a positive control (co-expression of WT *MtNFP-FP* and WT *MtLYK3-FP* constructs). Cell death induction upon separate expression or co-expression of each (pair of) constructs was analyzed between 24 and 72 hai in at least three independent experiments, every time using three different plants. In case of no macroscopic symptoms, three leaves were stained with Evans blue to confirm the lack of CD.

To confirm efficient accumulation of MtLRR1.1-YFP_C and AtBRI1-YFP_C fusions in *Nicotiana*, *Agrobacterium* transformants carrying the respective constructs were co-infiltrated at high optical densities (final OD₆₀₀ = 0.5) with *Agrobacterium* transformants carrying either *MtNFP-YFP_N* or *MtLYK3-YFP_N* constructs (final OD₆₀₀ = 0.5). The observed complementation of YFP

fluorescence reported on efficient accumulation, and even unspecific oligomerization of the respective encoded fusions.

Stereoscopic Analysis

Blue light-excitabile autofluorescence and far-red chlorophyll autofluorescence in intact *Nicotiana* leaves were imaged using 430/40 excitation and 485/50 emission BP filters, or 480/40 BP excitation and 510 LP emission filters, respectively. Images were captured using CMOS USB DCC1645C camera (THORlabs, Newton NJ, USA) implemented on a Leica MZ FLIII stereoscope. Evans blue staining was performed as described [42]. Leaves were cleared by boiling in acidic lactophenol/ethanol solution (10 g phenol in 10 ml lactic acid, mixed 2:1 with 96% ethanol) until the complete removal of chlorophyll (approximately 3 min per leaf). Ethanol-insoluble autofluorescence was excited with 312 nm wavelength. Images were captured using a Cool Snap CF camera (Photometrix, Tucson AZ, USA).

qRT-PCR Analysis

RNA extraction and qRT-PCR were performed as described [17] except that cDNA was prepared from 500 ng of total RNA (see Table S1 for primer sequences). Two technical replicates from two biological replicates were analyzed and results were collated.

Medicago Transformations

Complementation of *Mtlyk3-1* mutant seedlings was performed as described [20] using *MtLYK3-3xFLAG*, *MtLYK3*[K464A]-*3xFLAG*, and *MtLYK3*[T480A]-*sYFP2* constructs driven by the CaMV 35S promoter. Results were scored as: + (>75% of plants nodulated), reduced (<50% of plants nodulated) or - (0 plants nodulated).

Results

Co-production of MtNFP and MtLYK3 in *Nicotiana* Leaves Induces Cell Death

MtNFP and *MtLYK3* cDNA sequences were fused at their 3' ends to the sequence encoding a fluorescent protein (FP); either a super yellow fluorescent protein 2 (sYFP2) or mCherry [43–44], and were expressed from a CaMV 35S promoter in *Nicotiana* leaves, where they were delivered by *Agrobacterium*-mediated transformation. These and similar *MtNFP* and *MtLYK3* constructs were shown to complement *Mtnfp* and *Mtlyk3* mutants, respectively [17–18,22], and are therefore suitable for studying the encoded LysM-RLKs. Confocal laser-scanning microscopy analysis demonstrated co-localization of both MtNFP-sYFP2 and MtLYK3-sYFP2 fusions with a plasma membrane (PM) marker (the hyper-variable region [HVR] of maize [*Zea mays*] ROP7 fused to the C-terminus of mCherry; [20,22]), hence indicating PM localization of MtNFP and MtLYK3 fusions in *Nicotiana* leaf epidermal cells. Surprisingly, co-infiltration of *Agrobacterium* transformants carrying *MtNFP-FP* or *MtLYK3-FP* constructs, leading to the co-production of the encoded fusions, resulted in collapse and subsequent desiccation of the infiltrated region within 48 hours after infiltration (hai) (Fig. 1A), regardless of the *Agrobacterium* strain used (i.e. GV3101::pMP90 or LBA4404; unpublished data). This cell death (CD) response was not dependent on the tag attached to either protein, since an identical response was observed upon co-production of FP-tagged, 3xFLAG-tagged or untagged MtNFP and MtLYK3 (Fig. 1A, Table 1). Importantly, production of the separate MtNFP or MtLYK3 (fusions) did not induce CD as confirmed with an exclusion dye (Evans blue) staining (Fig. 1A), which reflects compromised membrane permeability attributed with cell death.

To investigate whether a similar CD response could be triggered by heterologous production of other plant RLKs, we analysed the *Nicotiana* response to expression of *Medicago Doesn't Make Infection 2* (*DMI2*; [45]), *MtLRR1.1* [41], and *Arabidopsis Brassinosteroid Insensitive 1* (*BRI1*; [46]), all driven by the CaMV 35S promoter. Notably, none of these RLKs, alone or in combination with either MtNFP or MtLYK3 fusions, induced CD (Fig. 1B, Table 1), despite being efficiently produced in *Nicotiana* leaves, as confirmed with fluorescence microscopy (see Materials & Methods). Thus, the *Nicotiana* CD response was not a general response to a heterologous production of RLKs but rather a specific response to MtNFP and MtLYK3 co-production.

Production of AtCERK1 also Induces Cell Death in *Nicotiana* Leaves

A rapid tissue collapse at the site of pathogen attack, termed the hypersensitive response (HR), is frequently observed in incompatible plant-pathogen interactions where it is thought to contribute to pathogen restriction and to generate a signal that activates systemic plant defence mechanisms [47–48]. The apparent phenotypic similarity of the CD response to MtNFP and MtLYK3 co-production with the HR elicited by various pathogen-derived components (MAMPs and so-called effectors) [49–50], prompted us to investigate whether co-production of the symbiotic LysM-RLKs might activate defence signalling similar to that mediated by LysM-RLKs functioning in innate immunity. AtCERK1 mediates signalling upon the perception of COs or PGN, although, to our knowledge, CD induction in response to these MAMPs has not been reported so far in any plant species. We therefore investigated the *Nicotiana* response to heterologous production of wild-type (WT) AtCERK1 or its kinase-inactive variant carrying a substitution in the catalytic lysine (Lys 349 in a kinase subdomain II). Both *AtCERK1* and *AtCERK1*[K349E] constructs were generated as fusions to the 5' end of the *sYFP2* sequence, and their expression in *Nicotiana* leaves was driven by the CaMV 35S promoter.

Notably, heterologous production of AtCERK1-sYFP2 fusion resulted in tissue collapse and desiccation of (nearly) the entire infiltrated region in 20 out of 22 infiltrations 36 hai (Fig. 2A). This CD induction abolished our attempts of precisely characterizing the subcellular localization of AtCERK1 fusion in *Nicotiana* leaf epidermal cells, although we could detect sYFP2 fluorescence at the cell boundary (unpublished data). On the contrary, we observed clear co-localization of AtCERK1[K349E]-sYFP2 fusion with the PM marker using confocal laser-scanning microscopy analysis (Fig. S1). The PM localization of AtCERK1[K349E] fusion in *Nicotiana* leaf epidermal cells is in agreement with the reported subcellular localization of AtCERK1 fluorescent fusion in onion (*Allium cepa*) epidermal cells [31]. Importantly, production of the kinase-inactive variant of AtCERK1 did not result in CD induction, as confirmed with Evans blue staining (Fig. 2B), indicating that biological activity of AtCERK1 in *Nicotiana* leaves was dependent on its kinase activity.

Cell Death Induction Upon MtNFP and MtLYK3 Co-production, and AtCERK1 Production in *Nicotiana* Leaves Requires an Influx of Extracellular Ca²⁺

An influx of extracellular Ca²⁺ causes an increase in the cytosolic [Ca²⁺] that is required for MAMP (including COs)-induced activation of a MAPK cascade, ROS production, and gene expression. Thus, Ca²⁺ influx is postulated to occur very early in the plant defence signalling pathway [51–52], possibly immediately upon the activation of the PM-localised MAMP




Figure 1. Co-production of MtNFP and MtLYK3 induces cell death in *Nicotiana* leaves. A, The following *MtNFP* and *MtLYK3* constructs were expressed alone or co-expressed in *Nicotiana* leaves: mock infiltration (1); *MtNFP* untagged+*MtLYK3* untagged (2); *MtNFP-sYFP2*+*MtLYK3-sYFP2* (3); *MtLYK3-sYFP2* (4); *MtLYK3* untagged (5); *MtNFP-sYFP2* (6); *MtNFP* untagged (7). Macroscopic observation (left panel) and subsequent Evans blue staining (right panel) are depicted 48 hai. Bar is 1 cm. B, *MtDMI2-sYFP2* construct was expressed alone or co-expressed with either *MtNFP-mCherry* or *MtLYK3-mCherry* construct in *Nicotiana* leaves. Macroscopic observations (left panel) and subsequent Evans blue stainings (right panel) are depicted 48 hai. Bars are 1 cm.

doi:10.1371/journal.pone.0065055.g001

receptors [53]. We wanted to know whether an influx of extracellular Ca^{2+} was similarly involved in CD induction upon *MtNFP* and *MtLYK3* co-production or separate production of *AtCERK1*. To this end, *MtNFP-3xFLAG* and *MtLYK3-3xFLAG* fusions or *AtCERK1-3xFLAG* fusion were (co-)produced in adjacent regions in *Nicotiana* leaves. Twelve hours later, parts of the infiltrated leaf regions were syringe-infiltrated with 5 mM lanthanum chloride (an established inhibitor of the PM-localized calcium channels) or water, and the CD development was monitored between 24 and 72 hours after the first infiltration (with *Agrobacterium*). Notably, in 24 out of 30 leaf regions co-producing *MtNFP* and *MtLYK3* fusions, compromised membrane permeability and tissue collapse were first (i.e. between 36 and 42 hai) localized only (or mostly) outside the lanthanum chloride-treated regions (Fig. 3A). Later on (i.e. 60 hai), 26 out of 30 parts of leaf regions treated with lanthanum chloride showed confluent death of the entire infiltrated region (unpublished data). Similar delay of the CD development was observed 33 hai in 11 out of 21 leaf regions producing *AtCERK1* fusion and treated with lanthanum chloride (Fig. 3C). On the contrary, control

treatment with water did not affect the development of confluent CD upon (co-)production of *MtNFP* and *MtLYK3* fusions or *AtCERK1* fusion (Fig. 3B, D).

Cell Death Upon *MtNFP* and *MtLYK3* Co-production, and *AtCERK1* Production in *Nicotiana* Leaves is Associated with an Induction of Defence-like Responses

Subsequently, we investigated whether co-production of *MtNFP* and *MtLYK3* or production of *AtCERK1* in *Nicotiana* leaves was associated with an accumulation of phenolic compounds and/or induction of defence-related gene expression, two established hallmarks of plant defence response, including that induced by COs and/or PGN [54–56]. We started by analysing the kinetics of CD development. To this end, *Agrobacterium* transformants carrying *MtNFP-3xFLAG*, *MtLYK3-3xFLAG* or *AtCERK1-3xFLAG* construct were (co-) infiltrated at different time-points in adjacent circles in *Nicotiana* leaves, and CD development was monitored between 24 and 48 hai. In case of co-production of *MtNFP* and *MtLYK3* fusions, macroscopic symptoms of CD were first observed around

Table 1. Cell death induction upon (co-)expression of various RLK-encoding genes in *Nicotiana* leaves.

Construct	Cell death induction		
	Separate expression	Co-expression with <i>MtNFP-FP</i> #	Co-expression with <i>MtLYK3-FP</i> #
<i>MtNFP-sYFP2</i>	0/12	Not applicable	20/22
<i>MtNFP-3xFLAG</i>	0/9	Not applicable	12/13*
<i>MtNFP</i>	0/9	Not applicable	8/9**
<i>MtLYK3-sYFP2</i>	0/12	20/22	Not applicable
<i>MtLYK3-3xFLAG</i>	0/9	12/13*	Not applicable
<i>MtLYK3</i>	0/9	8/9**	Not applicable
<i>MtDMI2-sYFP2</i>	0/12	0/12	0/12
<i>MtLRR1.1-YFP_C</i>	0/9	0/9***	0/9***
<i>AtBRI1-YFP_C</i>	0/7	0/9***	0/9***

– unless stated differently: with -3xFLAG (*) untagged (**), or -YFP_N (***) tagged construct.

Indicated constructs were expressed alone or co-expressed with either *MtNFP* or *MtLYK3* in *Nicotiana* leaves, and the infiltrated regions were marked. Macroscopic symptoms of cell death were scored 48 hai: only infiltrations that resulted in confluent death of (nearly) the entire infiltrated region were scored and are presented as a fraction of total infiltrations performed.

doi:10.1371/journal.pone.0065055.t001




Figure 2. Production of AtCERK1 in *Nicotiana* leaves induces cell death that requires AtCERK1 kinase activity. *AtCERK1-sYFP2* (A) and *AtCERK1*[K349E]-*sYFP2* (B) constructs were expressed in *Nicotiana* leaves. Macroscopic observations (left panel) and subsequent Evans blue stainings (right panel) are depicted 36 hai. Macroscopic symptoms of cell death were scored 36 hai: only infiltrations that resulted in confluent death of (nearly) the entire infiltrated region were scored and are presented (right panel) as a fraction of total infiltrations performed. Bars are 1 cm.

doi:10.1371/journal.pone.0065055.g002

36 hai (Fig. 4A) as a type of flaccidity and the appearance of small patches of collapsed tissue (these were more pronounced on the abaxial side of the leaf). Forty-eight hai, 30 out of 31 infiltrations showed pronounced tissue desiccation of the entire infiltrated region (Fig. 1A). Compromised membrane permeability preceded tissue collapse and often occurred over the entire infiltrated region approximately 33 hai (Fig. 4A). Compromised membrane permeability of leaf regions producing AtCERK1 fusion was observed already approximately 27–30 hai, and pronounced macroscopic symptoms of CD developed 36 hai (Fig. 2A, Fig. 3 C, D).

In addition, co-production of MtNFP-3xFLAG and MtLYK3-3xFLAG fusions resulted in accumulation of blue light-excitable autofluorescence (Fig. 4B) approximately 36 hai. This was not observed upon separate production of either fusion, or upon co-production of MtNFP-3xFLAG and kinase-inactive MtLYK3[G334E]-3xFLAG fusions (unpublished data). Accumulation of ethanol/lactophenol-inextractable and UV-excitable autofluorescence, indicative of phenolic compounds, was detected approximately 36 hai and 30 hai in leaf regions (co-)producing MtNFP-3xFLAG and MtLYK3-3xFLAG fusions or AtCERK1-3xFLAG fusion, respectively (Fig. 4C). Mock infiltration, separate production of MtNFP-3xFLAG or MtLYK3-3xFLAG fusion, production of kinase-inactive AtCERK1[K349E]-3xFLAG fusion or co-production of MtNFP-3xFLAG and kinase-inactive MtLYK3[G334E]-3xFLAG fusions did not result in the accumulation of similar autofluorescence (Fig. 4C).

Subsequently, we investigated induction of defence-related genes expression in *Nicotiana* leaves in response to: separate production and co-production of MtNFP-3xFLAG, MtLYK3-3xFLAG, MtLYK3[G334E]-3xFLAG, and AtCERK1-3xFLAG




Figure 3. Lanthanum chloride delays the cell death development upon MtNFP and MtLYK3, or AtCERK1 (co-)production.

Agrobacterium transformants carrying the following constructs were (co-)infiltrated at a final concentration: OD_{600} [MtNFP-3xFLAG] = 0.125 and OD_{600} [MtLYK3-3xFLAG] = 0.2 (A, B); OD_{600} [AtCERK1-3xFLAG] = 0.2 (C, D). Twelve hai parts of the infiltrated regions were syringe-infiltrated with 5 mM lanthanum chloride (circled in red) or water (circled in white). Macroscopic observations (left panel) and subsequent Evans blue stainings (right panel) are depicted 42 hai for leaf regions co-producing MtNFP and MtLYK3 fusions (A, B), and 33 hai for leaf regions producing AtCERK1 fusion (C, D). Cell death development was scored 42 hai (A, B) or 33 hai (C, D): only infiltrations that showed the lack of tissue collapse and no compromised membrane permeability in the lanthanum chloride- or water-treated region were scored and are presented (right panel) as a fraction of total infiltrations performed. Bars are 1 cm.

doi:10.1371/journal.pone.0065055.g003

fusion(s). Induction of: *NbHIN1*—a postulated marker gene for HR [50]; two *PR1* genes, i.e. *NbPR1a acidic* and *NbPR1 basic* [57]; and *NbACRE31*, *NbACRE132*, and *NbCYP71D20*—postulated marker genes for MAMP-triggered immunity [49] was analyzed 24 hai




Figure 4. MtNFP and MtLYK3, or AtCERK1 (co-)production in *Nicotiana* leaves induces defence-like responses. A, Kinetics of cell death development in *Nicotiana*. *Agrobacterium* transformants carrying either *MtNFP-3xFLAG* or *MtLYK3-3xFLAG* construct were co-infiltrated into *Nicotiana* leaves at five different time points (1–5). Macroscopic observation (left panel) and subsequent Evans blue staining (right panel) are depicted 42 hai (region 1), 39 hai (region 2), 36 hai (region 3), 33 hai (region 4) and 30 hai (region 5). Mock infiltration (region 6) was done concomitantly with the infiltration of region 1. Bar is 1 cm. B, Changes in leaf autofluorescence upon MtNFP and MtLYK3 co-production. Leaf regions co-producing MtNFP-3xFLAG and MtLYK3-3xFLAG fusions were analyzed between 24 and 48 hai (here depicted 36 hai) using a stereoscope. Note the decrease in chlorophyll content, as indicated by the decrease of far-red autofluorescence of chlorophyll (left panel), and enhanced accumulation of blue light-

excited autofluorescence (right panel) within the infiltrated region. Bar is 1 cm. C, Accumulation of phenolic compounds. The following fusions were (co-)produced in *Nicotiana* leaves: MtNFP-3xFLAG (1); MtLYK3-3xFLAG (2); MtNFP-3xFLAG+MtLYK3-3xFLAG (3); MtNFP-3xFLAG+MtLYK3[G334E]-3xFLAG (4); AtCERK1-3xFLAG (5); or AtCERK1[K349]-3xFLAG (6). Macroscopic observations (left panel) and subsequent UV-excited autofluorescence of ethanol/lactophenol-cleared (right panel) leaf regions are depicted 36 hai (except for 5–30 hai). Bars are 1 cm. D, Induction of *NbHIN1*, *NbPR1 basic*, *NbACRE31*, and *NbACRE132* expression in response to separate production or co-production of: MtNFP-3xFLAG (NFP), MtLYK3-3xFLAG (LYK3), MtLYK3[G334E]-3xFLAG (LYK3[G334E]), and AtCERK1-3xFLAG (CERK1). Leaf samples were collected 24 hai and induction of gene expression was analyzed using qRT-PCR. Histograms represent induction of *NbHIN1* (white columns), *NbPR1 basic* (grey columns), *NbACRE31* (hatched columns), and *NbACRE132* (black columns) normalized by one reference gene, *MtEF1 α* . Induction of each gene was normalized to that caused by mock infiltration, and then calculated as % induction relative to the induction observed upon co-production of MtNFP and MtLYK3 fusions. Bars represent standard deviation of the mean. At least two technical replicates from two biological replicates were analyzed.
doi:10.1371/journal.pone.0065055.g004

using quantitative reverse-transcriptase polymerase chain reaction (qRT-PCR). Co-production of MtNFP and MtLYK3 fusions, and separate production of AtCERK1 fusion resulted in an induction of *NbHIN1*, *NbPR1 basic*, *NbACRE31*, and *NbACRE132* gene expression that was significantly higher than that following co-production of MtNFP and MtLYK3[G334E] fusions or separate production of MtNFP, MtLYK3, and MtLYK3[G334E] fusions (Fig. 4D). The *NbPR1a acidic* and *NbCYP71D20* genes did not display significant induction upon (co-) production of any of the protein(s) tested (unpublished data).

Taken together, the indication of a localized accumulation of phenolic compounds and induction of defence-related gene expression suggested that the *Nicotiana* response to MtNFP and MtLYK3 co-production triggered responses that were qualitatively similar to the responses to heterologous production of AtCERK1. In addition, the fact that in both cases the impairment of Ca^{2+} influx delayed the CD development suggested that the apparent functional interaction of two symbiotic LysM-RLKs in *Nicotiana* leaves mimics the action of the MAMP receptor, AtCERK1, and triggers defence-like responses.

Cell Death Induced in *Nicotiana* Leaves Upon MtNFP and MtLYK3 Co-production is a NF-independent Response

Perception of NFs results in triggering host symbiotic programme mediated by MtNFP and/or MtLYK3 [2]. In contrast, co-production of these LysM-RLKs in *Nicotiana* leaves apparently triggered some signalling cascade in the absence of NFs. Therefore, we investigated the effect of NF produced by *Sinorhizobium meliloti*, a microsymbiont of *Medicago*, on this CD response. To this end, *Agrobacterium* transformants carrying either MtNFP-sYFP2 or MtLYK3-mCherry construct were co-infiltrated into *Nicotiana* leaves at varying concentrations (as measured with OD₆₀₀). Then, purified SmNF at 10⁻⁷ M (in diluted DMSO) or diluted DMSO alone was applied between 9 and 24 hai to parts of the leaf regions co-producing MtNFP and MtLYK3 fusions, and CD development was monitored between 24 and 72 hours after the first infiltration (with *Agrobacterium*) using Evans blue staining. For all bacterial concentrations and time-points of SmNF/DMSO application tested, compromised membrane permeability in leaf regions co-producing MtNFP and MtLYK3 fusions was observed at similar time irrespective of the SmNF or DMSO treatment (Fig. 5), indicating similar kinetics of CD development. Therefore, we did not obtain evidence for any stimulatory or inhibitory effect of the SmNF on the CD development upon MtNFP and MtLYK3 co-production.

The Intracellular Region of MtNFP and Kinase Activity of MtLYK3 are Required for Cell Death Induction in *Nicotiana* Leaves

The independence of CD induction upon MtNFP and MtLYK3 co-production from the SmNF perception prompted us to compare structural requirements of CD induction and nodulation with

regard to these LysM-RLKs. In case of MtNFP, a recent structure-function study in *Medicago* [22] showed that loss-of-function mutations located in the ExR could be attributed to retention of the mutated protein in the endoplasmic reticulum (ER), whereas most substitutions located in the InR were found not to have an effect on the MtNFP function in nodulation. Therefore, we decided to limit our analysis of MtNFP to three point-mutated variants carrying: Ser 67 Phe (encoded by the *Mtnfp-2* allele), Ser 67 Ala, and Gly 474 Glu substitution; and a truncated variant with almost the entire InR deleted, termed MtNFP[Δ InR] (amino acids: 1–283) (see Table 2). Based on structure-function studies on MtLYK3 and LjNFR1, respectively in *Medicago* and in *Lotus* ([12,20–21], Table 3 in this study), we decided to test the effect of 16 point mutations (listed in Table 3) on MtLYK3 ability to induce CD in *Nicotiana* leaves in the presence of MtNFP. These included: a Pro 87 Ser (encoded by the *Mtlyk3-3* allele) and a Gly 334 Glu (encoded by the *Mtlyk3-1* allele) mutations, and Ala substitutions of Thr 285, Ser 286, Thr 300, Thr 319, Lys 349, Glu 362, Thr 433, Asp 441, Lys 464, Ser 471, Thr 472, Thr 475, Thr 480, and Thr 512. With the exception of the P87S substitution located in the first LysM domain of MtLYK3, all the above mutations are located in the MtLYK3 InR but differ in their effect on MtLYK3 autophosphorylation activity *in vitro* ([20] and Fig. S2; see Table 3). All truncated/mutated variants were prepared as fusions to the N-terminus of sYFP2, and their production and correct PM localization in *Nicotiana* leaf epidermal cells was confirmed, except for two MtNFP variants: MtNFP[S67F]-sYFP2 fusion was retained in the ER, and MtNFP[G474E]-sYFP2 fusion showed a partial PM localization, ([20,22] and Fig. S1). Additionally, we ruled out a possibility that the presence of WT MtNFP-FP or WT MtLYK3-FP fusion might affect stability/localization of the truncated/mutated fusions by confirming their efficient production and PM localization in *Nicotiana* leaf epidermal cells also upon co-production with MtLYK3 or MtNFP fusions (unpublished data).

Subsequently, we analyzed the ability of truncated/mutated MtNFP and MtLYK3 variants to induce CD upon either their separate production or co-production with WT MtLYK3-mCherry or WT MtNFP-mCherry fusion, respectively. In order to compare the CD induction ability of truncated/mutated variants with WT proteins, concomitant co-infiltration with *Agrobacterium* transformants carrying either WT MtNFP-FP or WT MtLYK3-FP construct was done on every leaf. Development of CD was monitored between 36 and 72 hai, and in case of the absence of or weakly pronounced macroscopic symptoms, the occurrence of CD was further scrutinized with Evans blue staining. None of the truncated/mutated variants was able to induce CD in *Nicotiana* leaves on its own (Table 2, 3). Co-production of MtNFP[S67A]-sYFP2 and MtLYK3 fusions resulted in a confluent death of (nearly) the entire infiltrated region in 6 out of 9 infiltrations, and compromised membrane permeability that could be observed in the entire infiltrated region (Table 2). In contrast, co-production of MtLYK3 fusion with MtNFP[S67F]-sYFP2,




Figure 5. Cell death upon MtNFP and MtLYK3 co-production in *Nicotiana* leaves does not require *SmNF*. *Agrobacterium* transformants carrying either *MtNFP-3xFLAG* or *MtLYK3-3xFLAG* construct were co-infiltrated into *Nicotiana* leaves at a final concentration: OD₆₀₀ [*MtNFP*] = 0.25 and OD₆₀₀ [*MtLYK3*] = 0.4 (1); OD₆₀₀ [*MtNFP*] = 0.15 and OD₆₀₀ [*MtLYK3*] = 0.25 (2). Twelve hai parts of the transformed regions were syringe-infiltrated with 10⁻⁷ mM *SmNF* (circled in red) or DMSO diluted to the same concentration (circled in white). Macroscopic observation (left panel) and Evans blue staining (right panel) are depicted 33 hai. Bar is 1 cm. doi:10.1371/journal.pone.0065055.g005

MtNFP[G474E]-sYFP2, or MtNFP[ΔInR]-sYFP2 fusion did not induce CD in *Nicotiana* leaves (Table 2). In case of MtLYK3 mutated variants, co-production of MtLYK3[P87S]-sYFP2 and MtNFP fusions induced confluent CD in all infiltrated regions (Table 3). In contrast, co-production of MtNFP fusion with all seven MtLYK3-sYFP2 mutated variants affected for their autophosphorylation activity *in vitro* did not induce CD in *Nicotiana* leaves (Table 3). In case of mutations that do not affect autophosphorylation activity of MtLYK3 kinase, we found that MtLYK3[T285A S286A T300A]-sYFP2 and MtLYK3[S471A]-sYFP2 fusions were as active as WT MtLYK3-sYFP2 fusion for CD induction upon their co-production with MtNFP fusion, whereas MtLYK3[K464A]-sYFP2 fusion induced compromised membrane permeability (but no macroscopic symptoms of cell death) upon co-production with MtNFP fusion (Fig. S3, Table 3). Co-production of MtNFP fusion with MtLYK3[T433A]-sYFP2, MtLYK3[T472A]-sYFP2 or MtLYK3[T512A]-sYFP2 fusion resulted in a confluent death of (nearly) the entire infiltrated region in, respectively, 7 out of 20, 5 out of 11, and 12 out of 20

infiltrations, whereas the remaining leaf regions displayed only (a) small patch(es) of dead tissue (Fig. S3, Table 3).

Taken together, most of the structural requirements regarding the MtNFP and MtLYK3 InR, and the autophosphorylation activity of the MtLYK3 KD, appeared to be identical for biological activity of these LysM-RLKs in both *Medicago* and *Nicotiana*. More specifically, we found out that both nodulation and CD induction displayed the same requirements for 11 out of 15 residues located in the MtLYK3 InR. On the contrary, a single mutation in the MtLYK3 ExR tested (that does not affect the PM localization of the fusion) was found to be crucial for MtLYK3 function in nodulation but not in CD induction. In case of MtNFP, the substitution of Ser 67 similarly abolished (S67F) or did not have an effect (S67A) on MtNFP function in nodulation and CD induction, which seemed to correlate with, respectively, the absence or presence of MtNFP fusion at the PM.

Table 2. Cell death induction activity of MtNFP-sYFP2 truncated/mutated variants in *Nicotiana* leaves.

<i>MtNFP-sYFP2</i> construct	Subcellular localization*	Nodulation activity*	Cell death induction	
			Co-expression with <i>MtLYK3-mCherry</i>	Separate expression
WT	PM	+	28/30	0/12
S67F (<i>Mtnfp-2</i>)	ER	-	0/9	0/9
S67A	PM	+	6/9#	0/9
ΔInR	PM	-	0/20	0/9
G474E	partial PM	-	0/15	0/9

*-see [22], PM-plasma membrane, ER-endoplasmic reticulum.

The designated constructs were expressed alone or co-expressed with *MtLYK3-mCherry* construct in *Nicotiana* leaves. Macroscopic symptoms of cell death were scored 48 hai; only infiltrations that resulted in confluent death of (nearly) the entire infiltrated region were scored and are presented as a fraction of total infiltrations performed. # - In the 3 remaining leaf regions, co-expression of *MtNFP*[S67A]-sYFP2 and *MtLYK3-mCherry* constructs resulted in increased staining with Evans blue in the entire infiltrated region.

doi:10.1371/journal.pone.0065055.t002

Table 3. Cell death induction activity of MtLYK3-sYFP2 mutated variants in *Nicotiana* leaves.

MtLYK3-sYFP2 construct	Auto-phosphorylation activity*	Nodulation activity**	Cell death induction	
			Co-expression with MtNFP-mCherry	Separate expression
WT	+	+	28/30	0/12
P87S (Mtlyk3-3)	Not applicable	-	15/15	0/9
T319A	-	-	0/11	0/9
G334E (Mtlyk3-1)	-	-	0/20	0/9
K349A	-	-	0/16	0/9
E362A	-	-	0/15	0/9
D441A	-	-	0/16	0/9
T475A	-	-	0/13	0/9
T480A	-	-(0/24)	0/18	0/9
T285A/S286A/T300A	+	Reduced with T300A	15/16	0/9
T433A	+	Reduced	Reduced 7/20	0/10
K464A	Reduced	Reduced (7/18)	0/12#	0/9
S471A	+	Reduced	9/11	0/9
T472A	+	Reduced	Reduced 5/11	0/9
T512A	Reduced	-	Reduced 12/20	0/9

*-see [20], except for the T480A (Fig. S2), **-see [20], except for the P87S [12], K464A and T480A (this study; number of plants nodulated/number of plants tested). The designated constructs were expressed alone or co-expressed with MtNFP-mCherry construct in *Nicotiana* leaves. Macroscopic symptoms of cell death were scored 48 hai: only infiltrations that resulted in confluent death of (nearly) the entire infiltrated region were scored and are presented as a fraction of total infiltrations performed. # - despite the lack of pronounced macroscopic symptoms, the co-expression of MtLYK3[K464A]-sYFP2 and MtNFP-mCherry constructs resulted in increased staining with Evans blue in the entire infiltrated region.
doi:10.1371/journal.pone.0065055.t003

Discussion

Co-production of MtNFP and MtLYK3 in *Nicotiana* Induces Defence-like Responses that Resemble *Nicotiana* Responses to AtCERK1 Production

Efficient production of both MtNFP-FP and MtLYK3-FP fusions in *Nicotiana* leaves facilitated characterization of their subcellular localization [20,22] and oligomerization status *in vivo* (manuscript in preparation), and led to the surprising observation of a CD induction (Fig. 1A). This response phenotypically and kinetically (Fig. 4A) resembled the HR elicited in *Nicotiana* spp. by pathogen-derived components [49], or CD induced by (co-) production of certain defence-related proteins, e.g. Pto [58]. Notably, a similar CD response was observed in *Nicotiana* leaves upon production of AtCERK1 (Fig. 2A), a MAMP receptor. Although CD induction was not demonstrated in any plant species in response to COs or PGN [54–56], it has been observed upon deregulation of various stress/defence-related signalling components [59–60], including a mitogen-activated protein kinase kinase (OsMKK4) implicated in COs-induced signalling in rice [61]. In addition, expression of AtCERK1 from the CaMV 35S promoter (used also in our studies) in *Arabidopsis* was shown to result in a ligand-independent dimerization of AtCERK1 [38]. We hypothesize that overproduction of this LysM-RLK in *Nicotiana* analogously leads to its dimerization and enhanced (deregulated) kinase activity, which in turn is required and sufficient for triggering CD; a response not observed upon ligand-induced activation of AtCERK1 in *Arabidopsis*.

Cell death induction upon MtNFP and MtLYK3 co-production in *Nicotiana* is in agreement with the *Nicotiana* [21] and *Arabidopsis* [19] response to LjNFR1 and LjNFR5 co-production. However, in these studies the associated induction of putative defence-related

responses has not been investigated. We here showed that both MtNFP and MtLYK3 co-production and AtCERK1 production in *Nicotiana* leaves triggered local accumulation of phenolic compounds, and a similar induction of expression of 4 out of 6 tested defence-related genes (Fig. 4C, D). We speculate that the two other genes might display different kinetics of induced expression (here analyzed only 24 hai) or might undergo suppression by *Agrobacterium* [62]. Importantly, COs- and/or PGN-induced expression of *PRI*, *ACRE31*, *ACRE132* and a member of a *HLN1* gene family was reported previously [32,52,54], linking these genes to MAMP-induced gene regulation mediated by AtCERK1 in *Arabidopsis* and/or *NbCERK1* in *Nicotiana*. In addition, we found that the lanthanum chloride-induced impairment of a Ca²⁺ influx similarly delayed the CD development upon (co-)production of the LysM-RLKs in our study (Fig. 3A, C). Therefore, we speculate that the signalling triggered upon MtNFP and MtLYK3 co-production in *Nicotiana* mimics AtCERK1-mediated signalling and thereby results in an induction of defence-like responses.

Similarities and Differences between Symbiotic and Defence Signalling Mediated by LysM-RLKs

The similarity between *Nicotiana* response to MtNFP and MtLYK3 co-production and AtCERK1 production suggests a possible overlap in signalling mediated by these LysM-RLKs. Several NF-induced processes, such as: a transient increase of reactive oxygen species (ROS) production; activation of phospholipase C (PLC) and PLD; and prolonged oscillations of (peri)nuclear [Ca²⁺] are implicated in switching on the symbiotic programme [63–69], whereas a Ca²⁺ influx is postulated to act as a signal for infection thread formation [70]. Interestingly, (CERK1-mediated) COs- and/or PGN-induced responses also

involve a Ca^{2+} influx, an elevated ROS production, and PLC activation [33,35,37–38,52–53,71–73]. We speculate that these similar processes might be activated/regulated by related molecular components, hence allowing two *Medicago* LysM-RLKs to activate signalling components present in *Nicotiana* leaf. Remarkably, Nakagawa and associates [74] demonstrated that swapping of the AtCERK1 ExR and a certain part of the AtCERK1 InR for the corresponding regions from LjNFR1 conferred on AtCERK1 a competence, albeit inefficient, for symbiotic signalling during *Lotus-Mesorhizobium loti* interaction. Conversely, our results demonstrate that MtNFP and MtLYK3, when co-produced in *Nicotiana*, are capable of signalling in a similar manner to AtCERK1. We hypothesize that due to the absence of symbiosis-specific “decoders” or “modulators” in *Nicotiana*, MtNFP- and MtLYK3-mediated signalling might be differently interpreted in this species, resulting in the induction of defence-like responses.

Importantly, NF-induced host responses are partially contradictory. On one hand, NFs are postulated to suppress the production in legume roots of salicylic acid and ROS, two potent signals implicated in defence signalling, upon rhizobia perception [63,75]. This differential induction of the symbiotic competence or defence could be regulated quantitatively via NPR1 (*Non-expressor of Pathogenesis-related genes 1*)-mediated signalling [76]. On the other hand, even perception of compatible NFs leads to the induction of (some) defence-related gene expression and phosphorylation of defence-related proteins in the initial stage of symbiosis [7,11,68,74]. An even more striking example of a NF-induced defence-like response comes from *Sesbania rostrata*. NF produced by its microsymbiont, *Azorhizobium caulinodans*, triggers the production of hydrogen peroxide and ethylene that eventually lead to local CD required for the formation of cortical infection pockets [77]. Therefore, our results agree with the hypothesized stimulation of (initial) host's defence responses via LysM-RLKs that mediate NF-induced signalling.

Possible Co-functioning of MtNFP and MtLYK3 in *Nicotiana* and *Medicago*

To explain a possible signalling mechanism employed by the kinase-inactive MtNFP, it has been proposed to form a receptor complex with MtLYK3 and another LYK protein during, respectively, the infection thread growth and pre-infection stage of symbiosis [12,15]. In contrast, it is not known whether MtNFP functions alone during arbuscular mycorrhiza symbiosis [24–26] or resistance towards fungal and oomycete pathogens [27] or requires a similar co-functioning with (an)other RLK(s). We here demonstrated that MtNFP required MtLYK3 to induce CD in *Nicotiana*, and that neither of these LysM-RLKs could be substituted by an unrelated RLK (Fig. 1B, Table 1). We propose that this *Nicotiana* response reflects a functional interaction between these LysM-RLKs. A rather limited heteromerization of MtNFP and a kinase-inactive MtLYK3 variant observed in the PM of *Nicotiana* leaf epidermal cells (manuscript in preparation) does not exclude either hypothesized mechanism: a direct, phosphorylation-dependent physical interaction between MtNFP and WT MtLYK3; or indirect (functional) interaction between MtNFP and MtLYK3 that requires independent activation of different molecular components by either LysM-RLK, and a later convergence of such putative signalling pathways. Interestingly, MtLYK3, but not the homologous [78] AtCERK1, required the presence of MtNFP for CD induction in *Nicotiana* (Fig. 2A), indicating the specific requirement for MtNFP to potentiate the MtLYK3-mediated signalling. This observation agrees with the hypothesized specialization of the LysM-RLKs mediating NF-

induced signalling during the co-evolution of legumes with rhizobia [28].

Cell Death Induction in *Nicotiana* and Nodulation in *Medicago* Share Certain Structural Requirements Regarding MtNFP and MtLYK3

Curiously, CD induction in *Nicotiana* was independent from the NF perception (Fig. 5), and the presence of Pro 87 in the MtLYK3 ExR (Table 3), in contrast to MtNFP and MtLYK3 function in nodulation [12,15]. Further mapping of crucial amino acid residues, and detailed characterization of their exact role in signalling would be required to clarify in the future whether or not nodulation and CD induction indeed hold different structural requirements with regard to the ExRs of these LysM-RLKs. On the contrary, the biological activity of MtNFP in *Nicotiana* was dependent on its PM localization, the presence of its InR, and the Gly 474 (Table 2), thus mimicking the structural requirements of RL symbiosis regarding MtNFP [22], and proteins encoded by *MtNFP* orthologs in *Lotus* [79] and pea (*Pisum sativum*) [5]. The overlap between structural requirements of nodulation and CD induction was even more pronounced with respect to the MtLYK3 InR. Out of 16 residues whose role in nodulation was identified, 9 residues (Thr 319, Gly 334, Lys 349, Glu 362, Thr 433, Asp 441, Thr 472, Thr 475, and Thr 480) were found to be equally important, and 2 residues (Thr 285, Ser 286) were equally dispensable for MtLYK3 biological activity both in *Medicago* and in *Nicotiana* (Table 3). In addition, the K464A and T512A substitutions had a negative effect of MtLYK3 biological activity in both *Nicotiana* and *Medicago*, although this effect was more (K464A) or less (T512A) severely pronounced during CD induction assays than during nodulation (Table 3). Various mutations abolishing MtLYK3 autophosphorylation activity ([20] and Fig. S2) similarly abolished its biological activity in *Medicago* [20] and in *Nicotiana* (Table 3), supporting the hypothesis that autophosphorylation of MtLYK3 is crucial for its signalling function. Importantly, as the role of Thr 480 in nodulation has not been described so far, our results revealed its importance for MtLYK3 function *in vivo*. Notably, the shared structural requirements of nodulation and CD induction were also confirmed with regard to several (putative) phosphorylation sites that do not abolish MtLYK3 autophosphorylation activity *in vitro* (Table 3). Phosphorylation within the InR of a RLK is often required for activation and regulation of its catalytic activity, and for generation of docking sites for (downstream) signalling components [80–82]. The shared importance of three (Thr 433, Thr 472, and Thr 512) out of five such phosphorylation sites for MtLYK3 biological activity in both plant species suggests that some of these phosphorylation-dependent functions required for MtLYK3-mediated signalling are conserved in *Nicotiana* leaf.

Demonstrated significant overlap between structural requirements of nodulation and CD induction regarding the MtNFP and MtLYK3 InRs supports our notion of the relevance of the *Nicotiana* system for studies on these, and potentially other (symbiotic) LysM-RLKs. This system presents certain practical advantages over the legume root system, in terms of rapidity and ease of expression of multiple constructs. In view of hypothesized similarities between NF (i.e. lipo-chitooligosaccharide)-induced and COs-induced signalling, analyzing known molecular components/processes involved in the CERK1-mediated signalling [35–38,83] might provide information on the yet-unidentified players implicated in the perception and/or transduction of the NF signal. This would be especially important as still very little is known about the identity of interactors of these symbiotic LysM-RLKs [17–18,21,84]. Possible candidate signalling molecule(s) function-

ing in co-operation with, or downstream from the LysM-RLKs, and identified in this heterologous system should then be evaluated in legume root in order to confirm their involvement in symbiosis.

Supporting Information

Figure S1 Subcellular localization of various protein fusions in *Nicotiana* leaf epidermal cells. The plasma membrane marker, mCherry-HVR, was co-produced with the designated fusions in *Nicotiana* leaf epidermal cells, and the fluorescence (viewed from abaxial side) was imaged 24 hai using confocal laser scanning microscopy. From left to right: green fluorescence of sYFP2; orange fluorescence of mCherry; superimposition of green, orange, and far-red (chlorophyll) fluorescence with the differential interference contrast (DIC) image. Bars are 20 μm . Note 1: in case of subcellular localization of MtNFP[G474E]-sYFP2 fusion, strong fluorescent puncta (indicated with an arrowhead) at the cell boundary of many cells (sometimes in association with nuclei), and pronounced ER localization (indicated with an arrow) of the fusion were still visible at 48 hai. Nevertheless, some cells showed a more uniform pattern of fluorescence at the cell boundary, and this observation, together with a partial insensitivity of this mutated variant to the PNGaseF treatment [22], indicated that at least some MtNFP[G474E] fusion had reached the PM. Note 2: as all kinase-inactive MtLYK3 variants were produced and correctly localized to the plasma membrane in *Nicotiana* leaf epidermal cells, their lack of biological activity can be attributed to the general abolishment of kinase activity rather than to an individual effect of a particular mutation. (TIF)

Figure S2 Effect of the Thr 480 Ala substitution on MtLYK3 autophosphorylation activity *in vitro*. The puri-

fied intracellular regions of WT MtLYK3, MtLYK3[G334E], and MtLYK3[T480A], fused to the C terminus of GST, were analyzed for their autophosphorylation activity *in vitro* using radiolabeled ATP (γ - ^{32}P ATP) and phosphorimaging (PI). The coomassie blue staining (CB) shows the protein loading. (TIF)

Figure S3 Various (putative) phosphorylation sites are differentially required for MtLYK3 biological activity in *Nicotiana*. MtLYK3-sYFP2 mutated variants were co-produced with MtNFP-mCherry fusion in *Nicotiana* leaves: MtLYK3[T285A S286A T300A]+MtNFP (1); MtLYK3[T433A]+MtNFP (2); MtLYK3[T512A]+MtNFP (3); MtLYK3[T480A]+MtNFP (4). Macroscopic observation (left panel) and Evans blue staining (right panel) are depicted 48 hai. Bar is 1 cm. (TIF)

Table S1 Primer and linker sequences. (DOC)

Materials and Methods S1. (DOC)

Acknowledgments

We thank Dr. G. Morieri for providing us with purified *SinNod* factor, Dr. A. Streng for providing us with *AtCERK1* sequence and for critical reading of the manuscript, and Dr. H. van den Burg for helpful discussions.

Author Contributions

Conceived and designed the experiments: AP-B BL RG JVC TWJG. Performed the experiments: AP-B BL MAK DK-H JVC. Analyzed the data: AP-B BL FLWT RG JVC TWJG. Contributed reagents/materials/analysis tools: AP-B BL MAK DK-H FLWT RG JVC TWJG. Wrote the paper: AP-B BL MAK FLWT RG JVC TWJG.

References

- Masson-Boivin C, Giraud E, Perret X, Batut J (2009) Establishing nitrogen-fixing symbiosis with legumes: how many rhizobium recipients? *Trends Microbiol* 17: 458–466.
- Downie JA (2010) The roles of extracellular proteins, polysaccharides and signals in the interactions of rhizobia with legume roots. *FEMS Microbiol Rev* 34: 150–170.
- Gust AA, Willmann R, Desaki Y, Grabherr HM, Nürnberger T (2012) Plant LysM proteins: modules mediating symbiosis and immunity. *Trends Plant Sci* 17: 495–502.
- Oldroyd GED (2013) Speak friend, and enter: signalling systems that promote beneficial symbiotic associations in plants. *Nature* 11: 252–263.
- Madsen EB, Madsen LH, Radutoiu S, Olbryt M, Rakwalska M, et al. (2003) A receptor kinase gene of the LysM type is involved in legume perception of rhizobial signals. *Nature* 425: 637–640.
- Radutoiu S, Madsen LH, Madsen EB, Felle HH, Umehara Y, et al. (2003) Plant recognition of symbiotic bacteria requires two LysM receptor-like kinases. *Nature* 425: 585–592.
- El Yahyaoui FE, Küster H, Ben Amor B, Hohnjec N, Pühler A, et al. (2004) Expression profiling in *Medicago truncatula* identifies more than 750 genes differentially expressed during nodulation, including many potential regulators of the symbiotic program. *Plant Physiol* 136: 3159–3176.
- Mitra RM, Shaw SL, Long SR (2004) Six non-nodulating plant mutants defective for Nod factor-induced transcriptional changes associated with the legume-rhizobia symbiosis. *Proc Natl Acad Sci U S A* 101: 10217–10222.
- Arrighi JF, Barre A, Ben Amor B, Bersoult A, Soriano LC, et al. (2006) The *Medicago truncatula* Lysin [corrected] motif-Receptor-like kinase gene family includes NFP and new nodule-expressed genes. *Plant Physiol* 142: 265–279.
- Miwa H, Sun J, Oldroyd GED, Downie JA (2006) Analysis of Nod-factor-induced calcium signalling in root hairs of symbiotically defective mutants of *Lotus japonicus*. *Mol Plant Microbe Interact* 19: 914–923.
- Hogslund N, Radutoiu S, Krusell L, Voroshilova V, Hannah MA, et al. (2009) Dissection of symbiosis and organ development by integrated transcriptome analysis of *Lotus japonicus* mutant and wild-type plants. *PLoS ONE* 4: e6556.
- Smit P, Limpens E, Geurts R, Fedorova E, Dolgikh E, et al. (2007) *Medicago* LYK3, an entry receptor in rhizobial nodulation factor signalling. *Plant Physiol* 145: 183–191.
- Hayashi T, Banba M, Shimoda Y, Kouchi H, Hayashi M, et al. (2010) A dominant function of CcAMK in intracellular accommodation of bacterial and fungal endosymbionts. *Plant J* 63: 141–154.
- Madsen LH, Tirichine L, Jurkiewicz A, Sullivan JT, Heckmann AB, et al. (2010) The molecular network governing nodule organogenesis and infection in the model legume *Lotus japonicus*. *Nature* doi: 10.1038/ncomms1009.
- Bensmihen S, de Billy F, Gough C (2011) Contribution of NFP LysM domains to the recognition of Nod Factors during the *Medicago truncatula*/*Sinorhizobium meliloti* symbiosis. *PLoS ONE* 6: e26114.
- Limpens E, Mirabella R, Fedorova E, Franken C, Franssen H, et al. (2005) Formation of organelle-like N₂-fixing symbiosomes in legume root nodules is controlled by DMI2. *Proc Natl Acad Sci U S A* 102: 10375–10380.
- Mbengue M, Camut S, de Carvalho-Niebel F, Deslandes L, Froidure S, et al. (2010) The *Medicago truncatula* E3 ubiquitin ligase PUB1 interacts with the LYK3 symbiotic receptor and negatively regulates infection and nodulation. *Plant Cell* 22: 3474–3488.
- Haney CH, Riely BK, Tricoli DM, Cook DR, Ehrhardt DW, et al. (2011) Symbiotic rhizobia bacteria trigger a change in localization and dynamics of the *Medicago truncatula* receptor kinase LYK3. *Plant Cell* 23: 2774–2787.
- Broghammer A, Krusell L, Blaise M, Sauer J, Sullivan JT, et al. (2012) Legume receptors perceive the rhizobial lipochitin oligosaccharide signal molecules by direct binding. *Proc Natl Acad Sci U S A* 109: 13859–13864.
- Klaus-Heisen D, Nurisso A, Pietraszewska-Bogiel A, Mbengue M, Camut S, et al. (2011) Structure-function similarities between a plant receptor-like kinase and the human Interleukin-1 Receptor-Associated Kinase-4. *J Biol Chem* 286: 11202–11210.
- Madsen EB, Antolin-Llovera M, Grossmann C, Ye J, Vieweg S, et al. (2011) Autophosphorylation is essential for *in vivo* function of the *Lotus japonicus* Nod Factor Receptor 1 and receptor mediated signalling in cooperation with Nod Factor Receptor 5. *Plant J* 65: 404–417.
- Lefebvre B, Klaus-Heisen D, Pietraszewska-Bogiel A, Hervé C, Camut S, et al. (2012) Role of N-glycosylation sites and CXC motifs in trafficking of *Medicago truncatula* Nod Factor Perception protein to the plasma membrane. *J Biol Chem* 287: 10812–10823.
- Radutoiu S, Madsen LH, Madsen EB, Jurkiewicz A, Fukai E, et al. (2007) LysM domains mediate lipochitin-oligosaccharide recognition and *Nfr* genes extend the symbiotic host range. *EMBO J* 26: 3923–3935.

24. Oláh B, Brière C, Bécard G, Déneris J, Gough C (2005) Nod factors and a diffusible factor from arbuscular mycorrhizal fungi stimulate lateral root formation in *Medicago truncatula* via the DMI1/DMI2 signalling pathway. *Plant J* 44: 195–207.
25. Maillet F, Poinso V, André O, Peuch-Pagès V, Haouy A, et al. (2011) Fungal lipochitoooligosaccharide symbiotic signals in arbuscular mycorrhiza. *Nature* 469: 58–64.
26. Czaja LF, Hogekamp C, Lamm P, Maillet F, Andres Martinez E, et al. (2012) Transcriptional responses toward diffusible signals from symbiotic microbes reveal *MtNFP*- and *MtDMI3*-dependent reprogramming of host gene expression by arbuscular mycorrhizal fungal lipochitoooligosaccharides. *Plant Physiol* 159: 1671–1685.
27. Rey T, Nars A, Bonhomme M, Bottin A, Huguet S, et al. (2013) NFP, a LysM protein controlling Nod Factor perception, also intervenes in *Medicago truncatula* resistance to pathogens. *New Phytol* 198: 875–886.
28. Streng A, Op den Camp R, Bisseling T, Geurts R (2011) Evolutionary origin of *Rhizobium* Nod factor signalling. *Plant Signal Behav* 6: 1510–1514.
29. Boller T, Felix G (2009) A renaissance of elicitors: perception of microbe-associated molecular patterns and danger signals by pattern-recognition receptors. *Annu Rev Plant Biol* 60: 379–406.
30. Thomma BPHJ, Nürnberger T, Joosten MHJ (2011) Of PAMPs and effectors: the blurred PTI-ETI dichotomy. *Plant Cell* 23: 4–15.
31. Miya A, Albert P, Shinya T, Desaki Y, Ichimura K, et al. (2007) CERK1, a LysM receptor kinase, is essential for chitin elicitor signalling in Arabidopsis. *Proc Natl Acad Sci U S A* 104: 19613–19618.
32. Wan J, Zhang XC, Neece D, Ramonell KM, Clough S, et al. (2008) A LysM receptor-like kinase plays a critical role in chitin signalling and fungal resistance in Arabidopsis. *Plant Cell* 20: 471–481.
33. Gimenez-Ibanez S, Hann DR, Ntoukakis V, Petutschnig E, Lipka V, et al. (2009) AvrPtoB targets the LysM receptor kinase CERK1 to promote bacterial virulence on plants. *Curr Biol* 19: 423–429.
34. Shimizu T, Nakano T, Takamizawa D, Desaki Y, Ishii-Minami N, et al. (2010) Two LysM receptor molecules, CEBlP and OsCERK1, cooperatively regulate chitin elicitor signalling in rice. *Plant J* 64: 204–214.
35. Willmann R, Lajunen HM, Erbs G, Newman MA, Kolb D, et al. (2011) Arabidopsis lysin-motif proteins LYM1 LYM3 and CERK1 mediate bacterial peptidoglycan sensing and immunity to bacterial infection. *Proc Natl Acad Sci U S A* 108:19824–19829.
36. Liu B, Li JF, Ao Y, Qu J, Li Z, et al. (2012) Lysin motif-containing proteins LYP4 and LYP6 play dual roles in peptidoglycan and chitin perception in rice innate immunity. *Plant Cell* 24: 3406–3419.
37. Petutschnig EK, Jones AME, Serazetdinova L, Lipka U, Lipka V (2010) The Lysin Motif receptor-like kinase (LysM-RLK) CERK1 is a major chitin-binding protein in *Arabidopsis thaliana* and subject to chitin-induced phosphorylation. *J Biol Chem* 285: 28902–28911.
38. Liu T, Liu Z, Song C, Hu Y, Han Z, et al. (2012) Chitin-induced dimerization activates a plant immune receptor. *Science* 336: 221–234.
39. Oh CS, Martin GB (2010) Effector-triggered immunity mediated by the Pto kinase. *Trends Plant Sci* 16: 1360–1385.
40. Nguyen HP, Chakravarthy S, Velásquez AC, McLane HL, Zeng L, et al. (2010) Methods to study PAMP-triggered immunity using tomato and *Nicotiana benthamiana*. *Mol Plant Microbe Interact* 23: 991–999.
41. Lefebvre B, Timmers T, Mbengue M, Moreau S, Hervé C, et al. (2010) A remorin protein interacts with symbiotic receptors and regulates bacterial infection. *Proc Natl Acad Sci U S A* 107: 2343–2348.
42. van Ooijen G, Mayr G, Kassem MM, Albrecht M, Cornelissen BJ, et al. (2008) Structure-function analysis of the NB-ARC domain of plant disease resistance proteins. *J Exp Bot* 59: 1383–1397.
43. Shaner NC, Campbell RE, Steinbach PA, Giepmans BN, Palmer AE, et al. (2004) Improved monomeric red, orange and yellow fluorescent proteins derived from *Discosoma* sp. red fluorescent protein. *Nature* 428: 1567–1572.
44. Kremers GJ, Goedhart J, van Munster EB, Gadella TW Jr (2006) Cyan and yellow super fluorescent proteins with improved brightness, protein folding, and FRET Förster radius. *Biochem* 45: 6570–6580.
45. Endre G, Kereszt A, Kevei Z, Mihacea S, Kaló P, et al. (2002) A receptor kinase gene regulating symbiotic nodule development. *Nature* 417: 962–966.
46. Li J, Chory J (1997) A putative leucine-rich repeat kinase involved in brassinosteroid signal transduction. *Cell* 90: 929–938.
47. Mur LA, Kenton P, Lloyd AJ, Ougham H, Prats E (2008) The hypersensitive response; the centenary is upon us but how much do we know? *J Exp Bot* 59: 501–520.
48. Coll NS, Epple P, Dangel JL (2011) Programmed cell death in the plant immune system. *Cell Death Differ* 18: 1247–1256.
49. Heese A, Hann DR, Gimenez-Ibanez S, Jones AM, He K, et al. (2007) The receptor-like kinase SERK3/BAK1 is a central regulator of innate immunity in plants. *Proc Natl Acad Sci U S A* 104: 12217–12222.
50. Taguchi F, Suzuki T, Takeuchi K, Inagaki Y, Toyoda K, et al. (2009) Glycosylation of flagellin from *Pseudomonas syringae* pv. *tabaci* 6605 contributes to evasion of host tobacco plant surveillance system. *Physiol Mol Plant Pathol* 74: 11–17.
51. Ranf S, Eschen-Lippold L, Pecher P, Lee J, Scheel D (2011) Interplay between calcium signalling and early signalling elements during defence responses to microbe- or damage-associated molecular patterns. *Plant J* 68: 100–113.
52. Segonzac C, Feike D, Gimenez-Ibanez S, Hann DR, Zipfel C, et al. (2011) Hierarchy and roles of pathogen-associated molecular pattern-induced responses in *Nicotiana benthamiana*. *Plant Physiol* 156: 687–699.
53. Frei dit Frey N, Mbengue M, Kwaaitaal M, Nitsch L, Altenbach D, et al. (2012) Plasma membrane calcium ATPases are important components of receptor-mediated signalling in plant immune responses and development. *Plant Physiol* 159: 798–809.
54. Gust AA, Biswas R, Lenz HD, Rauhut T, Ranf S, et al. (2007) Bacteria-derived peptidoglycans constitute pathogen-associated molecular patterns triggering innate immunity in Arabidopsis. *J Biol Chem* 282: 32338–32348.
55. Erbs G, Silipo A, Aslam S, De Castro C, Liparoti V, et al. (2008) Peptidoglycan and muropeptides from pathogens *Agrobacterium* and *Xanthomonas* elicit plant innate immunity: structure and activity. *Cell* 15: 438–448.
56. Hamel LP, Beaudoin N (2010) Chitoooligosaccharide sensing and downstream signalling: contrasted outcomes in pathogenic and beneficial plant-microbe interactions. *Planta* 232: 787–806.
57. Cornelissen B, Horowitz J, van Kan JA, Goldberg RB, Bol JF (1987) Structure of tobacco genes encoding pathogenesis-related proteins from the PR-1 group. *Nucl Acid Res* 15: 6799–6811.
58. Mucyn TS, Clemente A, Andriotis VME, Balmuth AL, Oldroyd GED, et al. (2006) The tomato NBARC-LRR protein Prf interacts with Pto kinase *in vivo* to regulate specific plant immunity. *Plant Cell* 18: 2792–2806.
59. Takahashi Y, Bin Nasir KH, Ito A, Kanzaki H, Matsumura H, et al. (2007) A high-throughput screen of cell-death-inducing factors in *Nicotiana benthamiana* identifies a novel MAPKK that mediates INF1-induced cell death signalling and non-host resistance to *Pseudomonas cichorii*. *Plant J* 49: 1030–1040.
60. Gao M, Wang X, Wang D, Xu F, Ding X, et al. (2009) Regulation of cell death and innate immunity by two receptor-like kinases in Arabidopsis. *Cell* 6: 34–44.
61. Kishi-Kaboshi M, Okada K, Kurimoto T, Murakami S, Umezawa T, et al. (2010) A rice fungal MAMP-responsive cascade regulates metabolic flow to antimicrobial metabolite synthesis. *Plant J* 63: 599–612.
62. Ditt RF, Kerr KF, de Figueiredo P, Delrow J, Comai L, et al. (2006) The *Arabidopsis thaliana* transcriptome in response to *Agrobacterium tumefaciens*. *Mol Plant Microbe Interact* 19: 665–681.
63. Bueno P, Soto MJ, Rodriguez-Rosales MP, Sanjuan J, Olivares J, et al. (2001) Time-course of lipoxygenase, antioxidant enzyme activities and H₂O₂ accumulation during the early stages of *Rhizobium*-legume symbiosis. *New Phytol* 152: 91–96.
64. den Hartog M, Musgrave A, Munnik T (2001) Nod factor-induced phosphatidic acid and diacylglycerol pyrophosphate formation: a role for phospholipase C and D in root hair deformation. *Plant J* 25: 55–65.
65. Charron D, Pingret JL, Chabaud M, Journet EP, Barker DG (2004) Pharmacological evidence that multiple phospholipid signalling pathways link rhizobium nodulation factor perception in *Medicago truncatula* root hairs to intracellular responses, including Ca²⁺ spiking and specific *ENOD* gene expression. *Plant Physiol* 136: 3582–3593.
66. Lohar DP, Sharopova N, Endre G, Peñaela S, Samac D, et al. (2006) Transcript analysis of early nodulation events in *Medicago truncatula*. *Plant Physiol* 140: 221–234.
67. Cárdenas L, Martínez A, Sánchez F, Quinto C (2008) Fast, transient and specific intracellular ROS changes in living root hair cells responding to Nod factors (NFs). *Plant J* 56: 802–813.
68. Serna-Sanz A, Parniske M, Peck SC (2011) Phosphoproteome analysis of *Lotus japonicus* roots reveals shared and distinct components of symbiosis and defense. *Mol Plant Microbe Interact* 24: 932–937.
69. Sieberer BJ, Chabaud M, Timmers AC, Monin A, Fournier J, et al. (2009) A nuclear-targetedameleon demonstrates intracellular Ca²⁺ spiking in *Medicago truncatula* root hairs in response to rhizobial Nodulation factors. *Plant Physiol* 151: 1197–1206.
70. Shaw SL, Long SR (2003) Nod Factor elicits two separable calcium responses in *Medicago truncatula* root hair cells. *Plant Physiol* 131: 976–984.
71. den Hartog M, Verhoef N, Munnik T (2003) Nod factor and elicitors activate different phospholipid signalling pathways in suspension-cultured alfalfa cells. *Plant Physiol* 132: 311–317.
72. Kwaaitaal M, Huisman R, Maintz J, Reinstädler A, Panstruga R (2011) Ionotropic glutamate receptor (iGluR)-like channels mediate MAMP-induced calcium influx in *Arabidopsis thaliana*. *Biochem J* 440: 355–365.
73. Wan J, Tanaka K, Zhang XC, Son GH, Brechenmacher L, et al. (2012) LYK4, a lysine motif receptor-like kinase, is important for chitin signalling and plant innate immunity in Arabidopsis. *Plant Physiol* 160: 396–406.
74. Nakagawa T, Kaku H, Shimoda Y, Sugiyama A, Shimamura M, et al. (2010) From defense to symbiosis: limited alterations in the kinase domain of LysM receptor-like kinases are crucial for evolution of legume-*Rhizobium* symbiosis. *Plant J* 65: 169–180.
75. Martínez-Abarca F, Herrera-Cervera JA, Bueno P, Sanjuan J, Bisseling T, et al. (1998) Involvement of salicylic acid in the establishment of the *Rhizobium meliloti*-alfalfa symbiosis. *Mol Plant Microbe Interact* 11: 153–155.
76. Peleg-Grossmann S, Golani Y, Kaye Y, Melamed-Book N, Levine A (2009) NPR1 protein regulates pathogenic and symbiotic interactions between *Rhizobium* and legumes and non-legumes. *PLoS ONE* 4: e8399.
77. Capoen W, Oldroyd G, Goormachtig S, Holsters M (2010) *Sesbania rostrata*: a case study of natural variation in legume nodulation. *New Phytol* 186: 340–345.
78. Zhang XC, Cannon SB, Stacey G (2009) Evolutionary genomics of LysM genes in land plants. *BMC Evol Biol* 9: doi:10.1186/1471-2148-9-183.

79. Murray J, Karas B, Ross L, Brachmann A, Wagg C, et al. (2006) Genetic suppressors of the *Lotus japonicus har1-1* hypermodulation phenotype. *Mol Plant Microbe Interact* 19: 1082–1091.
80. Wang X, Kota U, He K, Blackburn K, Li J, et al. (2008) Sequential transphosphorylation of the BRI1/BAK1 receptor kinase complex impacts early events in brassinosteroid signalling. *Cell* 15: 220–235.
81. Lemmon MA, Schlessinger J (2010) Cell signalling by receptor tyrosine kinases. *Cell* 141: 1117–1134.
82. Oh MH, Wang X, Clouse SD, Huber SC (2012) Deactivation of the *Arabidopsis* Brassinosteroid Insensitive 1 (BRI1) receptor kinase by autophosphorylation within the glycine-rich loop. *Proc Natl Acad Sci U S A* 109: 327–332.
83. Chen L, Hamada S, Fujiwara M, Zhu T, Thao NP, et al. (2010) The Hop/Sti1-Hsp90 chaperone complex facilitates the maturation and transport of a PAMP receptor in rice innate immunity. *Cell Host Microbe* 7: 185–196.
84. Ke D, Fang Q, Chen C, Zhu H, Chen T, et al. (2012) The small GTPase ROP6 interacts with NFR5 and is involved in nodule formation in *Lotus japonicus*. *Plant Physiol* 159: 131–143.