

Osteonecrosis of the Humeral Head Following Arthroscopic Repair of Rotator Cuff Tears: A Case Report

Karim Ben Aicha¹, Lassaad Hassini¹, Amir Mhiri¹, Aymen Hanafi¹, Mourad Mtaoumi¹, Karim Bouattour¹

Learning Point of the Article:

The management of humeral head osteonecrosis as a complication of rotator cuff repair and the treatment with a reverse total shoulder arthroplasty.

Abstract

Introduction: The osteonecrosis and the collapse of the humeral head may have many risk factors such as trauma, alcoholism, metabolic diseases, and corticosteroid therapy. Otherwise, it was described as a rare complication of shoulder arthroscopy in the past few years.

Case Report: We report the case of a 65-year-old right-handed woman who had a rotator cuff tear of the right shoulder. She underwent a double-row arthroscopic repair. Six months later she had an intense shoulder pain, with radiological and MRI signs of humeral head osteonecrosis. The patient had a reverse total shoulder arthroplasty.

Conclusion: The aim of this case report is to underline the diagnostic particularities of this complication, and to shed light on the pathogenesis of the interruption of blood supply in the humeral head following rotator cuff repair. We also discuss the management of this complication with reverse total shoulder arthroplasty.

Keywords: Shoulder, rotator cuff, arthroscopy, humeral head, osteonecrosis.

Introduction

Rotator cuff tears are a common pathology in the elderly, causing invalid pain and shoulder disability [1]. The surgical treatment consists in arthroscopic repair of the torn tendons using suture anchors. This treatment option has good outcomes with some complications such as infection, delayed wound healing, algodystrophy, stiffness, and deep venous thrombosis [2]. Osteonecrosis of the humeral head is a rare cause of shoulder pain. It has many traumatic and metabolic etiologies [3]. The disruption of the vascular supply of the humeral head was the common factor [4]. A few cases described this osteonecrosis as a complication of shoulder arthroscopy with a rapid progressive

severe pain and collapse of the humeral head within the 1st year following surgery [5, 6, 7]. In this report, we aim to describe this rare complication and to provide more information about the risk factors and the treatment options.

Case Report

A 65-year-old, right-hand dominant woman, presented with 1-year history of rotator cuff syndrome of the right shoulder. Plain radiographs showed a maintained acromiohumeral distance with no signs of osteoarthritis (Fig. 1). Magnetic resonance imaging (MRI) showed full-thickness tear of the supraspinatus muscle

Access this article online

Website:
www.jocr.co.in

DOI:
<https://doi.org/10.13107/jocr.2023.v13.i12.4074>

Author's Photo Gallery



Dr. Karim Ben Aicha



Dr. Lassaad Hassini



Dr. Amir Mhiri



Dr. Aymen Hanafi



Dr. Mourad Mtaoumi



Dr. Karim Bouattour

¹Department of Orthopaedic Surgery, Sahloul Hospital, 4054 Sousse, Tunisia.

Address of Correspondence:

Dr. Amir Mhiri,
Department of Orthopaedic Surgery, Sahloul Hospital, 4054 Sousse, Tunisia.
E-mail: amirmhiri1@gmail.com

Submitted: 21/09/2023; Review: 02/10/2023; Accepted: November 2023; Published: December 2023

DOI: <https://doi.org/10.13107/jocr.2023.v13.i12.4074>

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License <https://creativecommons.org/licenses/by-nc-sa/4.0/>, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms






Figure 1: Anteroposterior (a) and lateral (b) radiographs of the right shoulder showing a maintained acromiohumeral distance with no signs of osteoarthritis.

with Goutallier Stage 1 fatty degeneration, normal signals of the humeral head and the glenoid cavity were observed (Fig. 2). The patient underwent shoulder arthroscopy that confirmed the full-thickness tears of the supraspinatus with Patte Type 2 retraction. We performed a double-row repair for the tendon using two suture anchors. Concomitant biceps tenotomy and acromioplasty were also performed. The patient was allowed to do self-assisted passive motion in the first 4 weeks postoperatively. Full active motion was initiated 6 weeks after the surgery then muscle strengthening physical therapy was

prescribed 2 weeks later. At 3-month follow-up, the patient had a good range of motion in all planes without any complications.

Six months postoperatively, the patient presented with severe right shoulder pain and restriction of all ranges of motion. Physical examination found tenderness around the anterior aspect of the shoulder with only 30° of forward elevation and abduction. The white blood cell count and the C-reactive protein level were within the normal limits. Shoulder radiographs revealed Cruess Stage 3 osteonecrosis of the humeral head with osteopenia and a crescent sign (Fig. 3 and 4). MRI was consistent with humeral head osteonecrosis

(Fig. 5). Due to the severe pain and the disability, the patient was treated with a reverse total shoulder arthroplasty 2 months after the diagnosis of osteonecrosis (Fig. 6). At the 4-month follow-up, the shoulder was not painful with an acceptable range of motion. The patient had 90° of abduction, 100° of forward elevation and she was highly satisfied with the outcome (Fig. 7).

Discussion

Shoulder arthroscopy may have some complications such as



Figure 2: Coronal View of MRI scan of the right shoulder showing a full-thickness tear of the supraspinatus muscle with Grade 2 retraction.



Figure 3: Anteroposterior radiograph of the right shoulder showing Cruess Stage 3 osteonecrosis of the humeral head with osteopenia and a crescent sign.



Figure 4: Lateral radiograph of the right shoulder showing an osteonecrosis of the humeral head.




Figure 5: MRI scan of the right shoulder showing humeral head osteonecrosis.

infection, neurovascular injuries, stiffness, algodystrophy, and even deep venous thrombosis leading to death. These complications can occur in 5.8%–10.6% of cases [2]. Osteonecrosis of the humeral head can have many non-traumatic causes, with the interruption of the blood supply as a common point [4]. These non-traumatic etiologies include corticosteroid therapy, alcoholism, Gaucher's disease, dysbarism, and pregnancy [3]. The understanding of vascular

anatomy and its variations is crucial for surgeons before proximal humerus procedures or shoulder arthroscopy [8]. The pathogenesis of humeral head osteonecrosis after shoulder arthroscopy was referred by some authors to a perioperative injury of the anterolateral branch of the anterior circumflex humeral artery by a poor placement of suture anchor or the use of multiple anchors [7]. However, Dilisio et al. [6] could not conclude that aberrant use of suture anchors was linked to the




Figure 6: Postoperative anteroposterior radiograph of reverse total shoulder arthroplasty.



Figure 7: Four-month follow-up anteroposterior radiograph of reverse total shoulder arthroplasty.

osteonecrosis because no hardware was used in two of the three patients in their study. Other suggested that additional damage might occur at the level of the biceps tenotomy or tenodesis [9]. In fact, the anterior circumflex humeral artery and its anterolateral branch may present anatomical variations leading to a higher risk of injury during biceps tenodesis tenotomy or placement of suture anchors [10]. Therefore, post-arthroscopic humeral head osteonecrosis can result from a damage to the anterolateral branch at the level of the transverse humeral ligament and the biceps tendon, or a damage to the intraosseous blood supply in the great tubercle, or both concurrently [10]. However, the poor intraosseous blood supply of the superomedial aspect of the humeral head makes it the initial starting location of osteonecrosis [9, 10]. The coagulation of blood vessels using radiofrequency, the increasing pressure and temperature of intra-articular fluid, embolism, and prolonged traction seem to be involved in the mechanisms of osteonecrosis. Possible allergic reactions or hypersensitivity due to the bioabsorbable suture anchors can also be involved in combination with the other mentioned factors [11]. Due to the severe pain and the disability, reverse total shoulder arthroplasty was the treatment of choice in many reported

cases, with satisfying functional results [11].

Conclusion

The knowledge of the anatomy of humeral head vascularization is necessary for shoulder surgeons, in a way to avoid this rare complication. Variations of the anterior circumflex humeral artery and its anterolateral branch should be taken in consideration. Thus, humeral head osteonecrosis leads to inevitable reinterventions for shoulder arthroplasty which can be risky in elderly patients with comorbidities. Further studies are required to establish a clear relationship between shoulder arthroscopy and rapid progressive osteonecrosis of the humeral head.

Clinical Message

To underline the diagnostic particularities of humeral head osteonecrosis following a rotator cuff repair using an anchor, and to shed light on the pathogenesis of the interruption of blood supply in the humeral head following rotator cuff repair.

Declaration of patient consent: The authors certify that they have obtained all appropriate patient consent forms. In the form, the patient has given the consent for his/ her images and other clinical information to be reported in the journal. The patient understands that his/ her names and initials will not be published and due efforts will be made to conceal their identity, but anonymity cannot be guaranteed.

Conflict of interest: Nil **Source of support:** None

References

1. Moosmayer S, Smith HJ, Tariq R, Larmo A. Prevalence and characteristics of asymptomatic tears of the rotator cuff: An ultrasonographic and clinical study. *J Bone Joint Surg Br* 2009;91:196-200.
2. Brislin KJ, Field LD, Savoie FH 3rd. Complications after arthroscopic rotator cuff repair. *Arthroscopy* 2007;23:124-8.
3. Hasan SS, Romeo AA. Nontraumatic osteonecrosis of the humeral head. *J Shoulder Elbow Surg* 2002;11:281-98.
4. Harreld KL, Marker DR, Wiesler ER, Shafiq B, Mont MA. Osteonecrosis of the humeral head. *J Am Acad Orthop Surg* 2009;17:345-55.
5. Beauthier V, Sanghavi S, Roulot E, Hardy P. Humeral head osteonecrosis following arthroscopic rotator cuff repair. *Knee Surg Sports Traumatol Arthrosc* 2010;18:1432-4.
6. Dilisio MF, Noble JS, Bell RH, Noel CR. Postarthroscopic humeral head osteonecrosis treated with reverse total shoulder arthroplasty. *Orthopedics* 2013;36:e377-80.
7. Goto M, Gotoh M, Mitsui Y, Okawa T, Higuchi F, Nagata K. Rapid collapse of the humeral head after arthroscopic rotator cuff repair. *Knee Surg Sports Traumatol Arthrosc* 2015;23:514-6.
8. Fontes EB, Precht BL, Andrade RC, Fernandes RM, Cisne R. Output relations of humeral circumflex arteries and its variations. *Int J Morphol* 2015;33:1171-5.
9. Keough N, de Beer T, Uys A, Hohmann E. An anatomical investigation into the blood supply of the proximal humerus: Surgical considerations for rotator cuff repair. *JSES Open Access* 2019;3:320-7.

10. Keough N, Lorke DE. The humeral head: A review of the blood supply and possible link to osteonecrosis following rotator cuff repair. *J Anat* 2021;239:973-82.
11. Kim JK, Jeong HJ, Shin SJ, Yoo JC, Rhie TY, Park KJ, et al. Rapid progressive osteonecrosis of the humeral head after arthroscopic rotator cuff surgery. *Arthroscopy* 2018;34:41-7.

Conflict of Interest: Nil

Source of Support: Nil

Consent: The authors confirm that informed consent was obtained from the patient for publication of this case report

How to Cite this Article

Aicha KB, Hassini L, Mhiri A, Hanafi A, Mtaoumi M, Bouattour K. Osteonecrosis of the Humeral Head Following Arthroscopic Repair of Rotator Cuff Tears: A Case Report. *Journal of Orthopaedic Case Reports* 2023 December;13(12): 48-52.

