



An Overview of Mycoviral Curing Strategies Used in Evaluating Fungal Host Fitness

Haris Ahmed Khan^{1,2} · Danish Ilyas Baig¹ · Muhammad Faraz Bhatti¹

Received: 4 November 2022 / Accepted: 11 February 2023

© The Author(s), under exclusive licence to Springer Science+Business Media, LLC, part of Springer Nature 2023

Abstract

The number of novel mycoviruses is increasing at a high pace due to advancements in sequencing technologies. As a result, an uncountable number of mycoviral sequences are available in public sequence repositories. However, only genomic information is not sufficient to understand the impact of mycoviruses on their host biology. Biological characterization is required to determine the nature of mycoviruses (cryptic, hypervirulent, or hypovirulent) and to search for mycoviruses with biocontrol and therapeutic potential. Currently, no particular selective method is used as the gold standard against these mycoviral infections. Given the importance of curing, we present an overview of procedures used in preparation of isogenic lines, along with their benefits and drawbacks. We concluded that a combination of single-spore isolation and hyphal tipping is the best fit for preparation of isogenic lines. Furthermore, recent bioinformatic approaches should be introduced in the field of mycovirolgy to predict virus-specific antivirals to get robust results.

✉ Haris Ahmed Khan
hkhan.phdabs13asab@asab.nust.edu.pk;
hakhan@umw.edu.pk


✉ Muhammad Faraz Bhatti
mfbhatti@asab.nust.edu.pk

Danish Ilyas Baig
danishbaig802@gmail.com

¹ Atta-ur-Rahman School of Applied Biosciences (ASAB),
National University of Sciences and Technology (NUST),
H-12, Islamabad 44000, Pakistan

² Department of Biotechnology, University of Mianwali,
Punjab 42200, Pakistan

Graphical Abstract



Keywords Mycovirus · Virus-infected/free isogenic lines · Curing · Single-spore isolation · Hyphal tipping

Introduction

Mycoviruses infecting phytopathogenic fungi have attracted huge interest from researchers in the fields of virology and pathology due to their variety of impacts on host biology [1]. Nearly all of the reported fungal groups contain mycoviruses [2, 3]. These viruses are classified into 23 different families by the International Committee on Taxonomy of Viruses (ICTV) (Table 1). Mycovirus transmission is facilitated by a number of mechanisms, including hyphal anastomosis. Mycoviruses have the power to alter fungal growth and completely halt the

virulence [4]. The potential of these viruses against fungal infections have previously exploited against fungal infections [5, 6]. A number of studies have shown the biocontrol potential of mycoviruses. The best example is Cryphonectria hypovirus 1 (CHV1), which has been used successfully in Europe to treat chestnut blight disease caused by *C. parasitica* [7]. This biocontrol was not commercialized due to the inability of CHV1 to infect distantly related fungal strains. A successful biocontrol agent should have the ability to transmit and replicate successfully in a distant fungal strain (by both extracellular and intracellular routes) [1]. Mycovirus-associated

Table 1 List of mycoviral families recognized by ICTV and their representative members

	Family	Genus	Representative member	Segment No.	Accession
ssDNA	<i>Genomoviridae</i>	Gemicirculavirus	Sclerotinia sclerotiorum hypovirulence-associated DNA virus 1 (SsHDV1)	1	GQ365709
		Gemytripvirus*	Gemytripvirus fungal (FgGMTV1)	3	MK430076; MK430077; MK430078
Reverse Transcribing RNA viruses	<i>Pseudoviridae</i>	Pseudovirus	Saccharomyces cerevisiae Ty1 virus (SceTy1V)	1	M18706
		Hemivirus	Saccharomyces cerevisiae Ty5 virus (SceTy5V)	1	U19263
	<i>Metaviridae</i>	Metavirus	Saccharomyces cerevisiae Ty3 virus (SceTy3V)	1	M34549
Double-stranded RNA viruses	<i>Reoviridae</i>	Mycoreovirus	Mycoreovirus 1 (MyRV1)	11	AY277888; AY277889; AY277890; AB179636; AB179637; AB179638; AB179639; AB179640; AB179641; AB179642; AB179643
	<i>Totiviridae</i>	Totivirus	Saccharomyces cerevisiae virus L-A (ScV-L-)A	1	J04692; X13426
		Victorivirus	Helminthosporium victoriae 190SV(Hv190SV)	1	U41345
	<i>Chrysoviridae</i>	Alphachrysovirus	Penicillium chrysogenum virus (PcV)	4	AF296439; AF296440; AF296441; AF296442
		Betachrysovirus	Botryosphaeria dothidea chrysovirus (BdCV1)	4	KF688736; KF688737; KF688738; KF688739
	<i>Megabirnaviridae</i>	Megabirnavirus	Rosellinia necatrix megabirnavirus 1 (RnMBV1)	2	AB512282; AB512283
	<i>Quadriviridae</i>	Quadrivirus	Rosellinia necatrix quadrivirus 1 (RnQV1)	4	AB620061; AB620062; AB620063; AB620064
	<i>Partitiviridae</i>	Alphapartitivirus	Rosellinia necatrix partitivirus 2 (RnPV2)	2	AB569997; AB569998
		Betapartitivirus	Atkinsonella hypoxylon virus (AhV)	2	L39125; L39126; L39127 (satellite)
		Gammapartitivirus	Penicillium stoloniferum virus S (PsV-S)	2	AY156521; AY156522
	<i>Amalgaviridae</i>	Zybavirus	Zygosaccharomyces bailii virus Z (ZbV-Z)	1	KU200450
	<i>Curvulaviridae</i> *	Orthocurvulavirus*	Curvularia orthocurvulavirus 1 (CThTV)	2	EF120984; EF120985
	<i>Polymycoviridae</i>	Polymycovirus	Aspergillus fumigatus polymycovirus 1 (AfuTmV1)	4	HG975302; HG975303; HG975304; HG975305
		Botybirnavirus	Botrytis porri botybirnavirus 1 (BpBBV1)	2	JF716350; JF716351
	Unassigned	Unassigned	Fusarium virguliforme dsRNA mycovirus 1 (FvV1)	1	JN671444
			Yado-nushi virus (YnV1)	1	LC061478
			Rosellinia necatrix megatotivirus 1 (RnMTV1)	1	LC333746
			Alternaria alternata virus 1 (AaV-1)	4	AB368492; AB438027; AB438028; AB438029
			Ustilagoidea virens RNA virus M (UvRV-M)	1	KJ101567

Table 1 (continued)

	Family	Genus	Representative member	Segment No.	Accession
Positive-sense RNA viruses	<i>Narnaviridae</i>	Narnavirus	Saccharomyces 20S narnavirus (ScNV-20S)	1	M63893
	<i>Mitoviridae</i>	Mitovirus	Cryphonectria mitovirus 1 (CpMV1)	1	L31849
	<i>Botourmiaviridae</i>	Botoulivirus	Botrytis botoulivirus (BOLV)	1	LN827955
		Magoulivirus	Magnaporthe magoulivirus 1 (MOLV1)	1	LT593139
		Scleroulivirus	Sclerotinia scleroulivirus 1 (SsOLV1)	1	KP900928
		Penoulivirus	Phaeoacremonium penoulivirus (PmOLV1)	1	MK584843
	<i>Barnaviridae</i>	Rhizoulivirus*	Rhizoctonia rhizoulivirus (RsOLV1)	1	KP900922
		Barnavirus	Mushroom bacilliform virus (MBV)	1	MBU07551
	<i>Hypoviridae</i>	Hypovirus	Cryphonectria hypovirus 1 (CHV1)	1	M57938
	<i>Endornaviridae</i>	Alphaendornavirus	Phytophthora alphaendornavirus 1 (PEV1)	1	AJ877914
		Betaendornavirus	Sclerotinia sclerotiorum betaendornavirus 1 (SsEV1)	1	KJ123645
		Botrexvirus	Botrytis virus X (BotVX)	1	AY055762
		Sclerodarnavirus	Sclerotinia sclerotiorum debilitation-associated RNA virus (SsDRV)	1	AY147260
	<i>Gammaflexiviridae</i>	Mycofflexivirus	Botrytis virus F (BVF)	1	AF238884
	<i>Deltaflexiviridae</i>	Deltaflexivirus	Sclerotinia sclerotiorum deltaflexivirus 1 (SsDFV1)	1	KT581451
	<i>Unassigned</i>	Unassigned	Diaporthe RNA virus (DRV)	1	AF142094
			Fusarium graminearum virus DK21 (FgV/DK21)	1	AY533037
			Oyster mushroom spherical virus (OMSV)	1	AY182001
			Hadaka virus 1 (HadV1)	11	LC519840; LC519841; LC519842; LC519843; LC519844; LC519845; LC519846; LC519847; LC519848; LC51984; LC519850

Table 1 (continued)

	Family	Genus	Representative member	Segment No.	Accession
Negative-sense RNA viruses	<i>Mymonaviridae</i>	<i>Sclerotimonavirus</i>	<i>Sclerotinia sclerotimonavirus</i> (SsNSRV-1)	1	KJ186782
		<i>Botrytimonavirus</i> *	<i>Botrytimonavirus botrytidis</i> (BcNSRV-7)	1	MT157413
		<i>Lentimonavirus</i> *	<i>Lentinula lentimonavirus</i> (LeNSRV-1)	1	LC466007
		<i>Phyllomonavirus</i> *	<i>Phyllomonavirus phyllosphaerae</i> (SLaNSRV-4)	1	KT598229
		<i>Auricularimonavirus</i> *	<i>Auricularimonavirus auriculariae</i> (AhNSRV-1)	1	MT259204
		<i>Penicillimonavirus</i> *	<i>Penicillimonavirus alphanenicillii</i> (PdNSRV-1)	1	MK584858
		<i>Plasmopamonavirus</i> *	<i>Plasmopamonavirus plasmoparae</i> (PvLAMV8)	1	MN557004
	<i>Phenuiviridae</i>	<i>Lentinivirus</i>	<i>Lentinula lentinivirus</i> (LeNSRV-2)	2	LC466008; LC466009
		<i>Entovirus</i>	<i>Entoleuca entovirus</i> (EnPLV-1)	2	MF375882; MK140653
	Unassigned	Unassigned	<i>Fusarium poae</i> negative-stranded RNA virus 1 (FpNSRV-1)	1	LC150618

*Taxa that are being suggested for creation or reclassification

hypovirulence has also been proposed to be used as a therapeutic against human pathogenic fungal infections [8–10]. Recent studies have reported that human pathogenic fungal strains are hosts to diverse mycoviruses [11]. The process of controlling human pathogenic fungi would be similar to that of phage therapy, where bacteriophages are used to target and cure selective bacterial infections [12]. Studies have reported that *Saccharomyces cerevisiae* (yeast) harboring mycoviruses are reported to produce killer toxins that are lethal to other sensitive fungal strains but have no influence on host fungi [13, 14]. Mycoviruses also induce interferon production in *Malassezia sympodialis* and *Penicillium stoloniferum* [15, 16]. The significant sequence similarity of *Sclerotinia sclerotiorum* RNA virus L (SsRV-L) to human Hepatitis E virus suggests the possibility of mycoviral replication in humans, but more research is needed in this scenario [9]. Mycovirus from *Aspergillus flavus* reproduced surface and genetic markers in the cells of patients suffering from acute lymphoblastic leukemia (ALL) disease. This character can be used as a test for the recognition of ALL patients in remission [17]. Mycovirus therapy against pulmonary aspergillosis has previously been reviewed in detail by Wendy et al. (van [18]. These properties of mycoviruses can also be utilized

to enhance agricultural, industrial, and pharmaceutical applications [19].

Extraction and gel electrophoresis of dsRNA elements are one of the most widely used techniques for detecting mycoviruses [20]. The only limitation is that it can solely detect ssDNA and dsRNA mycoviruses, while those viruses that do not accumulate dsRNA replicative form can only be detected by high-throughput sequencing [21]. The second step to determine the nature of mycoviruses is biological characterization, which is an important step toward determining the impact of mycoviruses on host fitness. The effect of mycoviruses on host (fungus) biology in natural conditions is a matter of great mystery. Mycoviruses can be classified as being either hypervirulent, latent (cryptic), or hypovirulent. These terms state the changes in pathogenicity of the fungal host and do not correlate with the virulence of mycoviruses, thus creating an unclear image of mycoviruses. *Cryphonectria parasitica* hypovirus 1 (CHV1) infecting American chestnut (*Castanea dentata*) was reported to decrease the pathogenicity, when compared to virus-free *Cryphonectria parasitica* and was classified as a hypovirulent mycovirus [7]. Similarly, *Nectria radicola* virus L1 (NrV-L1) induced greater pathogenicity in *Nectria radicola* fungus while infecting the plants of *Panax ginseng* (ginseng) and was reported as a hypervirulent mycovirus [22]. Many mycoviruses are

cryptic or latent mycoviruses as they do not show an obvious effect on the virulence of fungal hosts [23, 24]. It cannot be said with certainty that the effect of mycoviruses on fungal biology has a direct link to the fungus pathogenicity on its respective host. For instance, the growth rate of a *Monilinia fructicola* (the fundamental agent of brown rot on *Prunus* species) was increased (in vitro) via co-infection by three mycoviruses, but there were no visible effects on its lesions growth patterns on plums when infected by its isogenic lines [25]. To effectively investigate the impact of viruses on their hosts, virus-infected and virus-free isogenic lines are required. The communal method is to eliminate the virus from infected strains or introduce the virus into virus-free fungal isolates. To date, the available literature is lacking in collective information regarding the techniques for preparing isogenic lines. The purpose of this review is to highlight the gaps and to provide a deep insight into mycoviral curing strategies to researchers in the field of fungal virology. This review covers almost all information related to the available methods used in curing mycoviral infections. Furthermore, we also describe the processes, underlying mechanisms, and their pros and cons. The information will be helpful for researchers in selecting a suitable curing method for the understudied fungal strains to evaluate the effect of mycoviruses on host biology.

Drugs Used to Treat Mycoviral Infections

A variety of antiviral drugs are developed by the scientific community to limit the spread of viruses by halting their replication [26]. An antiviral drug is a substance (small or large molecule(s), natural or synthetic) that has the ability to lessen the spread of infectious disease triggered by a virus [27]. The use of already-available drugs and antibiotics to treat mycoviral infections has been widely documented, despite the fact that no specific drugs have been specifically designed to limit mycoviral infections. Curing mycoviral infections using drugs is achieved by inoculating dsRNA-positive mycelial plugs on PDA plates containing an optimized amount of drug (varies for different fungal hosts; needs optimization, no standard available) and incubating at optimized temperature and light conditions [28]. After incubating for a few days, small marginal mycelial plugs are transferred to new PDA plates and examined for the presence of mycoviruses through RT-PCR, dsRNA, or total RNA extraction.

Cycloheximide is a naturally occurring antibiotic produced by the *Streptomyces griseus* bacterium. It blocks eukaryotic translational elongation in protein synthesis by interfering with the translocation step, i.e., the movement of mRNA and two tRNA molecules with respect to the ribosome [29]. Therefore, the primary function is to

inhibit protein synthesis. Binding takes place at the ribosome and prevents eEF2-mediated translocation [30]. It is reported that the synthesis of dsRNAs in fungi is inhibited by cycloheximide treatment, which has been widely used previously. Cycloheximide prevents both the replication and translation of the viral genome from the negative-strand RNA by preventing the clearance of ribosomes from the viral genome (RNA) thus freezing ribosomes [31]. Successful elimination of mycoviruses has been reported from *Aspergillus fumigatus* [28] and *Lentinula edodes* [32]. It is also interesting to note that cycloheximide is not successful for curing all fungal strains, even at higher concentrations, e.g., *Metarhizium anisopliae* [33], *Aspergillus niger* [34], and *Pseudocercospora griseola* [35]. Partial success has been reported for *Pseudogymnoascus destructans* [36].

5'-Fluorouracil (FU) is a pyrimidine analog used in cancer chemotherapy. The basic mechanism of FU is considered to be the establishment of a covalently bound ternary complex achieved by the binding of the drugs deoxyribonucleotide (FdUMP) and N5,10-methylenetetrahydrofolate (the folate cofactor) to thymidylate synthase (TS) [37]. This inhibits the formation of thymidylate from uracil, leading to the inhibition of RNA and DNA synthesis and ultimately leading to cell death. Fluorouracil is also able to interfere with RNA processing and protein synthesis by generating a fraudulent RNA by replacing uridine triphosphate (UTP) in RNA [38]. FU can be broken down into 5-fluorouridine triphosphate, which is a substrate for viral RNA-dependent RNA polymerases. As a result, mutations are incorporated into viral genome, which are lethal for virus survival and result in loss of infectivity [39]. Ahmad et al. demonstrated that FU was ineffective against coronaviruses due to their ability to proofread. They also suggested that FU produces promising results when used in combination with deoxynucleosides [40]. The replication of *Penicillium stoloniferum* fast-moving virus (PsVf) was inhibited by FU treatment. FU was also successful in curing a virus from *Aspergillus flavus* [41]. FU is no longer used to prepare virus-free isogenic lines because it is ineffective against mycoviral infections.

Ribavirin (also known as tribavirin) is commonly used to treat hepatitis C, RSV infection, and some viral hemorrhagic fevers. Sometimes it is also used in combination with other drugs to get better results. Ribavirin is also reported to cure mycoviral infections. It accomplishes its antiviral activity by restraining the access of inosine monophosphate dehydrogenase (IMPDH) to its endogenous substrate, inosine-5-monophosphate through the inhibition of the enzyme [42]. This leads to a condition with minimized levels of intracellular guanosine triphosphate (GTP) pools that are required for viral replication. Studies have also indicated that the ribavirin antiviral activity might be due to inhibition of viral polymerase activity, capping of viral transcripts, or overwhelming the humoral and cellular immune responses.

Recent studies relate its primary antiviral mechanism to the deadly mutagenesis of viral RNA genomes. Being a nucleoside analog of guanosine, it also inhibits the viral polymerase [43]. A virus-free isogenic line of *Tolypocladium cylindrosporum* was prepared using ribavirin treatment [44]. *Lentinula edodes* was reported to carry a co-infection of two viruses: Lentinula edodes mycovirus HKB (LeV-HKB) and Lentinula edodes partitivirus 1 (LePV1) [44, 45]. Ribavirin treatment was only successful in curing LeV-HKB infection, while both viruses were lost during mycelial fragmentation [46]. In *Aspergillus* species, ribavirin treatment failed to eliminate mycoviruses belonging to *Fusariviridae*, *Mitoviridae*, and *Hypoviridae* [47].

Streptomycin and kanamycin have similar modes of action for viral elimination, i.e., halting protein synthesis by attaching to the ribosomal 30S subunit [48]. The infection of *Sclerotinia sclerotiorum* hypovirus 2 (SsHV2) and Botrytis virus F (BVF) in *Monilinia fructicola* fungus was eliminated by treatment with both of these antibiotics but was not successful in eliminating *Fusarium poae* virus 1 (FpV1) infection [25]. Partitiviruses are known to cause persistent infection and are difficult to eradicate [25, 49]. Rifampicin (a derivative of rifamycin) averts the attachment of RNA polymerase to DNA in order to act as an inhibitor of RNA viruses [50, 51]. Its curing mechanism is not clearly established, but in a minimal medium, a treatment of rifamycin with cAMP has been reported to cure a culture of the edible mushroom (*Pleurotus ostreatus*) from the RNA mycoviruses, i.e., oyster mushroom isometric virus (OMIV) and oyster mushroom spherical virus (OMSV) [52]. Multiple virus infections in *Ceratobasidium* sp. were eradicated using hyphal tipping and various antibiotics and drug treatments. Unexpectedly, the three mycoviruses reacted differently to each curing approach. The isolate containing *Ceratobasidium endornavirus* C (CbEVC) was eliminated by cycloheximide but retained by kanamycin or streptomycin. However, the isolate of CbEVD maintained stable replication with cycloheximide treatment. Interestingly, CbEVB was eliminated with all treatments [53, 54].

Single-Spore Isolation

Fungi produce an uncountable number of sexual (basidiospores and ascospores) and asexual spores (conidiophores) for propagation. Mycoviruses utilize these spores for their transmission (vertical transmission) to the next generation. Asexual (conidiophores) spores are most frequently used by mycoviruses for transmitting to progeny cells. The transmission rates vary among different mycoviruses, they range from 0% [23, 24] to 100% [55]. Single-spore isolation is commonly used to prepare virus-free and virus-infected isogenic lines for phenotype comparison and analyzing the

impact of mycoviruses. Spores from the mother plate are collected and a stock spore suspension is prepared by adding them to 1 ml of sterile distilled water. The spore suspension is serially diluted as required (based on haemocytometer spore count concentration) in sterilized double-distilled water and subsequently spread on PDA plates for growth. Several single conidiophores are picked from PDA media and analyzed via classical dsRNA extraction or one-step RT-PCR for detection of mycoviruses [56–58]. It is interesting to note that the virus vertical transmission rate decreases with the age of mycelia (in most cases) [23, 24, 59, 60], but this is not true for all fungal strains [23, 24]. Successful curing has been achieved in numerous studies, few examples include *Magnaporthe oryzae* [61], *Fusarium nygamai* [23, 24], *Fusarium oxysporum* [59], *Aspergillus fumigatus* [62], *Trichoderma harzianum* [63], and edible mushrooms [64]. In some cases, single-spore isolation is reported to be unsuccessful, i.e., *Aspergillus niger* [34], *Hypomyces chrysospermus* [65], *Alternaria alternata* [57, 66–69], and *Neofusicoccum parvum* [70, 71] and in some cases, partial success has been achieved, e.g., *Diplodia seriata* [71]. Some fungal strains produce asexual structures such as pycnidia, which yield pycnidiospores (asexual spores). These are macroscopic, hard, and solid round bodies, black or dark brown in color [72, 73]. Pycnidia have a diameter of 60–200 µm and have a sub-globose shape, with the presence of an ostiole either below or bulging through the stomatal pore [73, 74]. An estimated 10,000 pycnidiospores are produced by each pycnidium, which are projected through the ostiole in a gelatinous matrix, during high humidity conditions [75]. e.g., *Sclerotinia sclerotiorum* strain XG36-1 was allowed to grow on sterilized carrots for 30 days in order to produce pycnidia. Harvested pycnidia were soaked in water for hydration, causing spores to be released. These spores were utilized for single-spore isolation [76]. When compared to asexual spores in the same species, vertical mycovirus transmission through sexual spores appears to be much less effective. For example, in several species of *Aspergillus*, the majority of sexual spores are virus free, whereas asexual spores of the same species are infected [77, 78]. The exclusion of dsRNA segments from sexual spores was further observed in *Ophiostoma ulmi* [79], *Gaeumannomyces graminis* strains [80], and *Helicobasidium mompa* [81]. However, in some fungal strains the transmission through sexual spores is very efficient, e.g., *Ustilago maydis* [82] and *Heterobasidion annosum* [83], *Saccharomyces cerevisiae* [84], and *Fusarium graminearum* [85].

Hyphal Tipping

The growing tips of fungi sometimes do not contain mycoviral infection, similar to the apical meristem in plants [86, 87] but this is not true for all mycoviral infections (dependent on virus replication and titer). Being cytoplasmic residents, mycoviruses are expected to move in tandem with fungal nuclei via the cytoplasmic streaming movement, as reviewed by Xiang [88]. So it is possible to get virus-free fungal strains from marginal mycelia (tips). To obtain virus-free isogenic lines, fungal strains are grown on water agar medium and incubated at the optimum temperature. Water agar is deficient in nutrients and provides a kind of stress condition by depriving fungi of nutrients, due to which fungi spread branches in search of nutrition. After 2–4 days of growth, the growing tips are dissected (~1 mm) under a binocular microscope and transferred to fresh water agar plates [89]. For virus-free mycelial collection, all regenerated isolates are randomly shifted to potato dextrose broth (PDB) culture and then screened for mycoviral infection. This process can be repeated 4–5 times or until virus-free cultures are obtained [53]. *Alternaria alternata* virus 1 (AaV-1) was cured using a mixed approach of hyphal tipping and cycloheximide treatment [90]. Hyphal tipping was not successful for *Sclerotium sclerorum* and *Rhizoctonia solani* viruses [91, 92] but in most studies curing was unsuccessful [34, 93–97]. By inducing branching activities through antibiotic treatment, the number of hyphal tips should increase through rapid cell division, thus improving the chances of producing virus-free cultures from hyphal tips.

Protoplast Regeneration

The cell wall is a significant cellular structure in organisms, such as bacteria, fungi, algae, and plants. It serves a variety of purposes, including providing permeability barriers, assisting with cell design, and protecting cells in hazardous conditions [98]. The cell wall-deprived cells are termed “protoplasts” [99–101]. Similar to plant protoplasts fungal protoplasts also exhibit totipotency and can also be stimulated to produce new individuals by exposing them to appropriate external stimuli [102]. Different enzymatic methods can be applied to remove cell wall and produce protoplasts [99–101]. The cell wall of fungi are made up of polymers, i.e., dextran, mannose, chitin, and mannoproteins. Cell wall chemical composition varies from fungus to fungus and is dependent on cell development patterns and growth conditions [98]. The digestive enzymes required, such as cellulases, proteases, and chitinases, differ based on cell wall structure. As a result, the enzymatic mixture chosen from the suitable components and ratios is a critical aspect of

protoplast formation. Furthermore, during cell wall breakdown, protoplasts are extremely susceptible to osmotic pressure. To maintain protoplast integrity throughout the preparation procedure, an isotonic solution of particular sugars or salts is necessary. Osmotic stabilizers are chemicals that are used to stabilize osmotic pressure, maintain cell shape, and prevent membrane breakdown.

The protoplast fusion procedure has enabled researchers to overcome the vegetative incompatibility barrier and facilitate the transmission of viruses across different fungal species/strains. For example, dsRNA from *Fusarium boothii* was transferred via protoplast fusion to other *Fusarium* species as well as the model fungus *C. parasitica* [103]. Besides this, the transfection procedure serves the purpose of introducing virus into virus-free fungal strains for host range extension, as demonstrated by Rosellinia necatrix partitivirus 1 and mycoreovirus 3 in *C. parasitica*, *Diaporthe* sp., and *Valsa ceratosperma* [104]. Protoplast regeneration has been previously used to cure mycoviral infections. Virus-free isogenic lines were successfully prepared for *F. graminearum* strain HN1 using the protoplast regeneration method [105, 106]. No universal protocol is currently available as cell wall composition varies from fungus to fungus, so different digestion enzymes are required. Preparation of protoplasts (removing the cell wall) is chiefly achieved through enzymatic treatment. But physical methods like grinding and supersonic wave shocks have also been reported. However, due to practical inconvenience and the low yield of protoplasts, these methods are not widely used. A good quality protoplast can only be obtained from fresh mycelia [103]. Successful curing of mycoviral infection has been reported by [89]. An alternative way to transfer hypovirulence within a population of *C. parasitica* has been the construction of infectious cDNA clones of hypoviruses and the subsequent transgenic lines [107, 108]. An infectious clone of BVF successfully replicated in the protoplasts of *Botrytis cinerea* and maintained a stable infection [109]. CHV1 (EP713) and MyRV1 (9B21) were transferred to *Valsa mali* by fusing the protoplasts of *C. parasitica* (EP713 and 9B21) with *Valsa mali* protoplasts. The co-infection of both viruses reduced the growth and virulence of *Valsa mali* [110].

Mycelial Fragmentation

Mycelial fragmentation can be performed in a blender [111] or using screw-cap tubes and glass beads [112]. Scraped mycelia from PDA plate cultures can be transferred to bead-containing tubes, which were filled with sterile water and shaken for 10 s using a Mini-Bead Beater. Dilutions prepared from this solution can be spread on PDA plates to check for mycoviral infection using RNA extractions and




Fig. 1 A microscopic representation of **a** Spores of *Fusarium* sp. **b** Hyphal tips of *C. parasitica*. **c** Protoplast of *C. parasitica*. **d** Mycelial fragments of *Neofusicoccum parvum* (Khan et al., unpublished data)

PCR. The procedure is similar to that of single-spore isolation method. Successful curing of *Pleurotus ostreatus* virus (PoV) has been reported in *Pleurotus ostreatus* [113]. Prolonged incubation of mycelia results in mostly virus-free progenies. The use of mycelial fragmentation is reported in combination with drug treatment as has been reported for *Lentinula edodes* [114]. A microscopic representation of spores, hyphal tips, protoplasts, and mycelial fragments is shown in Fig. 1.

Polyethylene Glycol (PEG)-Mediated Stress

To generate water stress, water restrictors or osmotic stress inducers are widely utilized as media supplements. One of them is polyethylene glycol (PEG), which is a long-chain neutral polymer with impermeability and is soluble in water. It leads to stress in cell cultures that is similar to the drought stress seen in the cells of whole organisms [115–118]. PEG lowers the osmotic potential of the culture medium, inhibits water absorption, and is not digested by fungal strains. However, the appropriate concentrations of the inducers must be considered: the perfect concentration is one that is high enough to fulfill the experiment's goals while remaining low enough that no cultured organisms totally stop developing [119]. Polyethylene glycol (PEG) compounds are non-toxic, water-soluble polymer with different molecular weights and numerous applications [120]. PEG has been used in vivo and in vitro in different models of tissue injury and has shown many interesting biological properties; it has cytoprotective, anti-oxidant, immunosuppressive, and anti-inflammatory effects [121, 122]. PEG could create a

neutralization barrier that prevents coronavirus recognition of its receptors and thus inhibiting virus entry and invasion. PEG could inhibit virus adhesion by acting as an immunocamouflage. The mode of this protection is biophysical and depends on charge mass and steric hindrance induced by the polymer. Immunocamouflage depends on molecular weight: high molecular weight PEG of 10 kDa to 35 kDa are better absorbed and consequently, they are more effective [123, 124]. It also depends on the cell surface type: small molecular weight PEGs with a molecular weight of 2 kDa were effective at binding to respiratory syncytial virus (RSV), but completely ineffective in the host cell [124].

PEG-mediated stress is reported to cure mycoviral infections, but the underlying mechanism needs to be explored. Considering the example of *Pseudogymnoascus destructans*, which was infected with *Pseudogymnoascus destructans* partitivirus-pa (PdPV-pa), whose infection, was not cured by any available methods of curing (i.e., heat therapy, nutritional and chemical stress, including cycloheximide and ribavirin, hyphal tip culture, single-spore isolation, and protoplast culture). Finally, infection was cured using PEG-induced matric potential in MM (minimal nutrition media). Different matric potential gradients were used to check for PdPV-pa infection in *P. destructans*. No visible germination of *P. destructans* mycelial growth or conidia production was observed at 5 MPa and 6 MPa. The characteristic gray pigmentation of wild-type Pd isolates turned to white in PdPV-pa free isolates after PEG treatment. The virus-free isolate also produced ominously fewer conidia than the wild-type isolate [36]. *Phytophthora cactorum* bunya-like viruses 1 and 2 (PcBV1 and 2) infecting *Phytophthora cactorum* were cured using PEG 8000-mediated stress [125].

Thermotherapy: Heat or Cold Treatment

Plants cultured at high temperatures (40 °C) for an extended period of time may become virus free. At higher temperatures, the activity and synthesis of virus-encoded coat proteins (CPs) and mobility proteins (MPs) are disrupted [105, 106]. When combined with the tissue culture technique, thermotherapy is more effective [126]. Apparently, the detrimental effect of high temperatures on viral proliferation and dissemination in plants reduces their concentration following heat treatment. However, viruses that require elevated temperatures for reproduction and accumulation in plant tissues increase their concentration rather than being destroyed by heat treatment, i.e., Potato spindle tuber viroid (PSTVd) [127]. They can be eliminated by meristem excision at low temperatures (between 5 °C and 10 °C). The heat tolerance of the host and the type of virus determine the duration and intensity of thermotherapy [87]. For viral eradication in potatoes, meristem culture combined with thermotherapy is often utilized. This process is completed

by incubating virus-infected plants in a growth chamber for 2–6 weeks at a light intensity of 30–50 mol m² s⁻¹ and a temperature of 35 °C–37 °C. Following the corresponding duration of thermotherapy, the meristems are removed and grown on nutritional medium for regeneration [128, 129]. Virus elimination has also been observed when fungal samples are stored at lowered temperatures or when stored in glycerol stocks at – 80 °C [25].

Malassezia sympodialis harboring *Malassezia sympodialis* mycovirus 1 (MsMV1) was cured by exposing it to high temperatures (37 °C). At this temperature, virus replication and persistence may interfere with important biological functions. Furthermore, high temperatures are thought to increase heat shocks and misfolded proteins in the cell [130], leading to virus loss owing to overworked Hsp90 if complete Hsp90 activity is necessary for viral replication [15]. Golubev et al. failed to cure *Alternaria alternata* partitivirus 1 (AtPV1) infection using a compound approach of thermotherapy, chemotherapy, spheroplast regeneration, and PEG-induced matrix stress [131]. The same was observed for *Sclerotinia sclerotiorum* mitovirus 1 (SsMV1/KL-1) and SsMV2/KL-1 infecting *S. sclerotiorum* [132].

UltraViolet (UV) Light Exposure

Viruses are more vulnerable to UV radiation than red and white blood cells because they absorb significantly more energy. The killed viral fragments elicit a vaccination-like response, therefore boosting the immune system and producing resistance to the specific pathogen. UV radiation has a profound impact on airborne viruses. Far-UV radiation has the ability to destroy viruses. [133]. UV light has also been used to sterilize food items. The use of UV light for airborne disinfection was first shown more than eight decades ago [134]. Drug-sensitive, multidrug-resistant bacteria and different strains of viruses are inactivated by germicidal UV light [135]. Water is disinfected using germicidal lamps and UV radiation with a wavelength of 240 to 280 nm. Spectroscopy is often used to measure the average amounts of nucleic acids (DNA or RNA) present in a mixture, as well as their purity; the peak of DNA and RNA absorption is at 260 nm. UV germicidal activity is a well-known method for inactivating or killing microorganisms by causing nucleic acid damage. UV radiation with a wavelength of 300 to 320 nm is employed for light treatment in medical applications, e.g., 311 nm [136]. Ultraviolet B (UVB) narrowband lamps are used in phototherapy to treat T-cell cutaneous lymphoma and psoriasis, such as mycosis fungoides [137]. UV radiations have not been widely reported to be used for curing mycoviral infections, but a partial success was obtained by Castillo and colleagues [138]. It is not a widely used technique for eliminating mycoviruses as increasing UV light exposure may also kill fungi.

Co-Culture Transmission

Mycoviruses with persistent infections (from the families *Partitiviridae* and *Endornaviridae*) are difficult to eradicate using current curing methods. However, they can be transmitted to other compatible fungal strains to prepare virus-free and infected isogenic lines for evaluating their effect on host fitness. The dual-culture or co-culture technique is mostly used to transfer viruses to compatible fungal strains [128, 129, 139, 140]. However, depending on virus infectivity and host defense mechanisms, viruses may not be transferred to compatible strains [68, 131]. The virus-infected strain acts as a donor, while the other strain functions as a recipient. Both fungal strains are cultured side by side on the same PDA plate and allowed to grow under optimum conditions. After successful fusion of hyphae, three locations on the recipient side (i.e., near, far, and middle regions) are analyzed for the detection of mycovirus transmission. One-step RT-PCR or dsRNA extraction can be performed to check for the successful transmission of mycoviruses. The transmission of mycoviruses is hindered by vegetative incompatibility (VIC) among distant fungal strains. Allorecognition, or non-self-recognition, is a common phenomenon in the fungal kingdom, allowing them to distinguish one another [141]. This system is thought to have evolved to limit the spread of harmful organisms, like mycoviruses [142]. In many model organisms, this vegetative/heterokaryon incompatibility system is under the regulation of multiple allelic or non-allelic *vic* or *het* genes and most species seem to involve different genes [141, 143]. Non-self-recognition leads to compartmentalization between two isolates of dissimilar mycelial compatibility, leading to programmed cell death in order to interrupt fusion between hyphae and is known as heterokaryon incompatibility [144]. The mycovirus transmission between *Aspergillus* and *Cryphonectria* was unsuccessful [77, 145]. A recently reported virus, *Cryphonectria naterciae* fusagravirus (CnFGV1) infecting *Cryphonectria naterciae* was reported to cross vegetative compatibility barrier and was transferred to *Cryphonectria carpnicola* and *Cryphonectria radicalis* during co-culture [146]. Zinc chloride is previously reported to suppress the incompatible reaction (VIC) between two fungal strains and increase hyphal anastomosis, which can help in successful transmission of mycoviruses [147]. Zinc ions are also known to inhibit apoptosis by targeting caspase-3 activation in mammalian cells [148]. Some mycoviruses are found to be responsible for suppressing the fungal host's non-self-recognition, enabling the heterologous transmission of mycoviruses. *Sclerotinia sclerotiorum* mycoreovirus 4 (SsMYRV4), associated with hypovirulence in *Sclerotinia sclerotiorum*, has been reported to suppress non-self-recognition of the host as well as enable viral co-infection across vegetative

incompatible groups, through horizontal transmission of mycoviruses [149]. Vegetative incompatibility genes (*het* or *vic*) and proteins (heterotrimeric G proteins) are inhibited by SsMYRV4 [149]. In vitro horizontal transmission of mycoviruses is a commonly used technique in mycovirology. Carbone et al. found that transmission may occur in one direction, while it may not be as efficient in the reciprocal pairing [150].

Fungal Phenotyping and Pathogenicity Testing

After obtaining virus-infected and virus-free isogenic lines, fungal strains are evaluated by phenotype comparison and pathogenicity testing. Changes in colony diameter, pigmentation, and sporulation are usually assessed for determining the nature of mycovirus. The pros and cons of available methods are mentioned in Table 2. Mostly conidial suspension or mycelial plugs are used for pathogenicity testing [151, 152]. The lesion diameters of virus-infected and virus-free fungal isolates are compared in order to determine the impact of mycovirus on the pathogenicity of its host. There is a complex correlation between the occurrence of mycovirus and its effects on the fungal hosts, but overall, these

effects are categorized as hypovirulence, hypervirulence, and cryptic. It is still unclear how mycoviruses regulate hosts biology [153]. In the case of a single infection, assessing the effect of mycovirus is very simple but in multiple infections, it is very difficult to link a particular phenotype with understudied mycoviruses.

Challenges in Curing Mycoviral Infections

After curing attempts, mycoviruses can be found differently distributed in the mycelia due to which they are not detected with one-step RT-PCR and can reappear after subculturing or when antibiotic stress is removed. Mycoviruses have previously been reported to reappear in various fungal strains. The titer of a 12-kb dsRNA virus in *Rhizoctonia solani* went below the level of detection by RT-PCR using hyphal tipping and extended incubation on cycloheximide media. But the virus was detected again after subculturing [154]. In another case, mycovirus was detected again in a previously reported cured isolate [154, 155]. The infection of *Agaricus bisporus* endornavirus 1 (AbEV1) was cured with hygromycin B, but recovered after one month when the antibiotic stress was removed [156]. Similarly, a co-infection of two mycoviruses in

Table 2 Pros and Cons of techniques available for curing mycoviral infections

Technique	Pros	Cons
Drugs/Antibiotics	Easy to use High success rate when virus-specific drugs are used	Development of resistance Least focused in mycovirology Some drugs may kill fungal strain
Single-spore Isolation	High success rate Easy to use Not expensive Not harmful for fungal strain	Time consuming Cannot be used for fungal strains which do not produce asexual spores on growth media
Hyphal Tipping	Mostly used for those fungal strains which don't produce asexual spores on growth media Not expensive Easy to use	Time consuming
Protoplast Regeneration	cDNA infectious clones of mycovirus can easily infect fungal protoplast Protoplast fusion, sometimes produce virus-free fungal strain	Sensitive process, require careful handling Low success rate Required chemicals could be expensive
Polyethylene Glycol (PEG)	Easy to use	Low success rate Not widely used against mycoviruses
Thermotherapy	Easy to use Not expensive	Exposure to high temperatures may kill fungal strains
UltraViolet Light	Easy to use	Long exposure duration may kill fungal strain and small duration may be ineffective in curing Needs optimization
Mycelial Fragmentation	Is mostly used for those fungal strains which don't produce asexual spores on growth media	Time consuming Low success rate
Co-culture Transmission	Successful in transferring mycovirus to compatible fungal groups	Mycovirus cannot be transferred to compatible fungal strains

Pseudogymnoascus destructans was cured using cycloheximide and ribavirin, but they reappeared after removal of antibiotic stress [36]. Although it was hypothesized that the use of antibiotics reduced the concentration of dsRNA below the threshold for RT-PCR detection, no quantitative real-time qRT-PCR assays were conducted to assess the viral titers before and after treatments. The existence of integrated DNA copies of the dsRNA components might be a less reasonable explanation for the apparent reintroduction of dsRNA elements from fungus [154]. This argument is supported by the indication of sequence identity between *R. solani*'s dsRNA segments and portions of its genome [157]. More proof for this conceivable explanation would come from an analysis of the alignment between the mycovirus genome sequences and the fungal host genome.

The second problem is vegetative incompatibility, which we already discussed in detail. Some mycoviruses can successfully transfect protoplasts of distant fungal strains but are knocked out after subsequent subculturing. Here, we consider the example of *Botrytis gemydayirivirus* 1 (BGDaV1), whose virions successfully infected the protoplasts of *Botrytis cinerea* but were lost after the third subculture [158]. *Neofusicoccum parvum victorivirus* 3 (NpVV3) virions also successfully infected $\Delta dcl2$ mutant strain of *C. parasitica* but was lost after repeated subculturing [70, 71]. Similar results were observed for *Rosellinia necatrix victorivirus* 1 (RnVV1) [93]. These results suggested that host RNA silencing works as a counter defense against these viruses [159].

Conclusions

Mycoviruses are the most neglected group in the fields of virology and pathology, due to which our knowledge of global viral diversity in nature is still very incomplete and biased. Nevertheless, there has been a significant advancement in the taxonomy of mycoviruses based on genomic sequences and biological traits, including host antiviral responses. Recent studies have also highlighted the significance of mycoviruses as therapeutic and preventive agents against human and phytopathogenic fungal infections. Considering the importance of mycoviruses, it is very important to investigate the effects of mycoviruses that they induce in their hosts. Preparing virus-free and infected isogenic lines is the first step toward determining the nature of mycoviruses. Secondly, this process also results in the identification of mycoviruses with biocontrol potential. Previously, several methods were used to obtain virus-free isogenic lines of fungal strains. Cycloheximide and Ribavirin are among the most commonly used drugs against mycoviruses. Unfortunately, mycoviruses are developing resistance against available drugs. The unavailability of mycovirus-specific drugs is one

of the limiting factors in biological characterization. Thermotherapy, UV treatment, and PEG-induced matrix stress are all considered primitive techniques that are now rarely used and not very effective. Single-spore isolation, hyphal tipping, and mycelial fragmentation are among the most frequently used techniques and are considered more effective compared to other available techniques. Compared to drug treatment, the later techniques are more time consuming and labor intensive. The majority of studies revealed that combining these techniques yielded more effective results. Co-culture transmission assays are used to transfer viruses (with persistent infections) to other virus free fungal strains to investigate their impact on host biology by comparing the colony diameter. Pathogenicity tests are also performed to assess the pathogenicity of fungal strains that have and have not been infected with a virus. The size of the lesion will determine the cryptic hypo- and hypervirulent effects.

Future Directions

Recent advanced bioinformatics approaches, including molecular docking and simulation studies, should be introduced to design potential ligands or drugs to inhibit the replication of mycoviruses specifically by binding to their RNA-dependent RNA polymerase (RdRP), which is commonly present in all mycoviruses. Furthermore, suppressors of vegetative incompatibility genes (VIC) can also be designed, which will allow researchers to study the effect of viruses on a wide range of hosts. The field of nanotechnology can also be utilized for curing mycoviral infections using chemically coated nanoparticles that have antiviral activity and can specifically target mycoviruses. Polyethylene glycol-coated zinc oxide nanoparticles are already used as an effective nanoweapon to fight against herpes simplex virus type 1 [160]. Based on these suggestions, we will achieve a broader picture of viral cross-species transmission, which should in turn inform viral emergence studies. Progression in the field of mycovirolgy will lead to a tremendous change and soon mycoviruses will be used as bioweapons against fungal infections.

Acknowledgements We acknowledge National University of Sciences and Technology (NUST), Islamabad, Pakistan for providing research facilities.

Author Contributions Conceptualization: HAK. Literature search: HAK and DIB. Writing—original draft preparation: HAK and MFB. Critically revised the work: HAK and MFB.

Funding No particular funding was received from any public or private organization.

Data availability Data are available on request due to privacy or other restrictions.

Declarations

Conflict of interest The authors declares no conflicts of interest.

References

- García-Pedrajas, M. D., Cañizares, M. C., Sarmiento-Villamil, J. L., Jacquat, A. G., & Dambolena, J. S. (2019). Mycoviruses in biological control: From basic research to field implementation. *Phytopathology*, 109(11), 1828–1839. <https://doi.org/10.1094/phyto-05-19-0166-rvw>
- Kondo, H., Botella, L., & Suzuki, N. (2022). Mycovirus diversity and evolution revealed/inferred from recent studies. *Annual Review of Phytopathology*, 60, 307–336.
- Son, M., Yu, J., & Kim, K.-H. (2015). Five questions about mycoviruses. *PLOS Pathogens*, 11(11), e1005172. <https://doi.org/10.1371/journal.ppat.1005172>
- Kotta-Loizou, I., & Coutts, R. H. (2017). Mycoviruses in *Aspergilli*: A comprehensive review. *Frontiers in Microbiology*, 8, 1699.
- Ghabrial, S. A., Castón, J. R., Jiang, D., Nibert, M. L., & Suzuki, N. (2015). 50-plus years of fungal viruses. *Virology*, 479, 356–368.
- Umer, M., Qadeer, A., Razaq, Z., Anwar, N., & Kiptoo, J. J. (2023). Mycovirus: Biocontrol agent against *Sclerotium* of Rapeseed. *Phytopathogenomics and Disease Control*, 01, 97–108.
- Rigling, D., & Prospero, S. (2018). *Cryphonectria parasitica*, the causal agent of chestnut blight: Invasion history, population biology and disease control. *Molecular Plant Pathology*, 19(1), 7–20. <https://doi.org/10.1111/mpp.12542>
- Keçeli, S. A. (2017). Mycoviruses and importance in mycology. *Mikrobiyoloji Bülteni*, 51(4), 404–412. [https://doi.org/10.5578/mb.54128\(Mikovirüslervermikolojidekiönemi.\)](https://doi.org/10.5578/mb.54128(Mikovirüslervermikolojidekiönemi.))
- Liu, H., Fu, Y., Jiang, D., Li, G., Xie, J., Peng, Y., Yi, X., & Ghabrial, S. A. (2009). A novel mycovirus that is related to the human pathogen hepatitis E virus and Rubi-Like viruses. *Journal of Virology*, 83(4), 1981–1991. <https://doi.org/10.1128/JVI.01897-08>
- Van De Sande, W., Lo-Ten-Foe, J., van Belkum, A., Netea, M., Kullberg, B., & Vonk, A. (2010). Mycoviruses: Future therapeutic agents of invasive fungal infections in humans? *European Journal of Clinical Microbiology Infectious Diseases*, 29(7), 755–763.
- Kinsella, C. M., Deijs, M., Gittelbauer, H., van der Hoek, L., & van Dijk, K. (2022). Human clinical isolates of pathogenic fungi are host to diverse mycoviruses. *Microbiology Spectrum*, 10(5), e01610-01622.
- Lin, D. M., Koskella, B., & Lin, H. C. (2017). Phage therapy: An alternative to antibiotics in the age of multi-drug resistance. *World Journal of Gastrointestinal Pharmacology and Therapeutics*, 8(3), 162–173. <https://doi.org/10.4292/wjgpt.v8.i3.162>
- Schmitt, M. J., & Breinig, F. (2002). The viral killer system in yeast: From molecular biology to application. *FEMS Microbiology Reviews*, 26(3), 257–276. <https://doi.org/10.1111/j.1574-6976.2002.tb00614.x>
- Schmitt, M. J., & Breinig, F. (2006). Yeast viral killer toxins: Lethality and self-protection. *Nature Reviews Microbiology*, 4(3), 212–221. <https://doi.org/10.1038/nrmicro1347>
- Applen Clancey, S., Ruchti, F., LeibundGut-Landmann, S., Heitman, J., & Ianiri, G. (2020). A novel mycovirus evokes transcriptional rewiring in the fungus *Malassezia* and stimulates beta interferon production in macrophages. *MBio*, 11(5), e01534–e11520.
- Kotta-Loizou, I. (2021). Mycoviruses and their role in fungal pathogenesis. *Current Opinion in Microbiology*, 63, 10–18.
- Tebbi, C. K., Badiga, A., Sahakian, E., Powers, J. J., Achille, A. N., Patel, S., & Migone, F. (2021). Exposure to a mycovirus containing *Aspergillus flavus* reproduces acute lymphoblastic leukemia cell surface and genetic markers in cells from patients in remission and not controls. *Cancer Treatment and Research Communications*, 26, 100279. <https://doi.org/10.1016/j.ctarc.2020.100279>
- van de Sande, W. W., & Vonk, A. G. (2019). Mycovirus therapy for invasive pulmonary aspergillosis? *Medical Mycology*, 57(2), 179–188.
- Hussain, N., Baqar, Z., Mumtaz, M., El-Sappah, A. H., Show, P. L., Iqbal, H. M., & Bilal, M. (2022). Bioprospecting fungal-derived value-added bioproducts for sustainable pharmaceutical applications. *Sustainable Chemistry and Pharmacy*, 29, 100755. <https://doi.org/10.1016/j.scp.2022.100755>
- Okada, R., Kiyota, E., Moriyama, H., Fukuhara, T., & Natsuaki, T. (2015). A simple and rapid method to purify viral dsRNA from plant and fungal tissue. *Journal of General Plant Pathology*, 81(2), 103–107.
- Nerva, L., Ciuffo, M., Vallino, M., Margaria, P., Varese, G., Gnani, G., & Turina, M. (2016). Multiple approaches for the detection and characterization of viral and plasmid symbionts from a collection of marine fungi. *Virus Research*, 219, 22–38.
- Ahn, I.-P., & Lee, Y.-H. (2001). A viral double-stranded RNA up regulates the fungal virulence of *Nectria radicola*. *Molecular Plant-Microbe Interactions*, 14(4), 496–507.
- Khan, H. A., Sato, Y., Kondo, H., Jamal, A., Bhatti, M. F., & Suzuki, N. (2021). A second capsidless hadakavirus strain with 10 positive-sense single-stranded RNA genomic segments from *Fusarium nygamai*. *Archives of Virology*, 166(10), 2711–2722.
- Khan, H. A., Shamsi, W., Jamal, A., Javaied, M., Sadiq, M., Fatma, T., Ahmed, A., Arshad, M., Waseem, M., & Babar, S. (2021). Assessment of mycoviral diversity in Pakistani fungal isolates revealed infection by 11 novel viruses of a single strain of *Fusarium mangiferae* isolate SP1. *Journal of General Virology*, 102(12), 001690.
- Tran, T. T., Li, H., Nguyen, D. Q., Jones, M. G. K., & Wylie, S. J. (2019). Co-infection with three mycoviruses stimulates growth of a *Monilinia fructicola* isolate on nutrient medium, but does not induce hypervirulence in a natural host. *Viruses*, 11(1), 89. <https://doi.org/10.3390/v11010089>
- Tompa, D. R., Immanuel, A., Srikanth, S., & Kadhivel, S. (2021). Trends and strategies to combat viral infections: A review on FDA approved antiviral drugs. *International Journal of Biological Macromolecules*, 172, 524–541. <https://doi.org/10.1016/j.ijbiomac.2021.01.076>
- Paintsil, E., & Cheng, Y.-C. (2009). Antiviral agents. *Encyclopedia of Microbiology*. <https://doi.org/10.1016/B978-012373944-5.00178-4>
- Bhatti, M. F., Jamal, A., Petrou, M. A., Cairns, T. C., Bignell, E. M., & Coutts, R. H. A. (2011). The effects of dsRNA mycoviruses on growth and murine virulence of *Aspergillus fumigatus*. *Fungal Genetics and Biology*, 48(11), 1071–1075. <https://doi.org/10.1016/j.fgb.2011.07.008>
- Schneider-Poetsch, T., Ju, J., Eyler, D. E., Dang, Y., Bhat, S., Merrick, W. C., Green, R., Shen, B., & Liu, J. O. (2010). Inhibition of eukaryotic translation elongation by cycloheximide and lactimidomycin. *Nature Chemical Biology*, 6(3), 209–217. <https://doi.org/10.1038/nchembio.304>
- Young, C. W., Robinson, P. F., & Sacktor, B. (1963). Inhibition of the synthesis of protein in intact animals by

- acetoxycycloheximide and a metabolic derangement concomitant with this blockade. *Biochemical Pharmacology*, 12(8), 855–865.
31. Barton, D. J., Morasco, B. J., & Flanagan, J. B. (1999). Translating ribosomes inhibit poliovirus negative-strand RNA synthesis. *Journal of virology*, 73(12), 10104–10112.
 32. Kim, J.-M., Song, H.-Y., Choi, H.-J., Yun, S.-H., So, K.-K., Ko, H.-K., & Kim, D.-H. (2015). Changes in the mycovirus (LeV) titer and viral effect on the vegetative growth of the edible mushroom *Lentinula edodes*. *Virus Research*, 197, 8–12. <https://doi.org/10.1016/j.virusres.2014.11.016>
 33. Santos, V., Mascarin, G. M., da Silva Lopes, M., Alves, M. C. D. F., Rezende, J. M., Gatti, M. S. V., Dunlap, C. A., & Delalibera Júnior, Í. (2017). Identification of double-stranded RNA viruses in Brazilian strains of *Metarhizium anisopliae* and their effects on fungal biology and virulence. *Plant Gene*, 11, 49–58. <https://doi.org/10.1016/j.plgene.2017.01.001>
 34. van Diepeningen, A. D., Debets, A. J. M., & Hoekstra, R. F. (2006). Dynamics of dsRNA mycoviruses in black *Aspergillus* populations. *Fungal Genetics and Biology*, 43(6), 446–452. <https://doi.org/10.1016/j.fgb.2006.01.014>
 35. Lima, S. S., Abadio, A. K., Araújo, E. F., Kitajima, E. W., Sartorato, A., & de Queiroz, M. V. (2010). Mycovirus in *Pseudocercospora griseola*, the causal agent of angular leaf spot in common bean. *Canadian Journal of Microbiology*, 56(5), 359–365. <https://doi.org/10.1139/w10-022>
 36. Thapa, V., Turner, G. G., Hafenstein, S., Overton, B. E., Vanderwolf, K. J., & Roossinck, M. J. (2016). Using a novel partitivirus in *Pseudogymnoascus destructans* to understand the epidemiology of white-nose syndrome. *PLOS Pathogens*, 12(12), e1006076. <https://doi.org/10.1371/journal.ppat.1006076>
 37. Zhang, N., Yin, Y., Xu, S.-J., & Chen, W.-S. (2008). 5-Fluorouracil: Mechanisms of resistance and reversal strategies. *Molecules (Basel, Switzerland)*, 13(8), 1551–1569. <https://doi.org/10.3390/molecules13081551>
 38. Sethy, C., & Kundu, C. N. (2021). 5-Fluorouracil (5-FU) resistance and the new strategy to enhance the sensitivity against cancer: Implication of DNA repair inhibition. *Biomedicine & Pharmacotherapy*, 137, 111285. <https://doi.org/10.1016/j.biopha.2021.111285>
 39. Agudo, R., Arias, A., & Domingo, E. (2009). 5-fluorouracil in lethal mutagenesis of foot-and-mouth disease virus. *Future Medicinal Chemistry*, 1(3), 529–539. <https://doi.org/10.4155/fmc.09.26>
 40. Ahmad, S. I. (2020). 5-Fluorouracil in combination with deoxyribonucleosides and deoxyribose as possible therapeutic options for the Coronavirus, COVID-19 infection. *Medical Hypotheses*, 142, 109754–109754. <https://doi.org/10.1016/j.mehy.2020.109754>
 41. Schmidt, F. R., Lemke, P. A., & Esser, K. (1986). Viral influences on aflatoxin formation by *Aspergillus flavus*. *Applied Microbiology Biotechnology*, 24(3), 248–252.
 42. Thomas, E., Ghany, M. G., & Liang, T. J. (2012). The application and mechanism of action of ribavirin in therapy of hepatitis C. *Antiviral Chemistry & Chemotherapy*, 23(1), 1–12. <https://doi.org/10.3851/IMP2125>
 43. Tate, P. M., Mastrodomenico, V., & Mounce, B. C. (2019). Ribavirin induces polyamine depletion via nucleotide depletion to limit virus replication. *Cell Reports*, 28(10), 2620–2633. <https://doi.org/10.1016/j.celrep.2019.07.099>
 44. Herrero, N., & Zabalgoitia, I. (2011). Mycoviruses infecting the endophytic and entomopathogenic fungus *Tolypocladium cylindrosporum*. *Virus Research*, 160(1–2), 409–413. <https://doi.org/10.1016/j.virusres.2011.06.015>
 45. Niu, Y., Yuan, Y., Mao, J., Yang, Z., Cao, Q., Zhang, T., Wang, S., & Liu, D. (2018). Characterization of two novel mycoviruses from *Penicillium digitatum* and the related fungicide resistance analysis. *Scientific Reports*, 8(1), 5513. <https://doi.org/10.1038/s41598-018-23807-3>
 46. Sun, Y., Guo, M., Wang, J., Bian, Y., & Xu, Z. (2022). Curing two predominant viruses occurring in *Lentinula edodes* by chemotherapy and mycelial fragmentation methods. *Journal of Virological Methods*, 300, 114370. <https://doi.org/10.1016/j.jviromet.2021.114370>
 47. Ikeda, A., Chiba, Y., Kuroki, M., Urayama, S.-I., & Hagiwara, D. (2022). Efficient elimination of RNA mycoviruses in aspergillus species using RdRp-inhibitors ribavirin and 2'-C-methylribonucleoside derivatives. *Frontiers in Microbiology*, 13, 1–9. <https://doi.org/10.3389/fmicb.2022.1024933>
 48. Overington, J. P., Al-Lazikani, B., & Hopkins, A. L. (2006). How many drug targets are there? *Nature Reviews Drug Discovery*, 5(12), 993–996.
 49. Neang, S., Bincader, S., Rangsuwan, S., Keawmanee, P., Rin, S., Salaipeth, L., Das, S., Kondo, H., Suzuki, N., Sato, I., Takemoto, D., Rattanakreetakul, C., Pongpisutta, R., Arakawa, M., & Chiba, S. (2021). Omnipresence of partitiviruses in Rice aggregate sheath spot symptom-associated fungal isolates from Paddies in Thailand. *Viruses*, 13(11), 2269.
 50. Feklistov, A., Mekler, V., Jiang, Q., Westblade, L. F., Irschik, H., Jansen, R., Mustaev, A., Darst, S. A., & Ebright, R. H. (2008). Rifamycins do not function by allosteric modulation of binding of Mg²⁺ to the RNA polymerase active center. *Proceedings of the National Academy of Sciences USA*, 105(39), 14820–14825. <https://doi.org/10.1073/pnas.0802822105>
 51. Hartmann, G., Honikel, K. O., Knüsel, F., & Nüesch, J. (1967). The specific inhibition of the DNA-directed RNA synthesis by rifamycin. *Biochimica et Biophysica Acta*, 145(3), 843–844.
 52. Kwon, B. S., Kim, M.-N., Sung, H., Koh, Y., Kim, W.-S., Song, J.-W., Oh, Y.-M., Lee, S.-D., Lee, S. W., & Lee, J.-S. (2018). In vitro MIC values of rifampin and ethambutol and treatment outcome in *Mycobacterium avium* complex lung disease. *Antimicrobial Agents Chemotherapy*, 62(10), e00491–e1418.
 53. Cao, C., Li, H., Jones, M. G. K., & Wylie, S. J. (2019). Challenges to elucidating how endornaviruses influence fungal hosts: Creating mycovirus-free isogenic fungal lines and testing them. *Journal of Virological Methods*, 274, 113745. <https://doi.org/10.1016/j.jviromet.2019.113745>
 54. Cao, C. T. H. (2021). *Together forever: Molecular biology of a mycovirus-fungus symbiosis*. Murdoch University.
 55. Jacquat, A. G., Theumer, M. G., Cañizares, M. C., Debat, H. J., Iglesias, J., García Pedrajas, M. D., & Dambolena, J. S. (2020). A survey of mycoviral infection in *Fusarium* spp. isolated from maize and *Sorghum* in Argentina Identifies the first mycovirus from *Fusarium verticillioides*. *Viruses*, 12(10), 1161. <https://doi.org/10.3390/v12101161>
 56. Hansen, H. (1926). A simple method of obtaining single-spore cultures. *Science*, 64(1659), 384–384.
 57. Noman, E., Al-Gheethi, A. A., Rahman, N. K., Talip, B., Mohamed, R., & Kadir, O. A. (2018). Single spore isolation as a simple and efficient technique to obtain fungal pure culture. *IOP Conference Series: Earth and Environmental Science*, 140, 012055. <https://doi.org/10.1088/1755-1315/140/1/012055>
 58. Zhang, K., Yuan-Ying, S., & Cai, L. (2013). An optimized protocol of single spore isolation for fungi. *Cryptogamie, Mycologie*, 34(4), 349–356.
 59. Sato, Y., Shamsi, W., Jamal, A., Bhatti, M. F., Kondo, H., Suzuki, N., & Wickner, R. B. (2020). Hadaka virus 1: A Capsidless Eleven-segmented positive-sense single-stranded RNA virus from a phytopathogenic Fungus *Fusarium oxysporum*. *MBio*, 11(3), e00450–e1420. <https://doi.org/10.1128/mBio.00450-20>
 60. Yaegashi, H., Sawahata, T., Ito, T., & Kanematsu, S. (2011). A novel colony-print immunoassay reveals differential patterns of

- distribution and horizontal transmission of four unrelated mycoviruses in *Rosellinia necatrix*. *Virology*, 409(2), 280–289. <https://doi.org/10.1016/j.virol.2010.10.014>
61. Urayama, S., Kato, S., Suzuki, Y., Aoki, N., Le, M. T., Arie, T., Teraoka, T., Fukuhara, T., & Moriyama, H. (2010). Mycoviruses related to chrysovirus affect vegetative growth in the rice blast fungus *Magnaporthe oryzae*. *Journal of General Virology*, 91(12), 3085–3094.
 62. Takahashi-Nakaguchi, A., Shishido, E., Yahara, M., Urayama, S.-I., Sakai, K., Chibana, H., Kamei, K., Moriyama, H., & Gonoi, T. (2020). Analysis of an intrinsic mycovirus associated with reduced virulence of the human pathogenic fungus *Aspergillus fumigatus*. *Frontiers in Microbiology*, 10, 3045–3045. <https://doi.org/10.3389/fmicb.2019.03045>
 63. Chun, J., Yang, H.-E., & Kim, D.-H. (2018). Identification of a novel partitivirus of *Trichoderma harzianum* NCF319 and evidence for the related antifungal activity. *Frontiers in Plant Science*, 9, 1–10. <https://doi.org/10.3389/fpls.2018.01699>
 64. Zhang, C.-H., Liu, Y.-M., Qi, Y.-C., Gao, Y.-Q., Shen, J.-W., & Qiu, L.-Y. (2010). Comparisons of different methods for virus-elimination of edible fungi. *Chinese Journal of Virology*, 26(3), 249–254.
 65. Petrzik, K., & Siddique, A. B. (2019). A mycoparasitic and opportunistic fungus is inhabited by a mycovirus. *Archives of Virology*, 164(10), 2545–2549. <https://doi.org/10.1007/s00705-019-04359-x>
 66. Choi, Y.-W., Hyde, K. D., & Ho, W. (1999). Single spore isolation of fungi. *Fungal Diversity*, 10(43), 29–38.
 67. Ho, W.-C., & Ko, W.-H. (1997). A simple method for obtaining single-spore isolates of fungi. *Botanical Bulletin of Academia Sinica*, 38, 41–44.
 68. Shamsi, W., Sato, Y., Jamal, A., Shahi, S., Kondo, H., Suzuki, N., & Bhatti, M. F. (2019). Molecular and biological characterization of a novel botybirnavirus identified from a Pakistani isolate of *Alternaria alternata*. *Virus Research*, 263, 119–128. <https://doi.org/10.1016/j.virusres.2019.01.006>
 69. Singh, K. P., Kumar, D., & Bandyopadhyay, P. (2004). A new technique for single spore isolation of two predacious fungi forming constricting ring. *Mycobiology*, 32(4), 197–198.
 70. Khan, H. A., Sato, Y., Kondo, H., Jamal, A., Bhatti, M. F., & Suzuki, N. (2022). A novel victorivirus from the phytopathogenic fungus *Neofusicoccum parvum*. *Archives of Virology*, 167(3), 923–929. <https://doi.org/10.1007/s00705-021-05304-7>
 71. Khan, H. A., Telengech, P., Kondo, H., Bhatti, M. F., & Suzuki, N. (2022). Mycovirus hunting revealed the presence of diverse viruses in a single isolate of the phytopathogenic Fungus *Diplodia seriata* From Pakistan. *Frontiers in Cellular and Infection Microbiology*. <https://doi.org/10.3389/fcimb.2022.913619>
 72. Duncan, K. E., & Howard, R. J. (2000). Cytological analysis of wheat infection by the leaf blotch pathogen *Mycosphaerella graminicola*. *Mycological Research*, 104(9), 1074–1082.
 73. Kema, G., Annone, J., Sayoud, R., & Van Silfhout, C. (1996). Genetic variation for virulence and resistance in the wheat-*Mycosphaerella graminicola* pathosystem I. Interactions between pathogen isolates and host cultivars. *Genetics*, 16, 1–13.
 74. Eyal, Z., & van Ginkel, M. (1987). The Septoria Diseases of Wheat. Concepts and Methods of Disease Management. CIM-MYT. <http://hdl.handle.net/10883/1113>
 75. Palmer, C. L., & Skinner, W. (2002). *Mycosphaerella graminicola*: Latent infection, crop devastation and genomics. *Molecular Plant Pathology*, 3(2), 63–70.
 76. Zhang, L., Fu, Y., Xie, J., Jiang, D., Li, G., & Yi, X. (2009). A novel virus that infecting hypovirulent strain XG36-1 of plant fungal pathogen *Sclerotinia sclerotiorum*. *Virology Journal*, 6(1), 96. <https://doi.org/10.1186/1743-422X-6-96>
 77. Coenen, A., Kevei, F., & Hoekstra, R. F. (1997). Factors affecting the spread of double-stranded RNA viruses in *Aspergillus nidulans*. *Genetics Research*, 69(1), 1–10.
 78. Varga, J., Frisvad, J. C., & Samson, R. A. (2011). Two new aflatoxin producing species, and an overview of *Aspergillus* section Flavi. *Studies in Mycology*, 69, 57–80. <https://doi.org/10.3114/sim.2011.69.05>
 79. Brasier, C. (1983). A cytoplasmically transmitted disease of *Ceratocystis ulmi*. *Nature*, 305(5931), 220–223. <https://doi.org/10.1038/305220a0>
 80. McFadden, J., Buck, K., & Rawlinson, C. (1983). Infrequent transmission of double-stranded RNA virus particles but absence of DNA proviruses in single ascospore cultures of *Gaeumannomyces graminis*. *Journal of General Virology*, 64(4), 927–937.
 81. Ikeda, K.-I., Nakamura, H., Arakawa, M., & Matsumoto, N. (2004). Diversity and vertical transmission of double-stranded RNA elements in root rot pathogens of trees, *Helicobasidium mompa* and *Rosellinia necatrix*. *Mycological research*, 108(6), 626–634.
 82. Koltin, Y., & Day, P. (1976). Inheritance of killer phenotypes and double-stranded RNA in *Ustilago maydis*. *Proceedings of the National Academy of Sciences USA*, 73(2), 594–598.
 83. Ihrmark, K., Stenström, E., & Stenlid, J. (2004). Double-stranded RNA transmission through basidiospores of *Heterobasidium annosum*. *Mycological Research*, 108(2), 149–153.
 84. Brewer, B. J., & Fangman, W. L. (1980). Preferential inclusion of extrachromosomal genetic elements in yeast meiotic spores. *Proceedings of the National Academy of Sciences USA*, 77(9), 5380–5384. <https://doi.org/10.1073/pnas.77.9.5380>
 85. Chu, Y.-M., Lim, W.-S., Yea, S.-J., Cho, J.-D., Lee, Y.-W., & Kim, K.-H. (2004). Complexity of dsRNA mycovirus isolated from *Fusarium graminearum*. *Virus Genes*, 28(1), 135–143.
 86. Alam, I., Sharmin, S. A., Naher, M. K., Alam, M. J., Anisuzzaman, M., & Alam, M. F. (2013). Elimination and detection of viruses in meristem-derived plantlets of sweetpotato as a low-cost option toward commercialization. *3 Biotech*, 3(2), 153–164. <https://doi.org/10.1007/s13205-012-0080-6>
 87. Vivek, M., & Modgil, M. (2018). Elimination of viruses through thermotherapy and meristem culture in apple cultivar ‘Oregon Spur-II.’ *Virus Disease*, 29(1), 75–82. <https://doi.org/10.1007/s13337-018-0437-5>
 88. Xiang, X. (2018). Nuclear movement in fungi. *Seminars in cell & developmental biology*. <https://doi.org/10.1016/j.semcdb.2017.10.024>
 89. Kanematsu, S., Arakawa, M., Oikawa, Y., Onoue, M., Osaki, H., Nakamura, H., Ikeda, K., Kuga-Uetake, Y., Nitta, H., Sasaki, A., Suzuki, K., Yoshida, K., & Matsumoto, N. (2004). A Reovirus Causes Hypovirulence of *Rosellinia necatrix*. *Phytopathology*, 94(6), 561–568. <https://doi.org/10.1094/PHYTO.2004.94.6.561>
 90. Aoki, N., Moriyama, H., Kodama, M., Arie, T., Teraoka, T., & Fukuhara, T. (2009). A novel mycovirus associated with four double-stranded RNAs affects host fungal growth in *Alternaria alternata*. *Virus Research*, 140(1–2), 179–187.
 91. Liu, R., Cheng, J., Fu, Y., Jiang, D., & Xie, J. (2015). Molecular characterization of a novel positive-sense, single-stranded RNA mycovirus infecting the plant pathogenic Fungus *Sclerotinia sclerotiorum*. *Viruses*, 7(5), 2470–2484. <https://doi.org/10.3390/v7052470>
 92. Zheng, L., Zhang, M., Chen, Q., Zhu, M., & Zhou, E. (2014). A novel mycovirus closely related to viruses in the genus Alphapartitivirus confers hypovirulence in the phytopathogenic fungus *Rhizoctonia solani*. *Virology*, 456, 220–226.
 93. Chiba, S., Lin, Y.-H., Kondo, H., Kanematsu, S., & Suzuki, N. (2013). A novel victorivirus from a phytopathogenic fungus, *Rosellinia necatrix*, is infectious as particles and targeted by

- RNA silencing. *Journal of Virology*, 87(12), 6727–6738. <https://doi.org/10.1128/JVI.00557-13>
94. Kashif, M., Jurvansuu, J., Vainio, E. J., & Hantula, J. (2019). Alphapartitiviruses of *Heterobasidion* Wood decay fungi affect each other's transmission and host growth. *Frontiers in Cellular and Infection Microbiology*, 9(26), 64. <https://doi.org/10.3389/fcimb.2019.00064>
 95. Komatsu, A., Kondo, H., Sato, M., Kurahashi, A., Nishibori, K., Suzuki, N., & Fujimori, F. (2019). Isolation and characterization of a novel mycovirus infecting an edible mushroom *Grifola frondosa*. *Mycoscience*, 60(4), 211–220.
 96. Lin, Y.-H., Chiba, S., Tani, A., Kondo, H., Sasaki, A., Kanematsu, S., & Suzuki, N. (2012). A novel quadripartite dsRNA virus isolated from a phytopathogenic filamentous fungus *Rosellinia necatrix*. *Virology*, 426(1), 42–50. <https://doi.org/10.1016/j.virol.2012.01.013>
 97. Zhong, J., Chen, D., Zhu, H. J., Gao, B. D., & Zhou, Q. (2016). Hypovirulence of *Sclerotium rolfsii* caused by associated RNA mycovirus. *Frontiers in Microbiology*, 7, 1798.
 98. Garcia-Rubio, R., de Oliveira, H. C., Rivera, J., & Trevijano-Contador, N. (2020). The fungal cell wall: *Candida*, *Cryptococcus*, and *Aspergillus* species. *Frontiers in Microbiology*, 10(9), 2993. <https://doi.org/10.3389/fmicb.2019.02993>
 99. Cheng, Y., & Bélanger, R. R. (2000). Protoplast preparation and regeneration from spores of the biocontrol fungus *Pseudozyma flocculosa*. *FEMS Microbiology Letters*, 190(2), 287–291. <https://doi.org/10.1111/j.1574-6968.2000.tb09300.x>
 100. Necas, O. (1971). Cell wall synthesis in yeast protoplasts. *J Bacteriological reviews*, 35(2), 149–170.
 101. Wu, J.-D., & Chou, J.-C. (2019). Optimization of protoplast preparation and regeneration of a medicinal fungus *Antrodia cinnamomea*. *Mycobiology*, 47(4), 483–493. <https://doi.org/10.1080/12298093.2019.1687252>
 102. Roth, M. G., & Chilvers, M. I. (2019). A protoplast generation and transformation method for soybean sudden death syndrome causal agents *Fusarium virguliforme* and *F. brasiliense*. *Fungal Biology and Biotechnology*, 6(1), 7. <https://doi.org/10.1186/s40694-019-0070-0>
 103. Lee, K.-M., Yu, J., Son, M., Lee, Y.-W., & Kim, K.-H. (2011). Transmission of *Fusarium boothii* mycovirus via protoplast fusion causes hypovirulence in other phytopathogenic fungi. *PLoS ONE*, 6(6), e21629. <https://doi.org/10.1371/journal.pone.0021629>
 104. Kanematsu, S., Sasaki, A., Onoue, M., Oikawa, Y., & Ito, T. (2010). Extending the fungal host range of a partitivirus and a mycoreovirus from *Rosellinia necatrix* by inoculation of protoplasts with virus particles. *Phytopathology*, 100(9), 922–930. <https://doi.org/10.1094/PHYTO-100-9-0922>
 105. Wang, L., He, H., Wang, S., Chen, X., Qiu, D., Kondo, H., & Guo, L. (2018). Evidence for a novel negative-stranded RNA mycovirus isolated from the plant pathogenic fungus *Fusarium graminearum*. *Virology*, 518, 232–240. <https://doi.org/10.1016/j.virol.2018.03.008>
 106. Wang, M.-R., Cui, Z.-H., Li, J.-W., Hao, X.-Y., Zhao, L., & Wang, Q.-C. (2018). In vitro thermotherapy-based methods for plant virus eradication. *Plant Methods*, 14, 87–87. <https://doi.org/10.1186/s13007-018-0355-y>
 107. Chen, B., & Nuss, D. L. (1999). Infectious cDNA clone of hypovirus CHV1-Euro7: A comparative virology approach to investigate virus-mediated hypovirulence of the chestnut blight fungus *Cryphonectria parasitica*. *Journal of Virology*, 73(2), 985–992.
 108. Choi, G. H., & Nuss, D. L. (1992). Hypovirulence of chestnut blight fungus conferred by an infectious viral cDNA. *Science*, 257(5071), 800–803.
 109. Córdoba, L., Ruiz-Padilla, A., Rodríguez-Romero, J., & Ayllón, M. A. (2022). Construction and characterization of a *Botrytis* virus F infectious clone. *Journal of Fungi*, 8(5), 459.
 110. Yang, S., Dai, R., Salaipeth, L., Huang, L., Liu, J., Andika, I. B., & Sun, L. (2021). Infection of two heterologous mycoviruses reduces the virulence of *Valsa mali*, a fungal agent of apple Valsa canker disease. *Frontiers in Microbiology*, 25(12), 1092.
 111. Savage, G. M., & Brook, M. J. V. (1946). The fragmentation of the mycelium of *Penicillium notatum* and *Penicillium chrysogenum* by a high-speed blender and the evaluation of blended seed. *Journal of Bacteriology*, 52(3), 385–391. <https://doi.org/10.1128/jb.52.3.385-391.1946>
 112. Donzelli, B. G. G., & Churchill, A. C. (2007). A quantitative assay using mycelial fragments to assess virulence of *Mycosphaerella fijiensis*. *Phytopathology*, 97(8), 916–929.
 113. Song, H.-Y., Choi, H.-J., Jeong, H., Choi, D., Kim, D.-H., & Kim, J.-M. (2016). Viral effects of a dsRNA mycovirus (PoV-ASI2792) on the Vegetative growth of the edible mushroom *Pleurotus ostreatus*. *Mycobiology*, 44(4), 283–290. <https://doi.org/10.5941/MYCO.2016.44.4.283>
 114. Won, H.-K., Park, S.-J., Kim, D.-K., Shin, M. J., Kim, N.-R., Lee, S. H., Kwon, Y. C., Ko, H. K., Ro, H.-S., & Lee, H.-S. (2013). Isolation and characterization of a mycovirus in *Lentinula edodes*. *Journal of Microbiology*, 51, 118–122.
 115. Ali, L., Khalid, M., Asghar, H. N., & Asgher, M. (2017). Scrutinizing of rhizobacterial isolates for improving drought resilience in maize (*Zea mays*). *International Journal of Agriculture and Biology*, 19, 1054–1064.
 116. Bandeppa, P. S., & Kandpal, B. (2015). Evaluation of osmo-tolerant rhizobacteria for alleviation of water deficit stress in mustard. *Green Farming*, 6, 590.
 117. Karvembu, P., Gomathi, V., Anandham, R., & Mary, J. K. (2021). Isolation, screening and identification of moisture stress tolerant *Rhizobacteria* from xerophyte *Prosopis juliflora* (SW). *Journal of Pharmacognosy Phytochemistry*, 9, 605–609.
 118. Vurukonda, S. S. K. P., Vardharajula, S., Shrivastava, M., & SkZ, A. (2016). Multifunctional *Pseudomonas putida* strain FBKV2 from arid rhizosphere soil and its growth promotional effects on maize under drought stress. *Rhizosphere*, 1, 4–13.
 119. Rukundo, P., Carpentier, S., & Swennen, R. (2012). Development of in vitro technique to screen for drought tolerant banana varieties by sorbitol induced osmotic stress. *African Journal of Plant Science*, 6(15), 416–425.
 120. Bejaoui, M., Pantazi, E., Folch-Puy, E., Panisello, A., Calvo, M., Pasut, G., Rimola, A., Navasa, M., Adam, R., & Roselló-Catafau, J. (2015). Protective effect of intravenous high molecular weight polyethylene glycol on fatty liver preservation. *BioMed Research International*, 15, 1–10. <https://doi.org/10.1155/2015/794287>
 121. Lazar, H. L. (2015). High-molecular-weight polyethylene glycol: A new strategy to limit ischemia–reperfusion injury. *The Journal of Thoracic Cardiovascular Surgery*, 149(2), 594–595.
 122. Malhotra, M., Lane, C., Tomaro-Duchesneau, C., Saha, S., & Prakash, S. (2011). A novel method for synthesizing PEGylated chitosan nanoparticles: Strategy, preparation, and in vitro analysis. *International Journal of Nanomedicine*, 6, 485.
 123. Giraud, S., Codas, R., Hauet, T., Eugene, M., & Badet, L. (2014). Polyethylene glycols and organ protection against I/R injury. *Progress in Urology*, 24, 37–43.
 124. Kylvik, D. L., Sutton, T. C., Le, Y., & Scott, M. D. (2011). Polymer-mediated broad spectrum antiviral prophylaxis: Utility in high risk environments. *Progress in Molecular Environmental Bioengineering*, 1, 167–190.
 125. Poimala, A., Raco, M., Haikonen, T., Černý, M., Parikka, P., Hantula, J., & Vainio, E. J. (2022). Bunyaviruses affect growth,

- sporulation, and elicitor production in *Phytophthora cactorum*. *Viruses*, 14(12), 2596.
126. Wang, Q., Cuellar, W. J., Rajamäki, M.-L., Hirata, Y., & Valkonen, J. P. T. (2008). Combined thermotherapy and cryotherapy for efficient virus eradication: Relation of virus distribution, subcellular changes, cell survival and viral RNA degradation in shoot tips. *Molecular Plant Pathology*, 9(2), 237–250. <https://doi.org/10.1111/j.1364-3703.2007.00456.x>
 127. Owens, R. A. (2007). Potato spindle tuber viroid: The simplicity paradox resolved? *Molecular Plant Pathology*, 8(5), 549–560. <https://doi.org/10.1111/j.1364-3703.2007.00418.x>
 128. Zhang, T., Li, N., Yuan, Y., Cao, Q., Chen, Y., Tan, B., Li, G., & Liu, D. (2019). Blue-white colony selection of virus-infected isogenic recipients based on a *Chrysovirus* isolated from *Penicillium italicum*. *Virologica Sinica*, 34, 688–700. <https://doi.org/10.1007/s12250-019-00150-z>
 129. Zhang, Z., Wang, Q.-C., Spetz, C., & Blystad, D.-R. (2019). In vitro therapies for virus elimination of potato-valuable germplasm in Norway. *Scientia Horticulturae*, 249, 7–14. <https://doi.org/10.1016/j.scienta.2019.01.027>
 130. Parsell, D., & Lindquist, S. (1993). The function of heat-shock proteins in stress tolerance: Degradation and reactivation of damaged proteins. *Annual Review of Genetics*, 27, 437–497.
 131. da Silva Xavier, A., de Barros, A. P. O., Godinho, M. T., Zerbini, F. M., de Oliveira Souza, F., Bruckner, F. P., & Alfenas-Zerbini, P. (2018). A novel mycovirus associated to *Alternaria alternata* comprises a distinct lineage in Partitiviridae. *Virus Research*, 244, 21–26. <https://doi.org/10.1016/j.virusres.2017.10.007>
 132. Xie, J., & Ghabrial, S. A. (2012). Molecular characterizations of two mitoviruses co-infecting a hypovirulent isolate of the plant pathogenic fungus *Sclerotinia sclerotiorum*. *Virology*, 428(2), 77–85. <https://doi.org/10.1016/j.virol.2012.03.015>
 133. Welch, D., Buonanno, M., Grilj, V., Shuryak, I., Crickmore, C., Bigelow, A. W., Randers-Pehrson, G., Johnson, G. W., & Brenner, D. J. (2018). Far-UVC light: A new tool to control the spread of airborne-mediated microbial diseases. *Science and Reports*, 8(1), 2752. <https://doi.org/10.1038/s41598-018-21058-w>
 134. Reed, N. G. (2010). The history of ultraviolet germicidal irradiation for air disinfection. *Public Health Reports*, 125(1), 15–27. <https://doi.org/10.1177/003335491012500105>
 135. Yin, R., Dai, T., Avci, P., Jorge, A. E. S., de Melo, W. C. M. A., Vecchio, D., Huang, Y.-Y., Gupta, A., & Hamblin, M. R. (2013). Light based anti-infectives: Ultraviolet C irradiation, photodynamic therapy, blue light, and beyond. *Current Opinion in Pharmacology*, 13(5), 731–762. <https://doi.org/10.1016/j.coph.2013.08.009>
 136. Beck, S. E., Rodriguez, R. A., Hawkins, M. A., Hargy, T. M., Larson, T. C., & Linden, K. G. (2015). Comparison of UV-induced inactivation and RNA damage in MS2 Phage across the germicidal UV spectrum. *Applied and Environmental Microbiology*, 82(5), 1468–1474. <https://doi.org/10.1128/AEM.02773-15>
 137. Myers, E., Kheradmand, S., & Miller, R. (2021). An update on narrowband ultraviolet B therapy for the Treatment of skin diseases. *Cureus*, 13(11), e19182–e19182. <https://doi.org/10.7759/cureus.19182>
 138. Castillo, A., & Cifuentes, V. (1994). Presence of double-stranded RNA and virus-like particles in *Phaffia rhodozyma*. *Current genetics*, 26(4), 364–368.
 139. Lemus-Minor, C. G., Cañizares, M. C., García-Pedrajas, M. D., & Pérez-Artés, E. (2019). Horizontal and vertical transmission of the hypovirulence-associated mycovirus *Fusarium oxysporum* f. sp. dianthi virus 1. *European Journal of Plant Pathology*, 153(2), 645–650. <https://doi.org/10.1007/s10658-018-1554-0>
 140. Zhong, L., Shu, C., Zhang, M., Yang, M., & Zhou, E. (2019). Molecular characterization of a novel endornavirus conferring hypovirulence in rice sheath blight Fungus *Rhizoctonia solani* AG-1 IA strain GD-2. *Viruses*. <https://doi.org/10.3390/v11020178>
 141. Glass, N. L., & Dementhon, K. (2006). Non-self recognition and programmed cell death in filamentous fungi. *Current Opinion in Microbiology*, 9(6), 553–558. <https://doi.org/10.1016/j.mib.2006.09.001>
 142. Caten, C. E. (1972). Vegetative incompatibility and cytoplasmic infection in fungi. *Journal of General Microbiology*, 72(2), 221–229. <https://doi.org/10.1099/00221287-72-2-221>
 143. Saupe, S. J. (2000). Molecular genetics of heterokaryon incompatibility in filamentous ascomycetes. *Microbiology and Molecular Biology Reviews*, 64(3), 489–502.
 144. Glass, N. L., & Kaneko, I. (2003). Fatal attraction: Nonself recognition and heterokaryon incompatibility in Filamentous fungi. *Eukaryotic Cell*, 2(1), 1. <https://doi.org/10.1128/EC.2.1.1-8.2003>
 145. Milgroom, M. G., & Cortesi, P. (2004). Biological control of chestnut blight with hypovirulence: A critical analysis. *Annual Review of Phytopathology*, 42, 311.
 146. Cornejo, C., Hisano, S., Bragança, H., Suzuki, N., & Rigling, D. (2021). A new double-stranded RNA mycovirus in *Cryphonectria naterciae* is able to cross the species barrier and is deleterious to a new host. *Journal of Fungi*, 7(10), 861.
 147. Ikeda, K., Inoue, K., Kida, C., Uwamori, T., Sasaki, A., Kanematsu, S., & Park, P. (2013). Potentiation of mycovirus transmission by zinc compounds via attenuation of heterogenic incompatibility in *Rosellinia necatrix*. *Applied Environmental Microbiology*, 79(12), 3684–3691.
 148. Lambert, J. C., Zhou, Z., & Kang, Y. J. (2003). Suppression of Fas-mediated signaling pathway is involved in zinc inhibition of ethanol-induced liver apoptosis. *Experimental Biology Medicine*, 228(4), 406–412.
 149. Wu, S., Cheng, J., Fu, Y., Chen, T., Jiang, D., Ghabrial, S. A., & Xie, J. (2017). Virus-mediated suppression of host non-self recognition facilitates horizontal transmission of heterologous viruses. *PLOS Pathogens*, 13(3), e1006234. <https://doi.org/10.1371/journal.ppat.1006234>
 150. Carbone, I., Liu, Y.-C., Hillman, B. I., & Milgroom, M. G. (2004). Recombination and migration of *Cryphonectria hypovirus* 1 as inferred from gene genealogies and the coalescent. *Genetics*, 166(4), 1611–1629.
 151. Raimondo, M. L., & Carlucci, A. (2018). Characterization and pathogenicity assessment of *Plectosphaerella* species associated with stunting disease on tomato and pepper crops in Italy. *Plant Pathology*, 67(3), 626–641. <https://doi.org/10.1111/ppa.12766>
 152. Zhou, W., Shi, W., Xu, X.-W., Li, Z.-G., Yin, C.-F., Peng, J.-B., Pan, S., Chen, X.-L., Zhao, W.-S., Zhang, Y., Yang, J., & Peng, Y.-L. (2018). Glutamate synthase MoGlt1-mediated glutamate homeostasis is important for autophagy, virulence and conidiation in the rice blast fungus. *Molecular plant pathology*, 19(3), 564–578. <https://doi.org/10.1111/mpp.12541>
 153. Hyder, R., Pennanen, T., Hamberg, L., Vainio, E. J., Piri, T., & Hantula, J. (2013). Two viruses of *Heterobasidion* confer beneficial, cryptic or detrimental effects to their hosts in different situations. *Fungal Ecology*, 6(5), 387–396.
 154. Robinson, H. L., & Deacon, J. W. (2002). Double-stranded RNA elements in *Rhizoctonia solani* AG 3. *Mycological Research*, 106(1), 12–22.
 155. Koltin, Y., Finkler, A., & Ben-Zvi, B. (1987). Double-stranded RNA viruses of pathogenic fungi: virulence and plant protection. Symposium series-British Mycological Society,
 156. Maffettone, E. (2007). *Characterization of a novel virus associated with the MVX disease of Agaricus bisporus* [Doctoral dissertation, Cranfield University, Cranfield University].
 157. Tavantzis, S., & Bandy, B. (1988). Properties of a mycovirus from *Rhizoctonia solani* and its virion-associated RNA polymerase. *Journal of General Virology*, 69(7), 1465–1477.

158. Khalifa, M. E., & MacDiarmid, R. M. (2021). A mechanically transmitted DNA mycovirus is targeted by the defence machinery of its host *Botrytis cinerea*. *Viruses*, 13(7), 1315.
159. Bao, X., & Roossinck, M. J. (2013). Multiplexed interactions: Viruses of endophytic fungi. *Advances in Virus Research*, 86, 37–58.
160. Tavakoli, A., Ataei-Pirkooh, A., Mm Sadeghi, G., Bokharaei-Salim, F., Sahrapour, P., Kiani, S. J., Moghoofei, M., Farahmand, M., Javanmard, D., & Monavari, S. H. (2018). Polyethylene glycol-coated zinc oxide nanoparticle: An efficient nanoweapon to fight against herpes simplex virus type 1. *Nanomedicine*, 13(21), 2675–2690.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.