

Transcriptional Activity of an Estrogen Receptor β Subtype in the Medaka *Oryzias dancena*

Sejung Maeng¹, Sung Woo Yoon¹, Eun Jeong Kim², Yoon Kwon Nam², and [†]Young Chang Sohn¹

¹Dept. of Marine Molecular Biosciences, Gangneung-Wonju National University, Gangneung 25457, Korea

²Dept. of Marine Bio-Materials and Aquaculture, Pukyong National University, Busan 48513, Korea

ABSTRACT : In vertebrate reproductive system, estrogen receptor (ER) plays a pivotal role in mediation of estrogenic signaling pathways. In the present study, we report the cDNA cloning, expression analysis, and transcriptional activity of ER β 1 subtype from medaka *Oryzias dancena*. The deduced *O. dancena* ER β 1 (odER β 1; 519 amino acids) contained six characteristic A/B to E/F domains with very short activation function 2 region (called AF2). A phylogenetic analysis indicated that odER β 1 was highly conserved among teleost ER β 1 subgroup. A conventional RT-PCR revealed that the *odER β 1* transcripts were widely distributed in the multiple tissues, the ovary, brain, gill, intestine, kidney, and muscle. Further, the relatively higher *odER β 1* expressions in the ovary and brain were clearly reproduced in RT-qPCR assay. When HA-fused odER β 1 expression vector was transfected into HEK293 cells, an immunoreactivity for odER β 1 was mainly detected in the nucleus part. Finally, an estrogen responsive element driven luciferase reporter assays demonstrated that the transcriptional activity of odER β 1 significantly increased by estradiol-17 β (E2) in a dose dependent manner ($p < 0.05$). However, fold-activation of odER β 1 in the presence of E2 was markedly weak, when it compared with those of *O. latipes* ER β 1. Taken together, these data suggest that odER β 1 represents a functional variant of teleost ER β subtype and provides a basic tool allowing future studies examining the function of F domain of ER β 1 subtype and expanding our knowledge of ER β evolution.

Key words : Estrogen receptor beta, Esr2a, Estradiol, Marine medaka

INTRODUCTION

The importance of the role of estrogens in the sexual development and reproduction of vertebrates is well known, and in particular, estrogens broadly contribute to regulation of female reproductive system, secondary sex characteristics during puberty, and ovulation (Rasier et al., 2006). Biosynthesis of estrogens appears to occur via the cytochrome P450 aromatase enzyme complex in diverse vertebrates, and it is generally accepted that estrogens are effectors for the reproduction strategy, such as implantation,

pregnancy recognition, pregnancy maintenance, parturition, and lactation (Lange et al., 2002). Estradiol-17 β (E2) is an estrogen and the major female sex hormone involved in the regulation of the estrous and menstrual female reproductive cycles. The biological effects of E2 occur through binding with estrogen receptors (Ers; Evans, 1988).

ERs belong to the superfamily of nuclear receptors and specifically to the family of steroid hormone receptors that act as ligand-mediated transcription factors (Evans, 1988; Zhao et al., 2008). Two forms of the ER, ER α , and ER β , have been characterized up to date. Both ER α and ER β

Manuscript received September 10, 2019, Received in revised form September 20, 2019, Accepted September 29, 2019

[†] Corresponding Author : Young Chang Sohn, Ph.D., Dept. of Marine Molecular Biosciences, Gangneung-Wonju National University, Gangneung 25457, Korea. Tel: +82-33-640-2348, E-mail: ycsohn@gwnu.ac.kr



This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

consist of distinctive domains, such as N-terminal trans-activation domain, central DNA-binding domain (DBD), ligand-binding domain (LBD), and C-terminal coactivator/corepressor interacting domain (Zhao et al., 2008). Although the biochemical responses of ER α and ER β to estrogens are largely similar in their modes, the two ERs show different and specific gene expression pattern in some tissues such as breast cancer (Thomas & Gustafsson, 2011; Haldosén et al., 2014).

In teleost fishes, ER α and ER β are widely expressed in various tissues, and hepatic vitellogenesis which is a pivotal process for ovarian maturation provides a physiologically relevant endpoint of ER activation (Nelson & Habibi, 2013). In fact, hepatic vitellogenesis and the ER α gene expression are commonly used as bioassays for endocrine disruptors that have estrogen-like activity, whereas the biological significances of ER β genes are not well elucidated up to now. Furthermore, involvement of ER β in the control of estrogen-responsive genes in specific tissues is also poorly understood. In a previous report, E2-dependent induction of red fluorescent protein reporter gene expression driven by choriogenin H gene promoter was shown in transgenic marine medaka *Oryzias dancena* under a wide range of salinity conditions (Cho et al., 2013). To evaluate ER β -mediated E2 responsiveness in euryhaline teleosts, we firstly investigated the expression profile of ER β gene in several tissues and transcriptional activities of the ER β proteins were examined in marine medaka.

MATERIALS AND METHODS

1. cDNA cloning of *O. dancena* ER β

Total RNA was extracted from the whole body of adult medaka *O. dancena*, using the RNA Prep kit (Bioneer, Daejeon, Korea) according to the manufacturer's protocol. To obtain full-length ER β cDNA, the first-strand cDNA was synthesized with the extracted total RNA using a SMART rapid amplification of cDNA ends (RACE) cDNA

Amplification kit (Clontech Lab, Palo Alto, CA, USA). A cDNA fragment was amplified with a forward and reverse primer set (#1 and #2; Table 1) based on known *O. latipes* and *O. javanicus* ER β cDNA sequences (GenBank Accession nos. NM_001104702 and AY917148). PCR amplification was performed in a final volume of 20 μ L containing 1 μ g of the marine medaka whole-body cDNA, 2.5 mM dNTP 0.8 μ L, 10 pmole of each oligo primer, 2 \times GC LA buffer 10 μ L, 5U LA Taq DNA polymerase (Takara Bio, Shiga, Japan). The reactions were carried out under the following cycling conditions: 5 min denaturation at 94°C; 35 cycles of 30 s at 94°C, 30 s at 50°C, and 2 min at 72°C; and 7 min at 72°C. Full-length cDNAs for entire coding region of *O. dancena* ER β (odER β) were obtained by RACE PCR amplification in a final volume of 25 μ L including 1 μ g of 3'- or 5'-cDNA template, 2 pmole of gene-specific primers (#1 or #2; Table 1) and the Universal Primer A Mix (#3, Table 1), 10 \times Advantage 2 PCR buffer 2.5 μ L, 10 mM dNTP mix 0.4 μ L, and 50 \times advantage 2 polymerase mix (Clontech Lab) 0.4 μ L with the following cycles: 5 min denaturation at 94°C; 5 cycles of 30 s at 94°C and 3 min at 68°C; 5 cycles of 30 s at 94°C, 30 s at 66°C, and 3 min at 72°C; 25 cycles of 30 s at 94°C, 30 s at 64°C, and 3 min at 72°C; and 7 min at 72°C. The amplified PCR products were cloned into pGEM-T Easy vector (Promega, Madison, WI, USA). To construct the odER β expression vector, open reading frame of the odER β cDNA was amplified by a conventional PCR using a primer set (#4 and #5, Table 1) and cloned into the Eco RI-Xho I restriction sites in pcDNA3-HA-NLS vector (Zavacki et al., 1997). *Escherichia coli* competent cells were transformed with the vectors and all of the constructs were sequenced by the Sanger method.

2. Phylogenetic and sequence analysis of *O. dancena* ER β 1 cDNA

The amino acid alignment was carried out by CLUSTALW with MEGA6 (Tamura et al., 2013), and the phylogenetic tree was constructed using the maximum-likelihood (ML)

Table 1. Oligo primers used in the polymerase chain reactions

Target	Primer	Direction	Sequence (5'→3')	Application
<i>odERβ1</i>	#1	Sense	GCAATGGGAACCACTTTGGACTC	Partial cDNA cloning
<i>odERβ1</i>	#2	Antisense	GATGTGAGCGTCCAGCATCTC	Partial cDNA cloning
<i>odERβ1</i>	#3	Sense/ antisense	CTAATACGACTCACTATAGGGCAAGCA GTGGTCAACGCAGAGT	RACE PCR
<i>odERβ1</i>	#4	Sense	<u>GCGAATTC</u> ATGGGAACCACTTTGGACT CAGAG	Plasmid construction
<i>odERβ1</i>	#5	Antisense	GCCTCGAGTCAACGCAGAGTGGCCATT ACGG	Plasmid construction
<i>18S rRNA</i>	#6	Antisense	CAAGAATTTACCTCTAGCGGC	Reverse transcription
<i>odERβ1</i>	#7	Sense	TCCAAGCTGCTCAGCATTCA	RT-qPCR
<i>odERβ1</i>	#8	Antisense	TCTGTTCTGCAGACAAAGG	RT-qPCR
<i>18S rRNA</i>	#9	Sense	TCCAGCTCCAATAGCGTATC	RT-qPCR
<i>18S rRNA</i>	#10	Antisense	AGAACCGGAGTCCTATTCCA	RT-qPCR

Underlined sequences indicate restriction enzyme recognition sites.

RACE, rapid amplification of cDNA ends.

method based on the JTT matrix-based model (Jones et al., 1992). Branch supports were provided using 1,000 bootstrap replicates.

3. Tissue distribution assay of *O. dancena* ERβ1 transcripts

To examine tissue distribution pattern of ERβ1 transcripts, 12 kinds of tissues including brain, eye, fin, gill, heart, intestine, kidney, liver, muscle, spleen, ovary, and testis were surgically removed from mature marine medaka individuals (six males and six females; average body weight=2.3±0.3 g) that had been communally grown under the culture conditions: salinity at 15 ppt, temperature at 26±1°C and dissolved oxygen level at 6±1 ppm. Total RNA was extracted with Trizol reagent (Ambion, Thermo Fisher Scientific, Waltham, MA, USA) according to the manufacturer's instruction and further purified with RNeasy Plus Mini Kit (Qiagen, Hilden, Germany) including the DNase I treatment step. Integrity and purity of total

RNA sample was verified with 28S:18S rRNA ratio on formaldehyde/MOPS agarose gel and OD260/280 nm (260/230 nm) spectrophotometry, respectively. For end-point RT-PCR, equal amount of total RNA (1 µg) from each individual tissue was pooled within a tissue type, and an aliquot (2 µg) of the pooled RNA sample was reverse transcribed to cDNA using Omniscript RT kit (Qiagen) with oligo-dT₂₀ priming method according to the manufacturer's protocol. In order to prepare a normalized control (18S rRNA; GenBank accession number HM347347), an *O. dancena* 18S rRNA reverse primer (#6, Table 1) was also included in the RT reaction as described previously. RT product was 8-fold (for ERβ1) or 20-fold (for 18S rRNA) diluted with sterile water, and 2 µL of diluted cDNA was used as template for each PCR reaction. End-point RT-PCR was carried with a pair of primer (#7 and #8; Table 1) to amplify a 240-bp *odERβ1* cDNA fragment. Thermal cycling conditions are 33 cycles of 94°C for 20 s, 58°C for 20 s and 72°C for 20 s with an initial denaturation

step at 94°C for 2 min. From each cDNA template, a 253-bp fragment of 18S rRNA was also amplified as a normalization control with a pair of primer (#9 and #10; Table 1) under the same thermal cycling conditions described in Cho et al. (2011) except the reduction of cycle number to 22. Amplified products (5 μ L) for both odER β 1 and 18S rRNA were electrophoresed on 2% agarose gels, and visualized with ethidium bromide (EtBr)-staining.

Based on the end-point RT-PCR assay, tissues examined to show the clear amplification of odER β 1 transcripts were selected for real-time quantitative RT-PCR (RT-qPCR). Basal expression levels of odER β 1 among those tissues were assayed with individual cDNA samples (three males and three females), and each cDNA template was assessed in triplicates. Oligonucleotide primers and thermal cycling conditions for odER β 1 and 18S rRNA were same as those used in end-point RT-PCR above, except the extension of the cycle number to 45. PCR was conducted with LightCycler 480 SYBR Green I Master Mix and LightCycler 480 Real-Time PCR System (Roche Applied Sciences, Penzberg, Germany). PCR efficiencies of both primer pairs (i.e., for odER β 1 and 18S rRNA) were 0.96 and 0.94, respectively. Basal expression level of odER β 1 transcripts in each tissue was determined with $2^{-\Delta CT}$ method based on the normalization against 18S rRNA level (Schmittgen & Livak, 2008).

4. Reporter assay

HEK293 cells (5×10^4 cells/well) were routinely cultured in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% fetal bovine serum (FBS, HyClone, GE Healthcare, Chicago, IL, USA) and 1% antibiotic-antimycotic (Invitrogen, Grand Island, NY, USA) in a humidified atmosphere containing 5% CO₂ at 37°C. The cells were seeded in 24-well plates and transfected with odER β 1, *O. latipes* ER β 1 (oER β 1), and masu salmon ER α (msER α) expression vectors and reporter vectors including an estrogen-response element (ERE)-driven luciferase

reporter (ERE-Luc) (Maeng et al., 2005), and internal control vector for β -galactosidase (200 ng each) using Lipofectamine™ 2000 (Invitrogen). The oER β expression vector was kindly provided by Dr. Yoshitaka Nagahama, National Institute for Basic Biology, Okazaki, Japan. After 3 hr post-transfection, the cell medium was changed by DMEM with 10% charcoal-stripped FBS and 1% antibiotic-antimycotic. Detailed transfection, ligand treatment and luciferase assay were conducted as described previously (Maeng et al., 2005).

5. Subcellular localization

HEK293 cells were seeded on poly-D-lysine hydrobromide (Sigma-Aldrich, St. Louis, MO, USA)-coated coverslips and transiently transfected with HA-tagged ER expression vectors (200 ng each) as described before. After 30 h, the cells were fixed for 10 min in 4% paraformaldehyde in PBS (Wako, Osaka, Japan) and further incubated for 10 min with PBS containing 0.1% Triton X-100. To block non-specific binding, the cells were incubated with 1% bovine serum albumin in PBST (PBS+0.1% Tween 20) for 30 min at room temperature. The cells were incubated with a 1:5,000 dilution of monoclonal anti-HA antibody produced in mouse (H9658, Sigma-Aldrich) in PBST overnight at 4°C in a humidified chamber. The cells were further incubated with a 1:2,000 dilution of anti-mouse IgG (H+L), F(ab')₂ Fragment (AlexaFluor 488 conjugate, Cell Signaling, Danvers, MA, USA) in PBST for 1 h at room temperature in the dark. The coverslips were mounted with Fluoroshield Mounting Medium with DAPI (Abcam, Cambridge, UK) and the cells were observed using a fluorescence microscope (Eclipse E200, Nikon, Tokyo, Japan). Fluorescence signals were not detected in the control cells incubated without the primary antibody.

6. Statistical analysis

Differences in basal mRNA expression levels of *odER β* among adult tissues and luciferase activities

determined with reporter assays for ERs were assessed with ANOVA followed by Duncan's multiple range test, while differences between sexes given a tissue type with Student's *t*-test. Difference was considered to be significant when $p < 0.05$.

RESULTS

1. Isolation of *O. dancena* ERβ1 cDNA and sequence analysis

We cloned an ERβ1 cDNA from the whole body of adult marine medaka *O. dancena* based on *O. latipes* and *O. javanicus* ERβ cDNA sequences. The cDNA contained an open reading frame of 1,560 nucleotides encoding 519 amino acid residues. The overall protein sequence identities of the odERβ1 was relatively higher with ERβ1 subtype of *O. latipes*, *Paralichthys olivaceus*, and *Oncorhynchus mykiss* (93.8%, 81.5%, and 69.4%, respectively), but relatively lower with ERβ2 and ERα subtypes of teleosts and human (44.1%–57.0%; Fig. 1A). Amino acid sequences of N-terminal A/B and DNA binding C domains of odERβ1 were strikingly similar to those of olERβ1 (96.1% and 100%), while the sequences of hinge D domain and ligand binding E/F domains of odERβ1 showed relatively lower similarities with the corresponding regions of olERβ1 (82.9% and 92.3%). The C-terminal amino acid sequence of odERβ1 was short when compared with those of olERβ1 and human ERβ (Fig. 1B). A molecular phylogenetic analysis revealed that teleostean ERβ1 proteins including odERβ1 formed a monophyletic group that was distinct from a clade comprising tetrapodian ERβ and teleostean ERβ2 and ERα proteins (Fig. 2).

2. Tissue distribution pattern and basal expression levels of *O. dancena* ERβ1 transcripts

From end-point RT-PCR analysis, *odERβ1* mRNAs were predominantly expressed in the ovary (Fig. 3A). Besides the ovary, *odERβ1* transcripts were also detectable

in the brain, gill, intestine, kidney, and muscle, although expression levels in those tissues were significantly lower than that of the ovarian expression. Meanwhile, under the present RT-PCR conditions, *odERβ1* mRNAs were not clearly detected in the eye, fin, heart, liver, spleen and testis. The robust expression of *odERβ1* mRNAs in the ovary was clearly reproduced in RT-qPCR assay with female tissues, in which the ovarian expression level was 3.8–15.3 times greater than those in other tissues examined ($p < 0.05$; Fig. 3B). Among non-ovarian tissues, brain showed a higher expression compared to other tissues ($p < 0.05$). Expression patterns in male somatic tissues were not significantly dissimilar from those observed in female. Like in female tissues, male brains displayed a relatively higher expression than others did, although the difference was not highlighted as much as in female (Fig. 3C). In comparison between genders within a given tissue, females tended to show a higher expression level of *odERβ1* in the brain than males did, however, the difference was not statistically significant due to large individual variations. There was no gender-dependent difference in the expression level of the expression level of *odERβ1* for gill, intestine, muscle and kidney ($p > 0.05$).


3. Functional characterization of *O. dancena* ERβ1

The transcriptional activity of the odERβ1 was evaluated by cotransfection of a luciferase reporter, ERE-Luc, into a mammalian cell line, HEK293. The ERE-Luc reporter activities of odERβ1 were significantly elevated by E2 and the response of odERβ1 showed a dose dependent manner ($p < 0.05$). The transcriptional fold-activation of odERβ1 in the presence of E2 was markedly weak, when the activities of *O. latipes* ERβ1 (olERβ1) and *Oncorhynchus masou* ERα (msERα) were compared (Fig. 4B). The fold-transactivation of odERβ1 by E2 treatment showed a dose dependent manner. Transfected HA-tagged odERβ1, olERβ1, and msERα expression vectors were mainly expressed in the nucleus of HEK293 cells (Fig. 4C).

(A)

Entire coding region

<i>Oryzias latipes</i> ERβ1	93.8 (%)				
<i>Paralichthys olivaceus</i> ERβ1	81.5				
<i>Oncorhynchus mykiss</i> ERβ1	69.4				
<i>Oryzias latipes</i> ERβ2	53.4				
<i>Paralichthys olivaceus</i> ERβ2	57.0				
<i>Homo sapiens</i> ERβ	50.9				
<i>Oryzias latipes</i> ERα	43.9				
<i>Oncorhynchus masou</i> ERα	44.1				amino acids



(B)

1 80

O. dancena KGMDHLHCMK MKNVVFVLYDL LLEMLDAHIP AVMATLR

O. latipes KGMDHLHCMK MKNMVFLYDL LLEMLDAHIM HSSRLPRQFP PQDAADPTET SAQGQRSCCD VSKAWTTSSA GTAEEPQKSD

H. sapiens KGMEHLLNMK CKNVVFVYDL LLEMLNAHVL RGCKSSITGS ECSPAEDSKS KEGSQNPQSQ

Consensus KGM#HLhcmk mKNvVFVLYDL LLEML#AH!r.....a.....q.....

Fig. 1. Protein sequence identities of *Oryzias dancena* estrogen receptor (ER) β1 subtype with teleost and human ER orthologs. (A) Identities for the entire coding region are represented at the upper lines. ERs typically contain six distinct domains termed A to F extending from the N- to C-terminus. The four characteristic domains of the ERβ1 schematically represented with their corresponding identity percentages at the bottom lines. (B) Multiple sequence alignment of *Oryzias* ERβ1 and human ERβ C-terminal amino acids. Amino acids with 100 % and >60% conservation are shown in red and blue, respectively. The corresponding GenBank accession numbers for ER sequences are referred in the legend of Fig. 2.

DISCUSSION

In the present study, we cloned an ERβ1 subtype cDNA from *O. dancena* and examined the mRNA expression profile in various tissues. To date, two ERβ subtypes (ERβ1 and ERβ2) have been isolated from a variety of

teleost fish species, including Japanese medaka and largemouth bass (Sabo-Attwood et al., 2004; Chakraborty et al., 2011), although a single ERβ gene was reported in mammals (Haldosén et al., 2014). The overall sequence identity analysis of three subtypes of ER indicated that odERβ1 is the ortholog of teleost ERβ1. This classification




Fig. 2. Maximum-likelihood phylogenetic tree showing relationships between amino acid sequences of estrogen receptor (ER) subtypes. The bootstrap values from 1,000 replicates are given at each branch node. The scale bar indicates 0.1 amino acid replacements per site. GenBank accession numbers are as follows: ER β 1 subtype for *Oryzias dancena* (MN729292), *O. latipes* ER β 1 (NP_001098172), *Paralichthys olivaceus* (BAB85623), *Oreochromis aureus* (ACF75102), *Fundulus heteroclitus* (AAU44352), *Epinephelus coioides* (ADK90034), *Cyprinus carpio* (BAB91218), *Oncorhynchus mykiss* (P_001118225); ER β 2 subtype for *O. latipes* (NP_001121984), *O. aureus* (ACF75103), *F. heteroclitus* (NP_001296906), *P. olivaceus* (XP_019961638), *E. coioides* (ADK90035), *O. mykiss* (NP_001118042); ER β subtype for *Callorhinchus milii* (BAX07664), *Python bivittatus* (XP_007432491), *Xenopus tropicalis* (NP_001035101), *Gallus gallus* (NP_990125), *Homo sapiens* (CAA67555), *Mus musculus* (AAI41076); ER α subtype for *O. latipes* (P50241) and *O. masou* (AAS92970).

was supported by a phylogenetic analysis: ER β 1 and ER β 2 subtypes of teleost species and ER β of tetrapods. The estimated amino acid length of odER β 1 were 519 which is shorter than olER β 1 (562 amino acids; Chakraborty et al., 2011) and human ER β (530 amino acids; Ogawa et al.,

1998). The lacking of large portion of F domain known as a hormone-dependent activation function AF-2 (Glass & Rosenfeld, 2000) is the most different feature between odER β 1 and olER β 1. Taken together, odER β 1 is most likely a splicing variant of ER β 1 gene. In human, multiple




Fig. 3. Distribution and basal expression levels of ER β 1 transcripts in tissues of mature *Oryzias dancena*. (A) Representative end-point RT-PCR gels to show the RT-PCR products of ER β 1 and 18S rRNA control. (B, C) Relative expression levels of ER β 1 mRNAs in tissues selected from females (B) or males (C) determined with RT-qPCR assay based on the normalization against 18S rRNA levels. Only tissues proven to show the clear expression of ER β 1 in the end-point RT-PCR were subjected to RT-qPCR assay. Mean \pm SDs with different letters within a given gender (a–c) are significantly different based on ANOVA followed by Duncan's multiple range test at $p < 0.05$. ER, estrogen receptor.

ER β isoforms lacking or changing of F domain by alternative splicing of the last coding exons have been reported (Haldosén et al., 2014). The AF-2 region of E2-bound ERs mainly interact with transcriptional co-regulators, i.e., N-CoR, SMRT, and GRIP1, and then the ER complex may recruit the basal transcription machinery including RNA polymerase II (Glass & Rosenfeld, 2000; Webb et al., 2003). Although the function of AF-2 region of ER β is unclear in teleosts, AF2-deleted female mice were significantly less preferred by adult males than that of normal females in odor preference tests (Antal et al., 2012). Implication of ER β 1 variants in the reproductive axis of teleosts is needed to be elucidated.

Tissue-specific expression of ER subtypes was demonstrated during the life cycle of teleosts (Sabo-Attwood et al., 2004; Chakraborty et al., 2011). The *odERβ1* transcripts were strongly expressed in the ovary and significantly detectable in the brain than those of gill, intestine, kidney, and muscle, whereas *olERβ1* mRNA expression was observed as faint bands in the testis and the brain (Chakraborty et al., 2011). In rainbow trout, the mRNA for ER β 1 was detected in diverse tissues, e.g., the testis, ovary, brain, and kidney (Nagler et al., 2007). Interestingly, the rainbow trout ER β 1 transcript abundance in the ovary showed the highest expression at the beginning of the reproductive cycle, but then decreased afterward (Nagler et al., 2012). The expression of rockfish ER β 1 mRNA was also significantly higher in the ovary at the early-oocyte stage (Mu et al., 2013). The brain of female *O. dancena* showed relatively higher level of *odERβ1* transcript than other tissues examined. Recently, ER β 1-null female medaka showed apparently normal sexual behavior, but without fertility and oviposition in response to male courtship (Kayo et al., 2019). These observations suggest that ER β 1 expression may be important for estrogen signaling that starts the next cycle of ovarian development through brain-pituitary-gonad axis in teleosts. In fact, expression of *esr1* (ER α gene), *gnrhr1*, *fsh*, *lh* genes in the brain of red

Oryzias dancena ER β 1




Fig. 4. Transcriptional activities and localization of *Oryzias dancena* ER β 1 (odER β 1) in HEK-293 cells. (A) Transcriptional activity of odER β 1 with estradiol-17 β (E2, 1 μ M). HEK-293 cells were transfected with ERE-Luc reporter vector (200 ng) and β -galactosidase (β -gal) expression vector (100 ng), along with either pcDNA3-HA empty vector or odER β 1 expression vector as indicated. After 24 h of transfection, the cells were incubated in the presence of 1 μ M dosage of E2 or the same volume of DMSO for 18 h. All cells were lysed and the luciferase activities were measured and normalized against the β -gal expression as an internal control. Values are the average of at least four transfections (mean \pm SD). (B) HEK-293 cells were transfected with expression vectors for odER β 1, *O. latipes* ER β 1 (oER β 1), or masu salmon ER α (msER α) expression vector (200 ng each), together with ERE-Luc reporter vector and β -gal expression vector followed by treatment of E2 as indicated doses. Luciferase activity was measured 18 h after ligand addition and normalized as in (A). Mean \pm SDs with different letters are significantly different based on ANOVA followed by Duncan's multiple range test at $p < 0.05$. (C) HEK-293 cells were transfected with HA-tagged odER β 1, oER β 1, and msER α expression vector (200 ng each) and the cells were fixed and examined by immunohistochemistry with anti-HA antibody and DAPI. Magnification $\times 1,000$. ER, estrogen receptor.

spotted grouper was positively associated with the formation of ovarian cavity (Kim et al., 2016). More in-depth investigations of individual tissues during different reproductive stages of medaka *O. dancena* will be required to better understand the biological significance of *ERβ* isoforms.

The transcriptional activities of odERβ1 were significantly elevated by E2 concentration dependent manner. In addition, the nuclear localization and the E2-dependent transactivation mode of odERβ1 were evidently demonstrated like those of olERβ1 and msERα. These results are similar to a previous report that three medaka ER subtypes including olERβ1 are capable of initiating transactivation of vitellogenin genes via upstream EREs (Lee Pow et al., 2016). The transcriptional activity of steroid hormone receptor superfamily including ERs is mainly dependent upon C-terminal ligand-binding E/F domain and AF-2 region (Glass & Rosenfeld MG, 2000). When the amino acid sequence of odERβ1 was compared with those of olERβ1 and human ERβ, odERβ1 was estimated to be a splicing variant from ERβ1 gene that lacks F domain and/or AF2 region. The olERβ1 having a long F domain showed higher fold-activation of transcription of ERE-driven reporter gene in the presence of E2, whereas odERβ1 showed relatively weak fold-activation of the reporter gene in the presence of E2. These data indicate that the complete F domain and AF2 region are important for ligand binding and direct E2-dependent transcriptional activity of ERβ1. In fact, a mutant mammalian ERα deleted for the AF2 core domain impairs E2-induced transactivation by wild type ERα (Jung et al., 2001).

In summary, we cloned an *O. dancena* ERβ1 variant having a short C-terminal activation function 2 region. The *odERβ1* mRNAs were highly expressed in the ovary and brain. The transcriptional activity of odERβ1 increased by E2 in a dose dependent manner, although E2-dependent transactivation of odERβ1 was weaker than that of *O. latipes* ERβ1.

ORCID

Sejung Maeng

<https://orcid.org/0000-0002-0730-0300>

Sung Woo Yoon

<https://orcid.org/0000-0002-4104-9068>

Eun Jeong Kim

<https://orcid.org/0000-0001-8303-1731>

Yoon Kwon Nam

<https://orcid.org/0000-0001-8870-2098>

Young Chang Sohn

<https://orcid.org/0000-0002-4140-2669>

AUTHOR CONTRIBUTIONS

Conceptualization: Sohn YC.

Data curation: Maeng S, Yoon SW, Kim EJ.

Formal analysis: Nam YK, Sohn YC.

Methodology: Maeng S, Kim EJ.

Software: Maeng S, Yoon SW, Kim EJ.

Validation: Maeng S, Kim EJ.

Investigation: Maeng S, Yoon SW, Kim EJ.

Writing-original draft: Maeng S, Yoon SW, Kim EJ, Nam YK, Sohn YC.

Writing-review & editing: Maeng S, Yoon SW, Kim EJ, Nam YK, Sohn YC.

ETHICS APPROVAL

Experiment with living marine medaka samples was approved by the Animal Care and Use Committee of Pukyong National University (Approved number 201816).

REFERENCES

Antal MC, Petit-Demoulière B, Meziane H, Chambon P,

- Krust A (2012) Estrogen dependent activation function of ER β is essential for the sexual behavior of mouse females. *Proc Natl Acad Sci USA* 109:19822-19827.
- Chakraborty T, Shibata Y, Zhou LY, Katsu Y, Iguchi T, Nagahama Y (2011) Differential expression of three estrogen receptor subtype mRNAs in gonads and liver from embryos to adults of the medaka, *Oryzias latipes*. *Mol Cell Endocrinol* 333:47-54.
- Cho YS, Lee SY, Kim YK, Kim DS, Nam YK (2011) Functional ability of cytoskeletal β -actin regulator to drive constitutive and ubiquitous expression of a fluorescent reporter throughout the life cycle of transgenic marine medaka *Oryzias dancena*. *Transgenic Res* 20:1333-1355.
- Cho YS, Kim DS, Nam YK (2013) Characterization of estrogen-responsive transgenic marine medaka *Oryzias dancena* germlines harboring red fluorescent protein gene under the control by endogenous choriogenin H promoter. *Transgenic Res* 22:501-517.
- Evans RM (1988) The steroid and thyroid hormone receptor superfamily. *Science* 240:889-895.
- Glass CK, Rosenfeld MG (2000) The coregulator exchange in transcriptional functions of nuclear receptors. *Genes Dev* 14:121-141.
- Haldosén LA, Zhao C, Dahlman-Wright K (2014) Estrogen receptor beta in breast cancer. *Mol Cell Endocrinol* 382:665-672.
- Jones DT, Taylor WR, Thornton JM (1992) The rapid generation of mutation data matrices from protein sequences. *Comput Appl Biosci* 8:275-282.
- Jung DJ, Lee SK, Lee JW (2001) Agonist-dependent repression mediated by mutant estrogen receptor alpha that lacks the activation function 2 core domain. *J Biol Chem* 276:37280-37283.
- Kayo D, Zempo B, Tomihara S, Oka Y, Kanda S (2019) Gene knockout analysis reveals essentiality of estrogen receptor β 1 (Esr2a) for female reproduction in medaka. *Sci Rep* 9:8868.
- Kim HK, Kim JH, Baek HJ, Kwon JY (2016) Gene expression of aromatases, steroid receptor, GnRH and GTHs in the brain during the formation of ovarian cavity in red spotted grouper, *Epinephelus akaara*. *Dev Reprod* 20:367-377.
- Lange IG, Hartel A, Meyer HHD (2002) Evolution of oestrogen functions in vertebrates. *J Steroid Biochem Mol Biol* 83:219-226.
- Lee Pow CSD, Yost EE, Aday DD, Kullman SW (2016) Sharing the roles: An assessment of Japanese medaka estrogen receptors in vitellogenin induction. *Environ Sci Technol* 50:8886-8895.
- Maeng S, Jung Y, Choi E, Jeon JK, Kim S, Gen K, Sohn YC (2005) Expression of gonadotropin subunit genes following 4-nonylphenol exposure in masu salmon: Effects on transcript levels and promoter activities via estrogen receptor alpha. *Comp Biochem Physiol B Biochem Mol Biol* 142:383-390.
- Mu WJ, Wen HS, Shi D, Yang YP (2013) Molecular cloning and expression analysis of estrogen receptor betas (ER β 1 and ER β 2) during gonad development in the Korean rockfish, *Sebastes schlegeli*. *Gene* 523:39-49.
- Nagler JJ, Cavileer T, Sullivan J, Cyr DG, Rexroad C (2007) The complete nuclear estrogen receptor family in the rainbow trout: Discovery of the novel ERalpha2 and both ERbeta isoforms. *Gene* 392:164-173.
- Nagler JJ, Cavileer TD, Verducci JS, Schultz IR, Hook SE, Hayton WL (2012) Estrogen receptor mRNA expression patterns in the liver and ovary of female rainbow trout over a complete reproductive cycle. *Gen Comp Endocrinol* 178:556-561.
- Nelson ER, Habibi HR (2013) Estrogen receptor function and regulation in fish and other vertebrates. *Gen Comp Endocrinol* 192:15-24.
- Ogawa S, Inoue S, Watanabe T, Hiroi H, Orimo A, Hosoi T, Ouchi Y, Muramatsu M (1998) The complete primary structure of human estrogen receptor beta (hER beta)

- and its heterodimerization with ER alpha *in vivo* and *in vitro*. *Biochem Biophys Res Commun* 243:122-126.
- Rasier G, Toppari J, Parent AS, Bourguignon JP (2006) Female sexual maturation and reproduction after prepubertal exposure to estrogens and endocrine disrupting chemicals: A review of rodent and human data. *Mol Cell Endocrinol* 254-255:187-201.
- Sabo-Attwood T, Kroll KJ, Denslow ND (2004) Differential expression of largemouth bass (*Micropterus salmoides*) estrogen receptor isotypes alpha, beta, and gamma by estradiol. *Mol Cell Endocrinol* 218:107-118.
- Schmittgen TD, Livak KJ (2008) Analyzing real-time PCR data by the comparative C(T) method. *Nat Protoc* 3:1101-1108.
- Tamura K, Stecher G, Peterson D, Filipinski A, Kumar S (2013) MEGA6: Molecular evolutionary genetics analysis version 6.0. *Mol Biol Evol* 30:2725-2729.
- Thomas C, Gustafsson JA (2011) The different roles of ER subtypes in cancer biology and therapy. *Nat Rev Cancer* 11:597-608.
- Webb P, Valentine C, Nguyen P, Price RH, Marimuthu A, West BL, Baxter JD, Kushner PJ (2003) ERbeta binds N-CoR in the presence of estrogens via an LXXLL-like motif in the N-CoR C-terminus. *Nucl Recept* 1:4.
- Zavacki AM, Lehmann JM, Seol W, Willson TM, Klierer SA, Moore DD (1997) Activation of the orphan receptor RIP14 by retinoids. *Proc Natl Acad Sci USA* 94:7909-7914.
- Zhao C, Dahlman-Wright K, Gustafsson JA (2008) Estrogen receptor beta: An overview and update. *Nucl Recept Signal* 6:e003.