

Subdominant/Cryptic CD8 T Cell Epitopes Contribute to Resistance against Experimental Infection with a Human Protozoan Parasite

Mariana R. Dominguez^{1,2}, Eduardo L. V. Silveira^{1,2^{‡a}}, José Ronnie C. de Vasconcelos^{1,2}, Bruna C. G. de Alencar^{1,2^{‡b}}, Alexandre V. Machado³, Oscar Bruna-Romero⁴, Ricardo T. Gazzinelli^{4,5,6}, Mauricio M. Rodrigues^{1,2*}

1 Centro de Terapia Celular e Molecular (CTCMol), Universidade Federal de São Paulo-Escola Paulista de Medicina, São Paulo, Brazil, **2** Departamento de Microbiologia, Imunologia e Parasitologia, Universidade Federal de São Paulo-Escola Paulista de Medicina, São Paulo, Brazil, **3** Centro de Pesquisas René Rachou, FIOCRUZ, Belo Horizonte, Minas Gerais, Brazil, **4** Departamento de Microbiologia, Instituto de Ciências Biológicas, Universidade Federal de Minas Gerais, Belo Horizonte, Minas Gerais, Brazil, **5** Departamento de Bioquímica e Imunologia, Universidade Federal de Minas Gerais, Belo Horizonte, Minas Gerais, Brazil, **6** Division of Infectious Disease and Immunology, Department of Medicine, University of Massachusetts Medical School, Worcester, Massachusetts, United States of America

Abstract

During adaptive immune response, pathogen-specific CD8⁺ T cells recognize preferentially a small number of epitopes, a phenomenon known as immunodominance. Its biological implications during natural or vaccine-induced immune responses are still unclear. Earlier, we have shown that during experimental infection, the human intracellular pathogen *Trypanosoma cruzi* restricts the repertoire of CD8⁺ T cells generating strong immunodominance. We hypothesized that this phenomenon could be a mechanism used by the parasite to reduce the breadth and magnitude of the immune response, favoring parasitism, and thus that artificially broadening the T cell repertoire could favor the host. Here, we confirmed our previous observation by showing that CD8⁺ T cells of H-2^a infected mice recognized a single epitope of an immunodominant antigen of the *trans*-sialidase super-family. In sharp contrast, CD8⁺ T cells from mice immunized with recombinant genetic vaccines (plasmid DNA and adenovirus) expressing this same *T. cruzi* antigen recognized, in addition to the immunodominant epitope, two other subdominant epitopes. This unexpected observation allowed us to test the protective role of the immune response to subdominant epitopes. This was accomplished by genetic vaccination of mice with mutated genes that did not express a functional immunodominant epitope. We found that these mice developed immune responses directed solely to the subdominant/cryptic CD8 T cell epitopes and a significant degree of protective immunity against infection mediated by CD8⁺ T cells. We concluded that artificially broadening the T cell repertoire contributes to host resistance against infection, a finding that has implications for the host-parasite relationship and vaccine development.

Citation: Dominguez MR, Silveira ELV, de Vasconcelos JRC, de Alencar BCG, Machado AV, et al. (2011) Subdominant/Cryptic CD8 T Cell Epitopes Contribute to Resistance against Experimental Infection with a Human Protozoan Parasite. PLoS ONE 6(7): e22011. doi:10.1371/journal.pone.0022011

Editor: Georges Snounou, Université Pierre et Marie Curie, FRANCE

Received: March 29, 2011; **Accepted:** June 11, 2011; **Published:** July 14, 2011

Copyright: © 2011 Dominguez et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: Fundação de Amparo à Pesquisa do Estado de São Paulo (2009/06820-4), The National Institute for Vaccine Technology (INCTV-CNPq), The Millennium Institute for Vaccine Development and Technology (CNPq - 420067/2005-1) and The Millennium Institute for Gene Therapy (Brazil). EVLS, RTG and MMR are recipients of fellowships from CNPq. MRD and BCA are recipients of fellowships from FAPESP. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: mrodrigues@unifesp.br

^{‡a} Current address: Yerkes National Primate Research Center/Emory University, Atlanta, Georgia, United States of America

^{‡b} Current address: Institut Curie U932, Paris, France

Introduction

MHC class Ia-restricted CD8⁺ T cells are important mediators of the adaptive immune response against infections caused by intracellular microorganisms, including the digenetic intracellular protozoan parasite *Trypanosoma cruzi*, the causative agent of Chagas disease (American trypanosomiasis). During experimental infection, this T cell subpopulation has been shown to be critical for host survival even when small doses of parasites are used in challenges [1–5]. In spite of the CD8⁺ T-cell mediated immune response, the parasite survives within the host and establishes a life-long chronic infection. Parasite persistence is considered one of the critical factors in the development of the complex immuno-

pathology caused by *T. cruzi* that may occur years after the initial infection in ~30% of infected individuals [6–11]. Thus, understanding how the parasites escape the immune response and persist for such long periods may help us to find new means for interventions against Chagas disease that would improve quality of life for millions of infected individuals in Latin America

Recent studies on the CD8⁺ T-cell immune responses that occur during experimental *T. cruzi* infection in inbred mouse strains described a surprising immunodominance of certain epitopes expressed by members of a large family of *T. cruzi* surface antigens named *trans*-sialidases (TS) [1,5,12–20]. How and why this strong pattern of immunodominance is established is still a matter of debate. In general terms, immunodominance can emerge as a

result of different mechanisms that regulate the formation of the complex of MHC-I-peptide on the surface of antigen presenting cells (APC) such as antigen concentration, stability or epitope availability after processing and translocation to the endoplasmic reticulum, where the MHC-I-peptide complex is assembled to be transported to the APC surface [21–24].

After a stable MHC-I-peptide complex is formed on the surface of the APC, factors related to CD8⁺ T cells, such as the frequency of precursors, their TCR affinities, their capacity to proliferate in response to antigen and, thus, be incorporated into the pool of responder cells, are factors that shape immunodominance hierarchies. These factors transcend the MHC restriction element and may create T-cell competition for APCs and other resources, enabling certain CD8⁺ T cells to dominate and suppress others [23–29].

By comparing the specificity of CD8⁺ T cells of homozygous and heterozygous mouse strains, we observed that the immunodominance that occurs during experimental *T. cruzi* infection could be exerted not only on epitopes restricted by the same MHC molecules but also, unexpectedly, on the immune response to epitopes restricted by different MHC-I molecules. This phenomenon, termed cross-competition, represents a potent means by which T cells with a certain specificity may become immunodominant [30–34]. This strong and unusual phenomenon has been shown to be due to *T. cruzi* infection because following immunization with recombinant adenovirus expressing the same parasite antigens, this pattern of immunodominance was not observed [15].

Based on these observations, we hypothesized that this competition/immunodominance between T cells of different specificities could be a sophisticated strategy that *T. cruzi* developed to reduce the breadth and magnitude of CD8⁺ T-cell responses, suppressing the immune responses of these T cells with other specificities in order to escape complete elimination by host effector cells. Thus, we expected that artificially broadening the immune response to include T cells specific for subdominant or cryptic epitopes could favor the host, counteracting the restriction imposed by the infection. Here, we tested this hypothesis by using mice genetically immunized with a mutated form of the *amastigote surface protein (asp) - 2* gene in which the immunodominant CD8 T cell epitope is no longer functional. The CD8 T cell-mediated immune response of these mice was directed only to the newly described subdominant/cryptic CD8 T cell epitopes of ASP-2. Even in the absence of an immune response directed to the immunodominant epitope, these mice displayed a significant degree of protective immunity, albeit not as strong as the immune response elicited by the original gene expressing both the immunodominant and the subdominant epitopes. These results are compatible with our hypothesis that artificially broadening the immune response favors the host. Indirectly, we suggest that immunodominance may in fact be a mechanism to establish a chronic infection.

Materials and Methods

Ethics Statement

All experimental procedures were approved by the Ethics Committee for Animal Care of the Federal University of São Paulo (Id # CEP 0426/09).

Mice and parasites

Female 8-week-old H-2^a mice (B10.A and A/Sn) were purchased from CEDEME (Federal University of São Paulo). Bloodstream trypomastigotes of the Y strain of *T. cruzi* were obtained from A/Sn mice infected 7–8 days earlier [12]. Each

B10.A or A/Sn mouse was challenged i.p. with a final dose containing 10⁴ or 150 parasites, respectively, in a final volume of 0.2 mL. Parasite development was monitored by counting the number of bloodstream trypomastigotes in 5 µL of fresh blood collected from the tail vein [12].

Peptides

Peptides were purchased from Genscript (Piscataway, NJ). Purity was as follows: TEWETGQI (95%); PETLGHEI (97.4%); YEIVAGYI (99.40%); TPTAGLVGF (98.6%); GSRNGNDRL (97.1%); ESKSGDAPL (96.1%); HEHNLFGI (98.7%); ESSTP-TAGL (99.1%); ESEPKRPNM (98.7%); VSWGEPKSL (99.2%); YSDGALHLL (97.3%); AESWPSIV (96.5%); and RPNMSRHLF (99.4%).

Recombinant plasmids and adenoviruses

Plasmid pIgSPCL.9 and the human replication-defective adenovirus type 5 containing the *asp-2* gene were obtained as described previously [35,36]. Mutated *asp-2* was generated by a series of PCR reactions using DNA encoding the *asp-2 clone 9* gene as a template (Genbank Accession Number: AY186572). In the first reaction, the forward and reverse oligonucleotides were as follows:

- i) 5'-GGGGGTACCATGCTCTCACGTGTTGCT-3';
- ii) 5'-GAACGATCATGAGTGCTTGGCCCGTCTCC-CATGCGGTGATGCGGGGATC-3'

In the second reaction, they were as follows:

- i) 5'-GATCCCCGCATCACCGCATGGGAGACGGGA-CAAGCACTCATGATCGTTC-3'
- ii) 5'-GGGTCTAGATCAGACCATTTTGTAGTTCACC-3'.

PCR products were purified, mixed and subjected to a third PCR reaction. Forward and reverse oligonucleotides were respectively

- i) 5'-GGGGGTACCATGCTCTCACGTGTTGCT-3';
- ii) 5'-GGGTCTAGATCAGACCATTTTGTAGTTCACC-3'.

The final PCR product was completely sequenced. The only modifications found were in the nucleotide sequences encoding the immunodominant epitope TEWETGQI. The new sequence encoded the amino acids (AA) TAWETGQA. This plasmid is referred to as pIgSpTAWETGQA. The new gene was also subcloned into the pAdCMV shuttle vector, and the recombinant replication-defective adenovirus human adenovirus 5 was produced by Vectors BioLabs, Philadelphia, USA. This new recombinant adenovirus is referred to as AdTAWETGQA. Viruses and plasmids were purified as described previously [35–37]. Mice were inoculated intra-muscularly (i.m.) in each *tibialis anterioris* muscle with 50 µg of plasmid DNA 3 times every 3 weeks.

Heterologous prime-boost immunization consisted of priming i.m. with a total of 100 µg of plasmid DNA followed by a dose of viral suspension containing 2 × 10⁸ plaque forming units (pfu) of adenovirus twenty-one days later in the same locations. Immunological assays or challenges were performed 14 days after viral inoculation.

In vivo depletion of CD8⁺ T cells were performed by treating vaccinated A/Sn mice with 53.6.7 MAb. At days 2 and 3 before challenge with trypomastigotes, mice were injected i.p. with a dose of 1 mg of anti-CD8 or control Rat IgG. Seven days after challenge, each mouse received one more dose of 1 mg of anti-CD8 or Rat IgG. The efficacy of depletion of CD8⁺ spleen cells




Figure 1. Structure and localization of Amastigote Surface Protein-2 (ASP-2). A- Schematic view of the primary structure of *T. cruzi* ASP-2. B- HeLa cells were infected for 48 h with trypomastigotes of the Y strain. After fixation, indirect immunofluorescence or DAPI staining were performed as described using MAb K22 and imaged using fluorescence microscopy [39]. Bar, 14 μ M. doi:10.1371/journal.pone.0022011.g001

before challenge was more than 95% in anti-CD8 treated mice compared to Rat IgG treated ones.

Immunological T cell assays

Ex vivo ELISPOT (IFN- γ) or *in vivo* cytotoxic assays were performed exactly as described previously [12,15]. The surface mobilization of CD107a and the intracellular expression of cytokines (IFN- γ , TNF- α , IL-2 and IL-10) was evaluated after *in vitro* culture of splenocytes in the presence or absence of antigenic stimulus. Cells were washed three times in plain RPMI and re-suspended in cell culture medium consisting of RPMI 1640 medium, pH 7.4, supplemented with 10 mM Hepes, 0.2% sodium bicarbonate, 59 mg/l of penicillin, 133 mg/l of streptomycin, and 10% Hyclone fetal bovine sera (Hyclone, Logan, Utah). The viability of the cells was evaluated using 0.2% Trypan Blue exclusion dye to discriminate between live and dead cells. Cell concentration was adjusted to 5×10^6 cells/mL in cell culture medium containing anti-CD28 (2 μ g/mL), Brefeldin A (10 μ g/mL), Monensin (5 μ g/mL) and FITC-labeled anti-CD107a (Clone 1D4B, 2 μ g/mL, BD Pharmingen). In half of the cultures, a final concentration of 10 μ M of the VNHRFTLV peptide was added. The cells were cultivated in flat-bottom 96-well plates (Corning) in a final volume of 200 μ l in duplicate, at 37°C in a humid environment. After a 20-h incubation, cells were stained for surface markers with Per-CP or PE-labeled anti-CD8 on ice for 20 min. To detect IFN- γ , TNF- α ? IL-2 and IL-10 by intra-cellular staining (ICS), cells were then washed twice in buffer containing PBS, 0.5% BSA and 2 mM EDTA, fixed in 4% PBS-paraformaldehyde solution for 10 minutes and permeabilized for 15 minutes in a PBS, 0.1% BSA, 0.1% saponin solution. After being washed twice, cells were stained for intracellular markers using APC or PE-labeled anti-IFN- γ (Clone XMG1.2) and? PE- labeled anti-TNF- α (clone MP6-XT22), APC-labeled anti-IL-2 (clone JES6-5H4) or APC-labeled anti-IL-10 (JES5-16E3) for 20 minutes on ice. Finally, cells were washed twice and fixed in 1% PBS-paraformaldehyde. At least 300,000 cells were acquired on a BD FACS Canto flow cytometer and then analyzed with FlowJo.

Statistical analysis

Values were expressed as means \pm SD. These values were compared using one-way ANOVA followed by Tukey's HSD tests

(<http://faculty.vassar.edu/lowry/VassarStats.html>). The LogRank test was used to compare mouse survival rates after challenge with *T. cruzi* (<http://bioinf.wehi.edu.au/software/russell/logrank/>). The differences were considered significant when the *P* value was <0.05.

Results

During experimental infection of H-2^b or H-2^a inbred mouse strains with parasites of the Y strain of *T. cruzi*, two epitopes were identified within the ASP-2 antigen represented by the VNHRFTLV or TEWETGQI peptides. They were recognized by H-2K^b- or H-2K^k-restricted CD8⁺ cytotoxic T cells, respectively [12,14,15,37,38]. ASP-2 is a member of the large family of TS surface antigens and is abundantly expressed only in the intracellular forms (amastigotes) of *T. cruzi* [39]. Fig. 1A and

Table 1. Peptides used in the study.

Peptide	AA positions	Predicted H-2 restriction
<u>TEWETGQI</u>	320–327	K ^k
<u>PETLGHEI</u>	650–657	K ^k
<u>YEIVAGYI</u>	140–147	K ^k
<u>HEHNLFGI</u>	130–137	K ^k
<u>AESWPSIV</u>	121–128	K ^k
<u>TPTAGLVGF</u>	488–496	L ^d
<u>RPNMSRHLF</u>	36–44	L ^d
<u>GSRNGNDRL</u>	172–179	L ^d
<u>ESKSGDAPL</u>	69–77	L ^d
<u>ESEPKRPNM</u>	31–39	L ^d
<u>ESSTPTAGL</u>	485–493	L ^d
<u>VSWGEPKSL</u>	239–247	L ^d
<u>YSDGALHLL</u>	438–446	L ^d

Peptides were selected by the scores determined by programs available at the sites: <http://www.syfpeithi.de/> and http://www.bimas.cit.nih.gov/molbio/hla_bind/. Putative anchor residues are in bold and underlined. doi:10.1371/journal.pone.0022011.t001




Figure 2. CD8 T-cell epitope identification during immune responses of H-2^a mice infected with *T. cruzi* or genetically vaccinated with *asp-2* gene. B10.A mice were infected with 10⁴ *T. cruzi* blood parasites. A/S mice were immunized with plgSPCl.9 followed by AdASP-2 and injected i.m. at 0 and 3 weeks, respectively. Control mice were naïve or injected with pcDNA3 followed by pcDNA3/Adβ-gal. **A**- Two weeks after infection or the final immunizing dose, splenic cells were re-stimulated *in vitro* in the presence of medium or the indicated peptides at a final concentration of 10 μM. The number of splenic IFN-γ spot-forming cells (SFC) was estimated by *ex vivo* ELISPOT assay. **B**-

in vivo cytotoxic activity was estimated by injecting each mouse with syngeneic CFSE-labeled splenic cells coated with or without 2 μM of the indicated peptide. Results are expressed as means ± SD of 4 mice per group and are representative of experiments performed at least twice with similar results. Asterisks denote that the number of SFC or *in vivo* cytotoxicity were significantly higher when compared to naïve or pcDNA3/Adβ-gal injected mice (P<0.01, one-way ANOVA). doi:10.1371/journal.pone.0022011.g002

Fig. 1B show, respectively, some of the structural features of ASP-2 antigen and its expression by amastigotes.

To test whether the H-2K^k-restricted epitope TEWETGQI could be an immunodominant epitope, a series of synthetic peptides containing the predicted AA anchor motif for binding to the H-2K^k or H-2L^d alleles of mouse MHC haplotype H-2^a was employed. No epitope was predicted to bind to H-2D^d (Table 1). After infection of H-2^a mice, antigen-specific IFN-γ producing cells could only be detected in the presence of the peptide TEWETGQI (Fig. 2A). We concluded that this epitope was the immunodominant epitope of ASP-2 during infection of H-2^a mice with the Y strain of *T. cruzi*. H-2^a mice used in this experiment were B10.A because they are resistant to infection with *T. cruzi*.

We showed previously that the strong pattern of immunodominance observed following infection with *T. cruzi* was not duplicated in mice genetically immunized with a recombinant adenovirus expressing ASP-2 (AdASP-2, ref. 15). Then, we determined whether genetically immunized H-2^a mice could present a different pattern of immunodominance. We used a genetic immunization approach which consisted of a heterologous prime-boost regimen using plasmid DNA followed by a recombinant adenovirus both containing the same *asp-2* gene. This protocol provided strong and long lasting protective immunity against experimental infection mediated by CD8⁺ T cells [37,38]. In these experiments, we used H-2^a mice of the A/Sn strain. These mice are highly susceptible to infection with *T. cruzi*, allowing us to perform protective immunity studies [37,38]. Nevertheless, it is important to mention that the results were similar when we used B10.A mice.

After *ex vivo* stimulation with our synthetic peptides, as expected, IFN-γ producing cells were detected following stimulation with the TEWETGQI peptide. In addition to this epitope, two other epitopes (PETLGHEI and YEIVAGYI) induced IFN-γ production by immune cells (Fig. 2A). These peptide-specific IFN-γ producing cells were CD8⁺ T cells as determined by simultaneous staining of intra-cellular IFN-γ and the surface marker CD8 (see below).

These peptides were recognized by cytotoxic cells in H-2^a mice as determined by *in vivo* cytotoxicity assays using target cells coated with each of these peptides. B10.A mice infected with *T. cruzi* developed strong *in vivo* cytotoxicity against target cells coated with the peptide TEWETGQI. In contrast, very limited (if any) *in vivo* cytotoxicity was observed against target cells coated with peptides PETLGHEI and YEIVAGYI. These results were not due to different kinetics of the immune response because we observed the same results 14 or 28 days after an infectious challenge (Fig. 2B).

However, A/Sn (H-2^a) mice genetically vaccinated with a heterologous prime-boost vaccination regimen displayed easily detectable *in vivo* cytotoxic activity against target cells coated with any of these three peptides. The elimination of target cells coated with peptide TEWETGQI was always stronger than the two others, suggesting a pattern of immunodominance.

To determine whether other cytokines and/or effector molecules could be secreted by peptide-specific T cells, we performed staining to detect surface mobilization of CD107a (a marker for exocytosis) or intra-cellular accumulation of IFN-γ, TNF-α, IL-2




Figure 3. Surface mobilization of CD107a and expression of IFN-γ, TNF-α, IL-2 or IL-10 by specific CD8⁺ T cells from B10.A mice infected with *T. cruzi* or A/Sn mice immunized with plgSPCI.9/AdASP-2 vaccine. B10.A or A/Sn mice were infected or immunized as described in the legend of Fig. 2. Control mice were either naive mice or mice immunized with pcDNA3/Adβ-gal. Twenty one or fourteen days after infection or immunization, respectively, these mice had their splenic cells cultured in the presence of anti-CD107a and anti-CD28, with or without the peptides TEWETGQI, PETLGHEI or YEIVAGYI. After 12 h, cells were stained for CD8, IFN-γ, TNF-α, IL-2 and IL-10. Representative analyses (medians) are shown from four mice performed per experiment. **A**) Example of splenic CD8⁺ cells from B10.A naive mice cultivated *in vitro* in the presence of medium alone (Medium) or with the indicated peptides and stained for expression of IFN-γ and TNF-α. **B, C and D**) Examples of splenic CD8⁺ cells from B10.A infected mice cultivated *in vitro* in the presence of medium alone (Medium) or with the indicated peptides and stained for expression of: B) IFN-γ and TNF-α; C) IFN-γ and IL-2; D) IFN-γ and IL-10. **E and F**) Examples of splenic CD8⁺ cells from mice immunized with pcDNA3/Adβ-gal cultivated *in vitro* in the presence of medium alone (Medium) or with the indicated peptides and stained for surface mobilization of CD107a and expression IFN-γ (panel E) or expression of IFN-γ and TNF-α (panel F). **G and H**) Examples of splenic CD8⁺ cells from mice immunized with plgSPCI.9/AdASP-2 cultivated *in vitro* in the presence of medium alone (Medium) or with the indicated peptides and stained for surface mobilization of CD107a and expression IFN-γ (panel G) or expression of IFN-γ and TNF-α (panel H). **J**) Bar graph showing CD8 mobilization under different conditions. Legend: TEWETGQI (black), PETLGHEI (red), YEIVAGYI (green).

in vitro in the presence of medium alone (Medium) or with the indicated peptides and stained for surface mobilization of CD107a and expression IFN- γ (panel G) and expression IFN- γ and TNF- α (panel H). **I and J**) Determination of multifunctional CD8⁺ cells from mice immunized with pIgSPCL9/AdASP-2 cultivated *in vitro* in the presence of the indicated peptides and stained for surface mobilization of CD107a and expression IFN- γ and TNF- α . doi:10.1371/journal.pone.0022011.g003

or IL-10. When we used splenic cells from B10.A mice infected with *T. cruzi*, we observed that upon stimulation with TEWETGQI, a large fraction of CD8⁺ cells mobilize CD107a to the surface (data not shown) and accumulate intra-cellular IFN- γ and TNF- α (Fig. 3B). These CD8⁺ cells were therefore multifunctional CD107a⁺IFN- γ ⁺TNF- α ⁺ (~35%) or IFN- γ ⁺TNF- α ⁺ (~40%). We were unable to detect the presence of significant numbers of IL-2 or IL-10 expressing cells in these same samples (Fig. 3C and 3D). The expression of these cytokines was dependent on the infection because they were not detected in cells from naive mice (Fig. 3A).

In contrast, a relatively low frequency of CD8⁺ splenic cells stimulated with peptides PETLGHEI or YEIVAGYI mobilized CD107a to the surface (data not shown) or accumulated intra-cellular IFN- γ or TNF- α (Fig. 3B). We were also unable to detect the presence of significant numbers of IL-2 or IL-10 expressing cells in these same cells (Fig. 3C and 3D).

Splenic cells from A/Sn mice genetically vaccinated with heterologous prime-boost regimen were stimulated with peptides TEWETGQI, PETLGHEI or YEIVAGYI. The results shows that a large fraction of the CD8⁺ cells of pIgSPCL9/AdASP-2 immunized mice at the same time mobilize CD107a to the surface and expressed intra-cellular IFN- γ and TNF- α (Fig. 3G to 3J). These cells were therefore multifunctional CD8⁺ T cells as we have previously described [37]. We were unable to detect the presence of intra-cellular IL-2 or IL-10 in these same cells (data not shown). These results confirmed and extended the ones described in Fig. 2).

The description of these two new epitopes allowed us to test whether immunity to the subdominant/cryptic epitope could participate during protective immunity against *T. cruzi* infection, a phenomenon that has not previously been tested experimentally. For this purpose, we generated a plasmid DNA and a recombinant adenovirus containing a mutated form of the *asp-2* gene in which we modified the nucleotides encoding the anchor residues required for the immunodominant TEWETGQI epitope to bind to the H-2K^k molecule. The mutated gene expressed the AA sequence TAWETGQA, where the alanines (A) replaced glutamic acid (E) or isoleucine (I) of the original epitope. In preliminary experiments, we observed that the synthetic peptide TAWETGQA was not recognized by immune cells from genetically vaccinated H-2^a mice (data not shown). Details of the plasmid and recombinant adenovirus containing the mutated form of the *asp-2* gene are shown in Table 2.

Initially, we genetically immunized mice with plasmids containing the original gene (pIgSPCL9), the mutated gene (pIgSPTAWETGQA) or both plasmids simultaneously. Immune responses were estimated by ELISPOT after *ex vivo* stimulation

with the immunodominant epitope TEWETGQI or with the subdominant/cryptic epitopes PETLGHEI or YEIVAGYI two weeks after challenge with *T. cruzi*. We chose this protocol because, as the immune response following plasmid DNA vaccination is usually low, it is easier to detect the anamnestic immune responses after challenge [12]. We observed that all mice immunized with *asp-2* genes (mutated or not) presented specific IFN- γ producing cells when stimulated with subdominant/cryptic epitopes (PETLGHEI or YEIVAGYI, Fig. 4A). The immune response was specific because mice immunized with control plasmid pcDNA3 failed to recognize these peptides. However, we detected IFN- γ producing cells specific to TEWETGQI only in mice immunized with plasmid pIgSPCL9. It is noteworthy that the number of cells detected in mice immunized with pIgSPTAWETGQA was similar to the number of cells in pcDNA3 injected mice (Fig. 4A). These numbers reflect cells primed during infection. Analysis of *in vivo* cytotoxic activity also demonstrated that in mice immunized with the plasmid pIgSPTAWETGQA, the response to the immunodominant epitope TEWETGQI was not different from control mice immunized with pcDNA3 (Fig. 4B). These immunological analyses demonstrated that mice immunized with pIgSPTAWETGQA indeed lost the functional immunodominant TEWETGQI response but had an unaltered ability to elicit immune responses to the subdominant epitopes PETLGHEI and YEIVAGYI.

After challenge, mice immunized with plasmids containing the *asp-2* gene (mutated or not) presented significantly lower parasitemia than control mice immunized with pcDNA3 (Fig. 4C). Although the levels of parasitemia were not statistically different when compared to mice vaccinated with the plasmid containing the mutated *asp-2* gene, the mortality of these animals was significantly faster (Fig. 4D). We therefore concluded that broadening the CD8⁺ T cell immune response by vaccination with a plasmid containing epitopes that elicit immune responses to subdominant/cryptic epitopes of ASP-2 could provide some degree of protective immunity. Nevertheless, because protective immunity elicited by vaccination with pIgSPTAWETGQA was not as efficient, the presence of a functional immunodominant epitope was clearly important for effective protective immunity.

We then sought to test the same hypotheses described above using a distinct approach. For that purpose, H-2^a mice were primed with plasmid pIgSPCL9 followed by a booster immunization with AdASP-2 (heterologous prime-boost regimen). Alternatively, mice were primed with plasmid pIgTAWETGQA followed by a booster immunization with AdTAWETGQA. We consider this approach complementary to the one described above because plasmid or adenovirus may used distinct routes for stimulating CD8⁺ T cells.

Immune responses were estimated 14 days after the booster immunization by ELISPOT following *ex vivo* stimulation with synthetic peptides encoding the immunodominant epitope TEWETGQI, the subdominant/cryptic epitopes PETLGHEI or YEIVAGYI or the mutated epitope TAWETGQA. We chose this protocol because the immune responses following heterologous prime boost immunization generates strong immune responses that can be easily detected after boosting [37]. We observed that mice immunized with *asp-2* genes (mutated or not) had specific IFN- γ producing cells when stimulated with peptides PELTHGEI or YEIVAGYI. The fact that these responses were of similar magnitude strongly argued that the expression/immunogenicity of

Table 2. Genetic vectors used in the study.

Designation	Vector	ASP-2 CD8 epitopes		
pIgSPCL9	Plasmid	TEWETGQI	PETLGHEI	YEIVAGYI
pIgSPTAWETGQA.9	Plasmid	TAWETGQA	PETLGHEI	YEIVAGYI
AdASP-2	Adenovirus	TEWETGQI	PETLGHEI	YEIVAGYI
AdTAWETGQA	Adenovirus	TAWETGQA	PETLGHEI	YEIVAGYI

doi:10.1371/journal.pone.0022011.t002




Figure 4. CD8 immune responses and trypanosome-induced parasitemia and mortality in A/Sn mice genetically immunized with different plasmid DNA containing the *asp-2* gene. A/Sn mice were immunized with pcDNA3, plgSPCI.9, plgSPTAWETGQA or simultaneously with the last two (plgSPCI.9 and plgSPTAWETGQA). Immunization consisted of 3 doses of 100 μ g of DNA each given by the i.m. route in the *tibialis anterioris* three weeks apart. A- Two weeks after the final immunizing dose, mice were challenged i.p. with 150 bloodstream trypanosomes. Two weeks after infection, splenic cells were re-stimulated *in vitro* in the presence of medium only or the indicated peptides at a final concentration of 10 μ M. The number of splenic IFN- γ spot-forming cells (SFC) was estimated by *ex vivo* ELISPOT assay. B- *In vivo* cytotoxic activity was estimated by injecting each mouse with syngeneic CFSE-labeled splenic cells coated with or without 2 μ M of the indicated peptide. Results are expressed as means \pm SD of 4 mice per group and are representative of experiments performed at least twice with similar results. Asterisks denote that the number of SFC, or the *in vivo* cytotoxicity, were significantly higher when compared to SFC found in naive or pcDNA3/Ad β -gal-injected mice ($P < 0.01$). C- Parasitemia for each mouse group is represented as mean \pm SD ($n = 5-7$). Asterisks denote that mice from groups immunized with plgSPCI.9 or plgSPTAWETGQA or both had significantly lower parasitemia ($P < 0.01$) than animals injected with pcDNA3. The curves of parasitemia of animals immunized with plgSPCI.9 (squares) or plgSPTAWETGQA (triangles) are superimposed. D- Kaplan-Meier curves for the survival of mouse groups immunized and challenged as described above ($n = 5-7$). Mice from groups immunized with plgSPCI.9 or plgSPCI.9/plgSPTAWETGQA survived significantly longer than animals injected with pcDNA3 or plgSPTAWETGQA ($P < 0.05$, in all cases, LogRank test). Mice immunized with plgSPTAWETGQA also survived longer than animals injected with pcDNA3 ($P < 0.05$). No animals died after the 30th day until they were euthanized. Results are representative of two independent experiments. doi:10.1371/journal.pone.0022011.g004

both genes/antigens were very similar. We could detect IFN- γ producing cells specific for the TEWETGQI epitope only in mice immunized with plgSPCI.9/AdASP-2. In contrast, none of the immunized mice presented IFN- γ producing cells when stimulated with the TAWETGQA peptide (Fig. 5A). The analysis of *in vivo* cytotoxic activity showed a similar picture (Fig. 5B). We also

performed intra-cellular cytokine staining analysis for IFN- γ and TNF- α after *in vitro* peptide stimulation. As depicted in Fig. 5C, we detected CD8⁺ cells expressing IFN- γ and/or TNF- α specific for the TEWETGQI epitope only in mice immunized with plgSPCI.9/AdASP-2 (Fig. 5C). Similar analyses performed 14 days after an infectious challenge with *T. cruzi* exhibited the same




Figure 5. CD8 immune responses in A/Sn mice immunized with *asp-2* using the heterologous DNA prime-adenovirus boost vaccination regimen. A/Sn mice were primed i.m. with 100 μ g of plasmids pcDNA3, pIgSPCI.9 or pIgSPTAWETGQA. Three weeks later, these mice were boosted i.m. with 2×10^8 pfu Ad β -gal, AdASP-2 or AdTAWETGQA. A—Two weeks after the last dose, splenic cells were re-stimulated *in vitro* in the presence of medium only or the indicated peptides at a final concentration of 10 μ M. The number of splenic IFN- γ spot forming cells (SFC) was estimated by *ex vivo* ELISPOT assay. B—*In vivo* cytotoxic activity was estimated by injecting each mouse with syngeneic CFSE-labeled splenic cells coated with or without 2 μ M of the indicated peptide. Results are expressed as mean \pm SD of 4 mice per group and are representative of experiments performed at least twice with similar results. Asterisks denote that the number of SFC or *in vivo* cytotoxicity were significantly higher when compared to SFC found in naive or pcDNA3/Ad β -gal injected mice ($P < 0.01$). C—Fourteen days after the last dose, these mice had their splenic cells cultured in the presence of anti-CD28 and Medium or the indicated peptides. After 12 h, cells were stained for CD8, IFN- γ and TNF- α . Examples of splenic CD8⁺ cells from immunized mice. Representative analyses (medians) are shown from four mice performed per experiment. doi:10.1371/journal.pone.0022011.g005

pattern of response (data not shown). Together, these immunological analyses demonstrated that heterologous prime-boost immunization with pIgSPTAWETGQA/AdTAWETGQA failed to induce an immune response to the immunodominant epitope TEWETGQI but induced almost unaltered immune responses to the subdominant epitopes PETLGHEI or YEIVAGYI.

After challenge, parasitemia in mice immunized with the *asp-2* genes (mutated or not) was significantly lower than in control

mice injected with pcDNA3/Ad β -gal ($P < 0.01$, Fig. 6A). Although most of the mice immunized with pIgSPTAWETGQA/AdTAWETGQA died after challenge, they survived longer than control mice injected with pcDNA3/Ad β -gal (Fig. 6B, $P < 0.01$, LogRank test). In parallel, vaccinated and control mice were challenged by the s.c. route. We observed that the majority of the mice immunized with pIgSPTAWETGQA/AdTAWETGQA survived the infectious challenge (Fig. 6D). We therefore conclu-




Figure 6. Trypomastigote-induced parasitemia and mortality in A/Sn mice immunized with *asp-2* using the heterologous DNA prime-adenovirus boost vaccination regimen. A/Sn mice were immunized as depicted in the legend of Fig. 5. Two weeks after the final immunizing dose, mice were challenged i.p. (Panels A and B) or s.c. (Panels C and D) with 150 bloodstream trypomastigotes. Parasitemia for each mouse group is represented as mean \pm SD (n=10 or 11). Asterisks denote that mice from groups immunized with pIgSPC1.9/AdASP-2 or pIgSPTAWETGQA/AdTAWETGQA had significantly lower parasitemia ($P < 0.01$) than pcDNA3/Adβ-gal-injected animals. Panels B and D represent Kaplan-Meier curves for survival of the mouse groups immunized and challenged as described above (n=10 or 11). Mice immunized with pIgSPC1.9/AdASP-2 survived significantly longer than animals immunized with pIgSPTAWETGQA/AdTAWETGQA or pcDNA3/Adβ-gal ($P = 0.01$ or $P < 0.01$, respectively). Mice immunized with pIgSPTAWETGQA/AdTAWETGQA survived significantly longer than pcDNA3/Adβ-gal-injected animals ($P < 0.01$). Results are representative of two pooled experiments. No animals died after the 40th day. doi:10.1371/journal.pone.0022011.g006

ded that broadening the T cell immune response using a genetic vaccination that elicits immune responses to subdominant/cryptic epitopes of ASP-2 provides protective immunity against infection. Nevertheless, likewise in the case of the plasmid DNA, after an i.p. challenge, in terms of survival, immunization with only genes expressing subdominant epitopes was not as effective as immunization with genes expressing both the dominant and the subdominant epitopes.

Finally, to firmly establish that protective immunity was mediated by CD8⁺ T cells, we performed *in vivo* depletion experiments in mice vaccinated with the pIgSPC1.9/AdASP-2 or pIgSPTAWETGQA/AdTAWETGQA. Treatment with anti-CD8 MAb renders these mice more susceptible to infection. CD8 depleted mice presented higher parasitemia (Fig. 7A and B) and shorter survival times (Fig. 7C and D) when compared to A/Sn mice vaccinated with heterologous prime-boost regimen

(pIgSPC1.9/AdASP-2 or pIgSPTAWETGQA/AdTAWETGQA) and treated with rat IgG. CD8 depleted mice had their survival time reduced to the same time as control mice which were injected with pcDNA3/Adβ-gal.

Discussion

Here, we initially confirmed and extended our previous observation that experimental infection with the human intracellular pathogen *T. cruzi* restricted the repertoire of CD8⁺ T cells. While immune cells of infected H-2^a mice recognized a single immunodominant epitope of ASP-2, cells from mice immunized with recombinant genetic vaccines expressing this same *T. cruzi* antigen recognized, in addition to the immunodominant epitope, two other subdominant/cryptic epitopes. The sub-dominant epitopes not only failed to elicit IFN-γ and *in vivo* cytotoxicity




Figure 7. CD8 T cell dependence of protective immunity of A/Sn mice immunized with *asp-2* using the heterologous DNA prime-adenovirus boost vaccination regimen. A/Sn mice were immunized as described in the legend of Fig. 5. plgSPTAWETGQA/AdTAWETGQA had significantly lower parasitemia ($P < 0.01$) than pcDNA3/Adβ-gal-injected animals. Before and after challenge, mice were treated as described in Methods section with rat IgG (control) or anti-CD8 MAb. The parasitemia for each mouse group is represented as mean \pm SD ($n = 6$). Asterisks denote that mice from groups immunized with plgSPcI.9/AdASP-2 or plgSPTAWETGQA/AdTAWETGQA and treated with Rat IgG had significantly lower parasitemia ($P < 0.01$) than vaccinated mice treated with anti-CD8 (Panels A and B). Panels C and D represent Kaplan-Meier curves for survival of the mouse groups immunized and challenged as described above ($n = 6$). Mice immunized with plgSPcI.9/AdASP-2 or plgSPTAWETGQA/AdTAWETGQA and treated with Rat IgG survived significantly longer than vaccinated animals treated with anti-CD8 ($P < 0.01$ in both cases). doi:10.1371/journal.pone.0022011.g007

during infection (Fig. 2), but they also did not stimulate TNF- α , IL-2 or IL-10 secretion by CD8⁺ T cells (Fig. 3). The precise reason for this strong immunodominant pattern during *T. cruzi* infection is unknown at present. One possible explanation for this biased immune response could be to provide an advantage to the parasite by avoiding an even higher and broader immune response. This hypothesis is in agreement with earlier studies showing that immunity to subdominant epitopes can provide an important contribution to protective immunity against viral infection [34,40–43]. Our results also corroborated this hypothesis. Immunity to epitopes that are not commonly recognized during infection (cryptic) provided a significant degree of protective immunity. Nonetheless, immune responses against these cryptic epitopes did

not substitute completely for CD8⁺ T cells specific for the immunodominant TEWETGQI epitope. We observed that immunization with plasmids alone or in combination with a recombinant adenovirus that did not express the immunodominant epitope elicited immune responses to the subdominant/cryptic epitopes but failed to provide optimal protective immunity when compared to a plasmid that expresses both the immunodominant and subdominant epitopes (Fig. 4D, 6B and 7D). These results demonstrate that the response to the immunodominant epitope contributes to the immunity elicited by genetic vaccination and is might be required for highly efficient resistance.

In previous studies, we showed that immunization with short proteins in the presence of the TLR9 agonist CpG elicited CD8⁺

T-cell mediated immunity against *T. cruzi* infection in A/Sn mice [44]. These short proteins contained only the AA 261 to 500 or 261 to 380. In both cases, they did not express the subdominant/cryptic epitopes. Based on that, we concluded that immunization with the immunodominant CD8 epitope alone could provide a high degree of protective immunity even in the absence of the subdominant epitopes [44]. The fact that the immune response directed solely to the immunodominant epitope can provide significant degree of protective immunity against protozoan parasites has been established a long time ago by the use of adoptively transferred T cell clones or heterologous-prime boost vaccination regimen [45–49].

In earlier studies in which we depleted CD8⁺ T cells from genetically vaccinated mice, we observed that these mice were unable to control parasitemia and died at the same time as control unvaccinated animals [37]. Therefore, although genetic immunization elicits effector CD4⁺ T cells, these cells do not account for the protection we observed. This concept was further corroborated with experiments of CD8 T cell-depletion performed here (Fig. 7A to D).

Using a different approach, a similar conclusion was also reached by Rosenberg *et al.*, 2010 [17]. In their study, they induced simultaneous tolerance to two immunodominant *T. cruzi* epitopes in a resistant mouse strain. Following infection, an increased susceptibility to infection was observed. Nevertheless, they were still able to control and survive the experimental infection. This protective immunity was possibly mediated by CD8⁺ T cells specific to subdominant epitopes that substituted for the immunodominant ones. The AA sequences of these subdominant/cryptic epitopes have yet to be identified. Together with our study, they strongly support the notion that the immune responses to both dominant and subdominant/cryptic epitopes can be important for controlling experimental *T. cruzi* infection in inbred mouse strains. These results are in agreement with the observation with other parasites such as *Plasmodium*. Tolerance to or removal of the immunodominant CD8 epitope of *P. yoelii* led to the development of immunity to CD8 subdominant epitopes as well [50,51].

The mechanism operating during *T. cruzi* infection to restrict the immune response leading to immunodominance has yet to be characterized. We provide initial evidence that it could be explained by T cell competition for APCs by showing that in mice infected simultaneously with two different parasite strains containing different immunodominant epitopes, we could generate maximal responses to both epitopes without immunodominance or competition [15]. Our interpretation was that if the epitopes are presented by different APCs, then the immunodominant pattern is

disrupted. However, a more formal demonstration using bone marrow chimeric mice studies is lacking. At the molecular level, this strong immunodominance can be explained by the type of antigen presentation that predominates during *T. cruzi* infection. In recent studies, important evidence has been provided that subdominant epitopes can only be directly presented by the expressing cells, which might occur in the case of the recombinant adenovirus. However, during indirect priming (cross-priming), these epitopes would be at a disadvantage [52]. *T. cruzi* may use cross-priming as the dominant route, drastically reducing the priming of subdominant epitopes.

In addition to shedding some light on the host-parasite relationship, our results may have important implications for the development of T cell vaccines against parasitic diseases. In our earlier studies, we observed that genetic vaccination with a heterologous prime-boost regimen employing plasmid DNA and recombinant adenovirus elicited strong, long lasting CD8⁺ T cell-mediated protective immunity against experimental infection in a mouse strain highly susceptible to *T. cruzi* infection [37,38]. Here, we demonstrated that the CD8 T cell-mediated protective immunity observed was directed to three distinct epitopes, two of which are cryptic. The strategy of redirecting immunity to epitopes that are not usually targets of the naturally acquired immune response has been proposed as a possible means to improve immunity against viral infection [40–43]. Recently, this strategy was also proven useful to improve vaccination against *T. cruzi* infection [53].

Finally, our observations may have important implications regarding the basis for the strong immunodominance pattern observed after experimental infection with other intracellular parasites such as *Plasmodium*, *Toxoplasma gondii* and *Theileria parva* [50,51,54–57].

Acknowledgments

The authors are in debt with Dr. C. Claser for providing the immunofluorescence picture of Fig. 1. We are also in debt with Drs. Laurent Renia (Singapore Immunology Network), Dr. Chris Ibegbu (Emory Vaccine Center), Dr. Fanny Tzelepis (University of Ottawa) and Dr. Silvia B. Boscardin, for careful reviewing the manuscript.

Author Contributions

Conceived and designed the experiments: MRD ELVS JRCdV BCGdA MMR. Performed the experiments: MRD ELVS JRCdV BCGdA. Analyzed the data: MRD ELVS AVM OB-R RTG MMR. Contributed reagents/materials/analysis tools: AVM OB-R RTG. Wrote the paper: MMR.

References

1. Tarleton RL (2007) Immune system recognition of *Trypanosoma cruzi*. *Curr Opin Immunol* 19: 430–4.
2. Miyahira Y (2008) *Trypanosoma cruzi* infection from the view of CD8⁺ T cell immunity—an infection model for developing T cell vaccine. *Parasitol Int* 57: 38–48.
3. Padilla AM, Bustamante JM, Tarleton RL (2009) CD8⁺ T cells in *Trypanosoma cruzi* infection. *Curr Opin Immunol* 21: 385–90.
4. Boscardin SB, Torrecilhas AC, Manarin R, Revelli S, Rey EG, et al. (2010) Chagas' disease: an update on immune mechanisms and therapeutic strategies. *J Cell Mol Med* 14: 1373–84.
5. Junqueira C, Caetano B, Bartholomeu DC, Melo MB, Ropert C, et al. (2010) The endless race between *Trypanosoma cruzi* and host immunity: lessons for and beyond Chagas disease. *Expert Rev Mol Med* 12: e29.
6. Kierszenbaum F (2005) Where do we stand on the autoimmunity hypothesis of Chagas disease? *Trends Parasitol* 21: 513–6.
7. Marin-Neto JA, Cunha-Neto E, Maciel BC, Simões MV (2008) Pathogenesis of chronic Chagas heart disease. *Circulation* 115: 1109–23.
8. Bonney KM, Engman DM (2008) Chagas heart disease pathogenesis: one mechanism or many? *Curr Mol Med* 8: 510–8.
9. Gutierrez FR, Guedes PM, Gazzinelli RT, Silva JS (2009) The role of parasite persistence in pathogenesis of Chagas heart disease. *Parasite Immunol* 31: 673–85.
10. Scharfstein J, Gomes Jde A, Correa-Oliveira R (2009) Mem Inst Oswaldo Cruz. Back to the future in Chagas disease: from animal models to patient cohort studies, progress in immunopathogenesis research 104(Suppl 1): 187–98.
11. Lannes-Vieira J, Silverio JC, Pereira IR, Vinagre NF, Carvalho CM, et al. (2009) Chronic *Trypanosoma cruzi*-elicited cardiomyopathy: from the discovery to the proposal of rational therapeutic interventions targeting cell adhesion molecules and chemokine receptors—how to make a dream come true. *Mem Inst Oswaldo Cruz* 104(Suppl 1): 226–35.
12. Tzelepis F, de Alencar BC, Penido ML, Gazzinelli RT, Persechini PM, et al. (2006) Distinct kinetics of effector CD8⁺ cytotoxic T cells after infection with *Trypanosoma cruzi* in naive or vaccinated mice. *Infect Immun* 74: 2477–81.
13. Martin DL, Weatherly DB, Laucella SA, Cabinian MA, Crim MT, et al. (2006) CD8⁺ T-Cell responses to *Trypanosoma cruzi* are highly focused on strain-variant trans-sialidase epitopes. *PLoS Pathog* 2(8): e77.
14. Tzelepis F, Persechini PM, Rodrigues MM (2007) Modulation of CD4⁺ T cell-dependent specific cytotoxic CD8⁺ T cells differentiation and proliferation by the timing of increase in the pathogen load. *PLoS One* 2(4): e393.
15. Tzelepis F, de Alencar BC, Penido ML, Claser C, Machado AV, et al. (2008) Infection with *Trypanosoma cruzi* restricts the repertoire of parasite-specific CD8⁺ T cells leading to immunodominance. *J Immunol* 180: 1737–48.
16. Bixby LM, Tarleton RL (2008) Stable CD8⁺ T cell memory during persistent *Trypanosoma cruzi* infection. *J Immunol* 181: 2644–50.

17. Rosenberg CS, Martin DL, Tarleton RL (2010) CD8+ T cells specific for immunodominant trans-sialidase epitopes contribute to control of *Trypanosoma cruzi* infection but are not required for resistance. *J Immunol* 185: 560–8.
18. Freire-de-Lima L, Alisson-Silva F, Carvalho ST, Takiya CM, Rodrigues MM, et al. (2010) *Trypanosoma cruzi* subverts host cell sialylation and may compromise antigen-specific CD8+ T cell responses. *J Biol Chem* 285: 13388–96.
19. Oliveira AC, de Alencar BC, Tzelepis F, Klezewsky W, da Silva RN, et al. (2010) Impaired innate immunity in Tlr4(-/-) mice but preserved CD8+ T cell responses against *Trypanosoma cruzi* in Tlr4-, Tlr2-, Tlr9- or Myd88-deficient mice 29; 6(4): e1000870.
20. Rodrigues MM, de Alencar BC, Claser C, Tzelepis F (2009) Immunodominance: a new hypothesis to explain parasite escape and host/parasite equilibrium leading to the chronic phase of Chagas' disease? *Braz J Med Biol Res* 42: 220–3.
21. Chen W, Antón LC, Bennink JR, Yewdell JW (2000) Dissecting the multifactorial causes of immunodominance in class I-restricted T cell responses to viruses. *Immunity* 12: 83–93.
22. Tenzer S, Wee E, Burgevin A, Stewart-Jones G, Friis L, et al. (2009) Antigen processing influences HIV-specific cytotoxic T lymphocyte immunodominance. *Nat Immunol* 10: 636–46.
23. Yewdell JW (2006) Confronting complexity: real-world immunodominance in antiviral CD8+ T cell responses. *Immunity* 25: 533–43.
24. Irvine K, Bennink J (2006) Factors influencing immunodominance hierarchies in TCD8+ -mediated antiviral responses. *Expert Rev Clin Immunol* 2: 135–47.
25. La Gruta NL, Kedzierska K, Pang K, Webby R, Davenport M, et al. (2006) A virus-specific CD8+ T cell immunodominance hierarchy determined by antigen dose and precursor frequencies. *Proc Natl Acad Sci USA* 103: 994–9.
26. Kotturi MF, Scott I, Wolfe T, Peters B, Sidney J, et al. (2008) Naive precursor frequencies and MHC binding rather than the degree of epitope diversity shape CD8+ T cell immunodominance. *J Immunol* 181: 2124–33.
27. Sacha JB, Reynolds MR, Buechler MB, Chung C, Jonas AK, et al. (2008) Differential antigen presentation kinetics of CD8+ T-cell epitopes derived from the same viral protein. *J Virol* 82: 9293–8.
28. La Gruta NL, Rothwell WT, Cukalac T, Swan NG, Valkenburg SA, et al. (2010) Primary CTL response magnitude in mice is determined by the extent of naive T cell recruitment and subsequent clonal expansion. *J Clin Invest* 120: 1885–94.
29. Chen W, McCluskey J (2009) Immunodominance and immunodomination: critical factors in developing effective CD8+ T-cell-based cancer vaccines. *Adv Cancer Res* 95: 203–47.
30. Jenkins MR, Webby R, Doherty PC, Turner SJ (2009) Addition of a prominent epitope affects influenza A virus-specific CD8+ T cell immunodominance hierarchies when antigen is limiting. *J Immunol* 177: 2917–25.
31. Willis RA, Kappler JW, Marrack PC (2006) CD8 T cell competition for dendritic cells *in vivo* is an early event in activation. *Proc Natl Acad Sci USA* 103: 12063–8.
32. Newberg MH, McEvers KJ, Gorgone DA, Lifton MA, Baumeister SH, et al. (2006) Immunodomination in the evolution of dominant epitope-specific CD8+ T lymphocyte responses in simian immunodeficiency virus-infected rhesus monkeys. *J Immunol* 176: 319–28.
33. Kastenmuller W, Gasteiger G, Gronau JH, Baier R, Ljapoci R, et al. (2007) Cross-competition of CD8+ T cells shapes the immunodominance hierarchy during boost vaccination. *J Exp Med* 204: 2187–98.
34. Riedl P, Wieland A, Lamberth K, Buus S, Lemonnier F, et al. (2009) Elimination of immunodominant epitopes from multispecific DNA-based vaccines allows induction of CD8 T cells that have a striking antiviral potential. *J Immunol* 183: 370–80.
35. Vasconcelos JR, Hiyane MI, Marinho CR, Claser C, Machado AM, et al. (2004) Protective immunity against *trypanosoma cruzi* infection in a highly susceptible mouse strain after vaccination with genes encoding the amastigote surface protein-2 and trans-sialidase. *Hum Gene Ther* 15: 878–86.
36. Machado AV, Cardoso JE, Claser C, Rodrigues MM, Gazzinelli RT, et al. (2006) Long-term protective immunity induced against *Trypanosoma cruzi* infection after vaccination with recombinant adenoviruses encoding amastigote surface protein-2 and trans-sialidase. *Hum Gene Ther* 17: 898–908.
37. de Alencar BC, Persechini PM, Haolla FA, de Oliveira G, Silverio JC, et al. (2009) Perforin and gamma interferon expression are required for CD4+ and CD8+ T-cell-dependent protective immunity against a human parasite, *Trypanosoma cruzi*, elicited by heterologous plasmid DNA prime-recombinant adenovirus 5 boost vaccination. *Infect Immun* 77: 4383–95.
38. Haolla FA, Claser C, de Alencar BC, Tzelepis F, de Vasconcelos JR, et al. (2009) Strain-specific protective immunity following vaccination against experimental *Trypanosoma cruzi* infection. Strain-specific protective immunity following vaccination against experimental *Trypanosoma cruzi* infection. *Vaccine* 27: 5644–53.
39. Claser C, Espindola NM, Sasso G, Vaz AJ, Boscardin SB, et al. (2007) Immunologically relevant strain polymorphism in the Amastigote Surface Protein 2 of *Trypanosoma cruzi*. *Microbes Infect* 9: 1011–9.
40. Oukka M, Manuguerra JC, Livaditis N, Tourdot S, Riche N, et al. (1996) Protection against lethal viral infection by vaccination with nonimmunodominant peptides. *J Immunol* 157: 3039–45.
41. Friedrich TC, Valentine LE, Yant IJ, Rakasz EG, Piaskowski SM, et al. (2007) Subdominant CD8+ T-cell responses are involved in durable control of AIDS virus replication. *J Virol* 81: 3465–76.
42. Holtappels R, Simon CO, Munks MW, Thomas D, Deegen P, et al. (2008) Subdominant CD8 T-cell epitopes account for protection against cytomegalovirus independent of immunodomination. *J Virol* 82: 5781–96.
43. Ruckwardt TJ, Luongo C, Malloy AM, Liu J, Chen M, et al. (2010) Responses against a subdominant CD8+ T cell epitope protect against immunopathology caused by a dominant epitope. *J Immunol* 185: 4673–80.
44. Araújo AF, de Alencar BC, Vasconcelos JR, Hiyane MI, Marinho CR, et al. (2005) CD8+ T-cell-dependent control of *Trypanosoma cruzi* infection in a highly susceptible mouse strain after immunization with recombinant proteins based on amastigote surface protein 2. *Infect Immun* 73: 6017–25.
45. Rodrigues M, Nussenzweig RS, Romero P, Zavala F (1992) The *in vivo* cytotoxic activity of CD8+ T cell clones correlates with their levels of expression of adhesion molecules. *J Exp Med* 175: 895–905.
46. Li S, Rodrigues M, Rodriguez D, Rodriguez JR, Esteban M, et al. (1993) Priming with recombinant influenza virus followed by administration of recombinant vaccinia virus induces CD8+ T-cell-mediated protective immunity against malaria. *Proc Natl Acad Sci USA* 90: 5214–18.
47. Rodrigues M, Li S, Murata K, Rodriguez D, Rodriguez JR, et al. (1994) Influenza and vaccinia viruses expressing malaria CD8+ T and B cell epitopes. Comparison of their immunogenicity and capacity to induce protective immunity. *J Immunol* 153: 4636–48.
48. Sedegah M, Jones TR, Kaur M, Hedstrom R, Hobart P, et al. (1998) Boosting with recombinant vaccinia increases immunogenicity and protective efficacy of malaria DNA vaccine. *Proc Natl Acad Sci USA* 95: 7648–53.
49. Schneider J, Gilbert SC, Blanchard TJ, Hanke T, Robson KJ, et al. (1998) Enhanced immunogenicity for CD8+ T cell induction and complete protective efficacy of malaria DNA vaccination by boosting with modified vaccinia virus Ankara. *Nat Med* 4: 397–402.
50. Kumar KA, Sano G, Boscardin S, Nussenzweig RS, Nussenzweig MC, et al. (2006) The circumsporozoite protein is an immunodominant protective antigen in irradiated sporozoites. *Nature* 444: 937–40.
51. Mauduit M, Grüner AC, Tewari R, Depinay N, Kayibanda M, et al. (2009) A role for immune responses against non-CS components in the cross-species protection induced by immunization with irradiated malaria sporozoites. *PLoS One* 4(11): e7717.
52. Pavelic V, Matter MS, Mumprecht S, Breyer I, Ochsenbein AF (2009) CTL induction by cross-priming is restricted to immunodominant epitopes. *Eur J Immunol* 39: 704–16.
53. Cazorla SI, Frank FM, Becker PD, Arnaiz M, Mirkin GA, et al. (2010) Redirection of the immune response to the functional catalytic domain of the cysteine proteinase cruzipain improves protective immunity against *Trypanosoma cruzi* infection. *J Infect Dis* 202: 136–44.
54. Blanchard N, Gonzalez F, Schaeffer M, Joncker NT, Cheng T, et al. (2008) Immunodominant, protective response to the parasite *Toxoplasma gondii* requires antigen processing in the endoplasmic reticulum. *Nat Immunol* 9: 937–44.
55. Frickel EM, Sahoo N, Hopp J, Gubbels MJ, Craver MP, et al. (2008) Parasite stage-specific recognition of endogenous *Toxoplasma gondii*-derived CD8+ T cell epitopes. *J Infect Dis* 198: 1625–33.
56. Wilson DC, Grotenbreg GM, Liu K, Zhao Y, Frickel EM, et al. (2010) Differential regulation of effector- and central-memory responses to *Toxoplasma gondii* infection by IL-12 revealed by tracking of Tgd057-specific CD8+ T cells. *PLoS Pathog* 6(3): e1000815.
57. MacHugh ND, Connelley T, Graham SP, Pelle R, Formisano P, et al. (2009) CD8+ T-cell responses to *Theileria parva* are preferentially directed to a single dominant antigen: Implications for parasite strain-specific immunity. *Eur J Immunol* 39: 2459–69.