

(2*R*)-2-Benzene sulfonamido-2-phenyl-ethanoic acid: a new monoclinic polymorph

Islam Ullah Khan,* Shahzad Sharif, Muhammad Nadeem Arshad, Ejaz and Muhammad Idrees

Materials Chemistry Laboratory, Department of Chemistry, GC University, Lahore, Pakistan

Correspondence e-mail: iukhan.gcu@gmail.com


Received 1 September 2009; accepted 7 September 2009

Key indicators: single-crystal X-ray study; $T = 296\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$; R factor = 0.042; wR factor = 0.099; data-to-parameter ratio = 17.0.

In the title compound, $\text{C}_{14}\text{H}_{13}\text{NO}_4\text{S}$, a sulfonamide derivative of phenyl glycine, the aromatic rings are inclined at a dihedral angle of $28.03(12)^\circ$. In the crystal, $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds link the molecules into chains propagating in [100] and a weak $\text{C}-\text{H}\cdots\text{O}$ interaction cross-links the chains in the *c*-axis direction. In the previously published polymorph, the dihedral angle between the aromatic rings is $45.52(18)^\circ$ and the structure is stabilized by three different types of ring motif.

Related literature

For related sulfonamide structures see: Arshad *et al.* (2008a,*b*, 2009).



Experimental

Crystal data

$\text{C}_{14}\text{H}_{13}\text{NO}_4\text{S}$	$c = 15.3642(15)\text{ \AA}$
$M_r = 291.31$	$\beta = 100.384(3)^\circ$
Monoclinic, $P2_1$	$V = 663.64(10)\text{ \AA}^3$
$a = 8.2464(8)\text{ \AA}$	$Z = 2$
$b = 5.3251(4)\text{ \AA}$	Mo $K\alpha$ radiation

$\mu = 0.26\text{ mm}^{-1}$
 $T = 296\text{ K}$

$0.34 \times 0.19 \times 0.11\text{ mm}$

Data collection

Bruker Kappa APEXII CCD diffractometer
Absorption correction: multi-scan (*SADABS*; Bruker, 2007)
 $T_{\min} = 0.918$, $T_{\max} = 0.972$

7864 measured reflections
3174 independent reflections
2372 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.034$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.042$
 $wR(F^2) = 0.099$
 $S = 1.01$
3174 reflections
187 parameters
1 restraint

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.24\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.26\text{ e \AA}^{-3}$
Absolute structure: Flack (1983), 1360 Friedel pairs
Flack parameter: -0.04 (9)

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O4—H4O \cdots O2 ⁱ	0.83 (4)	1.98 (4)	2.727 (3)	150 (5)
C4—H4A \cdots O3 ⁱⁱ	0.93	2.56	3.317 (4)	139

Symmetry codes: (i) $x + 1, y, z$; (ii) $-x + 2, y + \frac{1}{2}, -z + 1$.

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997) and *PLATON* (Spek, 2009); software used to prepare material for publication: *WinGX* (Farrugia, 1999) and *PLATON*.

MNA acknowledges the Higher Education Commission of Pakistan for providing a PhD Scholarship under the Indigenous 5000 PhD fellowship program (PIN 042-120607-PS2-183).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HG2562).

References

- Arshad, M. N., Tahir, M. N., Khan, I. U., Ahmad, E. & Shafiq, M. (2008a). *Acta Cryst. E64*, o2380.
Arshad, M. N., Tahir, M. N., Khan, I. U., Shafiq, M. & Ahmad, S. (2009). *Acta Cryst. E65*, o940.
Arshad, M. N., Tahir, M. N., Khan, I. U., Siddiqui, W. A. & Shafiq, M. (2008b). *Acta Cryst. E64*, o2045.
Bruker (2007). *SADABS*, *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
Flack, H. D. (1983). *Acta Cryst. A39*, 876–881.
Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
Spek, A. L. (2009). *Acta Cryst. D65*, 148–155.

supplementary materials

Acta Cryst. (2009). E65, o2436 [doi:10.1107/S160053680903606X]

(2*R*)-2-Benzenesulfonamido-2-phenylethanoic acid: a new monoclinic polymorph

I. U. Khan, S. Sharif, M. N. Arshad, Ejaz and M. Idrees

Comment

We have already reported the crystal structures of sulfonamides (Arshad *et al.*, 2008a, b), (Arshad *et al.*, 2009). In continuation of our studies in this area, we report here a new polymorph of our previously published sulfonamide (Arshad *et al.*, 2009a), derivative (II).

The title compound (I) crystallizes in the monoclinic space group P21. The molecule has a chiral center at C₁₃ with a slightly distorted tetrahedral geometry. The dihedral angles between the two aromatic ring are 28.03 (12)^o in (I) and 45.52 (18)^o in (II). The crystal structure of I has no complex intermolecular interactions like II. There are only two types of hydrogen bonding interaction of O—H—O making a polymeric chain along the a-axes and C—H—O which linked these polymeric chain along c-axes (Fig. 2 and Table 1).

Experimental

Phenyl glycine (2 g, 13.2 mmol) was dissolved in distilled water (15 ml) in a round bottom flask (50 ml). The pH of the solution was maintained at 8–9 using 1M, Na₂CO₃ solution. Benzene sulfonyl chloride (2.32 g, 13.2 mmol) was then suspended to the solution, and stirred at room temperature until all the suspension had been disappeared. On completion of the reaction the pH was adjusted 1–2, using 1 M HCl with stirring. The precipitate formed was filtered off, washed with distilled water, dried and recrystallized in methanol.

Refinement

The H atoms for the C atoms were refined geometrically and treated as riding atoms: C—H = 0.93 for aromatic and C—H = 0.98 for the chiral carbon with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}$. The N—H and O—H were refined in calculated positions and treated as riding atoms: O—H = 0.83 (4) Å, N—H = 0.82 (3) Å, with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}$ (parent O atom) and = 1.2 U_{eq} (parent N atom)

Figures




Fig. 1. The labelled molecular structure diagram of the title compound with the 50% probability level of drawn thermal ellipsoids.

supplementary materials




Fig. 2. Unit cell packing diagram showing the intermolecular hydrogen bonding using dashed lines. The hydrogen atoms not involved in hydrogen bonding have been omitted.

(2*R*)-2-Benzensulfonamido-2-phenylethanoic acid

Crystal data

C ₁₄ H ₁₃ NO ₄ S	$F_{000} = 304$
$M_r = 291.31$	$D_x = 1.458 \text{ Mg m}^{-3}$
Monoclinic, $P2_1$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: P 2yb	Cell parameters from 1972 reflections
$a = 8.2464 (8) \text{ \AA}$	$\theta = 2.5\text{--}23.4^\circ$
$b = 5.3251 (4) \text{ \AA}$	$\mu = 0.26 \text{ mm}^{-1}$
$c = 15.3642 (15) \text{ \AA}$	$T = 296 \text{ K}$
$\beta = 100.384 (3)^\circ$	Needle, white
$V = 663.64 (10) \text{ \AA}^3$	$0.34 \times 0.19 \times 0.11 \text{ mm}$
$Z = 2$	

Data collection

Bruker Kappa APEXII CCD diffractometer	3174 independent reflections
Radiation source: fine-focus sealed tube	2372 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.034$
$T = 296 \text{ K}$	$\theta_{\text{max}} = 28.3^\circ$
φ and ω scans	$\theta_{\text{min}} = 1.4^\circ$
Absorption correction: multi-scan (SADABS; Bruker, 2007)	$h = -10 \rightarrow 11$
$T_{\text{min}} = 0.918$, $T_{\text{max}} = 0.972$	$k = -7 \rightarrow 6$
7864 measured reflections	$l = -20 \rightarrow 20$

Refinement

Refinement on F^2	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H atoms treated by a mixture of independent and constrained refinement
$R[F^2 > 2\sigma(F^2)] = 0.042$	$w = 1/[s^2(F_o^2) + (0.0379P)^2 + 0.1295P]$ where $P = (F_o^2 + 2F_c^2)/3$
$wR(F^2) = 0.099$	$(\Delta/\sigma)_{\text{max}} < 0.001$
$S = 1.01$	$\Delta\rho_{\text{max}} = 0.24 \text{ e \AA}^{-3}$
3174 reflections	$\Delta\rho_{\text{min}} = -0.26 \text{ e \AA}^{-3}$
187 parameters	Extinction correction: none

1 restraint
 Primary atom site location: structure-invariant direct methods
 Secondary atom site location: difference Fourier map

Absolute structure: Flack (1983), 1360 Friedel pairs
 Flack parameter: -0.04 (9)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.53130 (7)	0.77328 (15)	0.29076 (4)	0.03965 (18)
O1	0.5315 (3)	1.0355 (4)	0.27187 (13)	0.0569 (6)
O2	0.3791 (2)	0.6402 (4)	0.28841 (13)	0.0560 (6)
O3	0.9213 (3)	0.3901 (5)	0.25951 (16)	0.0728 (7)
O4	1.0525 (2)	0.7469 (6)	0.23666 (16)	0.0700 (7)
H4O	1.137 (5)	0.660 (9)	0.249 (3)	0.105*
N1	0.6206 (3)	0.6352 (5)	0.21858 (15)	0.0396 (6)
H1N	0.615 (4)	0.482 (6)	0.223 (2)	0.047*
C1	0.6493 (3)	0.7278 (5)	0.39752 (16)	0.0334 (6)
C2	0.6187 (3)	0.5204 (5)	0.44644 (19)	0.0411 (6)
H2	0.5406	0.4020	0.4222	0.049*
C3	0.7045 (3)	0.4914 (6)	0.53080 (19)	0.0448 (7)
H3	0.6846	0.3519	0.5637	0.054*
C4	0.8199 (3)	0.6657 (6)	0.56754 (19)	0.0454 (7)
H4A	0.8754	0.6469	0.6255	0.054*
C5	0.8524 (4)	0.8670 (6)	0.5181 (2)	0.0499 (8)
H5	0.9327	0.9823	0.5422	0.060*
C6	0.7673 (3)	0.9015 (6)	0.43262 (18)	0.0423 (6)
H6	0.7893	1.0393	0.3995	0.051*
C7	0.7556 (3)	0.7445 (5)	0.09274 (15)	0.0334 (5)
C8	0.6714 (3)	0.5591 (6)	0.03958 (18)	0.0433 (7)
H8	0.6165	0.4334	0.0647	0.052*
C9	0.6688 (4)	0.5605 (6)	-0.0501 (2)	0.0518 (8)
H9	0.6131	0.4341	-0.0852	0.062*
C10	0.7472 (4)	0.7454 (7)	-0.08849 (19)	0.0540 (8)
H10	0.7440	0.7458	-0.1493	0.065*
C11	0.8301 (4)	0.9294 (7)	-0.0367 (2)	0.0601 (9)
H11	0.8839	1.0552	-0.0624	0.072*
C12	0.8347 (4)	0.9302 (6)	0.0532 (2)	0.0492 (7)

supplementary materials

H12	0.8915	1.0566	0.0877	0.059*
C13	0.7667 (3)	0.7445 (6)	0.19245 (15)	0.0360 (6)
H13	0.7771	0.9186	0.2133	0.043*
C14	0.9209 (4)	0.6019 (6)	0.23465 (19)	0.0461 (7)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0295 (3)	0.0577 (4)	0.0306 (3)	0.0117 (3)	0.0025 (2)	-0.0032 (4)
O1	0.0695 (15)	0.0584 (14)	0.0409 (12)	0.0304 (11)	0.0050 (10)	0.0040 (10)
O2	0.0254 (9)	0.0997 (17)	0.0418 (12)	0.0027 (10)	0.0028 (8)	-0.0118 (11)
O3	0.0545 (14)	0.0755 (17)	0.0822 (18)	0.0204 (13)	-0.0038 (12)	0.0157 (14)
O4	0.0300 (10)	0.0916 (18)	0.0829 (17)	0.0032 (14)	-0.0042 (10)	-0.0082 (17)
N1	0.0373 (12)	0.0503 (13)	0.0324 (13)	0.0042 (11)	0.0096 (10)	-0.0076 (11)
C1	0.0263 (11)	0.0461 (17)	0.0274 (12)	0.0045 (12)	0.0038 (9)	-0.0056 (12)
C2	0.0339 (14)	0.0466 (16)	0.0425 (16)	-0.0035 (13)	0.0058 (12)	-0.0041 (13)
C3	0.0458 (16)	0.0509 (18)	0.0387 (16)	0.0017 (14)	0.0101 (13)	0.0087 (14)
C4	0.0434 (16)	0.0590 (19)	0.0317 (15)	0.0069 (14)	0.0013 (12)	-0.0001 (14)
C5	0.0445 (17)	0.0542 (18)	0.0465 (19)	-0.0095 (14)	-0.0036 (14)	-0.0098 (15)
C6	0.0412 (14)	0.0438 (14)	0.0405 (16)	-0.0008 (14)	0.0041 (12)	0.0043 (14)
C7	0.0277 (11)	0.0397 (15)	0.0324 (12)	0.0074 (12)	0.0041 (9)	-0.0005 (13)
C8	0.0457 (16)	0.0482 (16)	0.0344 (16)	-0.0072 (13)	0.0034 (13)	-0.0014 (13)
C9	0.0510 (18)	0.066 (2)	0.0370 (17)	-0.0071 (16)	0.0034 (14)	-0.0084 (16)
C10	0.0566 (16)	0.070 (2)	0.0360 (15)	0.0064 (19)	0.0107 (13)	0.0054 (17)
C11	0.064 (2)	0.066 (2)	0.056 (2)	-0.0083 (18)	0.0258 (17)	0.0109 (18)
C12	0.0486 (16)	0.0516 (18)	0.0479 (19)	-0.0066 (15)	0.0100 (14)	-0.0056 (15)
C13	0.0289 (11)	0.0439 (15)	0.0343 (13)	0.0034 (13)	0.0030 (9)	-0.0071 (13)
C14	0.0400 (16)	0.065 (2)	0.0310 (15)	0.0090 (15)	0.0002 (12)	-0.0056 (15)

Geometric parameters (\AA , $^\circ$)

S1—O1	1.426 (2)	C5—C6	1.385 (4)
S1—O2	1.436 (2)	C5—H5	0.9300
S1—N1	1.615 (2)	C6—H6	0.9300
S1—C1	1.766 (2)	C7—C12	1.384 (4)
O3—C14	1.191 (4)	C7—C8	1.385 (4)
O4—C14	1.328 (4)	C7—C13	1.518 (3)
O4—H4O	0.83 (4)	C8—C9	1.375 (4)
N1—C13	1.458 (3)	C8—H8	0.9300
N1—H1N	0.82 (3)	C9—C10	1.368 (4)
C1—C6	1.380 (4)	C9—H9	0.9300
C1—C2	1.384 (4)	C10—C11	1.365 (5)
C2—C3	1.369 (4)	C10—H10	0.9300
C2—H2	0.9300	C11—C12	1.374 (4)
C3—C4	1.376 (4)	C11—H11	0.9300
C3—H3	0.9300	C12—H12	0.9300
C4—C5	1.368 (4)	C13—C14	1.522 (4)
C4—H4A	0.9300	C13—H13	0.9800

O1—S1—O2	120.73 (14)	C12—C7—C8	118.4 (3)
O1—S1—N1	106.78 (13)	C12—C7—C13	119.7 (2)
O2—S1—N1	105.24 (13)	C8—C7—C13	121.9 (2)
O1—S1—C1	107.61 (13)	C9—C8—C7	120.2 (3)
O2—S1—C1	106.75 (12)	C9—C8—H8	119.9
N1—S1—C1	109.41 (12)	C7—C8—H8	119.9
C14—O4—H4O	109 (3)	C10—C9—C8	120.8 (3)
C13—N1—S1	120.6 (2)	C10—C9—H9	119.6
C13—N1—H1N	119 (2)	C8—C9—H9	119.6
S1—N1—H1N	111 (2)	C11—C10—C9	119.4 (3)
C6—C1—C2	120.5 (2)	C11—C10—H10	120.3
C6—C1—S1	120.2 (2)	C9—C10—H10	120.3
C2—C1—S1	119.33 (19)	C10—C11—C12	120.5 (3)
C3—C2—C1	119.4 (3)	C10—C11—H11	119.7
C3—C2—H2	120.3	C12—C11—H11	119.7
C1—C2—H2	120.3	C11—C12—C7	120.6 (3)
C2—C3—C4	120.9 (3)	C11—C12—H12	119.7
C2—C3—H3	119.6	C7—C12—H12	119.7
C4—C3—H3	119.6	N1—C13—C7	112.0 (2)
C5—C4—C3	119.4 (3)	N1—C13—C14	110.6 (2)
C5—C4—H4A	120.3	C7—C13—C14	108.9 (2)
C3—C4—H4A	120.3	N1—C13—H13	108.4
C4—C5—C6	120.9 (3)	C7—C13—H13	108.4
C4—C5—H5	119.5	C14—C13—H13	108.4
C6—C5—H5	119.5	O3—C14—O4	126.0 (3)
C1—C6—C5	118.9 (3)	O3—C14—C13	124.3 (3)
C1—C6—H6	120.6	O4—C14—C13	109.7 (3)
C5—C6—H6	120.6		
O1—S1—N1—C13	40.7 (2)	C13—C7—C8—C9	-177.7 (3)
O2—S1—N1—C13	170.1 (2)	C7—C8—C9—C10	-0.8 (5)
C1—S1—N1—C13	-75.5 (2)	C8—C9—C10—C11	0.6 (5)
O1—S1—C1—C6	-23.6 (2)	C9—C10—C11—C12	-0.2 (5)
O2—S1—C1—C6	-154.5 (2)	C10—C11—C12—C7	0.1 (5)
N1—S1—C1—C6	92.1 (2)	C8—C7—C12—C11	-0.2 (4)
O1—S1—C1—C2	154.5 (2)	C13—C7—C12—C11	178.1 (3)
O2—S1—C1—C2	23.6 (2)	S1—N1—C13—C7	-133.6 (2)
N1—S1—C1—C2	-89.8 (2)	S1—N1—C13—C14	104.7 (2)
C6—C1—C2—C3	1.2 (4)	C12—C7—C13—N1	150.2 (2)
S1—C1—C2—C3	-176.9 (2)	C8—C7—C13—N1	-31.6 (4)
C1—C2—C3—C4	0.3 (4)	C12—C7—C13—C14	-87.1 (3)
C2—C3—C4—C5	-1.8 (4)	C8—C7—C13—C14	91.1 (3)
C3—C4—C5—C6	1.9 (5)	N1—C13—C14—O3	23.4 (4)
C2—C1—C6—C5	-1.1 (4)	C7—C13—C14—O3	-100.1 (3)
S1—C1—C6—C5	177.0 (2)	N1—C13—C14—O4	-158.5 (2)
C4—C5—C6—C1	-0.5 (4)	C7—C13—C14—O4	78.0 (3)
C12—C7—C8—C9	0.6 (4)		


supplementary materials

Hydrogen-bond geometry (Å, °)

<i>D—H···A</i>	<i>D—H</i>	<i>H···A</i>	<i>D···A</i>	<i>D—H···A</i>
O4—H4O···O2 ⁱ	0.83 (4)	1.98 (4)	2.727 (3)	150 (5)
C4—H4A···O3 ⁱⁱ	0.93	2.56	3.317 (4)	139

Symmetry codes: (i) $x+1, y, z$; (ii) $-x+2, y+1/2, -z+1$.

Fig. 1



supplementary materials

Fig. 2

