

# Impact of Early Tracheostomy on Weaning From Ventilation and Sedation in COVID-19 Pregnant and Early Postpartum Patient: Two Case Reports

Review began 08/31/2022

Review ended 09/22/2022

Published 09/26/2022

© Copyright 2022

Karic et al. This is an open access article distributed under the terms of the Creative Commons Attribution License CC-BY 4.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Edin Karic <sup>1</sup>, Hassan Mitwally <sup>2</sup>, Lolwa M. Alansari <sup>5</sup>, Adel Ganaw <sup>1</sup>, Mohamed O. Saad <sup>2</sup>, Abdulhamid Azhaghdani <sup>3</sup>

1. Critical Care Medicine, Hamad Medical Corporation, Al Wakra, QAT 2. Pharmacy, Hamad Medical Corporation, Al Wakra, QAT 3. Obstetrics and Gynecology, Hamad Medical Corporation, Al Wakra, QAT

Corresponding author: Edin Karic, ekaric@hamad.qa

---

---

## Abstract

Pregnant women are at high risk of coronavirus disease 2019 (COVID-19) complications, including acute respiratory distress syndrome (ARDS) and the need for mechanical ventilation. There is no literature on the optimal strategy for the management of difficult-to-wean pregnant and early postpartum patients.

We report two cases of pregnant women with COVID-19 pneumonia and ARDS, who required mechanical ventilation and high doses of analgesia, and sedation with neuromuscular blocking agents to facilitate ventilation and oxygenation. Both patients had a tracheostomy procedure to facilitate weaning from mechanical ventilation and sedation. Shortly after tracheostomy, sedation and analgesia, along with ventilatory support were weaned off. Both patients were discharged home. These cases propose early tracheostomy as a strategy to facilitate weaning from mechanical ventilation and sedation in pregnant and early postpartum patients.

---

**Categories:** Anesthesiology, Obstetrics/Gynecology, Infectious Disease

**Keywords:** sedation, weaning from mechanical ventilator, covid-19, pregnancy, early tracheostomy

## Introduction

Since the World Health Organization (WHO) has defined the coronavirus disease 2019 (COVID-19) outbreak as a pandemic, COVID-2019 has triggered several waves in most countries around the globe. Pre-existing comorbidities, non-white ethnicity, chronic hypertension, pre-existing diabetes, high maternal age, and high body mass index are risk factors for severe COVID-19 outcomes in pregnancy, and pregnant women with COVID-19 are at higher risk for admission to the intensive care unit (ICU) and mechanical ventilation (MV) than non-pregnant women of reproductive age [1]. Additionally, during the second wave of the COVID-19 pandemic, pregnant and peripartum women experienced more severe illness than in the first wave [2]. A number of case reports of severe COVID-19 pregnant women requiring intensive care treatment and MV have been reported [3-6].

The recommendations for analgesia/sedation in mechanically ventilated pregnant patients are generally similar to nonpregnant individuals [7]. Many intubated patients with severe COVID-19 infection require high doses of analgesia/sedation with or without a neuromuscular blocking agent (NMBA) [8]. This usually increases the risk of delirium and impacts weaning from ventilation, length of stay, mortality, and costs [9-11].

Weaning from MV is the process of withdrawing ventilator support, which usually starts with spontaneous breathing trials (SBT). Weaning failure is defined as either the failure of SBT or the need for reintubation within 48 hours following extubation [12,13]. Weaning from sedation is the process of liberation from any sedative agent, which usually starts before or goes in line with weaning from ventilation. The application of weaning protocols for sedation and ventilation in ICU is a daily routine [14].

The conversion from an endotracheal tube to tracheostomy significantly improves the measured values of weaning parameters in difficult-to-wean patients [15]. Tracheostomy appears to facilitate weaning from sedation. The short duration of MV after tracheostomy suggests that sedation requirements and ventilator dys-synchrony may be primary barriers to weaning these patients from the ventilator [16]. A systematic review, which included eight randomized control trials with around 2000 patients, analyzed the timing of tracheostomy, suggesting early tracheostomy might be superior to late tracheostomy [17].

During the second wave, we observed that most pregnant and postpartum patients required excessive doses of multiple analgesic and sedation agents with or without NMBA to facilitate ventilation and oxygenation as life-saving management in severe acute respiratory distress syndrome (ARDS). To our knowledge, there is no literature that covers the regimen of sedation and the ideal time of tracheostomy in pregnant, critically ill

### How to cite this article

Karic E, Mitwally H, Alansari L M, et al. (September 26, 2022) Impact of Early Tracheostomy on Weaning From Ventilation and Sedation in COVID-19 Pregnant and Early Postpartum Patient: Two Case Reports. Cureus 14(9): e29633. DOI 10.7759/cureus.29633

COVID-19 patients. In these two case reports, we shared our experiences of how (early) tracheostomy in this patient population may impact the weaning from ventilation and sedation.

## Case Presentation

### Case one

A 32-year-old, gravida three (week 29 + three), para two with Middle Eastern background, with no chronic illness, hospitalized on the day of COVID-19 positive PCR test due to pneumonia. She was on metformin at home for gestational diabetes. The patient was admitted to the ICU, with a SOFA score of 3, after one week due to severe respiratory distress and deterioration on the obstetric ward with 91% oxygen saturation ( $SpO_2$ ) on 15 liters of oxygen on a non-nonbreathing mask (NRBM) with abnormal arterial blood gases (Table 1). Vital signs upon ICU admission: blood pressure (BP) 118/58 mmHg, heart rate (HR) 112/min, respiratory rate (RR) 31/min, and temperature 37.2 °C.

Parameter	Value	Normal range
pH	7.33 ↓	7.35 – 7.45
pO <sub>2</sub>	72 mmHg	65 – 105 mmHg
pCO <sub>2</sub>	22 mmHg ↓	35 – 45 mmHg
BE	-14.6 mmol/L ↓	-2 to +2 mmol/L
Bicarbonate	11.4 mmol/L ↓	22 – 26 mmol/L

**TABLE 1: Arterial blood gases analysis upon ICU admission**


pO<sub>2</sub> – Partial pressure of oxygen, pCO<sub>2</sub> – Partial pressure of carbon dioxide, BE – Base excess, pH – Potential of hydrogen

After admission, a high flow nasal cannula (HFNC) with 30 liters flow rate and 65% fraction of inspired oxygen (FiO<sub>2</sub>) was applied, and the patient got respiratory stabilized. In the evening of the next day, she suddenly deteriorated with desaturation and severe respiratory distress. Following the discussion in the multidisciplinary team, an emergency cesarean section (CS) was performed. Thereafter, the patient remained respiratory and hemodynamically stable and was transferred back to ICU on MV. The obstetric team regularly followed up with the patient. The ensuing two weeks spent at the ICU were accompanied by cytokine storm, hospital-acquired pneumonia (HAP), ARDS, and consequently difficult and prolonged weaning. COVID-19 infection, HAP, and ARDS were treated as per local protocol considering the international guidelines (Table 2). Weaning from sedation and accordingly from ventilation was the main challenge. The patient required simultaneously up to four analgesic and sedative agents at a high dose (Figure 1) plus NMBA cisatracurium up to 3 mcg/kg/min to maintain oxygenation and ventilation. On day 20 of ICU admission, when tracheostomy was applied, she required continuous infusions of fentanyl 5 mcg/kg/hour, midazolam 3 mcg/kg/min, and cisatracurium 2 mcg/kg/min. Ventilator settings on the day of tracheostomy were: adaptive pressure ventilation-controlled mechanical ventilation (APV-CMV), RR 33/min, tidal volume 350 mL, positive end-expiratory pressure (PEEP) 10 cmH<sub>2</sub>O, FiO<sub>2</sub> 45%. The bedside percutaneous dilatative tracheostomy under bronchoscopy guidance was performed without any complications, and tracheostomy tube size 8 was inserted. COVID-19 PCR status a day before the procedure was inconclusive. During the next 48 hours, most analgesic and sedative agents have been stopped (Figure 1), except dexmedetomidine, which reached up to 1.5 mcg/kg/hr and was stopped 72 hours after tracheostomy. The weaning from the ventilator went smoothly, and the patient was decannulated 10 days after the tracheostomy. On the following day, she was transferred to the ward after 30 days of ICU stay and discharged home after two more days, with 38 days of hospital stay. The baby has been discharged home with the mother in good condition. Follow-up by primary care physicians in the next two months did not show any issue.

Drug	Dose	Duration
Ceftriaxone	2 grams, daily	7 days
Azithromycin	500mg, daily	3 days
Enoxaparine	40 mg, twice daily	25 days
Dexamethason	8 mg, daily	10 days
Hydrochloroquine	400 mg, daily	4 days
Lopinavir/Ritonavir	200/50 mg, twice daily	4 days
Remdesivir	1 <sup>st</sup> day 200 mg, then 100 mg, daily	5 days
Tocilizumab	400 mg, once	on two occasions
Meropenem	2 g, q8hr	7 days

**TABLE 2: COVID-19, HAP and ARDS treatment**

HAP - hospital-acquired pneumonia, ARDS - acute respiratory distress syndrome



**FIGURE 1: Tracheostomy effect on sedation and analgesia weaning**

**Case two**

A 33-year-old, gravida three (week 23 + two), para two, with an Asian background, with no chronic illness or risk factors, was hospitalized on the day of COVID positive PCR test due to pneumonia. The patient was admitted to the ICU, with the SOFA score of 4, after one week, due to severe respiratory distress and deterioration in the emergency department, with 91% SpO<sub>2</sub> on 15 liters of oxygen on NRBM with abnormal arterial blood gases (Table 3). Vital signs upon ICU admission: BP 102/64 mmHg, HR 99/min, RR 30/min, and temp. 37.2 °C.

Parameter	Value	Normal range
pH	7.42	7.35 – 7.45
pO <sub>2</sub>	67 mmHg	65 – 105 mmHg
pCO <sub>2</sub>	26 mmHg ↓	35 – 45 mmHg
BE	-7.4 mmol/L ↓	-2 to +2 mmol/L
Bicarbonate	17.1 mmol/L ↓	22 – 26 mmol/L


**TABLE 3: Arterial blood gases analysis upon ICU admission**

pO<sub>2</sub> – Partial pressure of oxygen, pCO<sub>2</sub> – Partial pressure of carbon dioxide, BE – Base excess, pH – Potential of hydrogen

After admission, HFNC with 45 liters flow rate and 75% FiO<sub>2</sub> was applied, and the patient got respiratory stabilized. In the evening, due to deterioration and high oxygen requirements, the patient had to be intubated. The obstetric team regularly followed up with the patient. Clinically, the patient showed a full picture of severe ARDS. COVID-19 pneumonia has been treated as per local protocol (Table 4). The patient required simultaneously up to four analgesic and sedative agents at high doses (Figure 2) plus NMBA cisatracurium up to 3 mcg/kg/min to maintain oxygenation and ventilation. This challenged the weaning from sedation and ventilation accordingly. Taking into account the experience with the first case, a multidisciplinary decision for bedside tracheostomy has been made. On day 8 of ICU admission, when tracheostomy was applied, she required continuous infusions of fentanyl 5 mcg/kg/hour, propofol 45 mcg/kg/min, and midazolam two-point 5 mcg/kg/min to maintain Richmond Agitation Sedation Scale (RASS) of minus two. Ventilator settings early on the day of tracheostomy were APV-CMV, RR 25/min, tidal volume 360 mL, PEEP 12 cmH<sub>2</sub>O, FiO<sub>2</sub> 45%. The bedside percutaneous dilatative tracheostomy under bronchoscopy guidance was performed without any complications, and tracheostomy tube size eight was inserted. COVID-19 PCR status two days after the procedure was negative. On the second day after the tracheostomy, the patient developed paralytic ileus with severe abdominal pain, treated conservatively. The weaning from analgesia and sedation took about four days because of the ileus (Figure 2). Subsequent weaning from the ventilator went smoothly as well, and the patient was decannulated eight days after tracheostomy. On the following day, she was transferred to the ward after 17 days of ICU stay and discharged home after two more days, with in total of 20 days of hospital stay. Follow-up by primary care physician and obstetrician in the next two months did not show any serious concerns. Patient delivered per emergency CS in week 38 + one without any complications. The baby has been discharged home with the mother in good condition.

Drug	Dose	Duration
Piperacillin/Tazobactam	4.5 g, q8hr	7 days
Enoxaparine	40 mg, twice daily	10 days
Dexamethason	8 mg, daily	5 days
Mehtylprednisolon	40mg, q8hr	3 days
Remdesivir	1 <sup>st</sup> day 200 mg, then 100 mg, daily	4 days
Tocilizumab	400 mg, once	on two occasions
Meropenem	1 g, q8hr	5 days

**TABLE 4: COVID-19 pneumonia treatment**



**FIGURE 2: Tracheostomy effect on sedation, analgesia, and muscle relaxant weaning**

## Discussion

Sedation is paramount in all patients who require MV. Adequately managed sedation prevents patient-ventilator dyssynchrony, accidental extubation, and circuit disconnections. In addition, it allows to facilitate prone ventilation and considers paralysis in those who develop refractory hypoxia [18].

The physiological and pharmacokinetic changes during pregnancy, which include an increase in blood volume, cardiac output, and total body water, may explain the requirement for high doses of sedatives and analgesics [19,20]. These factors may lead to an increase in the volume of distribution (Vd). Additionally, glomerular filtration rate and total drug clearance (Cl) are increased in pregnancy. Both increases in Vd and Cl lead to a decrease in serum drug concentration [20,21].

In this paper, we report the clinical outcomes of two pregnant patients with severe COVID-19 pneumonia after tracheostomy. Our observation was a significant reduction in sedation and pain medication after tracheostomy. We were able to wean off the sedation and analgesia completely within 48 hours of tracheostomy in the first case and within 72 hours in the second case. Both our patients required high doses of analgesics, sedatives, and an NMBA. Drug safety, possible side effects, and unpredictable maximum dose due to physiology and pharmacokinetic changes during the pregnancy have contributed to our decision for the early tracheostomy in the second case in particular.

It has been reported that critically ill COVID-19 patients tend to require higher doses of sedation and analgesia [18]. This exposes them to serious side effects, especially in pregnant patients, such as hypotension and bradycardia, which may further affect placental perfusion, prolonged duration of MV, affect neurological assessment, cause paralytic ileus, delirium, nausea, and vomiting [22]. Furthermore, prolonged periods of sedation may lead to drug accumulation, tolerance, and tachyphylaxis.

In a prospective cohort study of 164 patients, tracheostomy was associated with significantly higher 30-day survival among COVID-19 pneumonia patients requiring MV [23]. Moreover, early tracheostomy (within 14 days from intubation) was associated with a decrease in both ventilator days and ICU length of stay, although no statistically significant difference in decannulation time was observed between early and late tracheostomy (13 versus 15 days, respectively). In this study, sedative agents were weaned off within 48 hours in 76% of the tracheostomized patients.

Carmichael et al. were able to liberate 81% of tracheostomized COVID-19 pneumonia patients from the ventilator an average of nine days post-tracheostomy. Additionally, sedation and pain medication requirements decreased significantly within one week after the procedure [16]. Although this study observed a significant decrease in sedation requirement and ventilator days, the authors did not compare these findings with non-tracheostomized patients. They did not mention if pregnant patients were included in this study.

During the COVID-19 surges, when most ICUs were fully occupied, reducing the patient's stay in the ICU is considered one of the most valuable outcomes to empty this bed for new admissions. Any intervention that contributes to this objective should be considered. In a recently published meta-analysis of 14 non-randomized studies, early tracheostomy was associated with a decrease in mortality risk and ICU stays [24]. These findings should be interpreted with caution, due to the high heterogeneity between the included studies ( $I^2 = 87\%$  and  $70\%$  in mortality, and ICU stay, respectively).

## Conclusions

Pregnant patients with severe COVID-19 pneumonia seem to require high doses of analgesics, sedatives, and NMBA. A prolonged period of sedation and ventilation, and possible drug side effects, are serious concerns that should be taken into account. Early tracheostomy can be considered a strategy to facilitate weaning from MV in intubated pregnant and early postpartum COVID-19 patients requiring high doses of sedation and analgesia. However, prospective studies are required to confirm these findings.

## Additional Information

### Disclosures

**Human subjects:** Consent was obtained or waived by all participants in this study. Medical Research Center, Hamad Medical Corporation issued approval MRC-04-21-737. **Conflicts of interest:** In compliance with the ICMJE uniform disclosure form, all authors declare the following: **Payment/services info:** All authors have declared that no financial support was received from any organization for the submitted work. **Financial relationships:** All authors have declared that they have no financial relationships at present or within the previous three years with any organizations that might have an interest in the submitted work. **Other relationships:** All authors have declared that there are no other relationships or activities that could appear to have influenced the submitted work.

## References

- Allotey J, Stallings E, Bonet M, et al.: Clinical manifestations, risk factors, and maternal and perinatal outcomes of coronavirus disease 2019 in pregnancy: living systematic review and meta-analysis. *BMJ*. 2020, 370:m3520. [10.1136/bmj.m3520](https://doi.org/10.1136/bmj.m3520)
- Kadiwar S, Smith JJ, Ledot S, et al.: Were pregnant women more affected by COVID-19 in the second wave of the pandemic?. *Lancet*. 2021, 397:1539-40. [10.1016/S0140-6736\(21\)00716-9](https://doi.org/10.1016/S0140-6736(21)00716-9)
- Hong L, Smith N, Keerthy M, Lee-Griffith M, Garcia R, Shaman M, Goyert G: Severe COVID-19 infection in pregnancy requiring intubation without preterm delivery: a case report. *Case Rep Womens Health*. 2020, 27:e00217. [10.1016/j.crwh.2020.e00217](https://doi.org/10.1016/j.crwh.2020.e00217)
- Kolkova Z, Bjurström MF, Länsberg JK, et al.: Obstetric and intensive-care strategies in a high-risk pregnancy with critical respiratory failure due to COVID-19: a case report. *Case Rep Womens Health*. 2020, 27:e00240. [10.1016/j.crwh.2020.e00240](https://doi.org/10.1016/j.crwh.2020.e00240)
- Mehta H, Ivanovic S, Cronin A, et al.: Novel coronavirus-related acute respiratory distress syndrome in a patient with twin pregnancy: a case report. *Case Rep Womens Health*. 2020, 27:e00220. [10.1016/j.crwh.2020.e00220](https://doi.org/10.1016/j.crwh.2020.e00220)
- Slayton-Milam S, Sheffels S, Chan D, Alkinj B: Induction of labor in an intubated patient with coronavirus disease 2019 (COVID-19). *Obstet Gynecol*. 2020, 136:962-4. [10.1097/AOG.0000000000004044](https://doi.org/10.1097/AOG.0000000000004044)
- Pacheco LD, Saade GR, Hankins GD: Mechanical ventilation during pregnancy: sedation, analgesia, and paralysis. *Clin Obstet Gynecol*. 2014, 57:844-50. [10.1097/GRF.0000000000000064](https://doi.org/10.1097/GRF.0000000000000064)
- Tapaskar N, Colon Hidalgo D, Koo G, et al.: Sedation usage in COVID-19 acute respiratory distress syndrome: a multicenter study. *Ann Pharmacother*. 2022, 56:117-23. [10.1177/10600280211021925](https://doi.org/10.1177/10600280211021925)
- Ely EW, Shintani A, Truman B, et al.: Delirium as a predictor of mortality in mechanically ventilated patients in the intensive care unit. *JAMA*. 2004, 291:1753-62. [10.1001/jama.291.14.1753](https://doi.org/10.1001/jama.291.14.1753)
- Hanidziar D, Bittner EA: Sedation of mechanically ventilated COVID-19 patients: challenges and special considerations. *Anesth Analg*. 2020, 131:e40-1. [10.1213/ANE.0000000000004887](https://doi.org/10.1213/ANE.0000000000004887)
- Kotfis K, Williams Roberson S, Wilson JE, Dabrowski W, Pun BT, Ely EW: COVID-19: ICU delirium management during SARS-CoV-2 pandemic. *Crit Care*. 2020, 24:176. [10.1186/s13054-020-02882-x](https://doi.org/10.1186/s13054-020-02882-x)
- Esteban A, Frutos F, Tobin MJ, et al.: A comparison of four methods of weaning patients from mechanical ventilation. Spanish Lung Failure Collaborative Group. *N Engl J Med*. 1995, 332:345-50. [10.1056/NEJM199502093320601](https://doi.org/10.1056/NEJM199502093320601)
- Vallverdú I, Calaf N, Subirana M, Net A, Benito S, Mancebo J: Clinical characteristics, respiratory functional parameters, and outcome of a two-hour T-piece trial in patients weaning from mechanical ventilation. *Am J Respir Crit Care Med*. 1998, 158:1855-62. [10.1164/ajrccm.158.6.9712135](https://doi.org/10.1164/ajrccm.158.6.9712135)
- Schmidt GA, Girard TD, Kress JP, et al.: Liberation from mechanical ventilation in critically ill adults: executive summary of an official American College of Chest Physicians/American Thoracic Society clinical practice guideline. *Chest*. 2017, 151:160-5. [10.1016/j.chest.2016.10.037](https://doi.org/10.1016/j.chest.2016.10.037)
- Lim CK, Ruan SY, Lin FC, et al.: Effect of tracheostomy on weaning parameters in difficult-to-wean mechanically ventilated patients: a prospective observational study. *PLoS One*. 2015, 10:e0138294. [10.1371/journal.pone.0138294](https://doi.org/10.1371/journal.pone.0138294)
- Carmichael H, Wright FL, McIntyre RC, et al.: Early ventilator liberation and decreased sedation needs after tracheostomy in patients with COVID-19 infection. *Trauma Surg Acute Care Open*. 2021, 6:e000591. [10.1136/tsaco-2020-000591](https://doi.org/10.1136/tsaco-2020-000591)
- Andriolo BN, Andriolo RB, Saconato H, Atallah ÁN, Valente O: Early versus late tracheostomy for critically ill patients. *Cochrane Database Syst Rev*. 2015, 1:CD007271. [10.1002/14651858.CD007271.pub3](https://doi.org/10.1002/14651858.CD007271.pub3)
- Karamchandani K, Dalal R, Patel J, Modgil P, Quintili A: Challenges in sedation management in critically ill patients with COVID-19: a brief review. *Curr Anesthesiol Rep*. 2021, 11:107-15. [10.1007/s40140-021-00440-x](https://doi.org/10.1007/s40140-021-00440-x)
- Blackburn S: Pharmacokinetic changes in the pregnant woman. *J Perinat Neonatal Nurs*. 2012, 26:13-4. [10.1097/JPN.0b013e318242fdf1](https://doi.org/10.1097/JPN.0b013e318242fdf1)
- Feghali M, Venkataramanan R, Caritis S: Pharmacokinetics of drugs in pregnancy. *Semin Perinatol*. 2015, 39:512-9. [10.1053/j.semperi.2015.08.003](https://doi.org/10.1053/j.semperi.2015.08.003)

21. Dawes M, Chowienczyk PJ: Drugs in pregnancy. Pharmacokinetics in pregnancy. *Best Pract Res Clin Obstet Gynaecol.* 2001, 15:819-26. [10.1053/beog.2001.0231](https://doi.org/10.1053/beog.2001.0231)
22. Barr J, Fraser GL, Puntillo K, et al.: Clinical practice guidelines for the management of pain, agitation, and delirium in adult patients in the intensive care unit. *Crit Care Med.* 2013, 41:263-306. [10.1097/CCM.0b013e3182783b72](https://doi.org/10.1097/CCM.0b013e3182783b72)
23. Breik O, Nankivell P, Sharma N, et al.: Safety and 30-day outcomes of tracheostomy for COVID-19: a prospective observational cohort study. *Br J Anaesth.* 2020, 125:872-9. [10.1016/j.bja.2020.08.023](https://doi.org/10.1016/j.bja.2020.08.023)
24. Staibano P, Levin M, McHugh T, Gupta M, Sommer DD: Association of tracheostomy with outcomes in patients with COVID-19 and SARS-CoV-2 transmission among health care professionals: a systematic review and meta-analysis. *JAMA Otolaryngol Head Neck Surg.* 2021, 147:646-55. [10.1001/jamaoto.2021.0930](https://doi.org/10.1001/jamaoto.2021.0930)