

# Discovering electrophysiology in photobiology

## A brief overview of several photobiological processes with an emphasis on electrophysiology

Vadim Volkov\*

Faculty of Life Sciences and Computing; London Metropolitan University; London, UK

**Keywords:** Photobiology, electrophysiology, polarized light, rod cells, dark noise, statistics of photons, optogenetics, quantum dots

The mini-review gives special attention to holistic approach and mechanisms of processes. The physical and chemical frames and background for visual perception and signaling are discussed. Perception of photons by retinal rod cells is described in more detail starting from photon absorption and culminating in ion currents. Dark noise and temperature-dependence of photocurrents in photoreceptor cells are analyzed. Perception of polarized light, its effects and informational importance are discussed based on underlying mechanisms and specialized morphological structures of biological organisms. Role of statistics of photons in photoreception is questioned. The review also pinpoints new and developing directions and raises questions for future research.

### Introduction

Life has evolved, developed and is flourishing nowadays in the permanent fluctuating fluxes of electromagnetic radiation of different frequencies and amplitudes. The electromagnetic waves are bringing information about the world and provide energy for the existence of living organisms. From the wide range of electromagnetic waves life chose narrow band of 400–800 nm for vision, light perception and photosynthesis. The reasons are in energy of photons and nature of chemical bonds. Much higher energies are harmful and break molecules; much lower energies are not distinguished from the thermal noise (see examples below).

The short review is focused on physico-chemical mechanisms of photobiological processes, conversion of photons into ion currents for further processing and on more specific aspects including perception of polarized light and light with unusual statistics, their role for biological systems and implications for

biological research. The general preface starts from storage of energy in photosynthesis moving to mechanisms of light perception and vision.

### Basic Preface: Plants and Light

Quanta of light (photons) are caught by pigments in photosynthetic membranes of plants, algae and some bacteria. The photosensitive pigments are mostly represented by chlorophylls and bacteriochlorophylls absorbing photons; the spectral properties of the molecules are naturally selected for solar radiation reaching earth. Some algae live in deep waters with lower proportion of larger wavelengths, spectrum of radiation is shifted to green-blue side. The algae usually possess accessory pigments called phycobilins to absorb photons within the range outside spectrum of chlorophylls and to transfer the energy to chlorophylls.

The next sequence of fast picosecond photochemical reactions results in electric charge transfer between the sides of the membranes. The generated electric potential is used for downhill charge transfer; the energy is converted to energy of chemical bonds, mostly in ATP (adenosine triphosphate), so the energy can be stored for further biochemical reactions and membrane transport processes.<sup>1</sup> The complex chain of reactions for ATP synthesis requires large protein macromolecular complexes. Usually protons are passing via subunits of ATP-synthase to change the conformation of protein complex; finally it is released in ATP synthesis from ADP and inorganic phosphate  $P_i$ .<sup>2,3</sup>

Energy  $E$  of photon (quantum of light) is determined by the equation:


$$E = h \nu = h c / \lambda \quad (1)$$

where  $\nu$  is the light wave frequency, which is reverse proportional to wavelength  $\lambda$  ( $c = \lambda \nu$ ),  $h$  is Planck's constant (about  $6.6 \cdot 10^{-34}$  J\*s) and  $c$  is the speed of light in vacuum (about  $3 \cdot 10^8$  m/s). The energies of single visible photons (400 – 700 nm, see below) will be then from 2.8 to  $5 \cdot 10^{-19}$  J (reverse proportional to wavelength). The values correspond to 170–300 kJ per mole of photons. For comparison, the standard energy of ATP hydrolysis is -30 kJ/mole and up to around -60 kJ/mole depending on pH, ATP, ADP and ion concentrations.<sup>4,5</sup> The energy of a single photon is quite large and sufficient to provide synthesis of ATP molecule or start the cascade of signaling events. Taking into account that 1) much less than 1% of photosynthetically active


\*Correspondence to: Vadim Volkov; Email: vadim.s.volkov@gmail.com and v.volkov@bio.gla.ac.uk

Submitted: 11/29/2013; Revised: 03/01/2014; Accepted: 03/03/2014; Published Online: 03/12/2014

Citation: Volkov V. Discovering electrophysiology in photobiology: A brief overview of several photobiological processes with an emphasis on electrophysiology. *Communicative & Integrative Biology* 2014; 7:e28423; <http://dx.doi.org/10.4161/cib.28423>



**Figure 1.** Channelrhodopsins are light-gated ion channels from green algae. After a flash of illumination the protein molecule of ion channel temporarily changes conformation due to isomerisation of molecule of retinal bound to lysine of the protein (compare with rhodopsin, see below). It allows selective passage of cations according to electrochemical gradient ( $\text{Na}^+$  selectivity is depicted); within a short time (milliseconds to seconds depending on the protein structure) the channel returns to the initial conformation and ion current stops.



**Figure 2.** Typical recording of membrane potential in cells of the emerged blade of the growing leaf 3 of barley, and the response of membrane potential to changes in illumination and further addition of NaCl (100 mM) to the root medium. Several phases of responses with different kinetics were observed upon changes in illumination (18 experiments with 9 plants). Light was supplied using fiber optics from the cold light source at the background of dim illumination in the electrophysiological rig. The figure is from<sup>14</sup> with permission from the Oxford University Press, extra information about the recordings is provided by V Volkov.

solar radiation is transformed to energy of photosynthetic products and 2) efficiency of photosynthesis is around 1–3%,<sup>1</sup> plants are not ideal biochemical convertors of light. Biosphere is rather wasting solar radiation and life is not limited by energy demands, having the other intrinsic and external (e.g., water resources, carbon and nitrogen availability, temperature etc.) regulators, limitations and reasons for development.

Apart from absorbing energy of light in photosynthesis, plants also have photoreceptor systems including several phytochromes, cryptochromes and phototropins, they respond to very low intensity of different light wavelengths and have regulatory functions.<sup>6</sup> Recently discovered plant receptor UVR8 for the range of UV-B light complements the earlier known receptors (reviewed in ref. 7).

An interesting phenomenon exhibited by some bacteria is direct transformation of light energy to electric fluxes of ions. Light-driven proton pump bacteriorhodopsin is an integral

membrane protein of some halobacteria (studied in detail for *Halobacterium halobium*); it pumps protons out of cell at the expenses of absorbed photons.<sup>8</sup> Green unicellular algae *Chlamydomonas reinhardtii* and some others possess ion channels channelrhodopsins, which are directly regulated by light.<sup>9,10</sup> Several channelrhodopsins were studied and characterized in detail (e.g., and reviewed in 11–13) (Fig. 1 and see later).

Light responses in photosynthesis and by photoreceptor cascades are registered by many ways and methods, from picosecond changes in spectral properties of photosynthetic pigments, conformational transitions of proteins, migration of small molecules, signaling events within and between cells, slow changes in membrane potentials of cells (Fig. 2) etc. to accumulation of new pools of synthesized organic molecules-photoassimilates.<sup>1</sup>



### Simple Scheme of Photoinduced Events in Retinal Cells

Vertebrate animals, invertebrates and the other non-photosynthetic organisms get visual information using specialized organs. Eyes of vertebrates are very sensitive. The estimate for the lower limit of human eye sensitivity is about 100 photons, which will correspond to about 10–20 photons reaching photoreceptor cells due to absorption and reflection.<sup>15</sup> Eyes contain retinal layer with photoreceptor cells, which includes 1) millions of more sensitive rods for lower illumination or night vision and may have 2) usually smaller number of less sensitive cones for color vision.<sup>16</sup> Rods are oblong cells (Fig. 3) with numerous stacked disks at the outer (distal) part; the disks are formed by membranes with photosensitive pigment rhodopsin.<sup>16</sup> The number of disks is quite large, for example about 1000 for a mouse rod; then about  $10^5$  rhodopsin molecules per disk will result in nearly hundred millions of rhodopsin molecules per a mouse rod (reviewed in 17).

The structure of rhodopsin is well studied: the protein part is called opsin, which is 30–50 kDa seven transmembrane G-protein coupled receptor (e.g., reviewed in 17,18); the chromophore is 11-*cis*-retinal, which photoisomerizes to all-*trans*-retinal after absorbing a photon.<sup>19</sup> Photochemistry of rhodopsin isomerization with picosecond rate constants and several intermediates is also well known: isomerization has quantum yield of over 0.6 and the energy barrier is nearly 190 kJ/mole for primary reaction (energy difference is slightly over 130 kJ/mole between rhodopsin and the first photoproduct bathorhodopsin), quantum yield is about 0.1 for one of the later steps (reviewed in ref. 20). Smaller values for activation energy around 100–120 kJ/mole were also reported, though there are variations between species and experimental conditions (reviewed in ref. 21).

Most adult insects have compound eyes, which are composed of thousands similar units called ommatidia. Each ommatidium has generally eight photoreceptor cells with rhodopsin.<sup>22</sup>

For an unaided human eye the visible wavelengths are limited by 400 nm – 700 nm,<sup>16</sup> from violet to red colors. Insect eyes have slightly different wavelengths of light perception, from below 300 nm to over 700 nm.<sup>23</sup> The difference in UV range is explained by the secondary UV-absorbing chromophore in




**Figure 3.** Microscope image of isolated *Xenopus laevis* retinal cells (A) and a rod cell from the preparation (B). Scale bar is 120  $\mu\text{m}$  (A) and 30  $\mu\text{m}$  (B).

some insects, the sensitizing pigment can transfer the energy further to the primary photopigment.<sup>22</sup>

The sequence of events from a photon hitting a rod cell to the registered photocurrent of the cell is deciphered in vertebrates (reviewed in ref. 24) (Fig. 4). Rhodopsin absorbs a photon and isomerizes to metarhodopsin; both proteins were crystallized and resolved structurally (refs. 25 and 26 correspondingly). Metarhodopsin has a short life half-time, so special approaches were used to crystallize this G-protein coupled receptor.<sup>26</sup> The next step is activation of transducin. Transducin is G-protein, which is composed of  $\alpha$ ,  $\beta$  and  $\gamma$  subunits. Transducin laterally diffuses on the surface of disk membrane, interacts with activated rhodopsin (the spectroscopic intermediate metarhodopsin II) and changes bound GDP for GTP (reviewed in refs. 24,27,28). Active form of transducin is  $\alpha$ -subunit-GTP, two activated subunits bind to phosphodiesterase PDE and activate it. PDE hydrolyses cGMP to GMP and decreases concentration of the cyclic nucleotide in a cell. Sharp drop in cGMP closes cyclic nucleotide gated channels, which are regulated (gated) by bound cGMP. Under low cGMP the channels are closed, while they are in an open state in darkness under higher micromolar concentration of cGMP.<sup>29</sup> Rod cell membrane hyperpolarizes having closed cyclic nucleotide gated channels, so the initial absorption of photon expresses finally in the change of the membrane potential and corresponding ion current, which is further passed to neurons (reviewed in refs. 24,27,28).


The amplification of signal in a rod cell is happening in the cascade: metarhodopsin  $R^*$  can activate up to hundreds of G-protein transducin molecules (G) molecules.<sup>24,30</sup> Rate of activation is around 125  $G^* s^{-1}$  per  $R^*$  for amphibian rods at room temperature and about 3 times higher in mammalian rods at body temperature.<sup>28</sup> Further on in darkness metarhodopsin, activated forms of transducin and phosphodiesterase are deactivated, concentration of cGMP is increasing and membrane depolarizes again; the processes add time components and kinetics to the development of the events (reviewed in ref. 31). Inactivation includes several steps, for  $R^*$  in mouse rods the activity is quenched with half-time about 50–80 ms<sup>32,33</sup> by




**Figure 4.** Signal transduction chain in a vertebrate retinal rod starting from a photon  $h\nu$  and leading to cyclic nucleotide gated channels. Closure of the ion channels after drop in cGMP results in membrane hyperpolarisation and stops inward ion current of sodium and calcium.

successive reactions of phosphorylation by rhodopsin kinase and binding of the protein arrestin (reviewed in ref. 31).

The kinetics of induced photocurrent in rods (Fig. 5) is reasonably modeled by several equations, which take into account kinetics of the reactions (rhodopsin to metarhodopsin, activation of transducin, phosphodiesterase, drop in cGMP and closure of ion channels), include diffusion, rod morphology and inactivation kinetics (e.g. refs. 31 and 35). Higher number of photons increases the photocurrent by activating more rhodopsin molecules; the response is nonlinear and reaches saturation at around eg 20,000 photons per light pulse for *Xenopus* rod.<sup>36</sup> Numerous mutations affecting components of the signal transduction chain for photon perception are known, some of them have an effect on photocurrent and its' kinetics (reviewed in ref. 31). Surprisingly stable and reproducible kinetics of photocurrent upon a given number of photons provides robust and reliable information about the light source. The rod photoreceptor could be considered like a natural example of engineering with numerous feedbacks; inactivation components are especially important for ensuring reproducible responses.<sup>37</sup> For example, C-terminus of rhodopsin has 6 phosphorylation sites, which are important for inactivation of activated  $R^*$  and reproducible kinetics of photocurrent;



**Figure 5.** Kinetics of retinal rod responses of *Xenopus laevis* to pulses of illumination. The color legend indicates the corresponding number of impinging photons. Reproduced with the permission of The Optical Society of America from.<sup>34</sup>



**Figure 6.** Light-activated ion channel channelrhodopsin-2 can restore basic visual function in photoreceptor-deficient mice with 6–9 orders of magnitude lower sensitivity (according to refs. 44 and 45).

decreasing the number of the sites in mouse mutants increased the duration of photocurrent and changed its shape.<sup>38</sup>

Invertebrates may have slightly different sequence of events during phototransduction. Fruitfly *Drosophila* is a well-known model in genetics; numerous mutants are useful for deciphering the light perception in the ommatidia of the insect. The difference in phototransduction from vertebrates is that in *Drosophila* 1) phospholipase C is present instead of phosphodiesterase, 2) signaling via inositol trisphosphate and diacylglycerol and probably polyunsaturated fatty acids without cGMP 3) results in opening of closed under darkness 4) transient receptor potential ion channels (reviewed in refs. 39 and 40). More differences could be revealed among species of numerous and strikingly unusual biological organisms.

### Artificial Receptors of Light and Means to Correct Impaired Vision

Obviously, the ion current is presumably the only converted measure of light intensity and it is further transferred to neurons of the neural system. Artificial electronic light receptors were proposed instead of absent or damaged retinal cells under

ophthalmological diseases; the electric response of the retinal prosthesis is passed to neurons restoring (at least partially) light sensing (reviewed in refs. 41 and 42). Compared with retinal photoreceptor cells, which are highly specialized and finely tuned systems, retinal prostheses are simpler. Another approach is to express light-sensitive proteins in neurons. Several genes from *Drosophila* including rhodopsin rendered light sensitivity to hippocampal neurons in primary culture.<sup>43</sup> Directly activated by light ion channel channelrhodopsin is also proposed for use in retinal prosthesis.<sup>42</sup> Model experiments using mouse model with degenerated retinas demonstrated that being genetically targeted to inner retinal neurons channelrhodopsin-2 can restore basic visual function with, however, 6–9 orders of magnitude lower sensitivity<sup>44,45</sup> (Fig. 6).

Here emerges an interesting area of discrimination between simple photic responses (difference light-dark and corresponding physiological reactions) and recognition of images derived from visual perception by specialized photoreceptor cells. The direction leads to neuroscience with different specialized types of neurons and is outside the scope of the present review. For example, photosensitive retinal ganglion cells were found in mice, the cells express melanopsin and confer simple photic responses in blind mice without photoreceptor rods and cones.<sup>46</sup> Human melanopsin from intrinsically photosensitive retinal ganglion cells was heterologously expressed in mouse paraneuronal cell line Neuro-2a and rendered light-induced ion currents.<sup>47</sup> Mouse melanopsin was heterologously expressed in *Xenopus* oocytes<sup>48</sup> and in human embryonic kidney HEK293 cells<sup>49</sup> making them light-sensitive. Visual perception is much more complicated.

### Dark Noise and Temperature-Dependence of Photoreceptors

Since each biological system is a physico-chemical system, it can be described in terms of energies, activation energies, kinetics of reactions and the other parameters. Numerous restrictions are implied by biological components (nature of interacting molecules; morphology of cells and tissues; pH, redox potential and chemical composition of cell medium). On the opposite, obvious predictions about behavior of biological systems could be done from the known parameters. The simplest prediction is about rhodopsin isomerization. High activation energy of photoisomerization for rhodopsin cannot prevent it from spontaneous isomerization without any photons, but due to temperature (around 300 K) and uneven distribution of energies of interacting molecules. One reason of so called dark noise of photoreceptor appears, it will depend (for the reason) on temperature and number of rhodopsin molecules in a rod cell. Dark noise of photoreceptor is expressed in spontaneous electric signals (voltage spikes or recorded ion currents), which are not distinguishable from the electric signals induced by photons. Indeed, dark noise of photoreceptor measured in events/(cell\*s) (varied from  $10^{-3}$  to 1) was linearly proportional to the number of rhodopsin molecules within the range of  $10^8$  –  $10^{10}$  rhodopsin molecules/cell (locust-human-toad-*Limulus*),<sup>21</sup> means around  $10^{-10}$  events/

second per a rhodopsin molecule. However, the situation looks much more complicated taking into account the sequence of events from activated rhodopsin to closure of ion channels. Each element (spontaneous activation of transducin or phosphodiesterase, drop in cGMP or stochastic closure of ion channels) has to be assessed using activation energies and stoichiometry of reactions. More convoluted models could be also used; for example, it was suggested to apply Hinshelwood distribution for complex molecules with many vibrational modes to thermal activation of rhodopsin.<sup>50</sup>

An interesting observation was done for photoreceptor cells of horseshoe crab *Limulus*: dark noise was about one bump of voltage per 5 min during evening and 25 times higher during midday.<sup>51</sup> On the opposite, the voltage response to illumination (gain) was much higher during night time. The model of two-step process for spontaneous temperature-dependent isomerization of rhodopsin was proposed.<sup>21,52</sup> The first step is deprotonation of protein-bound chromophore (form of retinal) with activation energy around 100 kJ/mole. The second step is isomerization of the chromophore with activation energy around the remaining 100 kJ/mole. The first step depends on pH and geometry of protein-chromophore complex, so changes in pH and protein microenvironment can strongly influence the noise. Temperature-dependent isomerization of chromophore at the second step can account for the observed temperature dependence of noise, which was the same during daytime and during night. Indeed, for photoreceptor cells of horseshoe crab the noise increased about twice with temperature increase by 6 °C (from 12 events/s at 14 °C to 28 events at 20 °C during daytime and from 3 events/s at 20 °C to 6.5 events/s at 26 °C during night) giving the estimated energy activation of 90–100 kJ/mole.<sup>52</sup>

However, the origin of photoreceptor noise and temperature dependence of photoreception could be much more complex. Dependence of thermal noise on pH of external medium was found for photoreceptor cells of horseshoe crab,<sup>52</sup> but not for cones of salamander<sup>53</sup> or toad rods.<sup>54</sup> Noise in salamander cones<sup>55</sup> had different origin (due to pigment or due to transduction species) for different types of cones.

Finally, physiological mechanisms and biological peculiarities are important for temperature-dependence of visual perception. For example, in swordfish (*Xiphias gladius*) the temporal resolution of retinograms (electrical activity recorded from isolated retinal preparations) was measured by the flicker fusion frequency. The parameter determines the frequency of pulses, which are distinct in retinograms. The flicker fusion frequency has very high temperature coefficient  $Q_{10}$  (change of measured parameter with changing temperature by 10°C) in swordfish: about 5; resolved frequency is around 30 Hz at 21 °C and just 6 Hz at 6 °C.<sup>56</sup> Swordfish has twice higher  $Q_{10}$  of flicker fusion frequency than bigeye and yellowfin tunas, but possesses special system for heating brain and eyes and, therefore, keeps good visual perception at depths of several hundred meters.<sup>56</sup>

At the molecular level it's reasonable to start from the point that a clear model for thermal isomerization of rhodopsin still has to be proposed, especially when rhodopsin is located in

membranes of rod outer segments and interacts with lipids, water molecules and proteins. For comparison, energy of gas molecules is described by Maxwell-Boltzmann distribution and statistics and in a simplified form can be expressed by:

$$N_i/N = e^{-\Delta E/kT} \quad (2)$$


where  $N$  is the total number of particles;  $N_i$  is the number of particles with energy by  $\Delta E$  higher than a certain energy (usually probabilities  $N_i/N$  are considered); Boltzmann constant  $k = 1.38 \cdot 10^{-23}$  J/K,  $T$  is temperature (e.g. ref. 57, p.203).

A simple estimate is useful to make an analogy to gas systems with the corresponding parameters: 1) the temperature is 300 K (27 °C), 2) activation energy  $E_a$  equals to 100 kJ/mole and for the known Avogadro number  $N_a = 6.02 \cdot 10^{23}$  molecules/mole gives  $E_a/N_a = 100\,000$  Joules /  $6.02 \cdot 10^{23} = 1.6 \cdot 10^{-19}$  Joules/molecule, then part of gas molecules with energy higher the energy barrier of  $1.6 \cdot 10^{-19}$  Joules/molecule will be:  $\exp(-1.6 \cdot 10^{-19} / (1.38 \cdot 10^{-23} \cdot 300)) \approx \exp(-39) \approx 3 \cdot 10^{-17}$ . Obviously, the number is far below the reasonable values of the observed dark noise ( $10^{-10}$  events/second per a rhodopsin molecule for  $10^{10}$  molecules per a photoreceptor rod while the activated rhodopsin molecule has a life half-time of about 50 ms); it proves that quantum effects of photochemical processes and interactions with surrounding molecules are involved to change the distribution.

Pondering the smart and ideal biological design, it's reasonable to think about photoreceptor without any dark noise. However, the biological systems are too noisy and have relatively high level of errors compared with technical or digital computer systems. Potentially the noise could be beneficial for the visual perception keeping a certain background activity of neurons. More general considerations propose limits for noise suppression in biological systems, otherwise negative feedbacks and the whole functioning of system are getting extremely expensive: the minimum standard deviation decreases with the quartic root of the number of events for Poisson communication channels.<sup>58</sup>

### Polarized Light, Occurrence in Nature and Mechanisms of Perception

Apart from wavelength and intensity electromagnetic waves have polarization. Electromagnetic wave consists of fluctuating electric and magnetic fields, which have perpendicular orientation to each other. Considering a single photon like a sort of wave packet and a pulse of light with thousands and millions of photons, which are not ordered and have random directions of individual vectors of electric field, it's simple to understand unpolarized light. However, depending on the light source and passed medium the orientation of electric vector in the wave may become ordered, the electromagnetic wave is becoming polarized. If the orientation of electric vector fluctuates in one plane, then the wave is linearly polarized; alternatively the vector can rotate and the wave is circularly polarized then. The degree of polarization characterizes the extent of ordered orientation of the electric vector. Polarization of light occurs naturally in atmosphere



**Figure 7.** Basic principle for perception of polarized light in biological organisms. Ordered orientation of pigment molecules in photoreceptor cell allows detection of electric vector E of light (B is the vector of magnetic field of electromagnetic wave). Molecules of photosensitive pigments are depicted in the form of blue ovals.



**Figure 8.** Underwater world with several photobiological phenomena and examples of perception. Light is reflected from the water surface and becoming partially polarized; light is also passing to water and getting partially linearly polarized; polarized light in water is sensed by fishes; fishes have multilayer reflecting skin with unusual properties; blonde, thorn-tail and undulate rays (*Raja brachyura*, *Raja clavata*, *Raja undulate*; size of specimens is around 1 m) possess electric sense. The picture is taken with the permission of staff of SEA LIFE London Aquarium.

during sunrise or sunset, after reflection of light from water surface, from Moon and also underwater.<sup>59-61</sup> So, the polarized light has essential information for biological organisms about time and surrounding environment. It cannot be directly digitized and translated to a fixed digital image.

Human beings have low sensitivity for polarized light and able to detect just very high polarization degrees, over 60%, while some insects and fishes can see polarization of 10% even.<sup>59</sup> Polarization of light is more important for invertebrates and fishes due to their habitat and size. Some insects are able to use polarized light for behavioral signaling. The metallic green elytra (forewings) of Japanese jewel beetle *Chrysochroa fulgidissima* reflects light, which is becoming highly polarized because of complex multi-layered surface of the elytra.<sup>62</sup> South American

beetle *Plusiotis boucardi* with metallic-looking forewings has dual-pitched helicoidal layer with bowl-shaped recesses, the forewings are special in reflecting circularly polarized light without changing its handedness.<sup>63</sup> The degree of polarization may depend on the angle of reflection and the polarized light reflected by the beetles may have function in recognition of the species. In butterfly *Heliconius* the reflected polarized light from the wings can be a preferential signal for mate recognition.<sup>64</sup>

In fish and underwater species, polarization pattern of light reflected from the skin or epidermal tissues can be also important for being less visible by predators. For describing the habitat, it is necessary to mention that many algae have iridescent surface layers (e.g., cuticle) and/or cellular structures,<sup>65-67</sup> so determining the underwater visual environment. Squid has iridescent skin with multilayer reflector cells; the cells contain plates of protein interspersed by cytoplasm and allow the animal to change color and polarization of reflected light and to send potential polarization signals to the other cephalopods with polarization-sensitive vision.<sup>68-70</sup> Several species of fish (sprat *Sprattus sprattus*, sardine *Sardina pilchardus*, and herring *Clupea harengus*) have similar cell structures: broadband guanine-cytoplasm 'silver' multilayer reflectors, which make, however, the reflected light non-polarized and provide a sort of cryptic camouflage for the situation.<sup>71</sup> The study on the structure of algal, insect and animal polarization surfaces (wings, forewings, skin etc.) may have important implications for material science

The perception of polarized light, obviously, requires ordered spatial positioning of rhodopsin/visual pigments and specialized cells for perceiving the direction and degree of polarization (Fig. 7). Much more is known about behavioral reactions of different species to polarized light than about mechanisms of perception of the light, though a few detailed studies on the perception exist and multiply. In continuation of the earlier experiments describing polarization-dependent dances of bees,<sup>72</sup> it was shown that bees possess receptors of polarized UV light, but not of the other wavelengths.<sup>73,74</sup> The specialized receptors at the dorsal rim of bee's compound eyes were examined electrophysiologically.<sup>75</sup> Very high polarization sensitivity up to 18 (ratio of maximal to minimal voltage responses to several directions of electric vector of light) was found in UV-responsive cells lacking sensitivity to green light.<sup>75</sup> In fish the perception of UV polarized light is associated with cones,<sup>76</sup> while the polarized light of another parts of spectrum is also sensed (according to electrical recordings from brain tectum).<sup>77,78</sup>

The ultrastructure of photoreceptors for polarized light in arthropods is studied for many species (e.g., and reviews: 79-82). Usually the pigments are packed orderly in microvilli within a photoreceptor cell; in each ommatidium the direction of microvilli is orthogonal (at 90°) in the photoreceptor cells. So, the direction of electric vector of light can be detected by different cells, the signal is periodically sent to neurons, which process the information about polarization. Periodical signal from different cells helps to exclude effect of light intensity, while the initial polarization of light is often amplified within an ommatidium or an arthropod eye by reflecting surfaces (reviewed in refs. 80-82). New approaches including methods of molecular biology and genetics help in dissecting and understanding the components of the polarization vision.<sup>83</sup>

It looks challenging (Fig. 8) to mix 1) perception of electric vector of polarized light with the electric sense of several sea inhabitants including sharks and rays<sup>84-86</sup> and 2) magnetic vector of polarized light with magnetoreception of many living organisms (e.g. reviewed in ref. 87). The electric sense of sharks and rays has resolution below 5 nV/cm<sup>84-86</sup> and seems comparable or even much lower than the electric field in light beams: e.g., sunlight has intensity of electric vector around 10 V/cm (88 p.10). Magnetic field vector of the light would correspond to about 3 mK (10 V/cm /  $c = 10^3 \text{ V/m} / (3 \cdot 10^8 \text{ m/s}) = 3.3 \cdot 10^{-6} \text{ T}$ ), lower than the magnetic field of Earth (25–65 mK).

However, the distinct sensual informational channels are determined by:

- 1) range of measured values and frequencies of electromagnetic fields (hundreds of Hz for electric sense and around  $10^{14} - 10^{15}$  Hz for visible light);
- 2) detailed possible mechanisms (often not studied yet) of perception;
- 3) morphological structures for specific sort of perception that are based on the possible mechanisms created and selected by nature (specialised photoreceptor cell types in eyes, ampullae of Lorenzini for electroreception in sharks and rays, magnetite-containing cells for magnetoperception etc.);
- 4) further numerous peculiarities.

Polarized light and direction of polarization may have an effect on development of biological organisms. For example, protonemata (primary germinating structures in development from spores) of ferns and mosses have preferential orientation according to electric vector of illumination.<sup>89-94</sup> Interesting experiments were performed with isolated and immobilized in agar protoplasts (single cells obtained from plants using cell wall degrading enzymes) of moss *Physcomitrella patens*.<sup>95</sup> The phragmoplast (scaffold for future cell wall between the new forming cells) of the first cell division was perpendicular to the electric vector of polarized light: nearly 50% of protoplasts had phragmoplasts at angles 80°-110° to electric vector compared with around 5% at e.g., angles 140°-170°. The polarization was induced by covering protoplasts in petri plates by plastic polarization filters, no ordered orientation of phragmoplasts was found at non-polarized light.<sup>95</sup> Phytochrome signaling is involved in the responses since effects of red light were reverted by far red light or were synergistic.<sup>90,93</sup> No effects of polarized UV or blue light were found for apical growth of fern *Adiantum capillus-veneris*,<sup>93</sup> but the effects of polarized blue and green light were present for growth of moss *Physcomitrella patens*.<sup>91</sup> Obviously, ordered localization of phytochromes (and probably of the other photoreceptors) is required for the reaction to polarized light, but phytochromes are generally known to be soluble proteins. The paradox is not solved yet; existence of membrane-bound fraction of phytochrome or/and interaction between phytochrome and membrane-associated phototropins was postulated (reviewed in ref. 94). Potential involvement of cytoskeleton is also suggested based on (1) experiments with cytoskeleton inhibitors for phototropic phytochrome-dependent responses (reviewed in ref. 94) and (2) involvement of actin-related protein ARPC4 in orientated growth of regenerating protoplasts of *Physcomitrella patens* under polarized white

light.<sup>96</sup> The development of biological polarity may imply further cytoskeleton reorganization and calcium waves/signaling, which are the typical elements in formation of polarity.

### Effects of Different Statistics of Photons on Light Perception

The main source of light existing in nature is solar radiation with properties of thermal light. Photons are emitted from numerous energy levels and the distribution of their number in a first approximation obeys Bose-Einstein statistics. The other well-known sources of light with different parameters are lasers. Distribution of photons emitted by lasers obeys Poisson statistics. Lasers provide a valuable tool for biological research and are widely used in technique due to their unique features (high possible power per a pulse, monochromatic light, high degree of polarization, low noise etc.). One of simple differences between the two statistics is that number of photons per unit of time is more uniform for lasers; functions for probabilities of distribution of photon number are different for thermal light and for lasers.<sup>97</sup>

Randomly occurring fluorescence, light from fires etc. usually are not discussed since having no known information/predictable effects for biological objects. Bioluminescence is widely spread (e.g., reviewed in ref. 98) though doesn't look different from the point of photon statistics and polarization. There are reports about super-Poisson statistics (typical for thermal light) of bacterial (*Photobacterium phosphoreum*) bioluminescence<sup>99</sup> and photon emission from cellular slime mold (*Dictyostelium discoideum*) during developmental processes,<sup>100</sup> which, however, need further investigation and confirmation.

An interesting question appears whether different sources of light may have different effects on biological systems due to statistics of number of photons. Potentially the same number of photons with the same energy, but distributed differently within the same time of illumination pulse may result in different effects. For example, conformations of proteins are subject to fast reversible and irreversible fluctuations due to thermal noise, interactions, changing microenvironment (at the scale of a few nanometers). The estimated upper limit for frequency of protein conformational changes is around  $10^6$  Hz.<sup>101</sup> Assuming just interaction of photons with one protein, it's conceivable to imagine several conformational levels for a protein and different effects after a multiphoton pulse of thermal or laser illumination.

A few experimental papers describe comparison of different sorts of illumination on visual perception paying attention to the statistics of photons (e.g. 102,103). Interesting results were obtained with retinal rods of *Xenopus*: the slope of response to Nd:YAG laser at 532 nm was steeper than for pseudothermal (the same statistics like thermal) light.<sup>36</sup> Photocurrent response of *Xenopus* rod cells was saturated by about 25,000 photons per a 30 ms pulse in the experiments; the difference between laser and pseudothermal light appeared after half-saturating amplitude of photocurrent. Relative photocurrent (normalized to saturating) was about 30–40% higher for laser source of photons in the range of illumination.<sup>36</sup> The observation is most likely connected to transduction chain species (the life of activated rhodopsin

is about 50–80 ms) and raises questions for most experiments with laser light. It might be possible that results with lasers and Poisson statistics require corrections when approximating for thermal sources of light.

## Outlook and Novel Technical Opportunities

The future directions in photobiology are bright and spread far outside the scope of the small review. Evident progress of optogenetics is expressed nowadays in potential medical applications. Further and deeper understanding of photobiological processes including leap to spatial nanoscale and temporal femtoscale in combination with new approaches of molecular biology and genetics needs also integrative and synthetic way of seeing. The new and more detailed picture with higher resolution will rise. More knowledge is gained from different species, so details of phototransduction may vary and leave plenty of space for future research.

Emerging new sources of light with different statistics of photons (lasers), light-emitting diodes with unusual spectral properties offer valuable tools for re-questioning old problems

and posing new ones. Recent interest in quantum dots was rewarded by opportunity to get single photons using quantum dots.<sup>104–106</sup> The source of single photons might be valuable for determining sensitivity of photoreception, for providing exact number of photons of certain energy (wavelength) and seems to be very promising for the future research.

## Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

## Acknowledgments

Unfortunately, I cannot number and thank here all colleagues who participated in numerous discussions within over 20 years of my fluctuating interest in the area, who inspired some ideas and thoughts, so I can just apologize for not citing all the relevant literature sources.

## Human and Animal Studies

All institutional and national guidelines for the care and use of laboratory animals were followed.

## References

- Govindjee, editor. Photosynthesis. Volume I. Energy Conversion by Plants and Bacteria. 1982. 799. Volume II. Development, Carbon Metabolism and Plant Productivity. Academic Press, NY. 1982. 580.
- Boyer PD. The ATP synthase—a splendid molecular machine. *Annu Rev Biochem* 1997; 66:717–49; PMID:9242922; <http://dx.doi.org/10.1146/annurev.biochem.66.1.717>
- McCarty RE, Evron Y, Johnson EA. THE CHLOROPLAST ATP SYNTHASE: A Rotary Enzyme? *Annu Rev Plant Physiol Mol Biol* 2000; 51:83–109; PMID:15012187; <http://dx.doi.org/10.1146/annurev.arplant.51.1.83>
- Alberty RA. Standard Gibbs free energy, enthalpy, and entropy changes as a function of pH and pMg for several reactions involving adenosine phosphates. *J Biol Chem* 1969; 244:3290–302; <http://www.jbc.org/content/244/12/3290.long>; PMID:4307313.
- Kammermeier H, Schmidt P, Jüngling E. Free energy change of ATP-hydrolysis: a causal factor of early hypoxic failure of the myocardium? *J Mol Cell Cardiol* 1982; 14:267–77; <http://www.sciencedirect.com/science/article/pii/00222888290205X>; PMID:7131563; [http://dx.doi.org/10.1016/0022-2888\(82\)90205-X](http://dx.doi.org/10.1016/0022-2888(82)90205-X).
- Chen M, Chory J, Fankhauser C. Light signal transduction in higher plants. *Annu Rev Genet* 2004; 38:87–117; PMID:15568973; <http://dx.doi.org/10.1146/annurev.genet.38.072902.092259>
- Hejide M, Ulm R. UV-B photoreceptor-mediated signalling in plants. *Trends Plant Sci* 2012; 17:230–7; PMID:22326562; <http://dx.doi.org/10.1016/j.tplants.2012.01.007>
- Lozier RH, Bogomolni RA, Stoerkenius W. Bacteriorhodopsin: a light-driven proton pump in *Halobacterium Halobium*. *Biophys J* 1975; 15:955–62; <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1334761/>; PMID:1182271; [http://dx.doi.org/10.1016/S0006-3495\(75\)85875-9](http://dx.doi.org/10.1016/S0006-3495(75)85875-9).
- Sineshchekov OA, Jung KH, Spudich JL. Two rhodopsins mediate phototaxis to low- and high-intensity light in *Chlamydomonas reinhardtii*. *Proc Natl Acad Sci U S A* 2002; 99:8689–94; PMID:12060707; <http://dx.doi.org/10.1073/pnas.122243399>
- Nagel G, Ollig D, Fuhrmann M, Kateriya S, Musti AM, Bamberg E, Hegemann P. Channelrhodopsin-1: a light-gated proton channel in green algae. *Science* 2002; 296:2395–8; PMID:12089443; <http://dx.doi.org/10.1126/science.1072068>
- Hegemann P. Algal sensory photoreceptors. *Annu Rev Plant Biol* 2008; 59:167–89; PMID:18444900; <http://dx.doi.org/10.1146/annurev.arplant.59.032607.092847>
- Lin JY. A user's guide to channelrhodopsin variants: features, limitations and future developments. *Exp Physiol* 2011; 96:19–25; PMID:20621963; <http://dx.doi.org/10.1113/expphysiol.2009.051961>
- Watanabe HC, Welke K, Sindhikara DJ, Hegemann P, Elstner M. Towards an understanding of channelrhodopsin function: simulations lead to novel insights of the channel mechanism. *J Mol Biol* 2013; 425:1795–814; PMID:23376098; <http://dx.doi.org/10.1016/j.jmb.2013.01.033>
- Fricke W, Akhilarova G, Wei W, Alexandersson E, Miller A, Kjellbom PO, Richardson A, Wojciechowski T, Schreiber L, Veselov D, et al. The short-term growth response to salt of the developing barley leaf. *J Exp Bot* 2006; 57:1079–95; PMID:16513814; <http://dx.doi.org/10.1093/jxb/erj095>
- Hecht S, Shlaer S, Pirenne MH. Energy, quanta, and vision. *J Gen Physiol* 1942; 25:819–40; <http://jgp.rupress.org/content/25/6/819.long>; PMID:19873316; <http://dx.doi.org/10.1085/jgp.25.6.819>.
- Dowling JE. *The Retina: An Approachable Part of the Brain*. Cambridge, MA: Belknap of Harvard UP. Print. Revised Edition. 2012. 384.
- Palczewski K. G protein-coupled receptor rhodopsin. *Annu Rev Biochem* 2006; 75:743–67; PMID:16756510; <http://dx.doi.org/10.1146/annurev.biochem.75.103004.142743>
- Terakita A. The opsins. *Genome Biol* 2005; 6:213; PMID:15774036; <http://dx.doi.org/10.1186/gb-2005-6-3-213>
- Wald G. The molecular basis of visual excitation. *Nature* 1968; 219:800–7; PMID:4876934; <http://dx.doi.org/10.1038/219800a0>
- Birge RR. Nature of the primary photochemical events in rhodopsin and bacteriorhodopsin. *Biochim Biophys Acta* 1990; 1016:293–327; PMID:2184895; [http://dx.doi.org/10.1016/0005-2728\(90\)90163-X](http://dx.doi.org/10.1016/0005-2728(90)90163-X)
- Birge RR, Barlow RB. On the molecular origins of thermal noise in vertebrate and invertebrate photoreceptors. *Biophys Chem* 1995; 55:115–26; PMID:7632872; [http://dx.doi.org/10.1016/0301-4622\(94\)00145-A](http://dx.doi.org/10.1016/0301-4622(94)00145-A)
- Land M, Chittka L. Vision. In: *The Insects: Structure and Function*, 5th Edition (eds. Simpson, S. J. and Douglas, A. E.). Cambridge: Cambridge University Press. 2013. 708–737.
- Briscoe AD, Chittka L. The evolution of color vision in insects. *Annu Rev Entomol* 2001; 46:471–510; PMID:11112177; <http://dx.doi.org/10.1146/annurev.ento.46.1.471>
- Stryer L. Exploring light and life. *J Biol Chem* 2012; 287:15164–73; PMID:22411992; <http://dx.doi.org/10.1074/jbc.X112.361436>
- Palczewski K, Kumasaka T, Hori T, Behnke CA, Motoshima H, Fox BA, Le Trong I, Teller DC, Okada T, Stenkamp RE, et al. Crystal structure of rhodopsin: A G protein-coupled receptor. *Science* 2000; 289:739–45; PMID:10926528; <http://dx.doi.org/10.1126/science.289.5480.739>
- Choe HW, Kim YJ, Park JH, Morizumi T, Pai EF, Krauss N, Hofmann KP, Scheerer P, Ernst OP. Crystal structure of metarhodopsin II. *Nature* 2011; 471:651–5; PMID:21389988; <http://dx.doi.org/10.1038/nature09789>
- Lamb TD. Gain and kinetics of activation in the G-protein cascade of phototransduction. *Proc Natl Acad Sci U S A* 1996; 93:566–70; <http://www.pnas.org/content/93/2/566.long>; PMID:8570596; <http://dx.doi.org/10.1073/pnas.93.2.566>.
- Lamb TD, Pugh EN Jr. Phototransduction, dark adaptation, and rhodopsin regeneration the proctor lecture. *Invest Ophthalmol Vis Sci* 2006; 47:5138–52; PMID:17122096; <http://dx.doi.org/10.1167/iovs.06-0849>
- Fesenko EE, Kolesnikov SS, Lyubarsky AL. Induction by cyclic GMP of cationic conductance in plasma membrane of retinal rod outer segment. *Nature* 1985; 313:310–3; PMID:2578616; <http://dx.doi.org/10.1038/313310a0>
- Fung BK-K, Hurley JB, Stryer L. Flow of information in the light-triggered cyclic nucleotide cascade of vision. *Proc Natl Acad Sci U S A* 1981; 78:152–6; <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC319009/>; PMID:6264430; <http://dx.doi.org/10.1073/pnas.78.1.152>.

31. Burns ME, Pugh EN Jr. Lessons from photoreceptors: turning off g-protein signaling in living cells. *Physiology* (Bethesda) 2010; 25:72-84; PMID:20430952; <http://dx.doi.org/10.1152/physiol.00001.2010>
32. Krispel CM, Chen D, Mellinger N, Chen YJ, Martemyanov KA, Quillinan N, Arshavsky VY, Wensel TG, Chen CK, Burns ME. RGS expression rate-limits recovery of rod photoreceptors. *Neuron* 2006; 51:409-16; PMID:16908407; <http://dx.doi.org/10.1016/j.neuron.2006.07.010>
33. Chen CK, Woodruff ML, Chen FS, Chen D, Fain GL. Background light produces a recoverin-dependent modulation of activated-rhodopsin lifetime in mouse rods. *J Neurosci* 2010; 30:1213-20; PMID:20107049; <http://dx.doi.org/10.1523/JNEUROSCI.4353-09.2010>
34. Sim N, Bessarab D, Jones CM, Krivitsky L. Method of targeted delivery of laser beam to isolated retinal rods by fiber optics. *Biomicroscop Opt Express* 2011; 2:2926-33; PMID:22076256; <http://dx.doi.org/10.1364/BOE.2.002926>
35. Cobbs WH, Pugh EN Jr. Kinetics and components of the flash photocurrent of isolated retinal rods of the larval salamander, *Ambystoma tigrinum*. *J Physiol* 1987; 394:529-72; <http://jpp.physoc.org/content/394/1/529.long>; PMID:2832596.
36. Sim N, Cheng MF, Bessarab D, Jones CM, Krivitsky LA. Measurement of photon statistics with live photoreceptor cells. *Phys Rev Lett* 2012; 109:113601; PMID:23005626; <http://dx.doi.org/10.1103/PhysRevLett.109.113601>
37. Rieke F, Baylor DA. Origin of reproducibility in the responses of retinal rods to single photons. *Biophys J* 1998; 75:1836-57; PMID:9746525; [http://dx.doi.org/10.1016/S0006-3495\(98\)77625-8](http://dx.doi.org/10.1016/S0006-3495(98)77625-8)
38. Doan T, Mendez A, Detwiler PB, Chen J, Rieke F. Multiple phosphorylation sites confer reproducibility of the rod's single-photon responses. *Science* 2006; 313:530-3; PMID:16873665; <http://dx.doi.org/10.1126/science.1126612>
39. Montell C. Visual transduction in *Drosophila*. *Annu Rev Cell Dev Biol* 1999; 15:231-68; PMID:10611962; <http://dx.doi.org/10.1146/annurev.cellbio.15.1.231>
40. Montell C. *Drosophila* visual transduction. *Trends Neurosci* 2012; 35:356-63; PMID:22498302; <http://dx.doi.org/10.1016/j.tins.2012.03.004>
41. Chader GJ, Weiland J, Humayun MS. Artificial vision: needs, functioning, and testing of a retinal electronic prosthesis. *Prog Brain Res* 2009; 175:317-32; PMID:19660665; [http://dx.doi.org/10.1016/S0079-6123\(09\)17522-2](http://dx.doi.org/10.1016/S0079-6123(09)17522-2)
42. Nirenberg S, Pandarinath C. Retinal prosthetic strategy with the capacity to restore normal vision. *Proc Natl Acad Sci U S A* 2012; 109:15012-7; PMID:22891310; <http://dx.doi.org/10.1073/pnas.1207035109>
43. Zemelman BV, Lee GA, Ng M, Miesenböck G. Selective photostimulation of genetically chARGed neurons. *Neuron* 2002; 33:15-22; PMID:11779476; [http://dx.doi.org/10.1016/S0896-6273\(01\)00574-8](http://dx.doi.org/10.1016/S0896-6273(01)00574-8)
44. Bi A, Cui J, Ma YP, Olshevskaya E, Pu M, Dizhoor AM, Pan ZH. Ectopic expression of a microbial-type rhodopsin restores visual responses in mice with photoreceptor degeneration. *Neuron* 2006; 50:23-33; PMID:16600853; <http://dx.doi.org/10.1016/j.neuron.2006.02.026>
45. Lagali PS, Balya D, Awatramani GB, Münch TA, Kim DS, Busskamp V, Cepko CL, Roska B. Light-activated channels targeted to ON bipolar cells restore visual function in retinal degeneration. *Nat Neurosci* 2008; 11:667-75; PMID:18432197; <http://dx.doi.org/10.1038/nn.2117>
46. Panda S, Provencio I, Tu DC, Pires SS, Rollag MD, Castrucci AM, Pletcher MT, Sato TK, Wiltshire T, Andahazy M, et al. Melanopsin is required for non-image-forming photic responses in blind mice. *Science* 2003; 301:525-7; PMID:12829787; <http://dx.doi.org/10.1126/science.1086179>
47. Melyan Z, Tarrtelin EE, Bellingham J, Lucas RJ, Hankins MW. Addition of human melanopsin renders mammalian cells photoreceptive. *Nature* 2005; 433:741-5; PMID:15674244; <http://dx.doi.org/10.1038/nature03344>
48. Panda S, Nayak SK, Campo B, Walker JR, Hogenech JB, Jegla T. Illumination of the melanopsin signaling pathway. *Science* 2005; 307:600-4; PMID:15681390; <http://dx.doi.org/10.1126/science.1105121>
49. Qiu X, Kumbalasingi T, Carlson SM, Wong KY, Krishna V, Provencio I, Berson DM. Induction of photosensitivity by heterologous expression of melanopsin. *Nature* 2005; 433:745-9; PMID:15674243; <http://dx.doi.org/10.1038/nature03345>
50. Ala-Laurila P, Donner K, Koskelainen A. Thermal activation and photoactivation of visual pigments. *Biophys J* 2004; 86:3653-62; PMID:15189862; <http://dx.doi.org/10.1529/biophysj.103.035626>
51. Barlow RB Jr., Kaplan E, Renninger GH, Saito T. Circadian rhythms in *Limulus* photoreceptors. I. Intracellular studies. *J Gen Physiol* 1987; 89:353-78; <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2215907/pdf/jg893353.pdf>; PMID:3559515; <http://dx.doi.org/10.1085/jgp.89.3.353>
52. Barlow RB, Birge RR, Kaplan E, Tallent JR. On the molecular origin of photoreceptor noise. *Nature* 1993; 366:64-6; PMID:8232538; <http://dx.doi.org/10.1038/366064a0>
53. Sampath AP, Baylor DA. Molecular mechanism of spontaneous pigment activation in retinal cones. *Biophys J* 2002; 83:184-93; PMID:12080111; [http://dx.doi.org/10.1016/S0006-3495\(02\)75160-6](http://dx.doi.org/10.1016/S0006-3495(02)75160-6)
54. Firsov ML, Donner K, Govardovskii VI. pH and rate of "dark" events in toad retinal rods: test of a hypothesis on the molecular origin of photoreceptor noise. *J Physiol* 2002; 559:837-46; PMID:11897853; <http://dx.doi.org/10.1113/jphysiol.2001.013207>
55. Rieke F, Baylor DA. Origin and functional impact of dark noise in retinal cones. *Neuron* 2000; 26:181-6; PMID:10798402; [http://dx.doi.org/10.1016/S0896-6273\(00\)81148-4](http://dx.doi.org/10.1016/S0896-6273(00)81148-4)
56. Fritsches KA, Brill RW, Warrant EJ. Warm eyes provide superior vision in swordfishes. *Curr Biol* 2005; 15:55-8; PMID:15649365; <http://dx.doi.org/10.1016/j.cub.2004.12.064>
57. Chang R. *Physical chemistry with applications to biological systems*. 1977. Macmillan Publishing. 538 p. ISBN: 9780023210204.
58. Lestas I, Vinnicombe G, Paulsson J. Fundamental limits on the suppression of molecular fluctuations. *Nature* 2010; 467:174-8; PMID:20829788; <http://dx.doi.org/10.1038/nature09333>
59. Können GP. *Polarised light in nature*. Publisher: Cambridge University Press. 1985. 196.
60. Pye DJ. *Polarised light in science and nature*. CRC Press. 2010. 124.
61. Cronin TW, Marshall J. Patterns and properties of polarized light in air and water. *Philos Trans R Soc Lond B Biol Sci* 2011; 366:619-26; PMID:21282165; <http://dx.doi.org/10.1098/rstb.2010.0201>
62. Stavenga DG, Wilts BD, Leertouwer HL, Hariyama T. Polarized iridescence of the multilayered elytra of the Japanese jewel beetle, *Chrysochroa fulgidissima*. *Philos Trans R Soc Lond B Biol Sci* 2011; 366:709-23; PMID:21282175; <http://dx.doi.org/10.1098/rstb.2010.0197>
63. Jewell SA, Vukusic P, Roberts NW. Circularly polarized colour reflection from helicoidal structures in the beetle *Plusiotis boucardi*. *New J Phys* 2007; 9:99; <http://dx.doi.org/10.1088/1367-2630/9/4/099>
64. Sweeney A, Jiggins C, Johnsen S. Insect communication: Polarized light as a butterfly mating signal. *Nature* 2003; 423:31-2; PMID:12721616; <http://dx.doi.org/10.1038/423031a>
65. Gerwick WH, Lang NJ. Structural, chemical and ecological studies on iridescence in *Iridaea* (Rhodophyta). *J Phycol* 1977; 13:121-7; <http://dx.doi.org/10.1111/j.1529-8817.1977.tb00567.x>
66. Pedersen M, Roomans GM, Hofsten A. von Blue iridescence and bromine in the cuticle of the red alga *Chondrus crispus* Stackh. *Bot Mar* 1980; 23:193-6
67. Pellegrini L, Pellegrini M. Iridescent bodies of *Cystoseira stricta* Sauvageau (Phaeophyta, Fucales): their fine structure, development and nature in vegetative cells. *Phycologia* 1982; 21:34-46; <http://dx.doi.org/10.2216/10031-8884-21-1-34.1>
68. Moody MF, Parriss JR. Discrimination of polarized light by octopus. *Nature* 1960; 186:839-40; PMID:14423793; <http://dx.doi.org/10.1038/186839a0>
69. Mäthger LM, Hanlon RT. Anatomical basis for camouflaged polarized light communication in squid. *Biol Lett* 2006; 2:494-6; PMID:17148271; <http://dx.doi.org/10.1098/rsbl.2006.0542>
70. Mäthger LM, Shashar N, Hanlon RT. Do cephalopods communicate using polarized light reflections from their skin? *J Exp Biol* 2009; 212:2133-40; PMID:19561202; <http://dx.doi.org/10.1242/jeb.020800>
71. Jordan TM, Partridge JC, Roberts NW. Non-polarizing broadband multilayer reflectors in fish. *Nat Photonics* 2012; 6:759-63; PMID:23160173; <http://dx.doi.org/10.1038/nphoton.2012.260>
72. Frisch K v. Die Polarisation des Himmelslichtes als orientierender Faktor bei den Tänzern der Bienen. *Experientia* (Basel) 1949; 5:142-8; <http://dx.doi.org/10.1007/BF02174424>
73. von Helversen O, Edrich W. Der Polarisationsempfänger im Bienenauge: ein Ultraviolettzeptor. *J Comp Physiol* 1974; 94:33-47; <http://dx.doi.org/10.1007/BF00610156>
74. Sakura M, Okada R, Aonuma H. Evidence for instantaneous e-vector detection in the honeybee using an associative learning paradigm. *Proc Biol Sci* 2012; 279:535-42; PMID:21733901; <http://dx.doi.org/10.1098/rspb.2011.0929>
75. Labhart T. Specialized photoreceptors at the dorsal rim of the honeybee's compound eye: polarization and angular sensitivity. *J Comp Physiol A Neuroethol Sens Neural Behav Physiol* 1980; 141:19-30; <http://dx.doi.org/10.1007/BF00611874>
76. Hawryshyn CW. Ultraviolet polarization vision in fishes: possible mechanisms for coding e-vector. *Philos Trans R Soc Lond B Biol Sci* 2000; 355:1187-90; PMID:11079395; <http://dx.doi.org/10.1098/rstb.2000.0664>
77. Waterman TH, Aoki K. E-vector sensitivity patterns in the goldfish optic tectum. *J Comp Physiol* 1974; 95:13-27; <http://dx.doi.org/10.1007/BF00624348>
78. Waterman TH, Hashimoto H. E-vector discrimination by the goldfish optic tectum. *J Comp Physiol* 1974; 95:1-12; <http://dx.doi.org/10.1007/BF00624347>
79. Nilsson DE, Labhart T, Meyer E. Photoreceptor design and optical properties affecting polarization sensitivity in ants and crickets. *J Comp Physiol A Neuroethol Sens Neural Behav Physiol* 1987; 161:645-58; <http://dx.doi.org/10.1007/BF00605006>
80. Labhart T, Meyer EP. Neural mechanisms in insect navigation: polarization compass and odometer. *Curr Opin Neurobiol* 2002; 12:707-14; PMID:12490263; [http://dx.doi.org/10.1016/S0959-4388\(02\)00384-7](http://dx.doi.org/10.1016/S0959-4388(02)00384-7)
81. Mueller KP, Labhart T. Polarizing optics in a spider eye. *J Comp Physiol A Neuroethol Sens Neural Behav Physiol* 2010; 196:335-48; PMID:20229246; <http://dx.doi.org/10.1007/s00359-010-0516-6>

82. Roberts NW, Porter ML, Cronin TW. The molecular basis of mechanisms underlying polarization vision. *Philos Trans R Soc Lond B Biol Sci* 2011; 366:627-37; PMID:21282166; <http://dx.doi.org/10.1098/rstb.2010.0206>
83. Wernet MF, Velez MM, Clark DA, Baumann-Klausener F, Brown JR, Klovstad M, Labhart T, Clandinin TR. Genetic dissection reveals two separate retinal substrates for polarization vision in *Drosophila*. *Curr Biol* 2012; 22:12-20; PMID:22177904; <http://dx.doi.org/10.1016/j.cub.2011.11.028>
84. Kalmijn AJ. The electric sense of sharks and rays. *J Exp Biol* 1971; 55:371-83; <http://jeb.biologists.org/content/55/2/371.long>; PMID:5114029.
85. Antipin NP, Krylov BV, Cherepnov VL. Topography of the ampullary system of *Raja clavata* and its role in electroorientation. *Neurophysiology* 1984; 16:628-33; <http://dx.doi.org/10.1007/BF01052682>
86. Gonzalez IF. The Electric Sense of the Thornback Ray, *Platyrhinoidis triseriata*: Linear Dynamic Range in Single-Unit Electrophysiological Recordings *in vivo* from the Afferent Nerve Fibers of the Ampullae of Lorenzini. Series: UC San Diego Electronic Theses and Dissertations; Degree: Ph. D., UC San Diego 2008; 104 p.; permalink: <https://escholarship.org/uc/item/8v03g541>
87. Johnsen S, Lohmann KJ. The physics and neurobiology of magnetoreception. *Nat Rev Neurosci* 2005; 6:703-12; PMID:16100517; <http://dx.doi.org/10.1038/nrn1745>
88. Singh J. Modern physics for engineers. John Wiley & Sons. 1999. 400.
89. Bünning E, Etzold H. Über die Wirkung von polarisiertem Licht auf keimende Sporen von Pilzen, Moosen und Farnen. *Ber Dtsch Bot Ges* 1958; 71:304-6;
90. Nebel BJ. Responses of moss protonemata to red and far-red polarized light: evidence for disc-shaped phytochrome photoreceptors. *Planta* 1969; 87:170-9; PMID:24504725; <http://dx.doi.org/10.1007/BF00386974>
91. Jenkins GI, Cove DJ. Phototropism and polarotropism of primary chloronemata of the moss *Physcomitrella patens*: responses of the wild-type. *Planta* 1983a; 158:357-64; PMID:24264756; <http://dx.doi.org/10.1007/BF00397338>
92. Jenkins GI, Cove DJ. Phototropism and polarotropism of primary chloronemata of the moss *Physcomitrella patens*: responses of mutant strains. *Planta* 1983b; 159:432-8; PMID:24258296; <http://dx.doi.org/10.1007/BF00392079>
93. Kadota A, Koyama M, Wada M, Furuya M. Action spectra for polarotropism and phototropism in protonemata of the fern *Adiantum capillus-veneris*. *Physiol Plant* 1984; 61:327-30; <http://dx.doi.org/10.1111/j.1399-3054.1984.tb06335.x>
94. Jaedicke JK. Molekulare Charakterisierung der cytoplasmatischen Phytochrom Funktion in *Physcomitrella patens*. Inaugural Dissertation Zur Erlangung des akademischen Grades eines Doktor der Naturwissenschaft (Dr. rer. nat.). 2012. Giessen. 158 p. [http://geb.uni-giessen.de/geb/volltexte/2012/8907/pdf/JaedickeKatharina\\_2012\\_05\\_04.pdf](http://geb.uni-giessen.de/geb/volltexte/2012/8907/pdf/JaedickeKatharina_2012_05_04.pdf)
95. Skripnikov AY, Shefer DD, Zrid Zh-P. Orientation of Cyto-Kinetic Apparatus of Moss Protoplasts *PHYSCOMITRELLA PATENS* under Direct Beam Influence of Plane-Polarized Light: New Model of Plant Development Biophysics. (In Russian, English abstract). *Problems of Contemporary Science and Practice, Vernadsky University* 2009; 6(20); <http://vernadsky.tstu.ru/en/vjpusk/2009/vjpusk-06.php> [http://vernadsky.tstu.ru/pdf/2009/06/rus\\_02\\_2009\\_06.pdf](http://vernadsky.tstu.ru/pdf/2009/06/rus_02_2009_06.pdf)
96. Perroud PF, Quatrano RS. The role of ARPC4 in tip growth and alignment of the polar axis in filaments of *Physcomitrella patens*. *Cell Motil Cytoskeleton* 2006; 63:162-71; PMID:16450411; <http://dx.doi.org/10.1002/cm.20114>
97. Mandel L, Wolf E. Optical coherence and quantum optics. Cambridge University Press. 1995. 1166.
98. Haddock SH, Moline MA, Case JF. Bioluminescence in the sea. *Ann Rev Mar Sci* 2010; 2:443-93; PMID:21141672; <http://dx.doi.org/10.1146/annurev-marine-120308-081028>
99. Kobayashi M, Devaraj B, Inaba H. Observation of super-Poisson statistics from bacterial (*Photobacterium phosphoreum*) bioluminescence during the early stage of cell proliferation. *Phys Rev E Stat Phys Plasmas Fluids Relat Interdiscip Topics* 1998; 64:2129-33; <http://dx.doi.org/10.1103/PhysRevE.57.2129>
100. Kobayashi M, Inaba H. Photon statistics and correlation analysis of ultraweak light originating from living organisms for extraction of biological information. *Appl Opt* 2000; 39:183-92; PMID:18337887; <http://dx.doi.org/10.1364/AO.39.000183>
101. Chakrapani S, Auerbach A. A speed limit for conformational change of an allosteric membrane protein. *Proc Natl Acad Sci U S A* 2005; 102:87-92; PMID:15618401; <http://dx.doi.org/10.1073/pnas.0406777102>
102. Teich MC, Prucnal PR, Vannucci G, Breton ME, McGill WJ. Multiplication noise in the human visual system at threshold: 1. Quantum fluctuations and minimum detectable energy. *J Opt Soc Am* 1982a; 72:419-31; PMID:7077429; <http://dx.doi.org/10.1364/JOSA.72.000419>
103. Teich MC, Prucnal PR, Vannucci G, Breton ME, McGill WJ. Multiplication noise in the human visual system at threshold. 3. The role of non-Poisson quantum fluctuations. *Biol Cybern* 1982b; 44:157-65; PMID:7115793; <http://dx.doi.org/10.1007/BF00344271>
104. Santori C, Fattal D, Vuckovic J, Solomon GS, Yamamoto Y. Single-photon generation with InAs quantum dots. *New J Phys* 2004; 6:89; <http://dx.doi.org/10.1088/1367-2630/6/1/089>
105. Ates S, Agha I, Gulinatti A, Rech I, Badolato A, Srinivasan K. Improving the performance of bright quantum dot single photon sources using temporal filtering via amplitude modulation. *Sci Rep* 2013; 3:1397; PMID:23466520; <http://dx.doi.org/10.1038/srep01397>
106. Holmes MJ, Choi K, Kako S, Arita M, Arakawa Y. Room-Temperature Triggered Single Photon Emission from a III-Nitride Site-Controlled Nanowire Quantum Dot. *Nano Lett* 2014; 14:982-6; PMID:24422516; <http://dx.doi.org/10.1021/nl404400d>