

MEETING ABSTRACT

Open Access

The neuroprotective effects of electrolyzed reduced water and its model water containing molecular hydrogen and Pt nanoparticles

Hanxu Yan¹, Taichi Kashiwaki², Takeki Hamasaki², Tomoya Kinjo¹, Kiichiro Teruya^{1,2}, Shigeru Kabayama³, Sanetaka Shirahata^{1,2*}

From 22nd European Society for Animal Cell Technology (ESACT) Meeting on Cell Based Technologies Vienna, Austria. 15-18 May 2011

Background

Human brain is the biggest energy consuming tissue in human body. Although it only represents 2% of the body weight, it receives 20% of total body oxygen consumption and 25% of total body glucose utilization. For that reason, brain is considered to be the most vulnerable part of human body against the reactive oxygen species (ROS), a by-product of aerobic respiration. Oxidative stress is directly related to a series of brain dysfunctional disease such as Alzheimer's disease, Parkinson's disease etc. Electrolyzed reduced water (ERW) is a functional drinking water containing a lot of molecular hydrogen and a small amount of platinum nanoparticles (Pt NPs, Table 1). ERW is known to scavenge ROS and protect DNA from oxidative damage [1]. We previously showed that ERW was capable of extending lifespan of *Caenorhabditis elegans* by scavenging ROS [2]. Molecular hydrogen could scavenge ROS and protected brain from oxidative stress [3]. Pt NPs are also a new type of multi-functional ROS scavenger [4].

Materials and methods

In this research, we used TI-200S ERW derived from 2 mM NaOH solution produced by a batch type electrolysis device and model waters containing molecular hydrogen and synthetic Pt NPs of 2-3 nm sizes as research models of ERW to examine the anti-oxidant capabilities of ERW on several kinds of neural cells such as PC12, N1E115, and serum free mouse embryo (SFME) cells. We pretreated the ERW and 200 μ M

H₂O₂ and examined the neuroprotective effects of ERW on PC12, N1E115 and SFME cells, using WST-8 method. We also examined the intracellular ROS scavenging effects of ERW on N1E115 cells after pretreated cells with ERW and H₂O₂ using DCFH-DA. We checked the protective effects of ERW on mitochondria and cytoplasm by Rh123 and Fuo-3 AM stain. We also examined the ATP production of SFME cells after pretreated with ERW and H₂O₂ by Bioluminescence Assay Kit. Finally, we used dissolved hydrogen (DH) and Pt NPs as research models to examine their neuroprotective effects.

Results

ERW significantly reduced the cell death induced by H₂O₂ pretreatment (Figure 1). ERW also scavenged the intracellular ROS and prevented the decrease of mitochondrial membrane potential and ATP production induced by ROS. We also examined the neuroprotective effects of molecular hydrogen and Pt NPs and showed that both molecular hydrogen and Pt NPs contributed to the neuroprotective effects of ERW.

Conclusion

The results suggest that ERW is beneficial for the prevention and alleviation of oxidative stress-induced human neurodegenerative diseases.

Author details

¹Graduate School of Systems Life Sciences, Kyushu University, 6-10-1 Hakozaki, Higashi-ku, Fukuoka 812-8581, Japan. ²Department of Bioscience and Biotechnology, Faculty of Agriculture, Kyushu University, Fukuoka 812-8581, Japan. ³Nihon Trim Co. Ltd., 1-8-34 Oyodonaka, Kita-ku, Osaka 531-0076, Japan.


* Correspondence: sirahata@grt.kyushu-u.ac.jp

¹Graduate School of Systems Life Sciences, Kyushu University, 6-10-1 Hakozaki, Higashi-ku, Fukuoka 812-8581, Japan

Full list of author information is available at the end of the article

Table 1 Characteristics of the water samples.The characteristics of water samples were determined immediately after the preparation of ERW. ERW, electrolyzed reduced water; CW, activated charcoal-treated water. The pH values were shown as average \pm standard deviation (N = 5). The values of DH, DO and Pt NPs were shown the minimum and maximum values after 5 independent measurements.

	MQ (NaOH)	TI-200 ERW	TI-9000 CW	TI-9000 ERW
pH	11.3 \pm 0.1	11.6 \pm 0.1	7.9 \pm 0.1	9.6 \pm 0.2
Dissolved Hydrogen (mM)	0	0.2– 0.45	0	0.1– 0.25
Dissolved Oxygen (μ M)	0	3.1– 78.1	0	0– 21.9
Pt NPs (nM)	0	0.5– 12.8	0	0– 3.6
Redox potential value (mV)	+ 350	-659	-	-



Published: 22 November 2011

References

- Shirahata S, Kabayama S, Nakano M, Miura T, Kusumoto K, Gotoh M, Hayashi H, Otsubo K, Morisawa S, Katakura Y: **Electrolyzed-reduced water scavenges active oxygen species and protects DNA from oxidative damage.** *Biophys Biochem Res Commun* 1997, **234**:269-274.
- Yan H, Tian H, Kinjo T, Hamasaki T, Tomimatsu K, Nakamichi N, Teruya K, Kabayama S, Shirahata S: **Extension of the lifespan of *Caenorhabditis elegans* by the use of electrolyzed reduced water.** *Biosci Biotech Biochem* 2010, **74**:2011-2015.
- Ohsawa I, Ishikawa M, Takahashi K, Watanabe M, Nishimaki K, Yamagata K, Katsura K, Katayama Y, Asoh S, Ohta S: **Hydrogen acts as a therapeutic antioxidant by selectively reducing cytotoxic oxygen radicals.** *Nature Med* 2007, **13**:688-694.
- Hamasaki T, Kashiwagi T, Imada T, Nakamichi N, Aramaki S, Toh K, Morisawa S, Shimakoshi H, Hisaeda Y, Shirahata S: **Kinetic analysis of superoxide anion radical-scavenging and hydroxyl radical-scavenging activities of platinum nanoparticles.** *Langmuir* 2008, **24**:7354-7364.

doi:10.1186/1753-6561-5-S8-P69

Cite this article as: Yan *et al.*: The neuroprotective effects of electrolyzed reduced water and its model water containing molecular hydrogen and Pt nanoparticles. *BMC Proceedings* 2011 **5**(Suppl 8):P69.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

