

Letter to the Editor

Comment on “Detecting Key Genes Regulated by miRNAs in Dysfunctional Crosstalk Pathway of Myasthenia Gravis”

Rozen Le Panse

Centre de Recherche en Myologie GH Pitié-Salpêtrière, 105 boulevard de l'Hôpital, 75013 Paris, France

Correspondence should be addressed to Rozen Le Panse; rozen.lepanse@upmc.fr

Received 7 June 2016; Accepted 6 December 2016; Published 8 February 2017

Academic Editor: Rituraj Purohit

Copyright © 2017 Rozen Le Panse. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

I am concerned by an article “Detecting Key Genes Regulated by miRNAs in Dysfunctional Crosstalk Pathway of Myasthenia Gravis” by Cao et al., 2015 [1].

In this study, the authors used data from public databases to compare microarray data from the thymus and miRNome data from peripheral blood mononuclear cells (PBMCs). They analyzed the regulation of mRNAs from the thymus by miRNAs from peripheral blood mononuclear cells: two different “tissues.” Of course, they ended up with results but one can really wonder about the meaning of these results.

Moreover, in their manuscript, Cao et al. compiled thymic microarray data altogether while data are from different categories of patients (seropositive (for anti-AChR antibodies) and seronegative patients) (Le Panse et al., *The Journal of Immunology*, 2006) [2]. In AChR-myasthenia gravis patients, the thymus is involved in the disease but not in seronegative myasthenia gravis patients.

and the role of CC chemokine ligand 21 in thymic hyperplasia,” *Journal of Immunology*, vol. 177, no. 11, pp. 7868–7879, 2006.

Competing Interests

The author declares no competing interests.

References

- [1] Y. Cao, J. Wang, H. Zhang et al., “Detecting key genes regulated by miRNAs in dysfunctional crosstalk pathway of myasthenia gravis,” *BioMed Research International*, vol. 2015, Article ID 724715, 10 pages, 2015.
- [2] R. Le Panse, G. Cizeron-Clairac, J. Bismuth, and S. Berrih-Aknin, “Microarrays reveal distinct gene signatures in the thymus of seropositive and seronegative myasthenia gravis patients