

IL-25 promotes Th2 bias by upregulating IL-4 and IL-10 expression of decidual $\gamma\delta$ T cells in early pregnancy

YUAN ZHANG^{1*}, YING WANG^{2*}, MING-QING LI¹, JIE DUAN¹, DENG-XUAN FAN¹ and LI-PING JIN^{1,3}

¹Laboratory for Reproductive Immunology, Hospital of Obstetrics and Gynecology, Fudan University, Shanghai Medical College, Shanghai 200011; ²Department of Obstetrics and Gynecology, Qilu Hospital of Shandong University, Jinan, Shandong 250012; ³Clinical and Translational Research Center, Shanghai First Maternity and Infant Hospital, Tongji University School of Medicine, Shanghai 201204, P.R. China

Received July 21, 2017; Accepted October 25, 2017

DOI: 10.3892/etm.2017.5638

Abstract. Decidual immune cells (DICs), consisting of both innate and adaptive immune cells, have a pivotal role in maintaining immune tolerance for normal pregnancy. Our previous study demonstrated that interleukin (IL)-25 stimulates the proliferation of decidual stromal cells (DSCs) in an autocrine manner. However, the role of IL-25 in functional regulation of DICs is largely unknown. Flow cytometry was used to analyze the expression of IL-25 and its receptor (IL-17RB) in DICs, and the effect of IL-25 on the expression of Ki-67, IL-4, IL-10, interferon (IFN)- γ and transforming growth factor (TGF)- β in decidual $\gamma\delta$ T cells. In addition, ELISA assays were performed to detect the secretion of IL-10 and TGF- β in decidual $\gamma\delta$ T cells. The present findings indicated that decidual CD56^{bright}CD16⁻ natural killer (NK) cells, natural killer T (NKT) cells, regulatory T (Treg) cells, CD3⁺ T cells, macrophages and $\gamma\delta$ T cells co-expressed IL-25 and IL-17RB, particularly $\gamma\delta$ T cells. Recombinant human (rh) IL-25 protein upregulated the expression of Ki-67, IL-4, and IL-10, but downregulated the expression of IFN- γ in $\gamma\delta$ T cells; however, anti-human IL-25 or IL-17RB neutralizing antibody reversed these effects. These data suggest that IL-25 may promote IL-10 production by $\gamma\delta$ T cells as well as the proliferation of $\gamma\delta$ T cells, and possibly

forms a positive feedback loop to maintain a T helper 2 cell bias at the maternal-fetal interface and further contributes to the maintenance of successful pregnancy.

Introduction

Normal pregnancy is equivalent to allogeneic transplantation that requires maternal immune tolerance for the development of embryo (1). At the maternal-fetal interface, apart from over 70% CD56^{bright}CD16⁻ natural killer (NK) cells, a few macrophages, CD3⁺ T cells, dendritic cells but not B cells are also enriched to compose the decidual immune cells (DICs), thereby modulating immune response during pregnancy. As has been determined, T helper 2 (Th2) bias at the maternal-fetal interface, characterized by interleukin (IL)-4, IL-5, and IL-10 secretion, is pivotal for immune tolerance. In contrast, Th1/Th2 imbalance can lead to pathological pregnancy, such as miscarriage or preeclampsia (2-4).

$\gamma\delta$ T cells, composed of γ and δ chains, are abundant in epithelial tissues, including reproductive tract, skin, and intestine, but only make up a fraction of the peripheral and organic lymphocytes. As a type of innate-like lymphocytes, unlike $\alpha\beta$ T cells $\gamma\delta$ T cells function without major histocompatibility complex restriction as well as CD4 or CD8 co-receptors, although a small proportion of human $\gamma\delta$ T cells express CD4 (5) or CD8 (6). Because of the widespread distribution of $\gamma\delta$ T cells, they are involved in a variety of disease conditions such as infection, autoimmunity, cancer, and miscarriage (7-9). Decidual $\gamma\delta$ T cells, expressing neither CD4 nor CD8 (double negative) markers, are increased significantly during pregnancy (10), and create a Th2 bias at the maternal-fetal interface by secreting IL-10 and transforming growth factor (TGF)- β (11-13). Moreover, a previous study has demonstrated that decidual $\gamma\delta$ T cells promote the proliferation and invasion of trophoblast cells as well as suppress the apoptosis via IL-10 (14).

As a member of the IL-17 family, IL-25, also known as IL-17E, seems to participate extensively in various immune-related diseases, such as allergy, asthma, and enteric nematode infection (15). On the one hand, IL-25 has a pathogenic effect on Th2-type diseases such as asthma; on the other hand, it plays a therapeutic role in Th1-type diseases such as

Correspondence to: Dr Deng-Xuan Fan, Laboratory for Reproductive Immunology, Hospital of Obstetrics and Gynecology, Fudan University, Shanghai Medical College, 413 Zhao Zhou Road, Shanghai 200011, P.R. China
E-mail: fdx_1228@163.com

Professor Li-Ping Jin, Clinical and Translational Research Center, Shanghai First Maternity and Infant Hospital, Tongji University School of Medicine, 2699 West Gaoke Road, Shanghai 201204, P.R. China
E-mail: jinlp01@163.com

*Contributed equally

Key words: interleukin-25, interleukin-17RB, $\gamma\delta$ T cells, type 2 T helper cells, proliferation, pregnancy

Crohn disease and Th17-type diseases such as autoimmune encephalomyelitis, thereby serving as a double-edged sword. For example, in asthma and airway high-reactivity disease, IL-25 secreted by airway epithelial cells or eosinophilic or basophilic granulocyte not only participates in the innate immune response to produce IL-4, IL-5, and IL-13, but also has a role in adaptive immune response to induce naïve T cells differentiation into Th2-type cells. In inflammatory bowel disease, antigen stimulates intestinal epithelial cells and macrophages to produce IL-25, which inhibits Th1 and Th17-type immune response by reducing the secretion of IL-12 and IL-23 by antigen-presenting cells, and protects the intestinal mucosal cells from damage. IL-17RB, a principal recognized IL-25 receptor, functions widely on diverse cells, including eosinophils, mast cells, monocytes, and T cells.

Moreover, our previous study has elucidated that human chorionic gonadotropin (hCG) derived from trophoblasts upregulates the expression of IL-25/IL-17RB in decidual stromal cells (DSCs), followed by further stimulating DSCs proliferation by activating c-Jun N-terminal kinase (JNK) and protein kinase B (AKT) signals, thus finally contributing to a successful pregnancy (16). However, whether DICs express IL-25/IL-17RB, and even whether IL-25 influences the local immune response at the maternal-fetal interface, are still unknown. Consequently, the present study was undertaken to investigate the expression of IL-25/IL-17RB in DICs and related functions in decidual $\gamma\delta$ T cells.

Materials and methods

Tissue collection. All tissue samples (n=42) were collected with informed consent according to the requirements of the Research Ethics Committee in the Obstetrics and Gynecology Institute, Fudan University Shanghai Medical College (Shanghai, China). All subjects completed informed consent forms for collection of tissue samples. In addition, this study was specifically approved by the Research Ethics Committee. Decidual samples were obtained from normal pregnant women (age 29.24±3.17 years; gestational age 8.11±1.37 weeks; mean ± standard deviation) whose pregnancies were terminated for non-medical reasons.

Cell isolation and culture. The tissues from the first-trimester pregnancy were placed immediately into cold Dulbecco's modified Eagle's medium (DMEM; Thermo Fisher Scientific, Inc., Waltham, MA, USA), transported to the laboratory within 1 h after surgery, and washed with Hank balanced salt solution for isolation of DICs. The DICs were isolated according to our previous procedures (17).

Isolation and culture of $\gamma\delta$ T cells. Decidual $\gamma\delta$ T cells were isolated using a magnetic activated cell-sorting human $\gamma\delta$ T cell positive selection isolation kit (Miltenyi Biotec, Bergisch Gladbach, Germany) in accordance with the manufacturer's instruction. The purity of $\gamma\delta$ T cells was evaluated by flow cytometry with fluorescein isothiocyanate- $\gamma\delta$ TCR (eBioscience; Thermo Fisher Scientific, Inc.) antibody.

Monoclonal antibodies. Phycoerythrin (PE)-conjugated anti-IL-25 monoclonal antibody (mAb) and allophycocyanin

(APC)-conjugated anti-IL-17RB mAb were purchased from R&D (R&D Systems, Inc., Minneapolis, MN, USA). Fluorescein-isothiocyanate-conjugated anti-CD56, CD14, CD25, and $\gamma\delta$ TCR mAbs, PE-conjugated anti-IL-4, TGF- β mAbs, PE-Cy5.5-conjugated anti-CD3, IL-10 mAbs, and APC-conjugated anti-interferon (IFN)- γ , Ki67 mAbs were purchased from eBioscience. APC-cy7-conjugated anti-CD45 and PE-cy7-conjugated anti-CD4 mAbs were purchased from BioLegend, Inc. (San Diego, CA, USA).

IL-25/IL-17RB expression in DICs. DICs were collected and washed with phosphate buffered saline. After blocking with 10% fetal bovine serum (Hyclone; GE healthcare, Logan, UT, USA), DICs were divided into several parts for detecting the percentage of IL-25 or IL-17RB positive cells in different immune cells, including CD45⁺CD3⁺CD56⁺ NK cells, CD45⁺CD3⁺CD56⁺ natural killer T (NKT) cells, CD45⁺CD3⁺ T cells, CD45⁺CD3⁺ $\gamma\delta$ T cells, CD45⁺CD14⁺ macrophages, and CD45⁺CD3⁺CD4⁺CD25⁺ regulatory T (Treg) cells. The recovered cells were immediately stained using a standard immunofluorescence assay with the different surface mAbs in darkness for 45 min at room temperature, followed by washing with phosphate buffered saline to remove unbound antibody, fixing, permeabilizing, and then incubated with anti-IL-25 antibody for another 1 h at room temperature. The isotype control was used as a negative control. After incubation, the cells were washed and analyzed immediately by flow cytometer (FACS Beckman; BD Biosciences, Franklin Lakes, NJ, USA).

Expression of cytokines in $\gamma\delta$ T cells. The isolated $\gamma\delta$ T cells were seeded in 24-well plates (1x10⁶ cells per well) and treated with different concentrations of recombinant human (rh) IL-25 (R&D Systems, Inc.) (0, 0.1, 1, 10 ng/ml), 1 ng/ml rhIL-25 plus 10 μ g/ml anti-IL-25 neutralizing antibody (anti-IL-25; R&D Systems, Inc.), or 1 ng/ml rhIL-25 plus 10 μ g/ml anti-IL-17RB neutralizing antibody (anti-IL-17RB; R&D Systems, Inc.) for 24 h. Then, after treatment with PMA+BFA+ionomycin for 4 h, we detected the expression of IL-10, TGF- β , IL-4, and IFN- γ in $\gamma\delta$ T cells by flow cytometry following the aforementioned staining procedure.

Ki-67 expression in $\gamma\delta$ T cells. $\gamma\delta$ T cells were seeded in 24-well plates (1x10⁶ cells per well) and treated with different concentrations of rhIL-25 (0, 0.1, 1, 10 ng/ml), 10 ng/ml rhIL-25 plus 10 μ g/ml anti-IL-25, or 10 ng/ml rhIL-25 plus 10 μ g/ml anti-IL-17RB for 24 h. Subsequently, we used flow cytometry assay to detect the expression of Ki-67 in $\gamma\delta$ T cells following the aforementioned intracellular cytokine staining procedure.

Enzyme-linked immunosorbent assay (ELISA). $\gamma\delta$ T cells were seeded in 24-well plates (1x10⁶ cells per well) and treated with 1 ng/ml rhIL-25, 1 ng/ml rhIL-25 plus 10 μ g/ml anti-IL-25, 1 ng/ml rhIL-25 plus 10 μ g/ml anti-IL-17RB for 48 h. The culture supernatants were then harvested, centrifuged to remove cellular debris, and stored at -80°C. The concentrations of IL-10 and TGF- β were detected by ELISA according to the manufacturer's instructions (eBioscience; Thermo Fisher Scientific, Inc.).




Figure 1. (A and B) Decidual $\gamma\delta$ T cells highly co-express IL-25 and IL-17RB. We used (A) flow cytometry to detect the percentage of (B) IL-25 or IL-17RB positive cells in different immune cells at the maternal-fetal interface (n=12). Data are presented as mean \pm standard deviation. Experiments were repeated three times. IL, interleukin; Treg, decidual regulatory T cells; NK, decidual natural killer cells; NKT, decidual natural killer T cells; $\gamma\delta$ T, decidual $\gamma\delta$ TCR cells; CD3, decidual CD3⁺ T cells; M, decidual macrophages.

Statistical analysis. The data were analyzed using the least significant difference test after one-way analysis of variance (SPSS version 17.0; SPSS, Inc., Chicago, IL, USA). All values are shown as mean \pm standard deviation. $P < 0.05$ was considered to indicate a statistically significant difference.

Results

Decidual $\gamma\delta$ T cells highly co-express IL-25 and IL-17RB. Because of the complex components of DICs, we examined the expression of IL-25/IL-17RB in different types of cells by flow cytometry, including NK cells, NKT cells, CD3⁺ T cells, $\gamma\delta$ T cells, macrophages, and Treg cells. The results of flow cytometer showed that 15.7 \pm 1.6% NK cells, 20.6 \pm 3.2% NKT cells, 29.1 \pm 2.6% Treg cells, 21.3 \pm 2.8% CD3⁺ T cells,

32.9 \pm 3.4% macrophages, and 32.0 \pm 3.1% $\gamma\delta$ T cells expressed IL-25, whereas 40.4 \pm 2.0% NK cells, 66.8 \pm 6.2% NKT cells, 71.3 \pm 2.4% Treg cells, 33.0 \pm 3.6% CD3⁺ T cells, 19.7 \pm 2.2% macrophages, and 70.9 \pm 1.6% $\gamma\delta$ T cells expressed IL-17RB (Fig. 1). Taken together, the expression levels of IL-25/IL-17RB in $\gamma\delta$ T cells were higher than other cells.

IL-25 promotes the proliferation of decidual $\gamma\delta$ T cells. Because of the high expression level of IL-25/IL-17RB in $\gamma\delta$ T cells, we intended to determine the effects of IL-25 on the functions of $\gamma\delta$ T cells. First, we isolated $\gamma\delta$ T cells from DICs by magnetic activated cell sorting, and the purity was examined by flow cytometry, which was approximately 90% (Fig. 2A). Next, we treated $\gamma\delta$ T cells with different concentrations of rhIL-25, rhIL-25 plus anti-IL-25, or rhIL-25 plus anti-IL-17RB for 24 h,




Figure 2. IL-25 promotes the proliferation of decidual $\gamma\delta$ T cells. (A) Isolation and purity identification of $\gamma\delta$ T cells. DICs were collected and isolated for $\gamma\delta$ T cells by magnetic activated cell sorting, the purity of which was detected by flow cytometry. (B) After treatment with different concentrations of rhIL-25 (0, 0.1, 1, 10 ng/ml), 10 ng/ml rhIL-25 plus 10 μ g/ml anti-IL-25 or 10 ng/ml rhIL-25 plus 10 μ g/ml anti-IL-17RB for 24 h, we used flow cytometry assay to examine the expression of proliferation-related molecule Ki-67 in $\gamma\delta$ T cells (n=9). IL, interleukin; IL-25, recombinant human (rh) IL-25; anti-IL-25, anti-IL-25 neutralizing antibody; anti-IL-17RB, anti-IL-17RB neutralizing antibody. *P<0.05 compared with control. Data are presented as mean \pm standard deviation.

after which we detected the expression of proliferation-related molecule Ki-67 in $\gamma\delta$ T cells. As shown, IL-25 upregulated the percentage of Ki-67 positive $\gamma\delta$ T cells, whereas anti-IL-25 or anti-IL-17RB abrogated this effect (Fig. 2B). These results indicated that IL-25 stimulates the proliferation of $\gamma\delta$ T cells remarkably, and may lead to the accumulation of $\gamma\delta$ T cells in decidua.

IL-25 stimulates the production of IL-4 and IL-10, and suppresses the production of IFN- γ in $\gamma\delta$ T cells. To explore the effect of IL-25 on the expression of cytokine in decidual $\gamma\delta$ T cells, we treated these cells with rhIL-25, rhIL-25 plus anti-IL-25, and rhIL-25 plus anti-17RB for 24 h, and then detected the expression of IL-10, TGF- β , IL-4, and IFN- γ in $\gamma\delta$ T cells by flow cytometry (Fig. 3). As shown, IL-25 could upregulate the percentage of IL-10 positive $\gamma\delta$ T cells in a concentration-dependent manner, whereas anti-IL-25 or anti-IL-17RB could abrogate this effect (Fig. 3A and D). Meanwhile, IL-25 had no influence on the percentage of TGF- β positive $\gamma\delta$ T cells (Fig. 3A and E). Additionally, IL-25 also promoted the expression of IL-4 in $\gamma\delta$ T cells (Fig. 3B and F), while inhibiting the expression of IFN- γ (Fig. 3C and G). IL-25 obviously increased the ratio of IL-4 to IFN- γ (Fig. 3H). In addition, the ELISA results showed that IL-25 could facilitate the secretion of IL-10 by $\gamma\delta$ T cells, which was abrogated

by anti-IL-25 or anti-IL-17RB, but had no influence on that of TGF- β (Fig. 3I and J). Therefore, these results suggest that IL-25 significantly increases the expression of IL-10 and IL-4, decreases the level of IFN- γ in $\gamma\delta$ T cells, and thus is beneficial to form a Th2 immune bias at the maternal-fetal interface.

Discussion

Accounting for the majority of decidual CD3⁺ T cells, $\gamma\delta$ T cells, composed of V γ 9/V δ 1 chains, are functionally polarized to be Th2 cells, whereas human peripheral lymphocytes preferentially expressing V γ 9/V δ 2 chains function as Th1 cells (18-21). Based on previous research, chemokine CXCL16/CXCR6 at the maternal-fetal interface recruits $\gamma\delta$ T cells to the local decidua, the proportion of which increases significantly higher than that of the peripheral blood. In addition, the number of $\gamma\delta$ T cells in the peripheral blood and decidua of normal early pregnancy is remarkably more than that during nonpregnancy (22). Decidual $\gamma\delta$ T cells express high levels of IL-10 and TGF- β , low levels of TNF- α and IFN- γ , but almost no IL-2 and IL-4, contributing to normal pregnancy. Interestingly, we have previously discovered that decidual $\gamma\delta$ T cells promote the proliferation and invasion of trophoblast cells as well as inhibit apoptosis via IL-10, thus facilitating the formation of placenta (14). Currently, two classical mechanisms are




Figure 3. IL-25 stimulates the production of IL-4 and IL-10, and suppresses the production of IFN- γ in $\gamma\delta$ T cells. (A and B) Detection of the expression of several cytokines in $\gamma\delta$ T cells. Cells received treatment with different concentrations of rhIL-25 (0, 0.1, 1, 10 ng/ml), 1 ng/ml rhIL-25 plus 10 μ g/ml anti-IL-25, 1 ng/ml rhIL-25 plus 10 μ g/ml anti-IL-17RB for 24 h. IL, interleukin; IFN, interferon.




Figure 3. Continued. IL-25 stimulates the production of IL-4 and IL-10, and suppresses the production of IFN- γ in $\gamma\delta$ T cells. (C-H) Detection of the expression of several cytokines in $\gamma\delta$ T cells. After treatment with different concentrations of rhIL-25 (0, 0.1, 1, 10 ng/ml), 1 ng/ml rhIL-25 plus 10 μ g/ml anti-IL-25, 1 ng/ml rhIL-25 plus 10 μ g/ml anti-IL-17RB for 24 h, the expression of (D-G) IL-10, TGF- β , IL-4, and IFN- γ in $\gamma\delta$ T cells was detected. In addition, the ratio of (H) IL-4 to IFN- γ in $\gamma\delta$ T cells was also elevated obviously after treatment with rhIL-25, n=12; *P<0.05 compared with control. (I and J) The effect of IL-25 on the expression of IL-10 in $\gamma\delta$ T cells was also confirmed. We used 1 ng/ml rhIL-25, 1 ng/ml rhIL-25 plus 10 μ g/ml anti-IL-25, 1 ng/ml rhIL-25 plus 10 μ g/ml anti-IL-17RB to treat $\gamma\delta$ T cells for 48 h, after which the secretion levels of IL-10 and TGF- β by $\gamma\delta$ T cells were detected by enzyme-linked immunosorbent assay. n=9; *P<0.05 compared with rhIL-25 treatment group. Data are presented as mean \pm standard deviation. IL, interleukin; IFN, interferon; TGF, transforming growth factor.

thought to induce immune tolerance by decidual $\gamma\delta$ T cells, either exerting immunosuppression directly through cytokine production or inducing the differentiation of naïve CD4⁺ T cells into inhibitory/regulatory T cells via IL-10 and TGF- β indirectly (18).

Growing evidence has shown that IL-25 not only stimulates human umbilical vein endothelial cell proliferation and the length, number, and area of microvessel structures in a concentration-dependent manner, but also promotes angiogenesis (23,24). However, downregulation of IL-25 in villus from patients with recurrent miscarriages may alter the local Th1/Th2 ratio (25), giving IL-25 an important role in normal pregnancy. In this study, we found that IL-25/IL-17RB were widely expressed in both the innate immune cells and adaptive immune cells of DICs, such as NK cells, NKT cells, CD3⁺ T cells, macrophages, and Treg cells, especially $\gamma\delta$ T cells, indicating that decidual $\gamma\delta$ T cells are not only a source of IL-25 at the maternal-fetal interface, but also one of the target cells of IL-25. Then, we further investigated the effect of IL-25 on the function of decidual $\gamma\delta$ T cells, finding that IL-25 could promote the proliferation of decidual $\gamma\delta$ T cells as well as the secretion of Th2 cytokines, especially IL-10. However, the mechanism by which IL-25 promotes decidual $\gamma\delta$ T cells proliferation is currently unknown, which needs to be further explored in the future.

IL-10, an anti-inflammatory cytokine, is critical for preventing inflammatory and autoimmune pathologies. Because of the deficient IL-10 expression in CBA x DBA/2 abortion mice, supplementation of exogenous IL-10 could reduce the embryo absorption rate (26). Likewise, the expression of IL-4 and IL-10 in decidual T cells of patients who have had an abortion is lower than that during normal pregnancy, whereas the expression of IFN- γ and IL-2 is in contrast (27), indicating that Th2 bias at the maternal-fetal interface is critical for a successful pregnancy. The reduction of IL-4 and IL-10 during pregnancy also leads to preeclampsia, fetal growth restriction, premature labor, and other conditions (28). In addition, it has been discovered that trophoblast cells promote the proliferation of $\gamma\delta$ T cells to secrete IL-10 and TGF- β through CXCL16, thus in turn increasing the proliferation and invasion of trophoblasts and eventually contributing to the development of placenta and the maintenance of normal pregnancy.

In summary, based on the aforementioned and our previous study results, some molecules at the maternal-fetal interface, such as CXCL16, may recruit peripheral $\gamma\delta$ T cells to local decidua, which not only produce Th2 cytokines (e.g., IL-10, IL-4) but also reduce the production of Th1 cytokines (e.g., IFN- γ). In addition, IL-25 secreted by various cells (e.g., DSCs and DICs) accelerates the proliferation of $\gamma\delta$ T cells, whereas the apoptosis of which could be inhibited by thymic stromal lymphopoietin (TSLP) (29), resulting in a positive feedback loop and regulatory Th2 bias at the maternal-fetal interface.

Acknowledgements

This study was supported by the National Basic Research Program of China (no. 2015CB943300), the National Natural Science Foundation of China (NSFC) (no. 81671460), the Program for Shanghai leaders, the Program of Shanghai

Outstanding Academic Leader (no. 15XD1500900), and the Foundations from Shanghai Science and Technology Committee (no. 134119a4300) to Li-Ping Jin.

References

1. Wegmann TG, Lin H, Guilbert L and Mosmann TR: Bidirectional cytokine interactions in the maternal-fetal relationship: Is successful pregnancy a TH2 phenomenon? *Immunol Today* 14: 353-356, 1993.
2. Laird SM, Tuckerman EM, Cork BA, Linjawi S, Blakemore AI and Li TC: A review of immune cells and molecules in women with recurrent miscarriage. *Hum Reprod Update* 9: 163-174, 2003.
3. Yamada H, Morikawa M, Kato EH, Shimada S, Kobashi G and Minakami H: Pre-conceptual natural killer cell activity and percentage as predictors of biochemical pregnancy and spontaneous abortion with normal chromosome karyotype. *Am J Reprod Immunol* 50: 351-354, 2003.
4. Saito S and Sakai M: Th1/Th2 balance in preeclampsia. *J Reprod Immunol* 59: 161-173, 2003.
5. Spits H, Paliard X, Vandekerckhove Y, van Vlasselaer P and de Vries JE: Functional and phenotypic differences between CD4⁺ and CD4⁻ T cell receptor-gamma delta clones from peripheral blood. *J Immunol* 147: 1180-1188, 1991.
6. Carding SR and Egan PJ: Gamma delta T cells: Functional plasticity and heterogeneity. *Nat Rev Immunol* 2: 336-345, 2002.
7. Goodman T and Lefrançois L: Expression of the gamma-delta T-cell receptor on intestinal CD8⁺ intraepithelial lymphocytes. *Nature* 333: 855-858, 1988.
8. Chen ZW: Comparative biology of gamma delta T cells. *Sci Prog* 85: 347-358, 2002.
9. Hayday AC: Gammadelta T cells and the lymphoid stress-surveillance response. *Immunity* 31: 184-196, 2009.
10. Mincheva-Nilsson L, Baranov V, Yeung MM, Hammarström S and Hammarström ML: Immunomorphologic studies of human decidua-associated lymphoid cells in normal early pregnancy. *J Immunol* 152: 2020-2032, 1994.
11. Arck PC, Ferrick DA, Steele-Norwood D, Croitoru K and Clark DA: Regulation of abortion by gamma delta T cells. *Am J Reprod Immunol* 37: 87-93, 1997.
12. Arck PC, Ferrick DA, Steele-Norwood D, Croitoru K and Clark DA: Murine T cell determination of pregnancy outcome: I. Effects of strain, alphabeta T cell receptor, gammadelta T cell receptor, and gammadelta T cell subsets. *Am J Reprod Immunol* 37: 492-502, 1997.
13. Clark DA, Flanders KC, Hirte H, Dasch JR, Coker R, McAnulty RJ and Laurent GJ: Characterization of murine pregnancy decidua transforming growth factor beta. I. Transforming growth factor beta 2-like molecules of unusual molecular size released in bioactive form. *Biol Reprod* 52: 1380-1388, 1995.
14. Fan DX, Duan J, Li MQ, Xu B, Li DJ and Jin LP: The decidual gamma-delta T cells up-regulate the biological functions of trophoblasts via IL-10 secretion in early human pregnancy. *Clin Immunol* 141: 284-292, 2011.
15. Fort MM, Cheung J, Yen D, Li J, Zurawski SM, Lo S, Menon S, Clifford T, Hunte B, Lesley R, *et al*: IL-25 induces IL-4, IL-5, and IL-13 and Th2-associated pathologies in vivo. *Immunity* 15: 985-995, 2001.
16. Wang Y, Zhang Y, Li MQ, Fan DX, Wang XH, Li DJ and Jin LP: Interleukin-25 induced by human chorionic gonadotropin promotes the proliferation of decidual stromal cells by activation of JNK and AKT signal pathways. *Fertil Steril* 102: 257-263, 2014.
17. Fu Q, Tao Y, Piao H, Du MR and Li DJ: Trophoblasts and decidual stromal cells regulate decidual NK cell functions via interaction between collagen and LAIR-1. *Am J Reprod Immunol* 71: 368-378, 2014.
18. Nagaeva O, Jonsson L and Mincheva-Nilsson L: Dominant IL-10 and TGF-beta mRNA expression in gammadeltaT cells of human early pregnancy decidua suggests immunoregulatory potential. *Am J Reprod Immunol* 48: 9-17, 2002.
19. Mincheva-Nilsson L, Hammarström S and Hammarström ML: Human decidua leukocytes from early pregnancy contain high numbers of gamma delta+ cells and show selective down-regulation of alloreactivity. *J Immunol* 149: 2203-2211, 1992.
20. Szekeres-Bartho J, Barakonyi A, Miko E, Polgar B and Palkovics T: The role of gamma/delta T cells in the fetomaternal relationship. *Semin Immunol* 13: 229-233, 2001.

21. Barakonyi A, Kovacs KT, Miko E, Szereday L, Varga P and Szekeres-Bartho J: Recognition of nonclassical HLA class I antigens by gamma delta T cells during pregnancy. *J Immunol* 168: 2683-2688, 2002.
22. Heyborne KD, Cranfill RL, Carding SR, Born WK and O'Brien RL: Characterization of gamma delta T lymphocytes at the maternal-fetal interface. *J Immunol* 149: 2872-2878, 1992.
23. Corrigan CJ, Wang W, Meng Q, Fang C, Wu H, Reay V, Lv Z, Fan Y, An Y, Wang YH, *et al*: T-helper cell type 2 (Th2) memory T cell-potentiating cytokine IL-25 has the potential to promote angiogenesis in asthma. *Proc Natl Acad Sci USA* 108: 1579-1584, 2011.
24. Wang W, Fan YQ, Lv Z, Yao XJ, Wang W, Huang KW, Meng Q, Fang CL, Lee TH, Corrigan CJ, *et al*: Interleukin-25 promotes basic fibroblast growth factor expression by human endothelial cells through interaction with IL-17RB, but not IL-17RA. *Clin Exp Allergy* 42: 1604-1614, 2012.
25. Giannubilo SR, Landi B, Pozzi V, Sartini D, Cecati M, Stortoni P, Corradetti A, Saccucci F, Tranquilli AL and Emanuelli M: The involvement of inflammatory cytokines in the pathogenesis of recurrent miscarriage. *Cytokine* 58: 50-56, 2012.
26. Chaouat G, Assal Meliani A, Martal J, Raghupathy R, Elliott JF, Mosmann T and Wegmann TG: IL-10 prevents naturally occurring fetal loss in the CBA x DBA/2 mating combination, and local defect in IL-10 production in this abortion-prone combination is corrected by in vivo injection of IFN-tau. *J Immunol* 154: 4261-4268, 1995.
27. Jin LP, Fan DX, Zhang T, Guo PF and Li DJ: The costimulatory signal upregulation is associated with Th1 bias at the maternal-fetal interface in human miscarriage. *Am J Reprod Immunol* 66: 270-278, 2011.
28. Chatterjee P, Chiasson VL, Bounds KR and Mitchell BM: Regulation of the anti-inflammatory cytokines interleukin-4 and interleukin-10 during pregnancy. *Front Immunol* 5: 253, 2014.
29. Duan J, Jiang XP, Li MQ, Fan DX, Wang Y, Li DJ and Jin LP: Thymic stromal lymphopoietin suppresses the apoptosis of decidual gamma-delta T cells via regulation of the signal transduction and activation of transcription 3/caspase-3 signaling pathway. *Am J Reprod Immunol* 70: 464-471, 2013.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.