



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



## Experimental infection of IS/885/00-like infectious bronchitis virus in specific pathogen free and commercial broiler chicks



Faez Awad<sup>a,b</sup>, Rajesh Chhabra<sup>a</sup>, Anne Forrester<sup>a</sup>, Julian Chantrey<sup>a</sup>, Matthew Baylis<sup>a,c</sup>, Stephane Lemiere<sup>d</sup>, Hussein Aly Hussein<sup>e</sup>, & Kannan Ganapathy<sup>a,\*</sup>

<sup>a</sup> University of Liverpool, Leahurst Campus, Neston, Cheshire, CH64 7TE, UK

<sup>b</sup> University of Omar Al-Mukhtar, Faculty of Veterinary Medicine, Al-Bayda, Libya

<sup>c</sup> NIHR Health Protection Research Unit in Emerging and Zoonotic Infections, Liverpool, UK

<sup>d</sup> Merial S.A.S., 29 avenue Tony Garnier, Lyon cedex 07, France

<sup>e</sup> Cairo University, Department of Virology, Faculty of Veterinary Medicine, Cairo, Egypt

### ARTICLE INFO

#### Article history:

Received 19 May 2015

Received in revised form 19 December 2015

Accepted 3 January 2016

#### Keywords:

Infectious bronchitis virus

IBV IS/885/00-like

Cystic oviduct

Head swelling

Chicks

### ABSTRACT

Pathogenesis of an IS/885/00-like (IS/885) strain of variant infectious bronchitis virus (IBV) was examined in one day old specific pathogen free (SPF) and commercial broiler chicks. Chicks were humanely euthanized at 3, 6, 9, 12, 15, 21 and 28 days post infection (dpi) for necropsy examination, and tissues were collected for histopathology and virus detection by reverse transcription polymerase chain reaction (RT-PCR). Respiratory clinical signs and gross lesions consisting of tracheal caseous exudate and plugs, and swollen kidneys (with or without) urate deposits were observed in SPF and broiler chicks. The onset of disease developed more slowly and were of lesser severity in broiler compared to SPF chicks, reflecting the inhibitory effects of the IBV maternal-antibodies in the broiler chicks or genetic/strain susceptibility, or both. Head swelling was observed in one infected broiler chick at 15 dpi and the virus was recovered by RT-PCR and isolation. In the IS/885-infected SPF chicks, cystic oviducts were found in two female chicks. IS/885 was isolated from the cystic fluid. Using ELISA, low to moderate levels of the antibodies to IBV was detected in the SPF compared to broiler infected chicks.

© 2016 Elsevier Ltd. All rights reserved.

### 1. Introduction

Avian infectious bronchitis is a common, highly contagious and economically important viral disease of chickens caused by a coronavirus (Cavanagh, 2007). A large number of serotypes and genotypes of infectious bronchitis virus (IBV) strains have been reported worldwide (de Wit et al., 2011a; Jackwood, 2012). This appears to be due to constant changes in the spike-protein of IBV strains resulting in the emergence of new IBV variants (Cavanagh, 2007). In the winter of 2000, a severe outbreak of renal disease occurred in several broiler farms in Israel. An IBV strain, designated IS/885/00, was isolated from the kidneys of these flocks (Meir et al., 2004).

In the Middle East and North Africa, the circulation of Massachusetts, 793B, QX, Dutch strains (Abdel-Moneim et al., 2006; Bourogaa et al., 2009; Amin et al., 2012; Boroomand et al., 2012) and Q1 (Ganapathy et al., 2015) have been reported. In addition, an increasing number of strains closely related to IS/885/00 have been reported in Israel (Meir

et al., 2004), Egypt (Abdel-Moneim et al., 2012), Iraq (Mahmood et al., 2011), Libya (Awad et al., 2014a) and other Middle East countries (Ganapathy et al., 2015). To date, through our diagnostic services, we have detected this strain in samples received from France, Ukraine and Pakistan. These strains were detected from broiler and layer flocks experiencing respiratory distress, renal lesions and high mortality (Meir et al., 2004; Selim et al., 2013; Awad et al., 2014a).

The pathogenesis and host immune responses to several IBV strains are known, such as those of M41 (Crinion and Hofstad, 1972; Butcher et al., 1990), Beaudette (Geilhausen et al., 1973), Australian T-strain (Chong and Apostolov, 1982; Ignjatovic et al., 2002), Moroccan G strain (El-Houadfi et al., 1986; Ambali and Jones, 1990), 793B (Dhinakar Raj and Jones, 1996; Boroomand et al., 2012), QX (Wang et al., 1998; Terregino et al., 2008; Ganapathy et al., 2012), It-02 (Dolz et al., 2012), Q1 (Yu et al., 2001; Toffan et al., 2013) and more recently Brazilian IBV variant (USP-10 and USP-50) (Chacón et al., 2014).

Despite the high circulation of IS/885/00-like strains in the Middle East and North Africa, and further detection in France, Ukraine and Pakistan, to date, there is no published information on the pathogenesis of this increasingly important variant IBV. To better understand the IS/

\* Corresponding author.

E-mail address: [gana@liverpool.ac.uk](mailto:gana@liverpool.ac.uk) (K. Ganapathy).



Fig. 1. Periocular swelling of an IS/885-infected broiler chick at 15 dpi.

885 strain, a series of experiment was performed to investigate the pathogenesis in specific pathogen free (SPF) and commercial broiler chicks. Clinical signs, gross and microscopic lesions, virus detection and humoral antibody responses were evaluated.

## 2. Materials and methods

### 2.1. Virus

The virus used throughout this study (referred to here as IS/885) was received as third passage allantoic fluid (AF) from the Istituto Zooprofilattico Sperimentale delle Venezie, Padova, Italy. The virus had been isolated from a recent outbreak of high mortality and respiratory disease complex in broiler flocks in Egypt. Initial isolation was carried out in the virology laboratory at Cairo University, Egypt, and the AF was submitted to the Italian laboratory. There, the AF went through three further passages in embryonated chicken eggs (ECEs) and was shown to be negative for avian influenza virus (AIV) and Newcastle disease virus (NDV) and positive for IBV by RT-PCR. Positive IBV amplicons were submitted for commercial sequencing (Source BioScience) of a 393 bp region of the hypervariable S1 region (Cavanagh et al., 1999). Sequence (GenBank Accession: awaiting) analysis of the IS/885 inoculum showed 90% nucleotide identity to the Israeli strain IS/885/00 (Meir et al., 2004) using BLAST (NCBI database).

At the University of Liverpool, the virus received a further passage in 9 to 11 day-old SPF ECE. The viral titre was determined by titration in ECE and calculated as previously described (Reed and Muench, 1938) to provide  $10^{4.66}$  EID<sub>50</sub>/ml. The AF was free of NDV, AIV, avian metapneumovirus (aMPV), infectious laryngotracheitis virus (ILT), infectious bursal disease virus (IBDV) and fowl adenovirus (FAdV). The inoculum was also free of bacterial contamination when tested using blood and MacConkey agars, and no mycoplasmas were detected either by culture or PCR.

### 2.2. Eggs and chicks

Fertile eggs from SPF White Leghorn chickens (Lohmann Animal Health, Cuxhaven, Germany) were incubated and hatched in our facilities. Day-old commercial broiler chicks with IBV maternally-derived antibodies (MDA) were obtained from a commercial hatchery. The parent flock of the broiler chicks had been vaccinated with a live IBV

H120 + D274 vaccine at 3 weeks old and an inactivated IBV M41 vaccine was administered four weeks before transfer to laying farms. Chicks were kept in an isolation unit (University of Liverpool) throughout the experiment and reared on deep litter with water and feed provided *ad libitum*. The chicks were raised on wood shavings as is practiced in the commercial farms. Feed and water were free of any antibiotics. All of the experimental procedures were undertaken after the approval of the University of Liverpool ethical review committee and according to the UK legislation on the use of animals for experiments, as permitted under the project license PPL 112 40/3723.

## 3. Experimental design

### 3.1. Experiment 1 (Expt 1): infection of SPF chicks

Seventy one-day-old SPF chicks were randomly divided into two groups, consisting of 45 and 25 chicks in the infected and control group respectively. The chicks in the infected group were inoculated oculonasally with 0.1 ml of virus-positive AF and those in the control group with virus-free AF.

### 3.2. Experiment 2 (Expt 2): infection of commercial broiler chicks

Seventy one-day-old commercial broiler chicks were divided into two groups and inoculated as per Experiment 1 described above.

For both experiments, clinical signs were observed daily throughout the experimental period. At 3, 6, 9, 12, 15, 21 and 28 dpi, five infected and three control chicks were randomly selected and euthanized to evaluate the gross lesions. Tissue samples of trachea, lung, caecal tonsils and kidney were collected individually and frozen at  $-70^{\circ}\text{C}$  for virus detection by RT-PCR. In addition, pieces of trachea and kidney were fixed in 10% buffered formalin for histopathology. Blood samples were collected from 8 randomly selected chicks at 0, 15, 21 and 28 dpi from the SPF and at 0, 3, 6, 9, 15, 21 and 28 dpi from the broiler chicks to monitor antibody responses.

### 3.3. Gross and microscopic examinations

All euthanised and found dead chicks were necropsied and examined for gross lesions. The upper part of the trachea and kidney tissues





Fig. 2. Cystic dilation of the oviduct of an IS/885-infected SPF female chick at 28 dpi.



**Fig. 3.** Hematoxylin and eosin staining of the trachea and kidney (magnification at 400×). Panels (a) –trachea – and (d) kidney correspond to control chick tissues taken at 3 and 9 days of age respectively. (b), Extensive epithelial deciliation, with severe lymphocyte and heterophil infiltration of trachea of SPF chick at 3 dpi. (c), Mild epithelial deciliation, with moderate lymphocyte and heterophil infiltrate and mild epithelial hyperplasia of broiler chicks at 3 dpi. (e), Moderate to severe lymphocyte and mild heterophil interstitial infiltration in kidney of SPF chick at 9 dpi. (f), Mild to moderate lymphocyte and heterophil interstitial infiltration in kidney of broiler chick at 9 dpi.

were fixed in 10% buffered formalin, embedded in paraffin and sections were cut for hematoxylin and eosin (H&E) staining. Trachea and kidney lesions were examined and scored as follows: 0 = no change, 1 = mild, 2 = moderate, 3 = severe (Chen et al., 1996).

**3.4. Processing of tissue samples**

Individual tissue samples of the trachea (Tr), lung (L), caecal tonsil (CT) and kidney (Kid) were collected from five chicks in each group.

Each tissue was dipped in a sterile bijoux containing 1.5 ml of Eagles serum-free minimum essential medium with glutamine, streptomycin [50 mg/ml] and penicillin [50 IU/ml] and stored at –70 °C until required.

Tissues were homogenized with sterile sand and 1.5 ml of TOC medium using a pestle and mortar. Homogenates were subjected to freeze–thaw three times and clarified by centrifugation at 3000 g for 15 min (Awad et al., 2014b). Supernatant from the tissues was stored at –70 °C until required. RNA was extracted from tissue samples

**Table 1**  
Histopathology tracheal lesion scores in SPF chicks infected with IBV IS/885 strain.

| Tracheal lesions        | SPF (days post infection) |    |                        |                        |           |                        |           |  |
|-------------------------|---------------------------|----|------------------------|------------------------|-----------|------------------------|-----------|--|
|                         | 3                         | 6  | 9                      | 12                     | 15        | 21                     | 28        |  |
| Epithelial deciliation  | 3.0 ± 0.0 <sup>a</sup>    | ND | 3.0 ± 0.0              | 2.8 ± 0.2              | 0.5 ± 0.2 | 0.0 ± 0.0 <sup>a</sup> | 0.0 ± 0.0 |  |
| Epithelial degeneration | 0.5 ± 0.2                 | ND | 1.2 ± 0.2              | 1.0 ± 0.3              | 1.0 ± 0.4 | 0.5 ± 0.2              | 0.5 ± 0.5 |  |
| Decrease mucous cells   | 1.2 ± 0.2 <sup>a</sup>    | ND | 2.2 ± 0.2              | 2.6 ± 0.2              | 2.0 ± 0.4 | 1.2 ± 0.5              | 1.5 ± 0.5 |  |
| Heterophil infiltration | 2.0 ± 0.0 <sup>a</sup>    | ND | 0.5 ± 0.2              | 0.8 ± 0.3              | 0.0 ± 0.0 | 0.0 ± 0.0 <sup>a</sup> | 0.0 ± 0.0 |  |
| Epithelial hyperplasia  | 1.7 ± 0.2                 | ND | 2.7 ± 0.2 <sup>a</sup> | 2.6 ± 0.2 <sup>a</sup> | 1.2 ± 0.2 | 1.0 ± 0.0 <sup>a</sup> | 1.5 ± 0.5 |  |
| Lymphoid infiltration   | 2.0 ± 0.0 <sup>a</sup>    | ND | 1.5 ± 0.2              | 1.4 ± 0.2              | 2.0 ± 0.0 | 1.7 ± 0.2              | 2.0 ± 0.0 |  |

ND: not done; Data are expressed as the mean histopathological lesion score ± SEM (n = 5). Tracheal lesions scores as follows: no change (0), mild (1), moderate (2) or severe (3). For each time point and lesions, different superscripts for infected SPF (Table 2) and broiler (Table 3) chicks indicate significant differences between them (p < 0.05). Absence of a letter indicates that there were no significant differences (p > 0.05).

**Table 2**  
Histopathology tracheal lesion scores in broiler chicks infected with IBV IS/885 strain.

| Tracheal lesions        | Broiler (days post infection) |           |                        |                        |           |                        |           |
|-------------------------|-------------------------------|-----------|------------------------|------------------------|-----------|------------------------|-----------|
|                         | 3                             | 6         | 9                      | 12                     | 15        | 21                     | 28        |
| Epithelial deciliation  | 1.6 ± 0.4 <sup>b</sup>        | 2.8 ± 0.2 | 2.4 ± 0.2              | 2.4 ± 0.4              | 2.2 ± 0.4 | 2.0 ± 0.3 <sup>b</sup> | 0.0 ± 0.0 |
| Epithelial degeneration | 0.8 ± 0.2                     | 1.2 ± 0.2 | 1.4 ± 0.2              | 1.8 ± 0.3              | 1.0 ± 0.0 | 1.4 ± 0.2              | 0.2 ± 0.2 |
| Decrease mucous cells   | 0.2 ± 0.2 <sup>b</sup>        | 1.8 ± 0.5 | 1.8 ± 0.2              | 2.0 ± 0.3              | 2.8 ± 0.2 | 2.2 ± 0.3              | 0.5 ± 0.2 |
| Heterophil infiltration | 0.8 ± 0.3 <sup>b</sup>        | 0.8 ± 0.3 | 1.6 ± 0.5              | 1.0 ± 0.3              | 0.6 ± 0.2 | 1.2 ± 0.3 <sup>b</sup> | 0.2 ± 0.2 |
| Epithelial hyperplasia  | 1.2 ± 0.2                     | 1.4 ± 0.2 | 1.6 ± 0.2 <sup>b</sup> | 1.0 ± 0.0 <sup>b</sup> | 1.0 ± 0.4 | 2.0 ± 0.0 <sup>b</sup> | 1.5 ± 0.2 |
| Lymphoid infiltration   | 1.0 ± 0.0 <sup>b</sup>        | 1.2 ± 0.2 | 1.8 ± 0.2              | 1.6 ± 0.2              | 1.6 ± 0.4 | 2.2 ± 0.2              | 1.5 ± 0.2 |

ND: not done; Data are expressed as the mean histopathological lesion score ± SEM (n = 5). Tracheal lesions scores as follows: no change (0), mild (1), moderate (2) or severe (3). For each time point and lesions, different superscripts for infected SPF (Table 2) and broiler (Table 3) chicks indicate significant differences between them (p < 0.05). Absence of a letter indicates that there were no significant differences (p > 0.05).

using the QIAamp viral RNA Mini Kit following manufacturer's instructions (Qiagen, UK). RT-PCR was carried out as previously described (Cavanagh et al., 1999; Worthington et al., 2008). Briefly, detection of the IBV genome was achieved by amplifying a 393 bp region of the S1 gene using amplification and sequencing primers common for most of the known IBV strains. This protocol has been previously validated using reference IBV genotypes (Worthington et al., 2008). The initial PCR used primers SX1 + and SX2 -. The amplicon was further amplified in a second internal PCR that used primers SX3 +, SX4 -. The amplified DNA product from the IBV positive head swelling swab and cystic oviduct were treated with 0.15 µl Exonuclease 1 (EXO) and 0.99 µl shrimp alkaline phosphatase (SAP) at 37 °C for 30 min followed by 80 °C for 10 min to remove any extraneous material. The purified product, together with positive sense primer (forward direction using primer SX3 +), were submitted to an external laboratory (Source Bioscience sequencing, Nottingham, UK) for analysis of the partial S1 gene sequences.

### 3.5. Serological assay

Sera were tested using commercial enzyme-linked immunosorbent assay (ELISA) kits (IDEXX, Hoofddorp, The Netherlands) according to the manufacturer's instructions. Haemagglutination inhibition (HI) testing was performed using four haemagglutination units (HAU) as previously described (Alexander and Chettle, 1977). The virus strains used as antigens for the HI test were IBV M41 and 793B (Animal Health Service, Deventer, The Netherlands). Haemagglutination antigens of IS/885 were prepared in our laboratory based on methods previously described (King and Hopkins, 1983; Alexander and Chettle, 1977). IBV HI titres were expressed as log<sub>2</sub> values of the highest reciprocal of the dilution that showed HI. Titres equal to or greater than 3 log<sub>2</sub> were considered positive.

### 3.6. Statistical analysis

Data for histopathological lesions were analyzed using the Mann-Whitney U test. The mean ELISA and HI antibody titres of the sera of

infected SPF and broiler chicks tested with different IBV antigens were compared and analyzed using one way analysis of variance (ANOVA) to test for a significant overall effect, followed by Tukey's test to identify which means were significantly different from each other. All analyses were conducted using GraphPad Prism 6.0 (<http://www.graphpad.com/scientific-software/prism/>).

## 4. Results

### 4.1. Clinical signs

No clinical signs or mortalities were observed in the control groups of either experiment.

In the SPF chicks, clinical signs were first observed at 1 dpi, which included mild tracheal râles, sneezing, coughing, head shaking and eye scratching. Gaspings, wheezing and open mouth breathing were seen in 4–5 birds between 4 to 13 dpi. After 13 dpi, birds with mild respiratory signs and wet droppings (white or milky faeces) were observed. The clinical signs resolved by 18 dpi. Of 45 chicks, one bird died at 8 dpi (2.2%).

For the commercial broiler chicks, the clinical signs were similar to those observed in SPF chicks but with a lower severity and lasted for a longer period. Mild clinical signs of tracheal râles, sneezing, coughing and head shaking were first observed from 3 dpi and lasted until 22 dpi. Of the 45 birds, three died; one at 12 dpi and two at 19 dpi (6.7%). Apart from respiratory signs, one bird showed swelling of the head, with a foamy ocular and nasal discharge at 14 dpi. The swelling increased by 15 dpi and the periocular tissues were also affected (Fig 1). For welfare reasons, this bird was euthanized for necropsy and sampling was carried out. A sterile dissection was performed to assess any gross lesions underneath the skin. Subcutaneous swabs were taken for virus detection by RT-PCR, VI and for bacterial culture. Virus detection was performed in our laboratory while the bacterial culture, isolation and identification was carried out by a bacteriology diagnostic laboratory (School of Veterinary Science, University of Liverpool).

**Table 3**  
Histopathology kidney lesion scores in SPF chicks infected with IBV IS/885 strain.

| Kidney lesions          | SPF (days post infection) |                        |           |           |                        |           |           |
|-------------------------|---------------------------|------------------------|-----------|-----------|------------------------|-----------|-----------|
|                         | 3                         | 6                      | 9         | 12        | 15                     | 21        | 28        |
| Epithelial degeneration | 0.0 ± 0.0                 | 0.0 ± 0.0              | 0.6 ± 0.2 | 0.2 ± 0.2 | 0.0 ± 0.0              | 0.0 ± 0.0 | 0.0 ± 0.0 |
| Ducto-tubular dilation  | 0.4 ± 0.2                 | 0.4 ± 0.2              | 0.2 ± 0.2 | 0.0 ± 0.0 | 0.0 ± 0.0              | 0.0 ± 0.0 | 0.0 ± 0.0 |
| Heterophil infiltration | 0.2 ± 0.2                 | 1.0 ± 0.0 <sup>a</sup> | 0.2 ± 0.2 | 0.0 ± 0.0 | 0.0 ± 0.0              | 0.0 ± 0.0 | 0.0 ± 0.0 |
| Lymphoid infiltration   | 0.0 ± 0.0                 | 0.0 ± 0.0              | 1.0 ± 0.0 | 1.6 ± 0.2 | 1.0 ± 0.0              | 0.6 ± 0.2 | 0.4 ± 0.2 |
| Epithelial regeneration | 0.0 ± 0.0                 | 0.0 ± 0.0              | 0.2 ± 0.2 | 0.2 ± 0.2 | 0.8 ± 0.2              | 0.0 ± 0.0 | 0.0 ± 0.0 |
| Epithelial hyperplasia  | 0.2 ± 0.2                 | 0.0 ± 0.0              | 0.6 ± 0.2 | 0.2 ± 0.2 | 1.0 ± 0.0 <sup>a</sup> | 0.0 ± 0.0 | 0.0 ± 0.0 |
| Lymphoid nodules        | 0.0 ± 0.0                 | 0.0 ± 0.0              | 0.6 ± 0.2 | 0.8 ± 0.2 | 1.0 ± 0.0 <sup>a</sup> | 0.4 ± 0.2 | 0.8 ± 0.2 |

Data are expressed as the mean histopathological lesion score ± SEM (n = 5). Kidney lesions scores as follows: no change (0), mild (1), moderate (2) or severe (3). For each time point and lesions, different superscripts for infected SPF (Table 3) and broiler (Table 4) chicks indicate significant differences between them (p < 0.05). Absence of a letter indicates that there were no significant differences (p > 0.05).

**Table 4**  
Histopathology kidney lesion scores in broiler chicks infected with IBV IS/885 strain.

| Kidney lesions          | Broiler (days post infection) |                        |           |           |                        |           |           |
|-------------------------|-------------------------------|------------------------|-----------|-----------|------------------------|-----------|-----------|
|                         | 3                             | 6                      | 9         | 12        | 15                     | 21        | 28        |
| Epithelial degeneration | 0.0 ± 0.0                     | 0.4 ± 0.2              | 0.0 ± 0.0 | 0.0 ± 0.0 | 0.2 ± 0.2              | 0.2 ± 0.2 | 0.0 ± 0.0 |
| Ducto-tubular dilation  | 0.0 ± 0.0                     | 0.0 ± 0.0              | 0.2 ± 0.2 | 0.2 ± 0.2 | 0.0 ± 0.0              | 0.0 ± 0.0 | 0.0 ± 0.0 |
| Heterophil infiltration | 0.0 ± 0.0                     | 0.0 ± 0.0 <sup>a</sup> | 0.2 ± 0.2 | 0.0 ± 0.0 | 0.2 ± 0.2              | 0.0 ± 0.0 | 0.4 ± 0.2 |
| Lymphoid infiltration   | 0.0 ± 0.0                     | 0.0 ± 0.0              | 1.0 ± 0.0 | 1.2 ± 0.2 | 0.6 ± 0.2              | 0.6 ± 0.2 | 0.0 ± 0.0 |
| Epithelial regeneration | 0.0 ± 0.0                     | 0.8 ± 0.3              | 0.2 ± 0.0 | 0.0 ± 0.0 | 0.0 ± 0.0              | 0.0 ± 0.0 | 0.0 ± 0.0 |
| Epithelial hyperplasia  | 0.0 ± 0.0                     | 0.2 ± 0.2              | 0.0 ± 0.0 | 0.0 ± 0.0 | 0.2 ± 0.2 <sup>b</sup> | 0.2 ± 0.2 | 0.0 ± 0.0 |
| Lymphoid nodules        | 0.0 ± 0.0                     | 0.0 ± 0.0              | 0.0 ± 0.0 | 0.6 ± 0.4 | 0.0 ± 0.0 <sup>b</sup> | 0.4 ± 0.4 | 1.0 ± 0.0 |

Data are expressed as the mean histopathological lesion score ± SEM (n = 5). Kidney lesions scores as follows: no change (0), mild (1), moderate (2) or severe (3). For each time point and lesions, different superscripts for infected SPF (Table 3) and broiler (Table 4) chicks indicate significant differences between them (p < 0.05). Absence of a letter indicates that there were no significant differences (p > 0.05).

#### 4.2. Gross necropsy findings

No detectable gross lesions were witnessed in the two control groups. In the SPF chicks (Expt 1), tracheal congestion and excess mucus were found in all the chicks that were necropsied at 3 dpi and such lesions disappeared by 12 dpi. Kidney lesions were first observed at 6 dpi and all the necropsied chicks showed large swelling, pale with tubules and/or ureters distended with urates, and these lesions were observed until 12 dpi. At 28 dpi, (during the *post mortem* examination) serous fluid accumulation was found in the left oviduct of two female SPF chicks (Fig 2). Samples of the fluid and tissues of the cystic oviduct were collected for virus detection and the oviduct was also collected for histopathology. The cystic fluid was positive for IBV by RT-PCR and VI, while the tissue was negative. Part-sequencing of the hypervariable region of the S1 gene of this isolate revealed 86% homology with the challenge inoculum. Necropsy of the bird that died at 8 dpi showed large, swollen, pale kidneys with tubules and ureters distended with urates.

In the broiler chicks (Expt 2), the gross lesions of the trachea and kidney were similar to those observed in the SPF chicks but were less severe. However, the lesions persisted for a longer time than in the SPF chicks. Tracheal lesions such as yellowish caseous plugs were observed at 15 dpi. Paleness and swelling of kidneys were observed from 6 to 21 dpi.

For the chick with the head swelling, the swab was positive for IBV and was negative for bacteria. On sequencing, the virus detected was 100% identical to the IS/885 strain (the inoculum). The virus was successfully isolated in SPF ECEs and typical IBV lesions were found in the embryo (e.g. dwarfing and curling of the embryos). Necropsy of birds that died showed emaciation, dehydration, caseous exudate and plugs in the upper trachea and pale, swollen kidneys.

#### 4.3. Histopathology

No significant histological abnormalities were observed in the trachea, kidneys or oviduct of the control groups in either experiment.

##### 4.3.1. Trachea of infected chicks

Details of histological lesions in the tracheas for SPF and broiler chicks are summarized in Table 1 and Table 2 respectively. In both lines of chicks (SPF and broiler), lesions were similar but the severity differed throughout the study periods.

In the SPF chicks, severe changes in the trachea were observed at 3 dpi. The most consistent lesions in the SPF chick were loss of cilia and heterophil infiltration, decreased mucous cells and an occasional heterophilic exudate in the tracheal lumen (Fig 3b). Mild to moderate lesions were witnessed until 15 dpi. Ciliated epithelium repaired by 21 dpi. Areas of severe lymphoid infiltration in the lamina propria or submucosa persisted and were found until the end of the experiment (Table 1).

In the broilers chicks, at an early stage of infection, each lesion type was less severe than those observed in the SPF chicks but persisted for longer period (Fig 3c). At 21 dpi, the histopathological changes were significantly greater (p < 0.05) than those observed in the SPF chicks (Table 1).

##### 4.3.2. Kidney of the infected chicks

A summary of the histopathological changes in the kidneys of infected SPF (Expt 1) and boiler (Expt 2) chicks are given in Table 3 and 4 respectively. In the SPF chicks, kidney lesions developed by 3 dpi, which included ducto-tubular dilation, interstitial heterophilic infiltration and epithelial hyperplasia. The main histological lesions consisted of interstitial lymphoid infiltration with mild lymphoid nodules observed throughout the study (Fig 3e). Most of the kidney lesions had cleared by 21 dpi, apart from mild lymphoid infiltration and lymphoid nodules that were present until the end of the experiment (Table 3).

In the broiler chicks, kidney lesions were first observed at 6 dpi, which included tubular degeneration and hyperplasia of the epithelium. Lymphoid infiltration was the main lesion throughout the observation period (Fig 3f). No significant differences were observed among the histological scores of the lesions between the SPF and broiler chicks (Table 4).

##### 4.3.3. Oviduct of infected SPF chicks

The oviduct lumen was multifocally partially lined by compact epithelial cells devoid of cilia (Fig 4). Many epithelial cells are shed into the lumen leaving sections without mucosa. The underlying submucosa was multifocally infiltrated by both moderate numbers of degenerate and viable lymphocytes and plasma cells with lesser numbers of macrophages.

#### 4.4. IBV RT-PCR

No virus was detected in the control groups of either experiment throughout the study.

**Table 5**  
Virus detection by RT-PCR in tissue of SPF or broiler chicks infected with IBV IS/885 strain.

| Days post infection | SPF            |    |    |     | Broiler |   |    |     |
|---------------------|----------------|----|----|-----|---------|---|----|-----|
|                     | Tr             | L  | CT | Kid | Tr      | L | CT | Kid |
| 3                   | 5 <sup>a</sup> | 5  | 4  | 4   | 3       | 1 | 5  | 5   |
| 6                   | 5              | 5  | 5  | 5   | 5       | 2 | 5  | 2   |
| 9                   | 5              | 5  | 5  | 5   | 2       | 0 | 4  | 0   |
| 12                  | 5              | 5  | 4  | 5   | 4       | 1 | 2  | 1   |
| 15                  | 5              | 5  | 5  | 5   | 4       | 2 | 4  | 1   |
| 21                  | 2              | 3  | 4  | 3   | 0       | 0 | 3  | 1   |
| 28                  | 1              | 3  | 3  | 0   | 1       | 0 | 5  | 2   |
| Total <sup>b</sup>  | 28             | 31 | 30 | 27  | 19      | 6 | 28 | 12  |

Tr = Trachea, L = lung, CT = caecal tonsil, Kid = kidney.

<sup>a</sup> Values presented are number of IBV-positive out of five birds examined.

<sup>b</sup> Total number of tissues that was positive for IBV.

**Table 6**  
Mean IBV ELISA antibody titres in the SPF and broiler chicks infected with IBV IS/885 strain.

| Days post infection | SPF                    |                        | Broiler                 |                       |
|---------------------|------------------------|------------------------|-------------------------|-----------------------|
|                     | Infected               | Control                | Infected                | Control               |
| 0                   | 21 ± 9.0               | 21 ± 9.0               | 2867 ± 229              | 2867 ± 229            |
| 3                   | ND                     | ND                     | 1813 ± 343              | 1809 ± 544            |
| 6                   | ND                     | ND                     | 531 ± 153               | 501 ± 64              |
| 9                   | ND                     | ND                     | 311 ± 85                | 282 ± 75              |
| 15                  | 160 ± 64 <sup>a</sup>  | 4.0 ± 2.0 <sup>b</sup> | 378 ± 175               | 75 ± 19               |
| 21                  | 551 ± 146 <sup>a</sup> | 12 ± 5.0 <sup>b</sup>  | 1116 ± 129 <sup>a</sup> | 19 ± 8.0 <sup>b</sup> |
| 28                  | 481 ± 194 <sup>a</sup> | 7.0 ± 4.0 <sup>b</sup> | 623 ± 35 <sup>a</sup>   | 10 ± 8.0 <sup>b</sup> |

ND: not done. Data are expressed as the mean values ± SEM (n = 8). Data in the same row with different superscript letters are significantly different in antibody titres (p < 0.05). Absence of a letter indicates that there were no significant differences (p > 0.05) between the infected and control groups at each sampling days. The cut-off values for ELISA = 396.

In the SPF chicks, the virus was detected in the trachea, lung, caecal tonsil and kidney in most of the tested samples for up to 15 dpi. Beyond that, the virus was infrequently detected in tissues (Table 5). In the broiler chicks, detection of the virus was lower than that observed in the SPF chicks throughout the study. IBV was found most frequently in the caecal tonsil, followed by the trachea (Table 5).

#### 4.5. Antibody detection

##### 4.5.1. Enzyme-linked immunosorbent assay

Analysis of the antibody titres of both experiments using ELISA is summarized in Table 6. In the SPF chicks, sera collected prior to experimental infection were free of IBV antibodies. Chicks which received virus-free AF had ELISA titres lower than the detectable level throughout the experimental period. In the IS/885-infected chicks, an increase of antibody titres occurred at 21 dpi which decreased by 28 dpi (Table 6). In broiler chicks, high levels of IBV MDA were detected in all birds at 1-day-old. By 3 dpi, the MDA had declined in the control and infected groups and showed no significant differences (Table 6). At 9 and 15 dpi, neither the infected nor control group had antibody titres above the cut-off point. However, at 21 and 28 dpi, highly significant antibody titres were observed in the infected group compared to the control group (Table 6).

##### 4.5.2. Haemagglutination inhibition test

Analysis of the HI antibody titres of both experiments using different IBV antigens are summarized in Tables 7 and 8. In the SPF chicks, all sera collected prior to experimental infection and from the control group were free of IBV HI antibodies (Table 7). At 15 dpi, the HI titres against the heterologous antigens of M41 and 793B were low (<3 log<sub>2</sub>) compared to the homologous antigen (3.8 log<sub>2</sub>). At 21 dpi, chicks showed high antibody titres (≥3 log<sub>2</sub>) compared to control birds. The birds showed high significant level (p < 0.05) of IBV HI antibody titres when IS/885 was used as antigen throughout the sampling time (Table 7). In

the broiler chicks, sera collected at one day old showed significant antibody titer when either M41 or 793B were used as the antigen (Table 8). Titres had started to decline significantly by 3 dpi in both the infected and control groups. At 6 and 9 dpi, there were no detectable titres in either (<3 log<sub>2</sub>). When all IBV antigens were compared, higher levels of IBV antibody titres were seen when IS/885 was used as the antigen from 15 dpi onwards.

## 5. Discussion

The results presented in this study demonstrate that IS/885/00 is a virulent IBV, as extensive disease was produced following infection in both types of chicks. In addition to the respiratory signs, gross lesions comprising of tracheal caseous exudate and plugs, and swollen kidneys (with or without) urate deposits were observed. Such clinical signs have been observed following infection with virulent IBV M41 (Butcher et al., 1990), Moroccan G strain (Ambali and Jones, 1990), 793B (Boroomand et al., 2012), It-02 (Dolz et al., 2012) and QX (Ganapathy et al., 2012). These clinical signs and lesions were more severe in SPF than in the broiler chicks, indicating differences in the susceptibility to IBV infection in these different types of birds. This could be due to the genetic line of the birds or the IBV MDAs in the broiler chicks which probably neutralized certain amounts of IBVs (Ignjatovic et al., 2003). Previous studies using virulent IBV M41 demonstrated a considerable difference between two white leghorn chick lines in terms of the severity and duration of respiratory signs (Otsuki et al., 1990).

It has been reported that chicks with high antibody titres had a better protection against IBV M41 infection at one day old but not at seven days old (Mondal and Naqi, 2001) and that MDA does not prevent the viral infection but does reduce the pathogenic effects of the IBV infection in young chicks (Klieve and Cumming, 1988). Our findings suggest that MDA is an important factor in determining the severity of the disease caused by IS/885 in young chicks.

Following IS/885 infection of the broiler chicks, clear periorbital swelling was noted in one bird at 15 dpi. The swab collected from this site was positive for IBV by RT-PCR and VI, and sequencing demonstrated 100% nucleotide level identity to the inoculated IS/885 strain. Neither aMPV nor bacteria (*E. coli* or *Mycoplasma*) were detected by PCR or cultured from swabbing, suggesting that the swelling was most likely caused the IBV infection. This finding is similar to the first report of swollen head syndrom (SHS) where an untyped coronavirus and *E. coli* were isolated from a broiler chicken in Southern Africa (Morley and Thomson, 1984). IBV M41 and *E. coli* were isolated from a broiler flock in the USA that experienced SHS (Droual and Woolcock, 1994). aMPV (Picault et al., 1987; Aung et al., 2008), FAdV (Droual and Woolcock, 1994; Georgiades et al., 2001) and secondary infection (such as *E. coli*) have been implicated as the cause of SHS in chickens (Nakamura et al., 1997; Nakamura et al., 1998).

In the IS/885-infected SPF chicks, necropsy examination at 28 dpi revealed dilation of the oviduct with fluid content (cystic oviduct) in two female chicks. Cystic oviduct formation following infection of virulent

**Table 7**  
Mean HI antibody titres in IS/885-infected SPF chicks using homologous (IS/885) and heterologous (M41, 793B) antigens.

| Days post infection | Infected                |                         |                         | Control   |           |           |
|---------------------|-------------------------|-------------------------|-------------------------|-----------|-----------|-----------|
|                     | M41                     | 793B                    | IS/885                  | M41       | 793B      | IS/885    |
| 0                   | 0.6 ± 0.2               | 1.1 ± 0.2               | 0.7 ± 0.2               | 0.6 ± 0.2 | 1.1 ± 0.2 | 0.6 ± 0.2 |
| 3                   | ND                      | ND                      | ND                      | ND        | ND        | ND        |
| 6                   | ND                      | ND                      | ND                      | ND        | ND        | ND        |
| 9                   | ND                      | ND                      | ND                      | ND        | ND        | ND        |
| 15                  | 1.5 ± 0.3 <sup>ab</sup> | 0.6 ± 0.1 <sup>a</sup>  | 3.8 ± 0.2 <sup>c*</sup> | 1.1 ± 0.1 | 0.6 ± 0.1 | 0.6 ± 0.1 |
| 21                  | 4.0 ± 0.2 <sup>a*</sup> | 3.6 ± 0.3 <sup>a*</sup> | 5.1 ± 0.3 <sup>b*</sup> | 0.8 ± 0.3 | 0.7 ± 0.2 | 0.3 ± 0.1 |
| 28                  | 2.5 ± 0.4 <sup>a*</sup> | 2.6 ± 0.1 <sup>a*</sup> | 3.9 ± 0.2 <sup>b*</sup> | 0.6 ± 0.1 | 0.2 ± 0.1 | 0.3 ± 0.2 |

ND: not done. Data are expressed as the mean values ± SEM (n = 8). At each sampling days, titers with different superscript letters within infected or control groups is significantly different (p < 0.05). Absence of a letter indicates that there were no significant differences (p > 0.05) between the IBV antigens at any of the time points. The cut-off point = >3log<sub>2</sub>.

\* Values differed significantly from each representative control group.

**Table 8**

Mean IBV HI antibody titres in IS/885-infected broiler chicks using homologous (IS/885) and heterologous (M41, 793B) antigens.

| Days post infection | Infected group          |                          |                         | Control                |                        |                        |
|---------------------|-------------------------|--------------------------|-------------------------|------------------------|------------------------|------------------------|
|                     | M41                     | 793B                     | IS/885                  | M41                    | 793B                   | IS/885                 |
| 0                   | 6.2 ± 0.6 <sup>a</sup>  | 5.8 ± 0.5 <sup>a</sup>   | 3.5 ± 0.2 <sup>b</sup>  | 6.2 ± 0.6 <sup>a</sup> | 5.8 ± 0.5 <sup>a</sup> | 3.5 ± 0.2 <sup>b</sup> |
| 3                   | 5.1 ± 0.7 <sup>a</sup>  | 5.0 ± 0.2 <sup>a</sup>   | 2.6 ± 0.1 <sup>a</sup>  | 5.1 ± 1.1 <sup>a</sup> | 5.5 ± 0.8 <sup>a</sup> | 3.1 ± 0.1 <sup>b</sup> |
| 6                   | 2.4 ± 0.6               | 1.3 ± 0.3                | 2.1 ± 0.2               | 3.0 ± 0.3 <sup>a</sup> | 3.3 ± 0.5 <sup>b</sup> | 1.7 ± 0.3 <sup>a</sup> |
| 9                   | 1.6 ± 0.8               | 1.4 ± 0.2                | 1.8 ± 0.2               | 1.6 ± 0.8              | 2.3 ± 0.6              | 1.1 ± 0.2              |
| 15                  | 3.5 ± 0.1 <sup>a*</sup> | 3.3 ± 0.1 <sup>a*</sup>  | 4.0 ± 0.2 <sup>a*</sup> | 1.3 ± 0.4              | 0.7 ± 0.1              | 1.0 ± 0.0              |
| 21                  | 5.0 ± 0.0 <sup>a*</sup> | 4.2 ± 0.4 <sup>ab*</sup> | 5.8 ± 0.2 <sup>c*</sup> | 1.1 ± 0.3              | 0.6 ± 0.1              | 0.3 ± 0.2              |
| 28                  | 3.7 ± 0.1 <sup>a*</sup> | 3.5 ± 0.1 <sup>a*</sup>  | 4.5 ± 0.1 <sup>c*</sup> | 0.7 ± 0.2              | 0.4 ± 0.1              | 0.1 ± 0.1              |


Data are expressed as the mean values ± SEM (n = 8). At each sampling days, titers with different superscript letters within infected or control groups is significantly different (p < 0.05). Absence of a letter indicates that there were no significant differences (p > 0.05). The cut-off point = >3log<sub>2</sub>.

\* Values differed significantly from each representative control group

IBVs at a young age has been reported for IBV M41 (Crinion et al., 1971; Crinion and Hofstad, 1972; Jones and Jordan, 1972) and, recently, QX (Benyeda et al., 2009; de Wit et al., 2011b; Ganapathy et al., 2012). We report, for the first time, similar pathological development in SPF chicks that received the IS/885 virus at 1-day old. In addition, virus with 86% part-S1 sequence identity to the original inoculum was isolated from the cystic fluid. The detection of IBV IS/885 in the oviduct fluid demonstrates both active replication of the virus in the epithelium and its abundance in the fluid and these were shed by epithelial cells. The epithelial cells of the oviduct are the primary cells infected by IBV (Crinion and Hofstad, 1972). The IS/885 infection has caused a diffuse loss in the cilia, epithelial necrosis and infiltration of lymphocytes and plasma cells into the oviduct which suggests that the virus has replicated in the epithelial lining of the oviduct.

The pathogenicity of IBV strains for the oviduct varies as some do not replicate in the oviduct neither causing lesions nor showing significant amounts of viral antigen in the epithelial cells (Crinion and Hofstad, 1972). IBV Conn and Iowa failed to produce any gross or histopathological change in the oviduct, in contrast, IBV M41 produces the greatest number of changes, followed by Australian T (Crinion and Hofstad, 1972). Benyeda et al. (2009) did not detect any abnormalities in the oviducts of 793B infected one-day-old SPF chicks, whereas a variable percentage of oviduct dilatations were detected following inoculation with five different QX-like strains. It appears that along with M41 and QX, IS/885 should be considered as a potential cause of cystic oviducts in female chickens.

The microscopic findings in the SPF and broiler chicks following IS/885 infection were similar to other virulent IBVs as reported (Albassam et al., 1986; Nakamura et al., 1991; Chen et al., 1996). In



**Fig. 4.** Hematoxylin and eosin stains of the cystic oviduct of IS/885-infected female SPF chicks at 28 dpi (magnification at 400×). Multifocal moderate chronic lymphocytic salpingitis with cystic dilation. Diffuse loss of the cilia (black arrow) and infiltration of lymphocytes and plasma cells in submucosa (blue arrow).

this work, the SPF chicks had greater necrosis in the trachea, indicating a more extensive virus replication than broiler chicks, which may have been due to extravasation of circulating maternal antibody in the latter. In both types of birds, severe and early onset of histopathological lesions was seen in the trachea, compared to relatively mild lesions in the kidney. Nephropathogenic IBVs have shown tropism for respiratory and kidney tissues although kidney lesions are more apparent (Albassam et al., 1986; Butcher et al., 1989). Based on these findings, the IS/885 strain appears to have an affinity for both respiratory and renal tissues.

In this study, persistence of the viral genome was observed in the selected tissues until the end of all experiments (28 dpi). Although no isolation was attempted, the results showed marked reductions in the detection in the trachea, lungs and kidneys of broilers in comparison to the SPF chicks. This again shows the advantage of having IBV maternal-antibodies in young chicks, which appears to inhibit the invasion and colonization of IS/885 in visceral tissues (Mondal and Naqi, 2001).

The immune response after infection with IBV was measured by ELISA and HI. With ELISA, a maximum mean antibody titer was found at 21 days post infection in SPF and broilers. For HI, sera were assayed against M41 and 793B since most producers use live and inactivated vaccines related to these serotypes. IS/885 antigen was also used as a homologous comparison. With HI, distinctive patterns were seen in broiler chicks; higher mean HI titres against M41 and 793B at day-old reflects the maternal antibodies derived from parent birds that were previously vaccinated using Massachusetts, 793B or D274 vaccines. For 3, 6 and 9 days post infection, though there were no significant differences between the IS/885-infected broiler chicks, the titres declined against all three HI antigens. From 15 days post infection, the mean HI titres against M41, 793B and IS/885 peaked by 21 days post infection and then declined by 28 days post infection. In contrast to day-old sera, the mean titer is highest for the IS/885 antigen, reflecting a better reactivity with the homologous antigen. These significant levels of HI antibodies against the IS/885 antigen were observed when compared to the level of antibody response against M41 and 793B antigens in IS/885-infected birds. This finding supports the use of the homologous virus as an antigen in the HI test, as it has been reported that the HI test is strain specific (Monreal et al., 1985; de Wit et al., 1997).

## 6. Conclusions

This study showed that under experimental conditions, the IBV strain, IS/885, is pathogenic for both SPF and broiler chicks, with lesser disease severity in the latter. The IS/885 showed tropism for both respiratory and renal tissues. In addition, this virus is associated with head-swelling and cystic oviduct development in young chicks.

## References

- Abdel-Moneim, A.S., El-Kady, M.F., Ladman, B.S., Gelb Jr., J., 2006. S1 gene sequence analysis of a nephropathogenic strain of avian infectious bronchitis virus in Egypt. *Viol. J.* 3, 78.



- Abdel-Moneim, A., Affi, M., El-Kady, M., 2012. Emergence of a novel genotype of avian infectious bronchitis virus in Egypt. *Arch. Virol.* 1–5.
- Albassam, M.A., Winterfield, R.W., Thacker, H.L., 1986. Comparison of the nephropathogenicity of four strains of infectious bronchitis virus. *Avian Dis.* 30, 468–476.
- Alexander, D.J., Chettle, N.J., 1977. Procedures for the haemagglutination and the haemagglutination inhibition tests for avian infectious bronchitis virus. *Avian Pathology* 6, 9–17.
- Ambali, A.G., Jones, R.C., 1990. Early pathogenesis in chicks of infection with an enterotropic strain of infectious bronchitis virus. *Avian Dis.* 34, 809–817.
- Amin, O.G.M., Valastro, V., Salviato, A., Drago, A., Cattoli, G., Monne, I., 2012. Circulation of QX-like infectious bronchitis virus in the Middle East. *Vet. Rec.* 171, 530.
- Aung, Y., Liman, M., Neumann, U., Rautenschlein, S., 2008. Reproducibility of swollen sinuses in broilers by experimental infection with avian metapneumovirus subtypes A and B of turkey origin and their comparative pathogenesis. *Avian Pathology* 37, 65–74.
- Awad, F., Baylis, M., Ganapathy, K., 2014a. Detection of variant infectious bronchitis viruses in broiler flocks in Libya. *International Journal of Veterinary Science and Medicine* 2, 78–82.
- Awad, F., Baylis, M., Jones, R.C., Ganapathy, K., 2014b. Evaluation of Flinders Technology Associates cards for storage and molecular detection of avian metapneumoviruses. *Avian Pathology* 43, 125–129.
- Benyeda, Z., Mató, T., Süveges, T., Szabó, É., Kardi, V., Abonyi-Tóth, Z., Rusvai, M., Palya, V., 2009. Comparison of the pathogenicity of QX-like, M41 and 793/B infectious bronchitis strains from different pathological conditions. *Avian Pathology* 38, 449–456.
- Boroomand, Z., Asasi, K., Mohammadi, A., 2012. Pathogenesis and tissue distribution of avian infectious bronchitis virus isolate IRFIBV32 (793/B serotype) in experimentally infected broiler chickens. *Sci. World J.* 6, 336–343.
- Bourogaa, H., Miled, K., Gribaa, L., El Behi, I., Ghram, A., 2009. Characterization of new variants of avian infectious bronchitis virus in Tunisia. *Avian Dis.* 53, 426–433.
- Butcher, G.D., Winterfield, R.W., Shapiro, D.P., 1989. An outbreak of nephropathogenic H13 infectious bronchitis in commercial broilers. *Avian Dis.* 33, 823–826.
- Butcher, G.D., Winterfield, R.W., Shapiro, D.P., 1990. Pathogenesis of H13 nephropathogenic infectious bronchitis virus. *Avian Dis.* 34, 916–921.
- Cavanagh, D., 2007. Coronavirus avian infectious bronchitis virus. *Vet. Res.* 38, 281–297.
- Cavanagh, D., Mawditt, K., Britton, P., Naylor, C.J., 1999. Longitudinal field studies of infectious bronchitis virus and avian pneumovirus in broilers using type-specific polymerase chain reactions. *Avian Pathol.* 28, 593–605.
- Chacón, J.L., Assayag, M.S., Revollo, L., Astolfi-Ferreira, C.S., Vejarano, M.P., Jones, R.C., Piantino Ferreira, A.J., 2014. Pathogenicity and molecular characteristics of infectious bronchitis virus (IBV) strains isolated from broilers showing diarrhoea and respiratory disease. *Br. Poult. Sci.* 55, 271–283.
- Chen, B.Y., Hosi, S., Nunoya, T., Itakura, C., 1996. Histopathology and immunohistochemistry of renal lesions due to infectious bronchitis virus in chicks. *Avian Pathology* 25, 269–283.
- Chong, K.T., Apostolov, K., 1982. The pathogenesis of nephritis in chickens induced by infectious bronchitis virus. *J. Comp. Pathol.* 92, 199–211.
- Crinion, R.A.P., Hofstad, M.S., 1972. Pathogenicity of four serotypes of avian infectious bronchitis virus for the oviduct of young chickens of various ages. *Avian Dis.* 16, 351–363.
- Crinion, R.A.P., Ball, R.A., Hofstad, M.S., 1971. Pathogenesis of oviduct lesions in immature chickens following exposure to infectious bronchitis virus at one day old. *Avian Dis.* 15, 32–41.
- de Wit, J.J., Mekkes, D.R., Kouwenhoven, B., Verheijden, J.H.M., 1997. Sensitivity and specificity of serological tests for infectious bronchitis virus antibodies in broilers. *Avian Pathology* 26, 105–118.
- de Wit, J.J., Cook, J.K.A., van der Heijden, H.M.J.F., 2011a. Infectious bronchitis virus variants: a review of the history, current situation and control measures. *Avian Pathology* 40, 223–235.
- de Wit, J.J., Nieuwenhuisen-van Wilgen, J., Hoogkamer, A., van de Sande, H., Zuidam, G.J., Fabri, T.H.F., 2011b. Induction of cystic oviducts and protection against early challenge with infectious bronchitis virus serotype D388 (genotype QX) by maternally derived antibodies and by early vaccination. *Avian Pathology* 40, 463–471.
- Dhinakar Raj, G., Jones, R.C., 1996. Immunopathogenesis of infection in SPF chicks and commercial broiler chickens of a variant infectious bronchitis virus of economic importance. *Avian Pathology* 25, 481–501.
- Dolz, R., Vergara-Alert, J., Perez, M., Pujols, J., Majo, N., 2012. New insights on infectious bronchitis virus pathogenesis: characterization of Italy 02 serotype in chicks and adult hens. *Vet. Microbiol.* 156, 256–264.
- Droual, R., Woolcock, P.R., 1994. Swollen head syndrome associated with *E. coli* and infectious bronchitis virus in the Central Valley of California. *Avian Pathology* 23, 733–742.
- El-Houadfi, M., Jones, R.C., Cook, J.K., Ambali, A.G., 1986. The isolation and characterisation of six avian infectious bronchitis viruses isolated in Morocco. *Avian Pathology* 15, 93–105.
- Ganapathy, K., Wilkins, M., Forrester, A., Lemiere, S., Cserep, T., McMullin, P., Jones, R.C., 2012. QX-like infectious bronchitis virus isolated from cases of proventriculitis in commercial broilers in England. *Vet. Rec.* 171, 597.
- Ganapathy, K., Ball, C., Forrester, A., 2015. Genotypes of infectious bronchitis viruses circulating in the Middle East between 2009 and 2014. *Virus Res.* 210, 198–204.
- Geilhausen, H.E., Ligon, F.B., Lukert, P.D., 1973. The pathogenesis of virulent and avirulent avian infectious bronchitis virus. *Archiv für die gesamte Virusforschung* 40, 285–290.
- Georgiades, G., Iordanidis, P., Koumbati, M., 2001. Cases of swollen head syndrome in broiler chickens in Greece. *Avian Dis.* 45, 745–750.
- Ignjatovic, J., Ashton, D.F., Reece, R., Scott, P., Hooper, P., 2002. Pathogenicity of Australian strains of avian infectious bronchitis virus. *J. Comp. Pathol.* 126, 115–123.
- Ignjatovic, J., Reece, R., Ashton, F., 2003. Susceptibility of three genetic lines of chicks to infection with a nephropathogenic T strain of avian infectious bronchitis virus. *J. Comp. Pathol.* 128, 92–98.
- Jackwood, M.W., 2012. Review of infectious bronchitis virus around the world. *Avian Dis.* 56, 634–641.
- Jones, R.C., Jordan, F.T., 1972. Persistence of virus in the tissues and development of the oviduct in the fowl following infection at day old with infectious bronchitis virus. *Res. Vet. Sci.* 13, 52–60.
- King, D.J., Hopkins, S.R., 1983. Evaluation of the hemagglutination-inhibition test for measuring the response of chickens to avian infectious bronchitis virus vaccination. *Avian Dis.* 27, 100–112.
- Klieve, A.V., Cumming, R.B., 1988. Immunity and cross-protection to nephritis produced by Australian infectious bronchitis viruses used as vaccines. *Avian Pathology* 17, 829–839.
- Mahmood, Z.H., Sleman, R.R., Uthman, A.U., 2011. Isolation and molecular characterization of Sul/01/09 avian infectious bronchitis virus, indicates the emergence of a new genotype in the Middle East. *Vet. Microbiol.* 150, 21–27.
- Meir, R., Rosenblut, E., Perl, S., Kass, N., Ayali, G., Perk, S., Hemsani, E., 2004. Identification of a novel nephropathogenic infectious bronchitis virus in Israel. *Avian Dis.* 48, 635–641.
- Mondal, S.P., Naqi, S.A., 2001. Maternal antibody to infectious bronchitis virus: its role in protection against infection and development of active immunity to vaccine. *Vet. Immunol. Immunopathol.* 79, 31–40.
- Monreal, G., Bauer, H.J., Wiegmann, J., 1985. Comparison of the enzyme-linked immunosorbent assay (ELISA), haemagglutination inhibition test and agar gel precipitation test for detection of antibodies to avian infectious bronchitis virus. *Avian Pathology* 14, 421–434.
- Morley, A.J., Thomson, D.K., 1984. Swollen-head syndrome in broiler chickens. *Avian Dis.* 28, 238–243.
- Nakamura, K., Cook, J.K., Otsuki, K., Huggins, M.B., Frazier, J.A., 1991. Comparative study of respiratory lesions in two chicken lines of different susceptibility infected with infectious bronchitis virus: histology, ultrastructure and immunohistochemistry. *Avian Pathology* 20, 241–257.
- Nakamura, K., Mase, M., Tanimura, N., Yamaguchi, S., Nakazawa, M., Yuasa, N., 1997. Swollen head syndrome in broiler chickens in Japan: its pathology, microbiology and biochemistry. *Avian Pathology* 26, 139–154.
- Nakamura, K., Mase, M., Tanimura, N., Yamaguchi, S., Yuasa, N., 1998. Attempts to reproduce swollen head syndrome in specific pathogen-free chickens by inoculating with *Escherichia coli* and/or turkey rhinotracheitis virus. *Avian Pathology* 27, 21–27.
- Otsuki, K., Huggins, M.B., Cook, J.K., 1990. Comparison of the susceptibility to avian infectious bronchitis virus infection of two inbred lines of white leghorn chickens. *Avian Pathology* 19, 467–475.
- Picault, J.P., Giraud, P., Drouin, P., Guittet, M., Bennejean, G., Lamande, J., Toquin, D., Gueguen, C., 1987. Isolation of a TRTV-like virus from chickens with swollen-head syndrome. *Vet. Rec.* 121, 135.
- Reed, L.J., Muench, H., 1938. A simple method of estimating fifty per cent endpoints. *Am. J. Epidemiol.* 27, 493–497.
- Selim, K., Arafa, A.S., Hussein, H.A., El-Sanousi, A.A., 2013. Molecular characterization of infectious bronchitis viruses isolated from broiler and layer chicken farms in Egypt during 2012. *International Journal of Veterinary Science and Medicine* 1, 102–108.
- Terregino, C., Toffan, A., Serena Beato, M., De Nardi, R., Vascellari, M., Meini, A., Ortali, G., Mancin, M., Capua, I., 2008. Pathogenicity of a QX strain of infectious bronchitis virus in specific pathogen free and commercial broiler chickens, and evaluation of protection induced by a vaccination programme based on the Ma5 and 4/91 serotypes. *Avian Pathology* 37, 487–493.
- Toffan, A., Bonci, M., Bano, L., Bano, L., Valastro, V., Vascellari, M., Capua, I., Terregino, C., 2013. Diagnostic and clinical observation on the infectious bronchitis virus strain Q1 in Italy. *Vet. Ital.* 49, 347–355.
- Wang, Y., Wang, Y., Zhang, Z., Fan, G., Jiang, Y., Liu Xiang, E., Ding, J., Wang, S., 1998. Isolation and identification of glandular stomach type IBV (QX IBV) in chickens. *Chinese Journal of Animal Quarantine* 15, 1–3.
- Worthington, K.J., Currie, R.J.W., Jones, R.C., 2008. A reverse transcriptase polymerase chain reaction survey of infectious bronchitis virus genotypes in Western Europe from 2002 to 2006. *Avian Pathology* 37, 247–257.
- Yu, L., Jiang, Y.H., Low, S., Wang, Z.L., Nam, S.J., Liu, W., Kwang, J., 2001. Characterization of three infectious bronchitis virus isolates from China associated with proventriculus in vaccinated chickens. *Avian Dis.* 45, 416–424.