

Research Article

5-Aminosalicylic Acid Inhibits Acute *Clostridium difficile* Toxin A-Induced Colitis in Rats

Steven R. Vigna^{1,2}

¹ Departments of Cell Biology & Medicine, Duke University Medical Center, P.O. Box 3011, Durham, NC 72210, USA

² VA Medical Center, Durham, NC 27705, USA

Correspondence should be addressed to Steven R. Vigna; steven.vigna@duke.edu

Received 13 January 2014; Revised 21 May 2014; Accepted 4 June 2014; Published 23 June 2014

Academic Editor: Jonathan Steven Alexander

Copyright © 2014 Steven R. Vigna. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We tested the hypothesis that 5-aminosalicylic acid (5-ASA) inhibits toxin A-induced generation of colonic leukotriene B₄ (LTB₄) and toxin A colitis in rats. Isolated colonic segments in anesthetized rats were treated intraluminally with toxin A for 3 hours with or without 30 minutes of pretreatment with either 5-ASA or sulfapyridine and then colonic tissue levels of LTB₄ were measured and inflammation was assessed. Separately, sulfasalazine was administered to rats in their drinking water for 5 days, isolated colonic segments were then prepared, toxin A was administered, and inflammation was assessed as before. Pretreatment with 5-ASA inhibited toxin A-induced increased tissue LTB₄ concentration in the colon. Sulfasalazine and 5-ASA but not sulfapyridine significantly inhibited toxin A colitis. However, pretreatment with 5-ASA did not protect against direct TRPV1-mediated colitis caused by capsaicin. Toxin A stimulated the release of substance P (SP), and this effect was also inhibited by sulfasalazine and 5-ASA but not by sulfapyridine. Thus, toxin A stimulates colonic LTB₄ resulting in activation of TRPV1, release of SP, and colitis. Inhibition of 5-LO by 5-ASA disrupts this pathway and supports the concept that LTB₄ activation of TRPV1 plays a role in toxin A colitis.

1. Introduction

Clostridium difficile produces exotoxins such as toxin A that cause acute colonic inflammation characterized clinically by watery diarrhea and cramps and pathologically by pseudomembranous colitis. The pathway by which toxin A causes colitis is incompletely known. We have shown that oral sulfasalazine inhibits toxin A colitis in rats [1]. Sulfasalazine is used to treat the chronic human inflammatory bowel diseases (IBDs), ulcerative colitis (UC), and Crohn's disease (CD). Sulfasalazine consists of one molecule of 5-aminosalicylic acid (5-ASA, mesalamine) coupled by an azo bond to one molecule of sulfapyridine. The azo bond allows uncoupling of the two constituent compounds in the lumen of the colon by the action of bacterial azo reductase enzymes resulting in topical delivery of the compounds [2]. It has been shown that the 5-ASA moiety of sulfasalazine is the therapeutically active component in UC and CD [3, 4] and that the sulfapyridine moiety is inactive and causes most of the allergic and intolerant effects of sulfasalazine [5, 6].

The mechanism of the therapeutic effect of 5-ASA in intestinal inflammation is unknown. Several possible mechanisms have been proposed including inhibition of 5-lipoxygenase (5-LO), the rate-limiting enzyme in the biosynthesis of leukotriene B₄ (LTB₄) [5, 7]. Inhibition of 5-LO is a promising candidate because of the demonstrated relationship between LTB₄ and human UC [5] and because of the efficacy of other 5-LO inhibitors in UC [8]. LTB₄ is also found in much higher concentrations in patients with IBD than in healthy controls [5]. In addition, LTB₄ has been shown to be increased in animal models of colitis [9–12] and inhibition of 5-LO caused reduction of tissue LTB₄ levels and inhibited colitis in these models [11, 13].

We have shown previously that LTB₄ activates the transient receptor potential vanilloid-1 (TRPV1) ion channel expressed by primary sensory neurons in the ileum resulting in neurogenic inflammation and that inhibition of TRPV1 reduces LTB₄- and toxin A-induced ileitis [14, 15]. The fact that LTB₄ is an endogenous TRPV1 agonist [16] coupled with




FIGURE 1: Effects of toxin A on colonic luminal fluid accumulation, MPO activity, and histopathology and inhibition of these effects by TRPV1 antagonism. (a) Toxin A-induced stimulation of colonic luminal fluid accumulation and the inhibition of this effect by pretreatment with 1 μg of I-RTX administered intraluminally 30 min before toxin A. (b) Toxin A-induced stimulation of colonic MPO activity and the inhibition of this effect by pretreatment with 1 μg of I-RTX administered intraluminally 30 min before toxin A. H&E-stained sections of rat colonic segments after they were treated intraluminally for 3 h with vehicle (c), 1 μg of I-RTX (d), 5 μg of toxin A (e), or 5 μg of toxin A after pretreatment with 1 μg of I-RTX 30 min before toxin A administration (f). Scale bar, 100 μm . * $P < 0.05$ versus toxin A(-)/I-RTX(-); ** $P < 0.001$ versus toxin A(-)/I-RTX(-); # $P < 0.05$ versus toxin A(+)/I-RTX(-).

our demonstration that inhibition of 5-LO inhibited both toxin A-induced ileal LTB_4 levels and toxin A- but not LTB_4 -induced ileitis led us to propose that *C. difficile* toxin A causes ileitis by stimulating mucosal LTB_4 production that

subsequently initiates neurogenic inflammation mediated by TRPV1.

We show here that 5-ASA decreases toxin A-stimulated colonic LTB_4 levels and SP release and that 5-ASA but not




FIGURE 2: Effects of 5-ASA, toxin A, and toxin A after pretreatment with 5-ASA on colonic tissue LTB₄ concentrations. * $P < 0.01$ versus no treatment.

sulfapyridine inhibits toxin A-induced colitis. However, 5-ASA does not inhibit direct TRPV1-mediated colitis caused by intraluminal capsaicin. Taken together, these results demonstrate that 5-ASA inhibits 5-LO and accounts for the protective effects of sulfasalazine in toxin A colitis.

2. Materials and Methods

2.1. Materials. Sulfasalazine, 5-aminosalicylic acid (5-ASA, mesalamine), sulfapyridine, and capsaicin were purchased from Sigma (St. Louis, MO). 5'-Iodoresiniferatoxin (IRTX) was purchased from Tocris Cookson (Ellisville, MO). *Clostridium difficile* toxin A was purchased from TechLab, Inc. (Blacksburg, VA).

2.2. Surgery. Isolated colonic segments were constructed in anesthetized male Sprague-Dawley rats (150–175 g) as previously described for construction of ileal segments [17, 18]. Isolated colonic segments 5 cm in length were constructed distal to the caecum by ligation with silk sutures.

2.3. Drug Administration. Toxin A was administered at a dose of 5 μ g in 400 μ L (or 200 μ L when given after other drugs) of PBS into the lumen of the isolated colonic segments. 5-ASA was dissolved in phosphate buffered saline (PBS) and injected intraluminally in volumes of 200 μ L 30 minutes prior to toxin A injection. Sulfapyridine was initially dissolved in dimethyl sulfoxide (DMSO) and then diluted to 10% DMSO in PBS. Capsaicin was administered at a dose of 4 mg in 400 μ L (or 200 μ L when given after other drugs) of 25% ethanol in saline. All intraluminal injections were made using a 27 ga syringe needle. Control rats were prepared identically and their isolated colonic segments were injected with the appropriate vehicle solutions. Sulfasalazine was initially dissolved in 0.1 N NaOH at 30 times the desired final concentration and then the pH was adjusted to 8.5 with 1N HCl and the solution was diluted 30-fold with distilled water. Sulfasalazine was administered chronically in this solution as drinking water for 5 days. The final concentration of sulfasalazine

used was calculated from preliminary data obtained on the amount of water that the animals drank per day. The drinking water for control animals was prepared identically with the omission of sulfasalazine. The animals drank the same amount of the sulfasalazine vehicle as tap water. These studies were approved by the Duke University and Durham VA Institutional Animal Care and Use Committees.

2.4. Luminal Fluid Accumulation. Luminal fluid accumulation was measured gravimetrically. After 3 hours of treatment, the isolated colonic segments were removed and weighed, and their lengths were measured. Luminal fluid accumulation is expressed as mg wet weight per cm length.

2.5. Myeloperoxidase Activity. Myeloperoxidase (MPO) activity was measured as described previously [19]. Briefly, pieces of control and treated colonic segments were homogenized in 0.5% hexadecyltrimethylammonium bromide in 50 mM KH₂PO₄ (pH 6), freeze-thawed three times, and centrifuged at 4°C for 2 minutes, and then the absorbance of each supernatant was read at 460 nm at 0, 30, and 60 seconds after the addition of 2.9 mL of *o*-dianisidine dihydrochloride to 0.1 mL supernatant. The maximal change in absorbance per minute was used to calculate the units of MPO activity based on the molar absorptivity index of oxidized *o*-dianisidine of $1.13 \times 10^4 \text{ M}^{-1} \text{ cm}^{-1}$. The results are expressed as MPO units of activity per gram of tissue wet weight.

2.6. Histopathology. After 3 hours of treatment, equivalent portions of the isolated colonic segments were fixed overnight in 10% formalin, dehydrated, and embedded in paraffin, and then sections of 5 μ m in thickness were cut, mounted on glass slides, and stained with hematoxylin and eosin.

2.7. LTB₄ Measurement. Colonic LTB₄ levels were measured by LTB₄ enzyme immunoassay (EIA) kits purchased from Cayman Chemical (Ann Arbor, MI) as previously described [15]. Briefly, samples of colon and colonic luminal fluid were collected after various treatments in 5 volumes of ice-cold 0.1M phosphate buffer, pH 7.4, containing 1 mM EDTA and 10 μ M indomethacin and homogenized for 15 sec on ice using a Tekmar Tissumizer (Tekmar, Cincinnati, OH) at a 50% power setting. Before homogenization, 10,000 cpm of ³H-LTB₄ (120–240 Ci/mmol; Perkin-Elmer Life Sciences, Boston, MA) was added to the buffer for later assessment of LTB₄ recovery. After homogenization, 2 volumes of ice-cold ethanol were added to each extract and the extracts were then incubated on ice for 5 min to precipitate proteins. After centrifugation at 3000 $\times g_{\text{max}}$ to remove the precipitated proteins, the ethanol in the supernatants was removed by vacuum centrifugation. The pH of the extracts was adjusted to ~4.0 by addition of 1M sodium acetate (pH 4.0). The resulting precipitate was removed by centrifugation and the supernatant was loaded onto C-18 solid phase extraction cartridges (Cayman Chemical, Ann Arbor, MI) previously washed with methanol and distilled water, washed with distilled water followed by hexane, and then eluted at unit gravity




FIGURE 3: Effects of sulfasalazine, 5-ASA, and sulfapyridine on toxin A-induced colonic luminal fluid accumulation and MPO activity. Sulfasalazine was administered in the drinking water at a dose of 100 mg/kg-day for 5 days. 5-ASA and sulfapyridine were administered as pretreatments for 30 minutes before toxin A was injected into isolated colonic segments. (a) Toxin A stimulation of colonic luminal fluid accumulation and the inhibition of this effect by chronic treatment with sulfasalazine or acute treatment with 5-ASA or sulfapyridine. (b) Toxin A stimulation of colonic MPO activity and the inhibition of this effect by chronic treatment with sulfasalazine or acute treatment with 5-ASA or sulfapyridine. * $P < 0.05$ versus no treatment; ** $P < 0.01$ versus no treatment; **** $P < 0.001$ versus no treatment; #### $P < 0.0001$ versus toxin A.

with 5 mL of 99% ethyl acetate: 1% methanol. The samples were then evaporated to dryness by vacuum centrifugation, reconstituted in LTB_4 EIA buffer, and assayed according to the instructions of the kit manufacturer. Because toxin A caused portions of the colonic mucosa to slough off into the intestinal lumen, luminal contents were collected by syringe from toxin A-treated colonic segments and assayed for LTB_4 content just as for colonic tissue, and the LTB_4 contents of the colonic tissue and corresponding luminal contents were added together for these samples. Because it seemed inappropriate to express the results as LTB_4 concentrations per unit wet weight, therefore, the results are expressed as LTB_4 concentrations per cm of colonic length.

2.8. Substance P Release. Substance P (SP) release was assessed by analysis of NK-1R endocytosis as described previously [14, 20] with modifications. Briefly, pieces of colonic segments taken from control, toxin A-treated, and capsaizepine pretreated/toxin A-treated rats were fixed in freshly depolymerized 4% paraformaldehyde overnight at 4°C. The tissue was then washed and embedded in Tissue Tek O.C.T. compound (Sakura, Torrance, CA), frozen, sectioned at 20 μ m, and mounted on Superfrost/Plus glass slides (Fisher, Pittsburgh, PA). After washing, the slides were stained using a rabbit antiserum (number 11886-5) specific for the C-terminal 15 amino acids of the rat NK-1R at a dilution of 1:3000 [21]. This was followed by incubation in a cyanine 3-conjugated donkey anti-rabbit IgG secondary antibody (Jackson ImmunoResearch, West Grove, PA) at a dilution of 1:600. The stained sections were analyzed using a Zeiss LSM-510 META inverted krypton-argon confocal laser scanning system coupled to a Zeiss Axiovert 200

MOT microscope. Images of 512 \times 512 pixels were obtained and processed using Adobe Photoshop. Quantification of NK-1R endocytosis was achieved by analyzing 20 NK-1R-immunoreactive (NK-1R-ir) myenteric plexus neuronal cell bodies per rat and determining the number of these cells containing more than 10 NK-1R-ir endosomes. Myenteric plexus neuronal cell bodies were identified by their large size, position between the internal and external layers of smooth muscle of the muscularis externa, encapsulation by a perineurium, and their morphology. Cytoplasmic NK-1R-ir endosomes were distinguished from NK-1R-ir plasma membranes or plasma membrane-associated endosomes by ensuring that the nucleus of the neurons was in the same optical section as the NK-1R-ir endosomes.

2.9. Statistical Analysis. Results are expressed as mean \pm SEM ($N = 5-15$). Mean differences among 2 groups were examined by the Student t -test and mean differences among several groups by one-way ANOVA with Dunnett's or Tukey-Kramer posttests, using GraphPad InStat version 3.05 for Windows (GraphPad Software, San Diego, CA). P values < 0.05 were considered significant.

3. Results

We previously demonstrated that TRPV1 activation plays a role in toxin A-induced inflammation in the rat ileum [14, 15, 20] but the role of TRPV1 has not been examined in toxin A-induced colitis. Therefore, we first sought to determine if pharmacological antagonism of TRPV1 inhibits toxin A colitis as it does toxin A ileitis in rats. Intraluminal administration of toxin A in the rat colon caused an intense




FIGURE 4: The effects of toxin A on colonic histopathology and protection against these effects by sulfasalazine and 5-ASA but not sulfapyridine. H&E-stained sections of rat colonic segments after they were treated intraluminally for 3 h with vehicle (a), 5 µg of toxin A (b), 5 µg of toxin A after chronic treatment with 100 mg/kg-day of sulfasalazine for 5 days (c), 5 µg of toxin A after acute pretreatment with 100 µg of 5-ASA for 30 minutes (d), and 5 µg of toxin A after acute pretreatment with 100 µg of sulfapyridine for 30 minutes (e). Scale bar, 100 µm.




FIGURE 5: The effects of sulfasalazine, 5-ASA, and sulfapyridine on toxin A-induced substance P release in the colon. Laser scanning confocal fluorescent micrographs of neurons immunostained for the NK-1R from colonic segments treated for 3 h with (a) vehicle, (b) 5 μ g toxin A, (c) 5 μ g toxin A after 5 days of treatment with sulfasalazine (100 mg/kg-day), (d) 5 μ g toxin A after 30 minutes of intraluminal pretreatment with 5-ASA (100 μ g), and (e) 5 μ g toxin A after 30 minutes of intraluminal pretreatment with sulfapyridine (100 μ g). Bar, 5 μ m.

inflammation similar to that previously observed in the rat ileum. Toxin A stimulated luminal fluid accumulation, increased MPO activity, and caused intense histopathology characterized by loss of mucosal folding, surface ulceration, and influx of neutrophils (Figure 1). In order to determine if TRPV1 mediates a portion of toxin A-induced colitis as it does ileitis, colonic segments were pretreated for 30 minutes with intraluminal administration of a TRPV1-defunctionalizing

concentration of the TRPV1 partial agonist, I-RTX [22], before toxin A injection. I-RTX significantly inhibited toxin A-induced luminal fluid accumulation, MPO activity, and histopathology in the colon (Figure 1) just as it does in the ileum.

We previously provided evidence that LTB₄ mediates the inflammatory effects of toxin A in the rat ileum [15]. Therefore, we examined the effects of toxin A, 5-ASA, and




FIGURE 6: Quantitation of toxin A-induced SP release and its inhibition by sulfasalazine and 5-ASA but not by sulfapyridine. The percentage of NK-1R-ir neuronal cell bodies in the myenteric plexus of the colon with >10 NK-1R-ir endosomes was determined in 20 cells per animal for each treatment as an index of endogenous SP release. Toxin A stimulated SP release and this effect was significantly inhibited by sulfasalazine and 5-ASA but not sulfapyridine. * $P < 0.05$ versus no treatment; ** $P < 0.001$ versus no treatment; # $P < 0.001$ versus toxin A alone treatment.

toxin A after 5-ASA pretreatment on levels of LTB_4 in the colon. Toxin A stimulated a significant increase in colonic tissue concentrations of LTB_4 (Figure 2). 5-ASA alone had no significant effect on tissue LTB_4 concentrations. Intraluminal pretreatment of the colonic segment for 30 minutes with 100 μg of 5-ASA inhibited the toxin A-induced increase in colonic LTB_4 levels (Figure 2).

When sulfasalazine was administered chronically to the rats for 5 days in their drinking water, it had no effect by itself but inhibited toxin A-stimulated luminal fluid accumulation and MPO activity (Figure 3). We next tested the effects of the two active constituents of sulfasalazine, 5-ASA and sulfapyridine. 5-ASA given alone caused a small but significant decrease in luminal fluid accumulation but had no effect on MPO activity. Sulfapyridine given alone caused a significant increase in MPO activity but had no effect on luminal fluid accumulation (Figure 3). Pretreatment of the colon for 30 minutes with 5-ASA before toxin A was administered resulted in highly significant inhibition of toxin A-induced luminal fluid accumulation and MPO activity (Figure 3). Pretreatment with sulfapyridine did not inhibit toxin A-induced colonic inflammation (Figure 3). When histopathology was examined, it was clear that both sulfasalazine and 5-ASA were highly protective of the structure of the colon against toxin A inflammation but sulfapyridine was not (Figure 4).

Because it has previously been shown that toxin A stimulates release of the proinflammatory neurotransmitter SP in the rat ileum via activation of TRPV1 [14, 17], we reasoned that if 5-ASA inhibits toxin A-induced colitis by blocking the generation of an endogenous TRPV1 agonist such as LTB_4 , then 5-ASA and sulfasalazine, but not sulfapyridine, should inhibit toxin A-induced SP release in

the colon. Therefore, we measured SP release in response to intraluminal toxin A and tested the effects of 5-ASA on SP release. We used immunocytochemical assessment of NK-1R endocytosis in myenteric plexus neuronal cell bodies as an index of endogenous SP release and as an index of the action of locally released SP as described previously [14, 17, 20, 23]. Intraluminal administration of toxin A caused NK-1R endocytosis, reflecting the action of endogenously released SP, and this effect was inhibited by chronic treatment with sulfasalazine and acute pretreatment with 5-ASA but not sulfapyridine (Figure 5). When quantitated, toxin A-induced SP release was statistically significant and sulfasalazine and 5-ASA significantly inhibited toxin A-induced SP release (Figure 6). Sulfapyridine did not significantly inhibit toxin A-induced SP release (Figure 6).

If 5-ASA blocks toxin A colitis by inhibiting the generation of an endogenous TRPV1 agonist such as LTB_4 , we reasoned that 5-ASA pretreatment should not inhibit colitis caused by an agent that directly stimulates TRPV1. Capsaicin, the pungent ingredient in chili peppers, is an activator of TRPV1 that has been shown to cause enteritis that is very similar to that caused by toxin A in the rat ileum [14]. Intraluminal injection of capsaicin in isolated rat colonic segments caused intense inflammation as assessed by luminal fluid accumulation and increased tissue MPO levels (Figure 7). Pretreatment of the segments for 30 minutes with 5-ASA had no effect on capsaicin-induced colitis, thus supporting the concept that 5-ASA protects against colitis by inhibiting toxin A-induced generation of an endogenous TRPV1 agonist such as LTB_4 . These results also demonstrate that 5-ASA is not itself a TRPV1 antagonist.

4. Discussion

Our first goal was to determine if TRPV1 activation plays a role in toxin A-induced colitis as it does in toxin A-induced ileitis in rats [14, 15, 20]. Intraluminal administration of toxin A into isolated segments of rat colon caused an intense inflammation similar to that previously described [24]. When pretreated for 30 minutes with a defunctionalizing dose of the specific TRPV1 partial agonist, I-RTX, toxin A-induced luminal fluid and MPO activity were significantly inhibited and normal colonic mucosal histology was preserved. The anti-inflammatory effects of TRPV1 inhibition in the rat colon are similar to those seen in the rat ileum previously and indicate that TRPV1 activation is important in toxin A colitis as well as in ileitis.

Toxin A also stimulated increased LTB_4 levels in the rat colon just as it does in the rabbit [25] and rat [15] ileum. We have previously demonstrated that LTB_4 causes ileitis similar to that seen after toxin A administration and that the inflammatory effects of both LTB_4 and toxin A are strongly inhibited by TRPV1 antagonism [15]. In addition, pharmacological inhibition of 5-LO activity inhibited both toxin A-induced ileal LTB_4 production and ileitis [15]. These findings coupled with the observations that LTB_4 is proinflammatory in human UC and that drugs containing 5-ASA inhibit LTB_4 production and colonic inflammation in UC led us to




FIGURE 7: Effects of 5-ASA on capsaicin-induced colonic luminal fluid accumulation and MPO activity. 5-ASA was administered intraluminally at a dose of 100 μg for 30 minutes before capsaicin. (a) Capsaicin stimulation of colonic luminal fluid accumulation and the lack of significant inhibition of this effect by acute pretreatment with 5-ASA. (b) Capsaicin stimulation of colonic MPO activity and the lack of significant inhibition of this effect by acute pretreatment with 5-ASA. H&E-stained sections of rat colonic segments after they were treated intraluminally for 3 h with vehicle (c), 100 μg of 5-ASA (d), 4 mg of capsaicin (e), or 4 mg of capsaicin after pretreatment with 100 μg of 5-ASA 30 min before toxin A administration (f). Scale bar, 100 μm . * $P < 0.001$ versus sulfapyridine-/toxin A-.

hypothesize that 5-ASA may also inhibit toxin A colitis by inhibiting 5-LO in the rat colon. It was previously shown that 5-ASA inhibits toxin A-induced LTB_4 release and mucosal permeability to mannose in the rabbit ileum but does not inhibit fluid secretion or morphological damage [25]. In the present study, however, 5-ASA significantly inhibited toxin A-induced luminal fluid accumulation and morphological

damage as well as MPO activity in the rat colon. These differing results may be due to species differences, organ differences, or other unknown factors.

In order to differentiate the anti-inflammatory mechanism of action of 5-ASA from sulfasalazine, which contains one molecule of 5-ASA coupled to one molecule of sulfapyridine via an azo bond, we separately tested the acute effects

of sulfapyridine in the toxin A model of rat colitis. This was important because 5-ASA has been shown to be the active moiety of sulfasalazine in preventing human UC [3] and proctitis [4] and because sulfasalazine but not 5-ASA has been shown to be an inhibitor of the proinflammatory nuclear factor, nuclear factor kappa B (NF- κ B) [26]. We found that sulfapyridine had no effect on toxin A colitis in rats, suggesting that 5-ASA is the active component of sulfasalazine in toxin A colitis just as it is in human UC and proctitis.

The present results suggest the possibility that sulfasalazine may be therapeutically useful in treating human *C. difficile* colitis. For example, although this disease is usually effectively treated by antibiotics [27], some cases of recurrent disease have proven difficult to treat [28] and new, more virulent strains of *C. difficile* have emerged [29] that may require new therapeutic approaches.

An interesting observation concerning the present results is that 5-ASA, a drug that has proven efficacious in treating chronic intestinal inflammatory diseases such as UC and CD, also strongly inhibits acute *C. difficile* toxin A-induced colitis. Although the causes of UC and CD are unknown, there is strong evidence that they are related to the commensal bacteria of the gut and depend on a T cell-mediated adaptive immune response. In contrast, toxin A colitis occurs in response to bacterial exotoxins, not the *C. difficile* organism itself, occurs too rapidly in the rat model to involve adaptive immune responses, and appears to be mediated largely by a neutrophilic innate immune response [24, 30]. Although it is possible that 5-ASA has different mechanisms of action in inhibiting chronic versus acute intestinal inflammation, it may be that its anti-inflammatory effect is due to inhibition of 5-LO in both cases. Support for this concept comes from animal studies showing that inhibition of 5-LO is also efficacious in blocking chronic colitis resulting from administration of trinitrobenzene sulfonic acid (TNBS) [11, 13, 31, 32]. Additionally, support for the involvement of TRPV1 in chronic colitis comes from animal studies of TNBS colitis [33, 34], dextran sulfate-induced colitis [35, 36], and colitis caused by adoptive transfer of CD4⁺/CD25⁻ T cells in SCID mice [37]. Based on these observations, it will be important in future studies to determine the role of LTB₄ activation of TRPV1 in the human IBDs.

Conflict of Interests

The author declares that there is no conflict of interests regarding the publication of this paper.

Acknowledgments

The author thanks Douglas C. McVey for technical support. This work was supported by a Veterans Affairs Merit Review Grant and National Institutes of Health Grant DK-50265.

References

- [1] D. C. McVey, R. A. Liddle, J. Riggs-Sauthier, N. Ekwuribe, and S. R. Vigna, "Inhibition of *Clostridium difficile* toxin A-induced colitis in rats by APAZA," *Digestive Diseases and Sciences*, vol. 50, no. 3, pp. 565–573, 2005.
- [2] H. Schröder and B. E. Gustafsson, "Azo reduction of salicyl azo sulphapyridine in germ free and conventional rats," *Xenobiotica*, vol. 3, no. 4, pp. 225–231, 1973.
- [3] A. K. A. Khan, J. Piris, and S. C. Truelove, "An experiment to determine the active therapeutic moiety of sulphasalazine," *The Lancet*, vol. 2, no. 8044, pp. 892–895, 1977.
- [4] P. A. M. Van Hees, J. H. Bakker, and J. H. M. van Tongeren, "Effect of sulphapyridine, 5-aminosalicylic acid, and placebo in patients with idiopathic proctitis: a study to determine the active therapeutic moiety of sulphasalazine," *Gut*, vol. 21, no. 7, pp. 632–635, 1980.
- [5] R. P. MacDermott, "Progress in understanding the mechanisms of action of 5-aminosalicylic acid," *The American Journal of Gastroenterology*, vol. 95, no. 12, pp. 3343–3345, 2000.
- [6] K. Nimmagadda and S. B. Hanauer, "Aminosalicylates in inflammatory bowel disease," *Clinical Perspectives in Gastroenterology*, vol. 5, no. 4, pp. 229–233, 2002.
- [7] S. M. Greenfield, N. A. Panchard, J. P. Teare, and R. P. H. Thompson, "Review article: the mode of action of the aminosalicylates in inflammatory bowel disease," *Alimentary Pharmacology and Therapeutics*, vol. 7, no. 4, pp. 369–383, 1993.
- [8] J. Rask-Madsen, K. Bukhave, L. S. Laursen, and K. Lauritsen, "5-Lipoxygenase inhibitors in the treatment of inflammatory bowel disease," *Advances in Prostaglandin, Thromboxane, and Leukotriene Research*, vol. 22, pp. 113–124, 1994.
- [9] L. G. Axelsson and S. Ahlstedt, "Actions of sulphasalazine and analogues in animal models of experimental colitis," *Inflammopharmacology*, vol. 2, no. 3, pp. 219–232, 1993.
- [10] P. Sharon and W. F. Stenson, "Enhanced synthesis of leukotriene B₄ by colonic mucosa in inflammatory bowel disease," *Gastroenterology*, vol. 86, no. 3, pp. 453–460, 1984.
- [11] J. L. Wallace, W. K. MacNaughton, G. P. Morris, and P. L. Beck, "Inhibition of leukotriene synthesis markedly accelerates healing in a rat model of inflammatory bowel disease," *Gastroenterology*, vol. 96, no. 1, pp. 29–36, 1989.
- [12] R. D. Zipser, C. C. Nast, M. Lee, H. W. Kao, and R. Duke, "In vivo production of leukotriene B₄ and leukotriene C₄ in rabbit colitis. Relationship to inflammation," *Gastroenterology*, vol. 92, no. 1, pp. 33–39, 1987.
- [13] J. L. Wallace and C. M. Keenan, "An orally active inhibitor of leukotriene synthesis accelerates healing in a rat model of colitis," *The American Journal of Physiology—Gastrointestinal and Liver Physiology*, vol. 258, no. 4, pp. G527–G534, 1990.
- [14] D. C. McVey and S. R. Vigna, "The capsaicin VR1 receptor mediates substance P release in toxin A-induced enteritis in rats," *Peptides*, vol. 22, no. 9, pp. 1439–1446, 2001.
- [15] D. C. McVey and S. R. Vigna, "The role of leukotriene B₄ in *Clostridium difficile* toxin A—induced ileitis in rats," *Gastroenterology*, vol. 128, no. 5, pp. 1306–1316, 2005.
- [16] S. W. Hwang, H. Cho, J. Kwak et al., "Direct activation of capsaicin receptors by products of lipoxygenases: endogenous capsaicin-like substances," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 97, no. 11, pp. 6155–6160, 2000.
- [17] C. R. Mantyh, T. N. Pappas, J. A. Lapp et al., "Substance P activation of enteric neurons in response to intraluminal *Clostridium difficile* toxin A in the rat ileum," *Gastroenterology*, vol. 111, no. 5, pp. 1272–1280, 1996.

- [18] C. Pothoulakis, I. Castagliuolo, J. T. Lamont et al., "CP-96,345, a substance P antagonist, inhibits rat intestinal responses to *Clostridium difficile* toxin A but not cholera toxin," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 91, no. 3, pp. 947–951, 1994.
- [19] P. P. Bradley, D. A. Priebe, R. D. Christensen, and G. Rothstein, "Measurement of cutaneous inflammation: estimation of neutrophil content with an enzyme marker," *Journal of Investigative Dermatology*, vol. 78, no. 3, pp. 206–209, 1982.
- [20] D. C. McVey, P. C. Schmid, H. H. O. Schmid, and S. R. Vigna, "Endocannabinoids induce ileitis in rats via the capsaicin receptor (VR1)," *Journal of Pharmacology and Experimental Therapeutics*, vol. 304, no. 2, pp. 713–722, 2003.
- [21] S. R. Vigna, J. J. Bowden, D. M. McDonald et al., "Characterization of antibodies to the rat substance P (NK-1) receptor and to a chimeric substance P receptor expressed in mammalian cells," *The Journal of Neuroscience*, vol. 14, no. 2, pp. 834–845, 1994.
- [22] I. Shimizu, T. Iida, N. Horiuchi, and M. J. Caterina, "5-Iodoresiniferatoxin evokes hypothermia in mice and is a partial transient receptor potential vanilloid 1 agonist in vitro," *Journal of Pharmacology and Experimental Therapeutics*, vol. 314, no. 3, pp. 1378–1385, 2005.
- [23] P. W. Mantyh, E. DeMaster, A. Malhotra et al., "Receptor endocytosis and dendrite reshaping in spinal neurons after somatosensory stimulation," *Science*, vol. 268, no. 5217, pp. 1629–1632, 1995.
- [24] C. Pothoulakis, "Pathogenesis of *Clostridium difficile*-associated diarrhoea," *European Journal of Gastroenterology & Hepatology*, vol. 8, no. 11, pp. 1041–1047, 1996.
- [25] G. Triadafilopoulos, C. Pothoulakis, R. Weiss, C. Giampaolo, and J. T. LaMont, "Comparative study of *Clostridium difficile* toxin A and cholera toxin in rabbit ileum," *Gastroenterology*, vol. 97, no. 5, pp. 1186–1192, 1989.
- [26] C. Wahl, S. Liptay, G. Adler, and R. M. Schmid, "Sulfasalazine: a potent and specific inhibitor of nuclear factor kappa B," *The Journal of Clinical Investigation*, vol. 101, no. 5, pp. 1163–1174, 1998.
- [27] C. P. Kelly, C. Pothoulakis, and J. T. Lamont, "*Clostridium difficile* colitis," *The New England Journal of Medicine*, vol. 330, no. 4, pp. 257–262, 1994.
- [28] L. Kyne and C. P. Kelly, "Recurrent *Clostridium difficile* diarrhoea," *Gut*, vol. 49, no. 1, pp. 152–153, 2001.
- [29] J. G. Bartlett and T. M. Perl, "The new *Clostridium difficile*—what does it mean?" *The New England Journal of Medicine*, vol. 353, no. 23, pp. 2503–2505, 2005.
- [30] C. Pothoulakis and J. T. Lamont, "Microbes and microbial toxins: paradigms for microbial-mucosal interactions II. The integrated response of the intestine to *Clostridium difficile* toxins," *American Journal of Physiology*, vol. 280, no. 2, pp. G178–G183, 2001.
- [31] C. Bobin-Dubigeon, X. Collin, N. Grimaud, J. Robert, G. Le Baut, and J. Petit, "Effects of tumour necrosis factor- α synthesis inhibitors on rat trinitrobenzene sulphonic acid-induced chronic colitis," *European Journal of Pharmacology*, vol. 431, no. 1, pp. 103–110, 2001.
- [32] B. Zingarelli, F. Squadrito, P. Graziani, R. Camcrini, and A. P. Caputi, "Effects of zileuton, a new 5-lipoxygenase inhibitor, in experimentally induced colitis in rats," *Agents and Actions*, vol. 39, no. 3-4, pp. 150–156, 1993.
- [33] K. Fujino, Y. Takami, S. G. de la Fuente et al., "Inhibition of the vanilloid receptor subtype-1 attenuates TNBS-colitis," *Journal of Gastrointestinal Surgery*, vol. 8, no. 7, pp. 842–848, 2004.
- [34] A. Miranda, E. Nordstrom, A. Mannem, C. Smith, B. Banerjee, and J. N. Sengupta, "The role of transient receptor potential vanilloid 1 in mechanical and chemical visceral hyperalgesia following experimental colitis," *Neuroscience*, vol. 148, no. 4, pp. 1021–1032, 2007.
- [35] N. Kihara, S. G. de la Fuente, K. Fujino, T. Takahashi, T. N. Pappas, and C. R. Mantyh, "Vanilloid receptor-1 containing primary sensory neurones mediate dextran sulphate sodium induced colitis in rats," *Gut*, vol. 52, no. 5, pp. 713–719, 2003.
- [36] I. Szitter, G. Pozsgai, K. Sandor et al., "The role of transient receptor potential vanilloid 1 (Trpv1) receptors in dextran sulfate-induced colitis in mice," *Journal of Molecular Neuroscience*, vol. 42, no. 1, pp. 80–88, 2010.
- [37] M. Gad, A. E. Pedersen, N. N. Kristensen, C. D. F. Fernandez, and M. H. Claesson, "Blockage of the neurokinin 1 receptor and capsaicin-induced ablation of the enteric afferent nerves protect SCID mice against T-cell-induced chronic colitis," *Inflammatory Bowel Diseases*, vol. 15, no. 8, pp. 1174–1182, 2009.