

Research article

Open Access

## Identification of novel light-induced genes in the suprachiasmatic nucleus

Veronica M Porterfield<sup>1</sup>, Helen Piontkivska<sup>1,2</sup> and Eric M Mintz\*<sup>1,2</sup>

Address: <sup>1</sup>School of Biomedical Sciences, Kent State University, Kent, OH 44242 USA and <sup>2</sup>Department of Biological Sciences, Kent State University, Kent, OH 44242 USA

Email: Veronica M Porterfield - vporterf@kent.edu; Helen Piontkivska - opiontki@kent.edu; Eric M Mintz\* - emintz@kent.edu

\* Corresponding author

Published: 19 November 2007

Received: 29 August 2007

BMC Neuroscience 2007, 8:98 doi:10.1186/1471-2202-8-98

Accepted: 19 November 2007

This article is available from: <http://www.biomedcentral.com/1471-2202/8/98>

© 2007 Porterfield et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### Abstract

**Background:** The transmission of information about the photic environment to the circadian clock involves a complex array of neurotransmitters, receptors, and second messenger systems. Exposure of an animal to light during the subjective night initiates rapid transcription of a number of immediate-early genes in the suprachiasmatic nucleus of the hypothalamus. Some of these genes have known roles in entraining the circadian clock, while others have unknown functions. Using laser capture microscopy, microarray analysis, and quantitative real-time PCR, we performed a comprehensive screen for changes in gene expression immediately following a 30 minute light pulse in suprachiasmatic nucleus of mice.

**Results:** The results of the microarray screen successfully identified previously known light-induced genes as well as several novel genes that may be important in the circadian clock. Newly identified light-induced genes include early growth response 2, proviral integration site 3, growth-arrest and DNA-damage-inducible 45 beta, and TCDD-inducible poly(ADP-ribose) polymerase. Comparative analysis of promoter sequences revealed the presence of evolutionarily conserved CRE and associated TATA box elements in most of the light-induced genes, while other core clock genes generally lack this combination of promoter elements.

**Conclusion:** The photic signalling cascade in the suprachiasmatic nucleus activates an array of immediate-early genes, most of which have unknown functions in the circadian clock. Detected evolutionary conservation of CRE and TATA box elements in promoters of light-induced genes suggest that the functional role of these elements has likely remained the same over evolutionary time across mammalian orders.

### Background

Circadian rhythms in mammals are driven by a clock located in the suprachiasmatic nucleus of the hypothalamus (SCN) [1]. The SCN is directly innervated by the retinas and photic information is transmitted to the SCN via a NMDA and pituitary adenylate cyclase activating peptide-dependent mechanism [2]. Photic input to the cir-

dian clock has differential effects on circadian rhythms depending on the timing of exposure to light. Light presented in the early night delays the phase of the clock, light during the late night advances the clock, and light presented during the subjective day has little or no effect on clock phase [3]. Photic signals during the subjective night activate a MAP kinase signaling pathway leading to

increased transcription of several immediate-early genes [4] and the core clock gene *Per1* [2].

The roles of light-induced genes in the circadian clock mechanism, other than the period genes, are not well understood. For example, the best characterized of these genes are *Fos*, *Egr1*, and *Nr4a1* (*nur77*). The expression of *c-fos* protein is commonly used as a marker for light-like activation of neurons in the SCN. *Fos* knockout mice still show behavioral circadian rhythms as well as a phase response curve to light, but the amplitude of the circadian rhythm of activity and the phase response curve are attenuated [5]. The threshold for photic induction of *c-fos* is similar to the threshold for behavioral phase shifts [6], however, expression of *c-fos* can be dissociated from behavioral shifts [7-9]. *Fos* appears to be a component of the circadian response to light, but not one that is absolutely necessary for phase shifts to occur, and the mechanism by which *Fos* protein is involved in photic signalling is unknown. In contrast, *Egr1* and *Nr4a1* knockout mice show normal entrainment patterns and no dysfunction in their response to light [10]. The expression of *egr1* occurs over a broader area of


the SCN than does *c-fos* [11], but the threshold for *egr1* induction is lower than for phase shifts of behavioral rhythms or *c-fos* induction. These data suggest that *Egr1* and *Nr4a1* may not be directly involved in entrainment pathways, but the lower threshold for stimulation of expression suggests that *egr1* may be involved in other SCN outputs, such as the one which regulates rhythmic melatonin secretion from the pineal gland, which has a lower light intensity threshold for photic regulation than behavioral phase shifts.

In order to better understand the molecular events governing the response of the SCN to photic input, we performed a microarray-based screen for genes rapidly induced by light, followed by a comparative evolutionary genomic analysis to identify common activation mechanisms among this gene population.

## Results

### **Light-induced immediate-early genes in the SCN**


We used laser capture microscopy (Figure 1) to isolate SCN tissue from mice immediately following either a 30 minute high-intensity light pulse or a sham light pulse.



**Figure 1**  
**Laser capture and gene expression in response to a light pulse.** (a) Twelve  $\mu\text{m}$  thick coronal section containing the SCN, indicated by arrows. Sections are stained with hematoxylin. (b) The same section after capture, showing the removal of the paired SCN.

The extent of the SCN was easily determined from the density of hemotoxylin-labeled nuclei. We examined a selected set of genes known to be markers of the SCN, but not the surrounding tissue, to confirm that we were measuring SCN gene expression. All 6 arrays displayed strong hybridization signals for the mRNA for vasoactive intestinal polypeptide (VIP), gastrin-releasing peptide, arginine vasopressin, calbindin 28 K, Clock, enkephalin, BMAL, and glutamic acid decarboxylase 67. VIP produced hybridization signals that were in the top 20 genes on 5 out of the 6 arrays (21<sup>st</sup> on the 6<sup>th</sup> array). In contrast, neural markers that are not expressed in the SCN, but are expressed in SCN-projecting neurons, received absent calls from all 6 arrays, including neuropeptide Y, preprohypocretin, tryptophan hydroxylase, and tyrosine hydroxylase.

The 22,626 probe sets on the Affymetrix arrays were reduced to 10,340 candidate transcripts for analysis based on the criteria that each probe set was identified as being present using the default settings of GCOS 1.1 software on at least 3 of the 6 arrays. The results of these probe sets are plotted in Figure 2. Differences between light pulse and sham light pulse groups were assessed according to p-values generated using a Bayesian t-test calculated using the logarithms of the signal intensities for each probe set. A total of six genes were measured as being differentially expressed using our most stringent criteria, which was the application of a Bonferroni correction to the critical p-



**Figure 2**  
**Gene expression after light pulse vs. sham light pulse.**  
 Summary comparison of gene expression after a light pulse vs. control dark pulse. Each point represents one gene that was present on at least half of the arrays. The boxes indicate data points that were subsequently confirmed as significantly different between conditions by qPCR.

value for significance ( $p < 4.84E-06$ ). A second group of 9 genes was identified as meeting a lesser criteria of  $p < 0.001$ . All 15 genes of these genes were then analyzed for differences in expression using quantitative real-time PCR (qPCR). All 6 of the genes in the first group were confirmed as having a more than 2-fold increase in expression following the light pulse, while 6 of the 9 genes in the second group were confirmed. In addition, we tested the fold-change of 3 additional genes that had not been identified in our microarray analysis but had previously been shown to be induced by a light pulse (*Egr3*, *Per1*, and *Per2*) [12,13]. Of this group, *Egr3* showed a significant increase in expression following the light pulse but no significant difference in *Per1* or *Per2* expression was detected. The identities of these genes and their fold-changes as calculated using the microarrays and qPCR are shown in Table 1 and Figure 3. The correlation coefficient between fold changes determined by microarray analysis and by real-time PCR was 0.87 (Spearman's rank-correlation,  $p = 0.00003$ ). There were no transcripts that showed a significant decrease in expression following the light pulse.

The failure to detect an increase in *Per1* expression was unexpected, as some previous reports detected an increase in *Per1* mRNA expression within 10 minutes of the onset of a nocturnal light pulse [14]. However, we hypothesized that the increase in *Per1* might be occurring later in our study. Therefore, we harvested SCN tissue from mice one hour after the start of a 30 minute light pulse or sham light pulse, 30 minutes later than in our initial study. qPCR assays of these captures revealed a significant increase in both *Per1* (7.0 $\times$  over dark pulse, 95% confidence interval of 2.8 – 17.8,  $n = 4$  per group) and *Per2* (1.9 $\times$  over dark pulse, 95% confidence interval of 1.1 – 3.4,  $n = 4$  per group) expression after the light pulse as compared to the sham light pulse.

#### **Analysis of promoter regions of light-induced genes**


We reasoned that if these light-induced genes were important to the basic mechanism of photic signalling to the SCN and were activated by a common mechanism, they may share common regulatory sequences.  $Ca^{2+}$ /cAMP response elements (CRE) have been shown to play a critical role in circadian rhythmicity and the photic entrainment system [15,16]. We therefore took a comparative evolutionary genomic approach to ask whether all of the light-inducible genes in the SCN that we detected contain conserved CRE elements in their promoters. We found at least one highly conserved CRE element within 2 kb upstream of the start codon in 12 of the 13 genes induced by light immediately following the light pulse (Figure 4, Table 2 Additional file 1). The only gene lacking a conserved CRE element within 2 kb of the start codon was *Nr4a1*. Most of these CRE elements were highly conserved not only at the nucleotide sequence level but also in their

**Table 1: Fold change in expression following a light pulse**

Gene Symbol	Gene Title	Entrez Gene ID	Fold-Array*	p-value**
Egr1	early growth response 1	13653	27.7	3.45E-13
Nr4a1	nuclear receptor subfamily 4, group A, member 1	15370	10.2	2.13E-09
Egr2	early growth response 2	13654	47.3	4.19E-08
Dusp1	dual specificity phosphatase 1	19252	5.8	6.18E-08
Rrad	Ras-related associated with diabetes	56437	26.2	7.89E-08
Pim3	proviral integration site 3	223775	2.9	1.32E-06
Klf4	Kruppel-like factor 4 (gut)	16600	3.4	2.07E-04
Fos	FBJ osteosarcoma oncogene	14281	29.3	2.18E-04
Gadd45b	growth arrest and DNA-damage-inducible 45 beta	17873	3.2	2.58E-04
Btg2	B-cell translocation gene 2, anti-proliferative	12227	3.3	5.84E-04
Tiparp	TCDD-inducible poly(ADP-ribose) polymerase	99929	2.4	6.02E-04
Jun	Jun oncogene	16476	3.5	6.99E-04

\* Fold-array indicates the fold-difference in gene expression following a light pulse as compared to a sham light pulse. \*\* Bayesian t-test using Cyber-T software (see Methods section for details)

relative positions within the promoter region in the five mammalian species examined (mouse, rat, human, cow and dog), suggesting that these elements are important to coordinate the regulation of this group of genes. Previous studies have also demonstrated that cAMP-dependent activation of transcription by p-CREB largely depends on the presence of TATA elements downstream of the CRE element [17]. We found that 8 out of the 12 genes with CRE elements also possess TATA elements in proximity to the CRE element, including the 4 genes that showed the largest increases in expression following the light pulse. By these same criteria, *Per1* has both a conserved CRE element and a conserved TATA box, while *Per2* has neither. In contrast, in 8 other clock genes (Table 3) that were not upregulated 30 minutes after the light pulse onset in this study, only one (*Csnk1d*) had both a conserved CRE element and a nearby TATA box. This difference (12/13 vs. 1/8) is statistically significant (Fisher's Exact Test,  $p < 0.001$ ).




**Figure 3**  
**Fold change of light-induced genes, measured by qPCR.** Fold change following a light pulse of genes significantly upregulated following the light pulse as compared to the sham light pulse, as measured by qPCR. Error bars represent 95% confidence intervals.

**Discussion**

The results of this study identify a total of 13 transcripts that show significant increases in expression 30 minutes after the onset of a light pulse during the early night. It is important to note that this study examined gene expression in only the earliest moments of the response to photic stimulation. It is likely that repeating this study at later time points would reveal additional light-induced genes.

Four of the differentially expressed transcripts identified in the microarray analysis have not previously been identified as being induced by light: *Egr2*, *Pim3*, *Gadd45b*, and *Tiparp*. *Egr2* is a zinc-finger transcription factor of the same family as *Egr1* and *Egr3*. There is insufficient information available to generate a clear hypothesis about the function of *Egr* genes in the SCN. *Pim3* is a serine/threonine protein kinase that has anti-apoptotic functions in a variety of tissues [18,19], but its role in neurons has not been investigated. In contrast, the known functions of *Gadd45b* and *Tiparp* suggest potential connections to SCN function. In beta cells, *Gadd45b* inhibits apoptosis as well as JNK and ERK activation [20]. *Gadd45b* has similar functions in hematopoietic cells [21]. *Gadd45b* is upregulated in the hippocampus in response to electroconvulsive shock, consistent with a neuroprotective function [22]. If *Gadd45b* acts to inhibit ERK activation in the SCN, then that would be consistent with the idea that some of these light-induced genes act to downregulate the sensitivity of the SCN to subsequent stimuli, and possibly contribute to the SCN's strong resistance to excitotoxicity [23].

*Tiparp* encodes TCDD-inducible poly [ADP-ribose] polymerase and is activated through an aryl hydrocarbon receptor-dependent pathway [24]. Activation of this pathway by tryptophan photoproducts alters the expression of clock genes and inhibits glutamate-induced phase shifts in SCN 2.2 cells [25]. The aryl hydrocarbon receptor



**Figure 4**  
**Conserved promoter elements in light-induced genes.** Conserved CRE and TATA-box elements in promoter regions of four immediate-early genes in five mammalian genomes: (A) *Egr1*, (B) *Fos*, (C) *Per1* and (D) *Klf4*. Translational start codon ATG begins at +1. Alignment gaps are shown as dashes (-), dots (.) indicate identity to the first sequence. For *Klf4*, genomic sequence of dog was not available. Predicted conserved CRE elements are outlined with shaded boxes; putative TATA-boxes are shown with white boxes.

shares significant structural similarity with *Bmal1*, a core clock gene, and may interact directly with *Bmal1*, providing a potential substrate for altering clock function [26]. The induction of *Tiparp* provides another potential mechanism for altering clock function, although the targets for ribosylation in the SCN are unknown.

Three genes identified in our study (*Rrad*, *Btg2*, *Klf4*) were recently reported as light-induced in a study using high-coverage expression profiling [27], and another gene, *Dusp1*, was also recently characterized as light-induced [28]. *Dusp1* encodes MAP kinase phosphatase 1 (MKP1), and is now thought to play a role in the termination of the photic signalling cascade. One of the critical steps in the response of the SCN to photic signaling is the phosphorylation of the extracellular signal regulated kinases (ERK), leading to phosphorylation of CREB and changes in transcription [29,30]. An increase in functional MKP1 would reduce the sensitivity of the SCN to photic stimuli, and shut down the transcriptional mechanisms turned on by light. It also might help explain the extraordinary resistance of SCN cells to excitotoxicity [23], by downregulating the responses of SCN cells during periods of extended stimulation.

*Rrad*, *Btg2*, and *Klf4* have not been examined in the context of circadian clock function. However, given what is known about the role of *Rrad* in other systems, increased *Rrad* protein would also serve to downregulate the response of SCN cells to subsequent stimulation. *Rrad*

encodes a small GTPase that binds to calmodulin and CaM Kinase II [31]. Increased *Rrad* expression is associated with the removal of voltage-gated calcium channels from the plasma membrane [32], which would reduce the response of SCN neurons to excitatory stimuli. In addition, CaM Kinase II has already been shown to play a role in the regulation of light-induced phase delays, through actions leading to the transcription of *Per1* [33,34]. Consistent with this theme of downregulating cellular responses to stimuli, *Btg2* and *Klf4* both have anti-apoptotic functions in other cellular systems [35,36], but their role in the SCN remains unknown.

The remaining five genes are well known light-induced genes: *Egr1*, *Egr3*, *Fos*, *Nr4a1*, and *Jun*. Investigations into the function of these genes in the response of the SCN to photic stimuli have been extremely limited. Mice lacking functional copies of *Egr1* and *Nr4a1* (*nur77*) do not show altered circadian rhythms, suggesting that these genes do not play an important role in entrainment [10], although there could be compensatory mechanisms in the knock-out mice that mask the role of these genes. In contrast, mice lacking *Fos* show an attenuated phase response curve to light [5], suggesting that this gene is involved in the signal transduction mechanism for conveying photic information to the molecular clock mechanism. The critical factor in the response of the circadian clock to light is the increase in expression of period genes, through a mechanism that does not appear to require *Fos*. One possibility is that mechanisms within the cell that are activated by

**Table 2: Conservation of CRE and TATA elements in light-induced genes**

Ensembl gene ID	Gene name	Number of conserved CRE elements*	Genomic positions of CRE elements**	Genomic positions of TATA boxes
ENSMUSG00000020423	Btg2	2	-277 to -257 (3)	No TATA
ENSMUSG00000024190c	Dusp1	2	-259 to -239 -349 to -329 -301 to -281 (4)	-180 to -175***
ENSMUSG00000038418	Egr1	3	-925 to -905 -424 to -404 -355 to -335	-363 to -360 -686 to -682
ENSMUSG00000037868	Egr2	1	-414 to -394	-302 to -298
ENSMUSG00000033730	Egr3	2	-645 to -625 (2)	-599 to -596
ENSMUSG00000021250	Fos	3	-592 to -572 -500 to -480 -452 to -432 (2)	-183 to -178
ENSMUSG00000015312	Gadd45b	1	-925 to -905	No TATA
ENSMUSG00000052684	Jun	2	-1261 to -1241 -1002 to -982	-505 to - 502****
ENSMUSG00000003032	Klf4	2	-1023 to - 1003 -986 to -966	-778 to -774
ENSMUSG00000023034	Nr4a1	N/A****	N/A	No TATA
ENSMUSG00000020893	Per1	1	-355 to -335 (2)	-500 to -497
ENSMUSG00000055866	Per2	N/A	N/A	No TATA
ENSMUSG00000035828	Pim3	1	-686 to -666 (2)	No TATA
ENSMUSG00000031880c	Rrad	1	-522 to -502	No TATA
ENSMUSG00000034640	Tiparp	1	-1863 to - 1843	-1615 to -1610

\* We defined CRE element as being evolutionary conserved if it was present in at least 3 out of 5 mammalian genomes (Mouse, Rat, Human, Cow and Dog). \*\* Number in parenthesis indicates how many overlapping CRE elements were identified in the promoter sequence, however, coordinates of only the first of the overlapping elements are given here. \*\*\* Mouse and rat promoters have non-canonical CATAAAA variants instead of canonical TATAA in human, cow and dog. However, this non-canonical variant is known to function in a TATA box-like fashion [56,57]. \*\*\*\* N/A = not applicable, because in Nr4a1 and Per2, no conserved CREs were detected within 2 kb of the promoter sequence. \*\*\*\*\* In human sequence TATA box was identified within 300 bp downstream of conserved CRE; however, it was not present in other genomes. See additional file 1 for additional notes for this table.

Fos protein are important for the communication of information from retinorecipient cells to other parts of the SCN. Another piece of evidence implicating both *Fos* and *Jun* in entrainment is that microinjection of antisense oligonucleotides for these transcripts into the brain inhibits the phase shifting effects of light [37]. In addition, the behavioral phase shifts are highly correlated with changes in *Fos* expression, but not with *Egr1* or *Nr4a1* expression [38]. There have been no functional studies regarding the role of *Egr3* in the circadian clock.

We found that nearly all of the light-induced genes identified in this study had evolutionary conserved CRE elements in their promoter regions, with the strongest light

inducible responses coming from the combination of CRE elements and associated TATA boxes. On the other hand, core clock components that are not induced by light lack this combination of promoter elements, with one exception (casein kinase 1 delta). The slower, longer-lasting increase in *Per2* relative to the other induced genes has been described elsewhere [39]. The loss of a conserved CRE element from the *Per2* promoter may be a component of its functional divergence from *Per1*. It is likely that while the CRE element helps drive transcription of light-induced genes, there are other important regulatory elements responsible for controlling the time course of expression in response to activation by p-CREB. However, these data strongly suggest that the presence of the CRE

**Table 3: CRE and TATA elements in selected clock genes**

Ensembl gene ID	Gene name	Number of conserved CRE elements within the promoter sequence *	Genomic positions of upstream CRE elements in mouse promoter relative to start codon (ATG starts at position +1)	Genomic positions of TATA boxes if any
ENSMUSG00000020038	Cry1	1	-600 to -580	No TATA
ENSMUSG00000068742	Cry2	1	-262 to -242	No TATA
ENSMUSG00000029238	Clock	No CRE ***	N/A	-472 to -469 ****
ENSMUSG00000055116	Bmal1	No CRE	N/A	No TATA
ENSMUSG00000020889	Rev-erb-alpha (Nr1d1)	No CRE	N/A	No TATA
ENSMUSG00000032238	Ror-alpha (Rora)	No CRE	N/A	No TATA
ENSMUSG00000022433	Casein kinase I epsilon (Csnk1e)	No CRE	N/A	No TATA
ENSMUSG00000025162	Casein kinase I delta (Csnk1d)	1	-330 to -310 (4) **	-570 to -567

\* We defined CRE element as being evolutionary conserved if it was present in at least 3 out of 5 mammalian genomes (Mouse, Rat, Human, Cow and Dog). \*\* Number in parenthesis indicates how many overlapping CRE elements were identified in the promoter sequence, however, coordinates of only the first of the overlapping elements are given here. \*\*\* Only mouse and human promoters had predicted CREs; however, these CREs were located in different positions and therefore we did not consider them as being evolutionary conserved. \*\*\*\* Multiple TATA boxes were identified in each species; however, not all of them were evolutionary conserved in all sequences. TATA box shown was shared between four out of five species because dog genomic sequence was not available in that region.

element drives transcription of a coordinated array of immediate-early genes in the SCN in response to photic stimulation. Further evidence of this is the finding that CREB specifically binds to the CRE sequences in the *Dusp1* promoter [28]. Observed evolutionary conservation of CRE elements among mammalian genomes that diverged at least 90 million years ago (MYA), such as primate-cattle divergence at about 90–98 MYA [40], and primate-rodent divergence at about 90 MYA [41], indicates that the functional role of these elements in regulation of clock genes has likely remained the same in different mammals. Although this kind of *in silico* analysis requires further experimental evidence, and while the role of many light-induced genes in the SCN remains unknown, these data support the idea that the response of the SCN to photic input is not limited to pathways involved directly in entrainment.

## Conclusion

The results from this study demonstrate the existence of previously unknown light-induced immediate-early genes in the SCN. Several of these genes have been previously shown to be involved in a reduction of cellular activity and/or the prevention of apoptosis. These data suggest that in addition to the responding to light by shifting the timing of the circadian clock, mechanisms exist to reduce the long-term sensitivity of the SCN to light during nocturnal light exposure. Most light-induced genes have evolutionary conserved CRE elements in their promoter regions, supporting a common mechanism for a coordinated transcriptional response to photic input. Further investigation into the functional role of light-induced

genes may yield new insight into the mechanisms of circadian clock function.

## Methods

### Animals

Adult male C57BL/6 mice were individually housed in a 14:10 light/dark cycle in their experimental room and cage for at least two weeks prior to the experiment. They were then exposed to a 30 minute light pulse (2400lux) or a dark pulse (sham) starting one hour after lights off. Previous research has shown that this duration and intensity of light is well below the threshold for induction of retinal apoptosis [42]. Immediately after treatment, mice were euthanized via cervical dislocation. Dark pulse mice were euthanized under dim red light, and both light and dark pulse animals' eyes were dissected out immediately to avoid excess exposure to light. The mice were then decapitated and the brains were quickly removed and frozen in isopentane cooled in dry ice. Brains were stored at -70°C until ready for further processing. Animals were handled in accordance with the guidelines of the PHS Guide to the Care and Use of Laboratory Animals and all NIH regulations.

### Laser Capture Microscopy/RNA purification

Brains were cut into 12 µm thick sections on a cryostat and directly mounted onto glass slides. Sections were stained using a quick protocol to allow for visual identification of the suprachiasmatic nucleus. First the sections were fixed in a 75% EtOH solution for 30 seconds, rinsed in water to remove excess EtOH from the slide, and then immersed in Hemotoxylin for 90 seconds. Slides were

then washed in molecular biology grade water. The slides then were taken through an alcohol dehydration series of 75%, 95% and 100% EtOH for 30 seconds each, followed by immersion in xylenes for 5 minutes. The slides were removed from the xylene, and once the remaining xylenes had evaporated the slides were placed into a laser capture microscope (Arcturus) and the SCN was identified and captured into CapSure<sup>®</sup> HS LCM Caps (Molecular Devices).

For microarrays and validation of microarrays by quantitative real-time PCR (qPCR), 6 consecutive SCN sections were captured from each mouse. The specificity of each capture for SCN tissue was confirmed after capture by examination of both pre and post-capture images of the tissue. While it is possible that a few cells were technically extra-SCN from any given sample, we estimate that this would comprise less than 1% of the total captured material. Samples from three mice were then pooled together within the same treatment, so in total there were 18 SCN sections pooled together. For the *Per1* and *Per2* time course experiment, 6 consecutive SCN sections from each mouse were used but multiple mice were not pooled for analysis. The pooled samples were purified using an RNA purification kit (Picopure from Molecular Devices) including a DNase treatment. A 1  $\mu$ l aliquot of each sample was removed and processed on an Agilent Bioanalyzer 2100 using the Agilent Lab-on-a-Chip Picochip RNA kit. Only samples with RNA integrity numbers about 6.8 were processed further. RNA was stored at -70°C until ready to proceed.

### Microarrays

For experiments involving microarrays, 1 ng of total RNA was processed through two rounds of linear amplification using RiboAmp HS kits (Molecular Devices). Amplified samples were labeled, fragmented, and hybridized to Affymetrix Mouse 430A 2.0 Genechips using the standard Affymetrix protocols according to the 2004 edition of the Affymetrix technical manual. Signal intensities for each gene were generated using the Microarray Suite 5.0 algorithm by Affymetrix GCOS software. In addition to the signal intensity, each gene was determined to be present, marginal, or absent using default software settings. Signal intensity and present/absent data were imported into Excel and filtered such that only genes that received a present call in at least 3 of the 6 arrays were included in the analysis. This reduced the total number of probe sets to be analyzed from 22,626 to 10,340. Signal intensities for the three light pulse and three dark pulse arrays were analyzed using Cyber-T software [43] using the default settings. This software generates p-values for each gene as a test of differences between light and dark pulse groups using a Bayesian t-test [44]. Two sets of candidate genes were selected based on the results of this test. The first set

was characterized by Bayesian t-test p-values that were below the critical value for a Bonferroni multiple-comparison correction ( $p < 4.8E-06$ ). The second set met a lesser criterion of  $p < 0.001$ . The false discovery rate was examined empirically by qPCR confirmation.

The data discussed in this publication have been deposited in NCBI's Gene Expression Omnibus (GEO, [45]) and are accessible through GEO Series accession number GSE6904.

### Real Time PCR

For experiments involving qPCR validation, Taqman 20 $\times$  gene expression probes were ordered from Applied Biosystems for the genes shown to be significantly upregulated from the microarray results. qPCR was performed on an Applied Biosystems Prism 7000 sequence detection system. Purified Total RNA from above was reversed transcribed using Taqman Reverse Transcription reagents kit, using standard protocol with random hexamers. qPCR was conducted using Taqman Universal Master Mix on the experimental samples, with all samples being assayed in triplicate. Each plate run included a subset of primers for mouse *gapdh* as a control gene. Analysis of relative gene expression in real-time PCR experiments were performed using the  $2^{-\Delta\Delta CT}$  method [46].

### Analysis of evolutionary conservation of CRE and TATA elements in the promoter sequences

Genomic sequences of promoter sequences for five mammalian species were downloaded from Ensembl Genome Browser [47], release 42. The following genomes were used: NCBI build m36 assembly of mouse (*Mus musculus*) genome, NCBI 36 assembly of human (*Homo sapiens*) genome, whole genome shotgun (WGS) assembly CanFam2.0 of dog (*Canis familiaris*) genome, WGS preliminary assembly Btau\_2.0 of cow (*Bos taurus*) genome, and RGSC 3.4 assembly of rat (*Rattus norvegicus*) genome. Promoters were defined as 2 kb upstream of the annotated translational start sites (ATG) of respective immediate-early and clock genes.

Putative CRE (cAMP-response) elements were identified using MatInspector [48,49] using vertebrate matrices of the Matrix Family Library Version 6.2 (October 2006) [50]. If multiple overlapping CREs were detected within 10 bp of each other, they were considered to form a single CRE. After CRE elements were identified in each sequence, a search for downstream TATA boxes was performed.

Multiple sequence alignments of the promoter sequences were constructed with the program BlastZ that is specifically fine-tuned to capture sequence similarity of large non-coding sequences such as those found in the promoter regions [51] using MultiPipMaker [52,53]. Putative



CRE elements and TATA boxes from mouse promoters were mapped onto the alignments. If the same elements were detected in the homologous position in at least two out of four other mammalian genomes, such CREs and TATA boxes were considered evolutionary conserved and likely to be functional [52,54,55].

### Authors' contributions

VMP carried out the laser capture, microarray hybridization, and qPCR analysis and helped to design the study and draft the manuscript. HP carried out the comparative genomic analysis and participated in drafting the manuscript. EMM conceived of the study, performed the statistical analyses, and drafted the manuscript. All authors read and approved the final manuscript.

### Additional material

#### Additional File 1

Supplementary notes for table 2. Additional information on the alignment of promoter sequences across species.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2202-8-98-S1.pdf>]

### Acknowledgements

The authors would like to thank Aaron Jasnow for his assistance with real-time PCR procedures. This research was supported by the National Science Foundation major research instrumentation grant DBI-0320858, an Ohio Board of Regents/Kent State research challenge grant, and the Kent State University Department of Biological Sciences.

### References

- Antle MC, Silver R: **Orchestrating time: arrangements of the brain circadian clock.** *Trends Neurosci* 2005, **28**:145-151.
- Butcher GQ, Lee B, Cheng HY, Obrietan K: **Light stimulates MSK1 activation in the suprachiasmatic nucleus via a PACAP-ERK/MAP kinase-dependent mechanism.** *J Neurosci* 2005, **25**:5305-5313.
- Daan S, Pittendrigh CS: **A Functional analysis of circadian pacemakers in nocturnal rodents II. The variability of phase response curves.** *Journal of Comparative Physiology A* 1976, **106**:253-266.
- Kornhauser JM, Mayo KE, Takahashi JS: **Light, immediate-early genes, and circadian rhythms.** *Behav Genet* 1996, **26**:221-240.
- Honrado GI, Johnson RS, Golombek DA, Spiegelman BM, Papaioannou VE, Ralph MR: **The circadian system of c-fos deficient mice.** *J Comp Physiol [A]* 1996, **178**:563-570.
- Travnickova Z, Sumova A, Peters R, Schwartz WJ, Illnerova H: **Photoperiod-dependent correlation between light-induced SCN c-fos expression and resetting of circadian phase.** *Am J Physiol* 1996, **271**:R825-R831.
- Edelstein K, de la Iglesia HO, Schwartz WJ, Mrosovsky N: **Behavioral arousal blocks light-induced phase advances in locomotor rhythmicity but not light-induced Per1 and Fos expression in the hamster suprachiasmatic nucleus.** *Neuroscience* 2003, **118**:253-261.
- Schwartz WJ, Peters RV, Aronin N, Bennett MR: **Unexpected c-fos gene expression in the suprachiasmatic nucleus of mice entrained to a skeleton photoperiod.** *J Biol Rhythms* 1996, **11**:35-44.
- Colwell CS, Kaufman CM, Menaker M: **Phase-shifting mechanisms in the mammalian circadian system: new light on the carbachol paradox.** *J Neurosci* 1993, **13**:1454-1459.
- Kilduff TS, Vugrinic C, Lee SL, Milbrandt JD, Mikkelsen JD, O'hara BF, Heller HC: **Characterization of the circadian system of NGFI-A and NGFI-A/NGFI-B deficient mice.** *J Biol Rhythms* 1998, **13**:347-357.
- Tanaka M, Amaya F, Tamada Y, Okamura H, Hisa Y, Ibata Y: **Induction of NGFI-A gene expression in the rat suprachiasmatic nucleus by photic stimulation.** *Brain Res* 1997, **756**:305-310.
- Morris ME, Viswanathan N, Kuhlman S, Davis FC, Weitz CJ: **A screen for genes induced in the suprachiasmatic nucleus by light.** *Science* 1998, **279**:1544-1547.
- Shearman LP, Zylka MJ, Weaver DR, Kolakowski LF Jr., Reppert SM: **Two period homologs: circadian expression and photic regulation in the suprachiasmatic nuclei.** *Neuron* 1997, **19**:1261-1269.
- Shigeyoshi Y, Taguchi K, Yamamoto S, Takekida S, Yan L, Tei H, Moriya T, Shibata S, Loros JJ, Dunlap JC, Okamura H: **Light-induced resetting of a mammalian circadian clock is associated with rapid induction of the mPer1 transcript.** *Cell* 1997, **91**:1043-1053.
- Obrietan K, Impey S, Smith D, Athos J, Storm DR: **Circadian regulation of cAMP response element-mediated gene expression in the suprachiasmatic nuclei.** *J Biol Chem* 1999, **274**:17748-17756.
- Tischkau SA, Mitchell JW, Tyan SH, Buchanan GF, Gillette MU: **Ca2+/cAMP response element-binding protein (CREB)-dependent activation of Per1 is required for light-induced signaling in the suprachiasmatic nucleus circadian clock.** *J Biol Chem* 2003, **278**:718-723.
- Conkright MD, Guzman E, Flechner L, Su AI, Hogenesch JB, Montminy M: **Genome-wide analysis of CREB target genes reveals a core promoter requirement for cAMP responsiveness.** *Mol Cell* 2003, **11**:1101-1108.
- Li YY, Popivanova BK, Nagai Y, Ishikura H, Fujii C, Mukaida N: **Pim-3, a proto-oncogene with serine/threonine kinase activity, is aberrantly expressed in human pancreatic cancer and phosphorylates bad to block bad-mediated apoptosis in human pancreatic cancer cell lines.** *Cancer Res* 2006, **66**:6741-6747.
- Popivanova BK, Li YY, Zheng H, Omura K, Fujii C, Tsuneyama K, Mukaida N: **Proto-oncogene, Pim-3 with serine/threonine kinase activity, is aberrantly expressed in human colon cancer cells and can prevent Bad-mediated apoptosis.** *Cancer Sci* 2007, **98**:321-328.
- Larsen CM, Dossing MG, Papa S, Franzoso G, Billestrup N, Mandrup-Poulsen T: **Growth arrest- and DNA-damage-inducible 45beta gene inhibits c-Jun N-terminal kinase and extracellular signal-regulated kinase and decreases IL-1beta-induced apoptosis in insulin-producing INS-1E cells.** *Diabetologia* 2006, **49**:980-989.
- Gupta M, Gupta SK, Hoffman B, Liebermann DA: **Gadd45a and Gadd45b protect hematopoietic cells from UV-induced apoptosis via distinct signaling pathways, including p38 activation and JNK inhibition.** *J Biol Chem* 2006, **281**:17552-17558.
- Ploski JE, Newton SS, Duman RS: **Electroconvulsive seizure-induced gene expression profile of the hippocampus dentate gyrus granule cell layer.** *J Neurochem* 2006, **99**:1122-1132.
- Colwell CS, Levine MS: **Glutamate receptor-induced toxicity in neostriatal cells.** *Brain Res* 1996, **724**:205-212.
- Ma Q: **Induction and superinduction of 2,3,7,8-tetrachlorodibenzo-rho-dioxin-inducible poly(ADP-ribose) polymerase: role of the aryl hydrocarbon receptor/aryl hydrocarbon receptor nuclear translocator transcription activation domains and a labile transcription repressor.** *Arch Biochem Biophys* 2002, **404**:309-316.
- Mukai M, Tischkau SA: **Effects of tryptophan photoproducts in the circadian timing system: searching for a physiological role for aryl hydrocarbon receptor.** *Toxicol Sci* 2007, **95**:172-181.
- Hogenesch JB, Chan WK, Jackiw VH, Brown RC, Gu YZ, Pray-Grant M, Perdev GH, Bradfield CA: **Characterization of a subset of the basic-helix-loop-helix-PAS superfamily that interacts with components of the dioxin signaling pathway.** *J Biol Chem* 1997, **272**:8581-8593.

27. Araki R, Nakahara M, Fukumura R, Takahashi H, Mori K, Umeda N, Sujino M, Inouye SI, Abe M: **Identification of genes that express in response to light exposure and express rhythmically in a circadian manner in the mouse suprachiasmatic nucleus.** *Brain Res* 2006, **1098**:9-18.
28. Doi M, Cho S, Yujnovsky I, Hirayama J, Cermakian N, Cato AC, Sassone-Corsi P: **Light-inducible and clock-controlled expression of MAP kinase phosphatase I in mouse central pacemaker neurons.** *J Biol Rhythms* 2007, **22**:127-139.
29. Butcher GQ, Lee B, Obrietan K: **Temporal regulation of light-induced extracellular signal-regulated kinase activation in the suprachiasmatic nucleus.** *J Neurophysiol* 2003, **90**:3854-3863.
30. Dziema H, Oatis B, Butcher GQ, Yates R, Hoyt KR, Obrietan K: **The ERK/MAP kinase pathway couples light to immediate-early gene expression in the suprachiasmatic nucleus.** *Eur J Neurosci* 2003, **17**:1617-1627.
31. Moyers JS, Bilan PJ, Zhu J, Kahn CR: **Rad and Rad-related GTPases interact with calmodulin and calmodulin-dependent protein kinase II.** *J Biol Chem* 1997, **272**:11832-11839.
32. Beguin P, Mahalakshmi RN, Nagashima K, Cher DH, Ikeda H, Yamada Y, Seino Y, Hunziker W: **Nuclear sequestration of beta-subunits by Rad and Rem is controlled by 14-3-3 and calmodulin and reveals a novel mechanism for Ca<sup>2+</sup> channel regulation.** *J Mol Biol* 2006, **355**:34-46.
33. Nomura K, Takeuchi Y, Fukunaga K: **MAP kinase additively activates the mouse *Per1* gene promoter with CaM kinase II.** *Brain Res* 2006, **1118**:25-33.
34. Nomura K, Takeuchi Y, Yamaguchi S, Okamura H, Fukunaga K: **Involvement of calcium/calmodulin-dependent protein kinase II in the induction of *mPer1*.** *J Neurosci Res* 2003, **72**:384-392.
35. Lim IK: **TIS21 (BTG2/PC3) as a link between ageing and cancer: cell cycle regulator and endogenous cell death molecule.** *J Cancer Res Clin Oncol* 2006, **132**:417-426.
36. Rowland BD, Peeper DS: **KLF4, p21 and context-dependent opposing forces in cancer.** *Nat Rev Cancer* 2006, **6**:11-23.
37. Wollnik F, Brysch W, Uhlmann E, Gillardon F, Bravo R, Zimmermann M, Schlingensiepen KH, Herdegen T: **Block of c-Fos and JunB expression by antisense oligonucleotides inhibits light-induced phase shifts of the mammalian circadian clock.** *Eur J Neurosci* 1995, **7**:388-393.
38. Lin JT, Kornhauser JM, Singh NP, Mayo KE, Takahashi JS: **Visual sensitivities of *nur77* (NGFI-B) and *zif268* (NGFI-A) induction in the suprachiasmatic nucleus are dissociated from c-fos induction and behavioral phase-shifting responses.** *Brain Res Mol Brain Res* 1997, **46**:303-310.
39. Travnickova-Bendova Z, Cermakian N, Reppert SM, Sassone-Corsi P: **Bimodal regulation of *mPeriod* promoters by CREB-dependent signaling and CLOCK/BMAL1 activity.** *Proc Natl Acad Sci U S A* 2002, **99**:7728-7733.
40. Blair HS, Kumar S: **Genomic clocks and evolutionary timescales.** *Trends Genet* 2003, **19**:200-206.
41. Kumar S, Hedges SB: **A molecular timescale for vertebrate evolution.** *Nature* 1998, **392**:917-920.
42. Marti A, Hafezi F, Linsel N, Hegi ME, Wenzel A, Grimm C, Niemeyer G, Reme CE: **Light-induced cell death of retinal photoreceptors in the absence of p53.** *Invest Ophthalmol Vis Sci* 1998, **39**:846-849.
43. **Welcome to Cyber-T** 2007 [<http://cybert.microarray.ics.uci.edu/>].
44. Baldi P, Long AD: **A Bayesian framework for the analysis of microarray expression data: regularized t-test and statistical inferences of gene changes.** *Bioinformatics* 2001, **17**:509-519.
45. **Gene Expression Omnibus** 2007 [<http://www.ncbi.nlm.nih.gov/geo/>].
46. Livak KJ, Schmittgen TD: **Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method.** *Methods* 2001, **25**:402-408.
47. Hubbard TJ, Aken BL, Beal K, Ballester B, Caccamo M, Chen Y, Clarke L, Coates G, Cunningham F, Cutts T, Down T, Dyer SC, Fitzgerald S, Fernandez-Banet J, Graf S, Haider S, Hammond M, Herrero J, Holland R, Howe K, Howe K, Johnson N, Kahari A, Keefe D, Kokocinski F, Kulesha E, Lawson D, Longden I, Melsopp C, Megy K, Meidl P, Ouverdin B, Parker A, Prlic A, Rice S, Rios D, Schuster M, Sealy I, Severin J, Slater G, Smedley D, Spudich G, Trevanion S, Vilella A, Vogel J, White S, Wood M, Cox T, Curwen V, Durbin R, Fernandez-Suarez XM, Flicek P, Kasprzyk A, Proctor G, Searle S, Smith J, Ureta-Vidal A, Birney E: **Ensembl** 2007. *Nucleic Acids Res* 2007, **35**:D610-D617.
48. Quandt K, Frech K, Karas H, Wingender E, Werner T: **MatInd and MatInspector: new fast and versatile tools for detection of consensus matches in nucleotide sequence data.** *Nucleic Acids Res* 1995, **23**:4878-4884.
49. Cartharius K, Frech K, Grote K, Klocke B, Haltmeier M, Klingenhoff A, Frisch M, Bayerlein M, Werner T: **MatInspector and beyond: promoter analysis based on transcription factor binding sites.** *Bioinformatics* 2005, **21**:2933-2942.
50. **Matrix Family Library Version 6.2** 2007 [<http://www.genom.atix.de/>].
51. Schwartz S, Kent WJ, Smit A, Zhang Z, Baertsch R, Hardison RC, Haussler D, Miller W: **Human-mouse alignments with BLASTZ.** *Genome Res* 2003, **13**:103-107.
52. Schwartz S, Elnitski L, Li M, Weirauch M, Riemer C, Smit A, Green ED, Hardison RC, Miller W: **MultiPipMaker and supporting tools: Alignments and analysis of multiple genomic DNA sequences.** *Nucleic Acids Res* 2003, **31**:3518-3524.
53. Schwartz S, Zhang Z, Frazer KA, Smit A, Riemer C, Bouck J, Gibbs R, Hardison R, Miller W: **PipMaker—a web server for aligning two genomic DNA sequences.** *Genome Res* 2000, **10**:577-586.
54. Dubchak I, Brudno M, Loots GG, Pachter L, Mayor C, Rubin EM, Frazer KA: **Active conservation of noncoding sequences revealed by three-way species comparisons.** *Genome Res* 2000, **10**:1304-1306.
55. Thomas JW, Touchman JW: **Vertebrate genome sequencing: building a backbone for comparative genomics.** *Trends Genet* 2002, **18**:104-108.
56. Stewart JJ, Fischbeck JA, Chen X, Stargell LA: **Non-optimal TATA elements exhibit diverse mechanistic consequences.** *J Biol Chem* 2006, **281**:22665-22673.
57. Stewart JJ, Stargell LA: **The stability of the TFIIA-TBP-DNA complex is dependent on the sequence of the TATAAA element.** *J Biol Chem* 2001, **276**:30078-30084.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
[http://www.biomedcentral.com/info/publishing\\_adv.asp](http://www.biomedcentral.com/info/publishing_adv.asp)

