

ORAL PRESENTATION

Open Access

Chikungunya, a new threat propagated by the cosmopolite *Aedes albopictus*

M Vazeille, E Martin, L Mousson, AB Failloux*

From Institut Pasteur International Network Annual Scientific Meeting
Hong Kong. 22-23 November 2010

The recent outbreaks of chikungunya (CHIK) were due to an East-Central-South African genotype harbouring a substitution from an Alanine (E1-226A) to a Valine (E1-226V) at the position 226 in the E1 gene. The new variant E1-226V is very efficiently transmitted by an unusual mosquito vector, *Aedes albopictus*. We found that (i) *Ae. albopictus* females ensure a high replication rate of E1-226V up to 10^9 viral particles per female, (ii) the mid-gut plays a key role in limiting viral dissemination of E1-226A in the mosquito, (iii) the virus is detectable in the saliva as soon as two days after the infectious blood-meal, and (iv) *Ae. albopictus* is able to deliver up to 3000 viral particles with its saliva. All these characteristics led to exacerbate CHIK transmission by *Ae. albopictus*. However, this species is affected by the viral infection. Indeed, CHIK infection reduces sharply the survival of *Ae. albopictus*, females laying their eggs just before dying. Females did not die from an excess of viral replication but more likely in attempting to mount an immune anti-viral response. Moreover, by removing the intracellular bacteria *Wolbachia* from *Ae. albopictus* through successive antibiotic treatments, we aimed to determine if *Wolbachia* interferes with CHIK replication in the mosquito. We found that *Ae. albopictus* cleaned of *Wolbachia* was not affected by CHIK infection. Thus, *Wolbachia* may regulate viral replication in *Ae. albopictus* with consequences on its survival. So, the response of a vector to a particular pathogen is also closely linked to the presence of other microorganisms. Finally, we showed that *Ae. albopictus* was able to be orally co-infected with CHIK and dengue (DEN) viruses and to deliver concomitantly infectious particles of both viruses in saliva. This finding is of particular concern as *Ae. albopictus* is still expanding its geographical range and as both CHIK and DEN

viruses can co-circulate in the same geographical regions. Indeed, reports of co-infections in patients with both viruses are increasing.

Published: 10 January 2011

doi:10.1186/1753-6561-5-S1-O8

Cite this article as: Vazeille et al.: Chikungunya, a new threat propagated by the cosmopolite *Aedes albopictus*. *BMC Proceedings* 2011 5(Suppl 1):O8.

Submit your next manuscript to BioMed Central
and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



Molecular Genetics of Bunyavirus, Institut Pasteur, F-75724 Paris Cedex 15,
France