

Physician's attitudes on pulmonary rehabilitation following COVID-19: a brief perspective from a developing country

Ivan Cherrez-Ojeda,^{1,2} Emanuel Vanegas,^{1,2} Miguel Felix,^{1,2} María José Farfán Bajaña,^{1,2} Azza Sarfraz,³ Zouina Sarfraz,^{2,4} Genesis Camacho,⁵ Alanna Barrios-Ruiz,^{6,7} Jack Michel⁶

¹Universidad Espíritu Santo, Samborondón, Ecuador

²Respiralab Research Group, Guayaquil, Ecuador

³Department of Pediatrics and Child Health, the Aga Khan University, Karachi, Pakistan

⁴Research and Publications, Fatima Jinnah Medical University, Lahore, Pakistan

⁵División de Estudios para Graduados, Facultad de Medicina, Universidad del Zulia, Maracaibo, Venezuela

⁶Division of Clinical and Translational Research, Larkin Community Hospital, South Miami, FL, USA

⁷Instituto Tecnológico y de Estudios Superiores de Monterrey, Monterrey, México

ABSTRACT

Despite the uncertainty about the follow up of COVID-19 survivors, there is a growing body of evidence supporting specific interventions including pulmonary rehabilitation, which may lead to a reduced hospital stay and improved overall respiratory function. The aim of this short report was to assess the attitudes toward pulmonary rehabilitation following COVID-19 among Ecuadorian physicians. A cross-sectional study was conducted, in which a 5-question survey was used to assess the level of agreement to specific statements with a 5-point Likert scale. Out of the 282 participants, 48.2% (n=136) were male, with a mean of 12.6 (SD=11.3) years of experience. More than half of physicians (63.8%, n=180; $\chi^2(2) = 139.224$, p=0.000) considered that diagnosis and treatment of patients with sub-acute and chronic COVID-19 pulmonary sequelae is not clear. Additionally, 94.3% (n=266; $\chi^2(2) = 497.331$, p=0.000) agreed that pulmonary rehabilitation must be considered as a relevant strategy in long-term care following an acute infection, with 92.6% (n=261; $\chi^2(2) = 449.772$, p=0.000) stating it will improve the likelihood of survival and return to baseline health. In conclusion, we found that considerable majority of physicians held positive attitudes to the role of pulmonary rehabilitation and considered it as a relevant strategy in long-term care following COVID-19. However, most of them also conveyed that the diagnosis and treatment of chronic pulmonary sequelae is unclear, and that guidelines for assessing pulmonary function should be established.

Key words: attitudes; developing countries; pulmonary rehabilitation; COVID-19.

Correspondence: Ivan Cherrez-Ojeda, Universidad Espíritu Santo, Km. 2.5 Vía La Puntilla Samborondón 0901-952, Ecuador. Tel. +593.999981769. E-mail: ivancherrez@gmail.com

Contributions: Authors have made substantial contributions to conception and design, acquisition, analysis and interpretation of data, have been involved in drafting the manuscript or revising it critically for important intellectual content, and given final approval of the version to be published. All the authors have read and approved the final version of the manuscript and agreed to be accountable for all aspects of the work.

Ethics approval and consent to participate: This study was conducted in accordance with the Declaration of Helsinki and was approved by the ethics committee: Comité de ética e Investigación en Seres Humanos (CEISH), Guayaquil-Ecuador (#HCK-CEISH-18-0060). Informed consent was obtained from all participants prior to their voluntary participation.

Consent for publication: Not applicable.

Availability of data and material: The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Conflict of interest: The authors declare that they have no competing interests, and all authors confirm accuracy.

Funding: This work was funded and supported by Universidad Espíritu Santo. The sponsor had no role in the study design, data recollection or statistical analyses.

Introduction

To date, millions of patients have recovered from COVID-19; however there are growing concerns related to potential long-term complications and persistent symptoms. A significant number of patients are reporting fatigue, dyspnea, anxiety, and depression even 2-3 months after recovery [1,2]. As documented in a previous report by the European Respiratory Society Task Force, these prolonged symptoms can affect daily life activities, and it appears that age and initial disease severity correlate with long-term sequelae [3]. Based on current evidence, these prolonged symptoms may impair daily life activities, thus requiring comprehensive evaluation and systematic follow up of COVID survivors with unresolved or new progressive symptoms [4].

Studies have also reported impairment in lung function, and reduced exercise capacity, which ultimately may contribute to the observed diminished quality of life [5,6]. In relation to changes in pulmonary function, a recent metanalysis found that roughly a third of patients presented abnormalities following initial infection, mainly a restrictive spirometry pattern, and a reduced diffusing capacity for carbon monoxide (DL_{CO}) [7]. Despite the uncertainty about the follow up of COVID-19 survivors, there is a growing body of evidence supporting specific interventions including pulmonary rehabilitation, which may lead to a reduced hospital stay and improved overall respiratory function [8]. With this in mind, our study aimed to assess the attitudes toward pulmonary rehabilitation following COVID-19 among Ecuadorian physicians.

Methods

Study design

We conducted a cross-sectional study involving 282 Ecuadorian physicians. To be recruited, participants were required to have an active medical practice including COVID-19 patients, regardless of specialization. Physicians filled a non-validated 5-question survey of attitudes and perceptions toward pulmonary rehabilitation in post-COVID-19 patients. Demographic and general characteristics were reported.

Questionnaire

The survey was comprised of two parts: i) questions regarding general characteristics, and ii) questions aimed to assess attitudes and perceptions. With respect to the latter, questions Q_1 , Q_{4-5} measured agreement on certain aspects of pulmonary rehabilitation through a Likert scale whose options were: “strongly disagree”, “disagree”, “neutral”, “agree” and “strongly agree”. Meanwhile, to assess perceived relevance, questions $Q_{2,3}$ employed a Likert scale as follows: “not relevant”, “slightly relevant”, “neutral”, “moderately relevant” and “extremely relevant”. For questions Q_{1-5} participants could only choose one option as an answer.

Ethical statement

This study was approved by the ethics committee Comité de ética e Investigación en Seres Humanos (IRB #HCK-CEISH-18-0060) in accordance with the principles established by the Declaration of Helsinki. All participants were informed of the study aims and gave informed consent prior to filling the survey.

Statistical analyses

Descriptive statistics were applied for general characteristics. Prior to be analyzed, the Likert scale categories were compiled in 3

groups. For questions Q_1 , Q_{4-5} answers were assembled as “agree” (agree and strongly agree), “neutral” or “disagree” (strongly disagree and disagree). Furthermore, for questions $Q_{2,3}$, answers were categorized as: “relevant” (moderately and extremely important), “neutral” or “irrelevant” (slightly important, not important). Chi square goodness of fit was applied to determine if the observed frequencies of the former categories differed with the expected ones.

Results

Out of the 282 participants, 48.2% ($n=136$) were male while 46.5% ($n=131$) were female. The majority of physicians (59.2%, $n=167$) were not specialized; mean years of experience was 12.6 ($SD=11.3$). More details about general characteristics are summarized in Table 1.

Attitudes toward pulmonary sequelae and guidelines

More than half of physicians (63.8%, $n=180$; $\chi^2(2) = 139.224$, $p=0.000$) considered that diagnosis and treatment of patients with sub-acute and chronic COVID-19 pulmonary sequelae is not clear (Figure 1A). As such, 92.2% ($n=260$; $\chi^2(2) = 463.818$, $p=0.000$) appraised it is relevant to establish standardized guidelines on pulmonary testing after COVID-19 pneumonia infection (Figure 1B).

Attitudes toward pulmonary rehabilitation following COVID-19

A large majority (94.3%, $n=266$; $\chi^2(2) = 497.331$, $p=0.000$) of physicians agreed that pulmonary rehabilitation must be considered as a relevant strategy in long-term care following an acute infection. In fact, 9 in 10 participants agreed pulmonary rehabilitation following COVID-19 pneumonia will improve the likelihood of survival and return to baseline health (92.6%, $n=261$; $\chi^2(2) = 449.772$, $p=0.000$) and that standardized pulmonary testing guidelines are required to prescribe pulmonary rehabilitation for these patients (90.8%, $n=256$; $\chi^2(2) = 420.511$, $p=0.000$).

Discussion

The COVID-19 pandemic has brought an unprecedented rate of scientific publication that has overwhelmed healthcare providers and the public health community [9]. Understandably, most participants in our study agreed that the diagnosis and treatment of chronic pulmonary sequelae is not clear and that guide-

Table 1. Demographic and general characteristics of surveyed population ($n=282$).

Characteristics	% (n)
Gender	
Male	48.2 (136)
Female	51.8 (146)
Years of practice (mean, SD)	12.6 (11.3)
Medical specialty	40.8 (115)
Pulmonary medicine and critical care	9.9 (28)
Internal medicine	8.5 (24)
Pediatrics	8.2 (23)
Other	73.4 (207)

lines for assessing pulmonary function following COVID-19 should be established. Regarding the latter, several expert organizations have already proposed recommendations detailing specific indications and procedures to be followed [8]. For example, in the U.K an expert panel recommends that respiratory complications should be considered in post COVID-19 patients, and that low-intensity exercise (≤ 3 METs) with increased intensity according to symptoms is indicated in patients on oxygen therapy [10]. Guidelines in Turkey stress the need to individualize the approach in patients with mild disease and recommend pulmonary rehabilitation either at a specialized center or at home for patients who experienced moderate disease [11].

Pulmonary rehabilitation includes thorax mobilizing exercise, expectoration therapy, and respiratory training to improve symptoms [8]. A previous study among patients in the post-acute phase of mild to severe COVID-19 found an improvement in the 6 minute walk test (6MWT), functional vital capacity (FVC), and the mental component of the SF-36 health survey among patients who completed a 3-week pulmonary rehabilitation program [12]. No adverse event was observed in the aforementioned study, indicating that it is a feasible, safe, and effective options in COVID-19 patients independent of disease severity [12]. In our study, a large majority of respondents agreed that pulmonary rehabilitation must be considered in post COVID-19 patients and that it will improve the likelihood of survival and return to baseline health. Thus, their attitude towards pulmonary rehabilitation reflects what it is known to date on the subject, which may prove useful when designing and implementing a local protocol for the follow up of COVID-19 patients. On a final note, despite the growing number of recommendations from expert panels we believe that guidelines should be adapted according to the limitations and resources of each region for them to be successful, and this is of utmost importance in developing countries.

Limitations

There are some limitations to our study. For instance, we used a non-validated survey to assess a limited set of circumstances regarding attitudes and perceptions towards pulmonary rehabilitation. In addition, about 60% of the participants had no specialization in any medical field, while 9.9% were specialized in pulmonary medicine or critical care. Accordingly, familiarity with pulmonary rehabilitation may have been limited in the sample. Consequently, the reported results may have undervalued the perceptions of pulmonary rehabilitation. However, to the best of our knowledge, our study is among the first to assess the attitudes to pulmonary rehabilitation among Ecuadorian physicians, providing valuable insights that might be useful to design future interventions.

Conclusions

We found that considerable majority of physicians held positive attitudes to the role of pulmonary rehabilitation and considered it as a relevant strategy in long-term care following COVID-19. However, most of them also conveyed that the diagnosis and treatment of chronic pulmonary sequelae is unclear, and that guidelines for assessing pulmonary function should be established.

Acknowledgement

Special thanks to all members of Respiralab Research Group for their initial input regarding this project, in particular to Dr. Karla Robles-Velasco and Matías Panchana. We also want to extend our gratitude to Dr. Arjola Musta Agolli, and Dr. Zeynep Yukselen for their collaboration. Finally, we want to express our gratitude to Universidad Espiritu Santo, Ecuador and Larkin Community Hospital, USA for their continuous support in our research endeavors.




Figure 1. Proportions of physicians expressing their attitudes and perceptions towards pulmonary rehabilitation in post-COVID-19 acute infection. A) Proportion of physicians according to their level of agreement regarding their attitudes and perceptions towards pulmonary rehabilitation in post-COVID-19 acute infection. B) Proportion of physicians according to their level of perception of relevance regarding pulmonary rehabilitation in post-COVID-19 acute infection. Q₁, diagnosis and treatment of patients with sub-acute and chronic COVID-19 pulmonary sequelae is not clear; Q₂, How relevant is it to establish standardized guidelines on pulmonary testing after COVID-19 pneumonia infection?; Q₃, How relevant is pulmonary rehabilitation after COVID-19 pneumonia infection?; Q₄, Pulmonary rehabilitation following COVID-19 pneumonia will improve likelihood of survival and return to baseline health; Q₅, Standardized pulmonary testing guidelines are required to prescribe pulmonary rehabilitation in COVID-19 pneumonia patients.

Abbreviations

DL_{CO}: diffusing capacity for carbon monoxide;
 6MWT: 6-minute walk test;
 FVC: functional vital capacity;
 SF-36: 36-item short form survey.

References

- Carfi A, Bernabei R, Landi F. Persistent symptoms in patients after acute COVID-19. *JAMA* 2020;324:603-5.
- Goërtz YM, Van Herck M, Delbressine JM, Vaes AW, Meys R, Machado FV, et al. Persistent symptoms 3 months after a SARS-CoV-2 infection: the post-COVID-19 syndrome? *ERJ Open Res* 2020;6:00542-2020.
- Antoniou KM, Vasarmidi E, Russell AM, Andrejak C, Crestani B, Delcroix M, et al. European Respiratory Society Statement on long COVID-19 follow-up. *Eur Respir J* 2022; 2102174. Online ahead of print.
- Spruit MA, Holland AE, Singh SJ, Tonia T, Wilson KC, Troosters T. COVID-19: interim guidance on rehabilitation in the hospital and post-hospital phase from a European Respiratory Society-and American Thoracic Society-coordinated international task force. *Eur Respir J* 2020;56:2002197.
- Sanchez-Ramirez DC, Normand K, Zhaoyun Y, Torres-Castro R. Long-term impact of COVID-19: A systematic review of the literature and meta-analysis. *Biomedicines* 2021;9:900.
- Huang C, Huang L, Wang Y, Li X, Ren L, Gu X, et al. 6-month consequences of COVID-19 in patients discharged from hospital: a cohort study. *Lancet* 2021;397:220-32.
- Torres-Castro R, Vasconcello-Castillo L, Alsina-Restoy X, Solis-Navarro L, Burgos F, Puppo H, et al. Respiratory function in patients post-infection by COVID-19: a systematic review and meta-analysis. *Pulmonology* 2021;27:328-37.
- Siddiq MAB, Rathore FA, Clegg D, Rasker JJ. Pulmonary rehabilitation in COVID-19 patients: A scoping review of current practice and its application during the pandemic. *Turk J Phys Med Rehabil* 2020;66:480-94.
- Kang M, Gurbani SS, Kempker JA. The published scientific literature on covid-19: An analysis of PubMed abstracts. *J Med Syst* 2021;45:3.
- Barker-Davies RM, O'Sullivan O, Senaratne KPP, Baker P, Cranley M, Dharm-Datta S, et al. The Stanford Hall consensus statement for post-COVID-19 rehabilitation. *Br J Sports Med* 2020;54:949-59.
- Aytür YK, Köseoğlu BF, Taşkıran ÖÖ, Ordu-Gökkaya NK, Delialioğlu SÜ, Tur BS, et al. Pulmonary rehabilitation principles in SARS-COV-2 infection (COVID-19): A guideline for the acute and subacute rehabilitation. *Turk J Phys Med Rehabil* 2020;66:104-20.
- Gloeckl R, Leitl D, Jarosch I, Schneeberger T, Nell C, Stenzel N, et al. Benefits of pulmonary rehabilitation in COVID-19: a prospective observational cohort study. *ERJ Open Res* 2021;7:00108-2021.

Received for publication: 9 February 2022. Accepted for publication: 22 March 2022.

This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0).

©Copyright: the Author(s), 2022

Licensee PAGEPress, Italy

Multidisciplinary Respiratory Medicine 2022; 17:837

doi:10.4081/mrm.2022.837