

SCIENTIFIC REPORTS



OPEN

Inhibition of dipeptidyl peptidase 8/9 impairs preadipocyte differentiation

Received: 10 February 2015

Accepted: 26 June 2015

Published: 05 August 2015

Ruijun Han^{1,*}, Xinying Wang^{1,*}, William Bachovchin², Zofia Zukowska^{3,†} & John W. Osborn¹

Adipocytes are the primary cells in adipose tissue, and adipocyte dysfunction causes lipodystrophy, obesity and diabetes. The dipeptidyl peptidase (DPP) 4 family includes four enzymes, DPP4, DPP8, DPP9 and fibroblast activation protein (FAP). DPP4 family inhibitors have been used for the treatment of type 2 diabetes patients, but their role in adipocyte formation are poorly understood. Here we demonstrate that the DPP8/9 selective inhibitor 1G244 blocks adipogenesis in preadipocyte 3T3-L1 and 3T3-F422A, while DPP4 and FAP inhibitors have no effect. In addition, knockdown of DPP8 or DPP9 significantly impairs adipocyte differentiation in preadipocytes. We further uncovered that blocking the expression or activities of DPP8 and DPP9 attenuates PPAR γ 2 induction during preadipocyte differentiation. Addition of PPAR γ agonist thiazolidinediones (TZDs), or ectopic expression of PPAR γ 2, is able to rescue the adipogenic defect caused by DPP8/9 inhibition in preadipocytes. These results indicate the importance of DPP8 and DPP9 on adipogenesis.

Over the last several decades, increase in the prevalence of maladies such as metabolic disease, cardiovascular diseases, and even some cancers is often attributed to the paralleled rise in obesity rate worldwide^{1–3}. The obesity phenotype is often characterized by the increase in fat mass, or adipose tissue accumulation. This adipose tissue is primarily comprised of adipocytes, which store excess energy as triglycerides and secrete various endocrine signals, which in turn and contributes to the regulation of total body energy homeostasis^{4–7}. While the role of adipose tissue in obesity has been thoroughly examined, the development and proliferation of adipocytes remain an important question in obesity research. To this end, further examination of adipocyte development is important for the treatment and prevention of obesity and associated diseases.

Much of our knowledge on adipogenesis comes from *in vitro* studies of fibroblasts or pre-adipocytes, such as the mouse cell lines 3T3-L1, 3T3-F442A⁴. Adipocyte differentiation is coordinated by the nuclear receptor peroxisome proliferator-activated receptor γ (PPAR γ) and members of the CCAAT enhancer binding proteins (CEBPs) family⁸. Many of the genes involved in adipogenesis are targets of PPAR γ and/or CEBP α ^{9–11}. Thiazolidinediones (TZDs), which are known to have potent adipogenic and antidiabetic effects, are agonists for PPAR γ . TZDs promote adipogenesis and adipokine production in adipose tissue^{8,12}.

Proteases of the S9B/dipeptidyl peptidase (DPP) 4 family are serine amino peptidases. They comprise of four enzymes: DPP4, fibroblast activation protein (FAP), DPP8 and DPP9. Active members of the DPP4 family preferentially cleave Xaa-Pro- and Xaa-Ala- dipeptides (where Xaa is any amino acid except proline) from the N-terminus of proteins¹³. Further, DPP4 and FAP are cell surface peptidases, and substrates of DPP4 include numerous neuropeptides, hormones and chemokines, such as: fibronectin, substance P, neuropeptide Y, peptide YY, glucagon-like-peptides 1 and 2 (GLP-1, GLP-2) and glucose-dependent insulinotropic peptide (GIP)¹³. Particularly, two substrates of DPP4, GLP-1 and GIP, are released from the intestinal mucosa and account for \sim 60% of postprandial insulin secretion^{14,15}.

¹Department of Integrative Biology and Physiology, University of Minnesota, Minneapolis, MN, USA. ²Sackler School of Biomedical Sciences, Tufts University School of Medicine, Boston, MA, USA. ³These authors contributed equally to this work. [†]Deceased. Correspondence and requests for materials should be addressed to R.H. (email: rhan@umn.edu)

Inhibition of DPP4 prolongs the activity of GLP and GIP, playing an important role in insulin secretion and blood glucose regulation^{16,17}. Therefore, DPP4 inhibitors have drawn increased attention, and are in clinical use as antidiabetic drugs^{13,17}.

Mice lacking DPP4 or mice treated with DPP4 inhibitors are resistant to high fat diet (HFD) induced obesity and insulin resistance^{18,19}. These results were attributed to the reduced food intake and increased energy expenditure in the HFD-treated mice^{18,19}. Human diabetic subjects treated with DPP4 inhibitors show a significant decrease in HbA1c, postprandial glucose, and circulating triglyceride levels compared to placebo treatment group²⁰. Interestingly, despite their wide use in treating type 2 diabetes, the importance of DPP4 family inhibitors on adipogenesis is yet to be identified.

The two cytosolic members of the DPP4 family are DPP8 and DPP9 that share approximately 60% homology^{21,22}. *In vitro*, DPP8 and DPP9 are similar in their biochemical properties, including enzyme kinetics and substrate specificity^{23,24}. However, the importance of DPP8 and DPP9 in obesity and diabetes remain undetermined.

In the present study, we examined the effect of three DPP specific inhibitors on adipocyte differentiation in preadipocytes: DPP4-selective inhibitor MK-0431, DPP8/9-selective inhibitor 1G244, and the FAP-selective inhibitor 3099. We found that only the DPP8/9 inhibitor 1G244 blocked adipogenesis in preadipocytes. We further discovered that inhibition of DPP8 and DPP9 attenuated the expression of PPAR γ 2 in basal level and during preadipocyte differentiation.

Results

DPP4 family inhibitor P32/98 impairs adipocyte differentiation. To address whether the DPP4 family (DPP4, DPP8, DPP9 and FAP) play a role in preadipocyte differentiation, we added the non-selective competitive DPP4 family inhibitor isoleucine-thiazolidide (P32/98) to the classic 3T3-L1 differentiation cocktail: dexamethasone, methylxanthine and insulin (DMI). We found that P32/98 blocked adipogenesis in a dose dependent manner, starting at the concentration of 100 μ M, as assessed by oil red O staining (Fig. 1A). At 500 μ M concentration, P32/98 completely blocked adipogenesis in 3T3-L1 cell line. However, 500 μ M concentration of a chemically similar but non-inhibitory compound P34/98 had no effect (Fig. 1A). We further confirmed the inhibitory effects of P32/98 by detecting the expression of adipocyte markers at the end of differentiation. The expression of FABP4, adiponectin and leptin was markedly decreased in 3T3-L1 adipocytes when treated with P32/98 during adipogenesis (Fig. 1B). A similar inhibitory effect of P32/98 on adipocyte differentiation was observed in 3T3-F442A preadipocytes (supplementary Fig. 1). Notably, only the high dose (over 100 μ M) of P32/98 blocked adipogenesis in preadipocytes. Since P32/98 has IC₅₀ values of 460 nM for DPP4, 2180 nM for DPP8 and 1600 nM for DPP9²⁵, we suggest that the activities of DPP4, DPP8 and DPP9 were all inhibited in preadipocytes.

DPP8/9 inhibition blocks adipocyte differentiation. To further investigate which DPP contributes to the adipogenic defect as described above, specific DPP inhibitors such as a DPP4-selective inhibitor MK-0431/Sitagliptin, DPP8/9-selective inhibitor 1G244, and FAP-selective inhibitor 3099, were added during 3T3-L1 differentiation. We found that the DPP4 inhibitor MK-0431 and the FAP inhibitor 3099 had no effect on adipocyte differentiation, while the DPP8/9 inhibitor 1G244 blocked adipogenesis in 3T3-L1 cells as measured by Oil red O staining (Fig. 2A). At the end of differentiation, the expression of the adipocyte markers, FABP4, adiponectin and leptin, were markedly decreased (Fig. 2B) after treatment with the DPP8/9 inhibitor 1G244. No difference in the expression of these markers was found with either DPP4 inhibitor MK-0431 or FAP inhibitor 3099 (Fig. 2B). A similar inhibitory effect of the DPP8/9 inhibitor 1G244 on adipogenesis was observed in 3T3-F442A preadipocytes (supplementary Fig. 1). Taken together, these data indicate that DPP8/9, but not DPP4 or FAP, is essential for adipogenesis.

Time dependent expression of DPP4 family during 3T3-L1 differentiation. We next assessed the gene expression and enzyme activity of DPPs during adipocyte differentiation. The mRNA level of DPP4 and FAP were low after 48 hours of DMI treatment, whereas the DPP8 and DPP9 mRNA levels were much higher (supplementary Fig. 2A). Evidence suggest that majority of the DPPs activity during differentiation is through DPP8/9 as the DPP8/9 inhibitor 1G244 blocked 82% of total DPPs activity in 3T3-L1 cells; while MK-0431 blocked only 15% and 3099 blocked 20%, respectively (supplementary Fig. 2B). We tested whether DPP8/9 was a regulated component of the differentiation program and if its expression changed over the course of differentiation. Indeed, analysis of a time course of 3T3-L1 cell differentiation revealed that DPP8/9 mRNA expression was significantly increased during DMI treatment (Fig. 3A). Conversely, DPP4 and FAP mRNA expression were decreased (supplementary Fig. 3).

DPP8 and DPP9 are required for 3T3-L1 differentiation. To further validate the adipogenic action of DPP8 and DPP9, we generated stable cell lines with knockdown of DPP8, DPP9 or both using shRNAs. The efficiency and specificity of DPP8 or DPP9 knockdown was confirmed by real time PCR and western blot (Supplementary Fig. 4). Stable 3T3-L1 cell lines expressing shRNA sequences targeting DPP8 or DPP9 exhibited reduced differentiation capacity, compared to those expressing a control shRNA, as assessed by oil red O staining (Fig. 3B). Knockdown of both DPP8 and DPP9 completely blocked adipocyte differentiation (Fig. 3B). In agreement with the morphological differentiation, the




Figure 1. The non-specific DPP4 family inhibitor P32/98 impairs adipocyte differentiation. (A) Oil Red O staining of control 3T3-L1 cells (DMI) and 3T3-L1 cells treated with DPP4 family inhibitor P32/98 (P32/98) or inactive inhibitor (P34/98) at day 8 of differentiation. (B) Adipocyte markers, FABP4, adiponectin (AdipoQ) and leptin, were measured by real time PCR at day 8 of differentiation. β -actin expression was used as an internal control.

expression of the adipocyte markers FABP4, adiponectin and leptin were also impaired in DPP8, DPP9 and DPP8/9 knockdown cells (Fig. 3C).

Next we sought to elucidate the mechanism of action of DPP8/9 in preadipocyte differentiation. PPAR γ is the master regulator of adipocyte differentiation. Activation of PPAR γ at an early stage is necessary and sufficient for adipogenesis⁴. We hypothesized that the effect of DPP8/9 inhibitor on adipogenesis was mediated by PPAR γ . The expression PPAR γ 2 was significantly decreased in stable 3T3-L1 cell expressing shRNA sequences targeting both DPP8 and DPP9, compared to those expressing a control shRNA at basal level with no differentiation cocktail (DMI) (Fig. 4A). When treated with a differentiation cocktail DMI, PPAR γ 2 was markedly increased after 48 hours in 3T3-L1 cells (Fig. 4B). DMI induced expression of PPAR γ 2 was partly blocked with depletion of either DPP8 or DPP9. Since PPAR γ gene is regulated by CEBPs⁴, we determined the effect of DPP8 or DPP9 depletion on the expression of CEBP α , CEBP β and CEBP δ . We found that depletion of DPP8 or DPP9 also attenuated basal and DMI-induced CEBP α mRNA levels (Fig. 4A,B) but had no effect on expression of CEBP β and CEBP δ (Fig. 4A,B). Knockdown of both DPP8 and DPP9 completely blocked the DMI induced expression of PPAR γ 2 and CEBP α (Fig. 4B) in 3T3-L1. In addition, the DPP8/9 inhibitor 1G244 attenuated DMI-induced PPAR γ 2 and CEBP α mRNA (Fig. 4C), but had no effect on the expression of CEBP β and CEBP δ (Fig. 4C). On




Figure 2. DPP8/9 inhibitor impairs adipocyte differentiation. (A) Oil Red O staining of 3T3-L1 control cells (DMI) and cells treated with specific DPP inhibitors like DPP4 inhibitor MK-0431(DMI+ MK-0431), the DPP8/9 inhibitor 1G244 (DMI+1G244) and FAP inhibitor 3099 (DMI+3099) at day 8 of differentiation. (B) Adipocyte markers, FABP4, adiponectin (AdipoQ) and leptin were measured by real time PCR at day 8 of differentiation. β -actin expression was used as an internal control.

the other hand, DPP4 inhibitor MK-0431 and FAP inhibitor 3099 had no effect on DMI induced expression of CEBP α , CEBP β , CEBP δ and PPAR γ 2 during 3T3-L1 differentiation (Fig. 4C).

TZDs or ectopic PPAR γ 2 rescues inhibition of DPP8/9 induced adipogenic defects. To determine whether PPAR γ 2 can rescue the adipogenic defect caused by DPP8/9 inhibition, we generated stable PPAR γ 2 transduced 3T3-L1 cells (Fig. 5A). Ectopic expression of PPAR γ 2 was able to rescue P32/98 or 1G244 caused adipogenic defect in preadipocytes (Fig. 5B). The TZDs, rosiglitazone or troglitazone, are known ligands of PPAR γ . To investigate whether rosiglitazone or troglitazone were able to rescue the adipogenic defect in DPP8/9 inactivated preadipocytes, we added the DPP8/9 inhibitor 1G244 with either rosiglitazone or troglitazone during 3T3-L1 differentiation. In the absence of rosiglitazone or troglitazone, 1G244 markedly inhibited the adipogenesis in 3T3-L1 cells. However, administration of the rosiglitazone or troglitazone completely rescued the adipogenesis defect caused by 1G244, as assessed by oil red O staining (Fig. 5C) and confirmed by the expression of adipocyte markers FABP4, adiponectin and leptin (Fig. 5D).

Discussion

Compared to the well-studied DPP4, the functions of DPP8 and DPP9 are mostly unclear. To date, there are several studies indicating that DPP8 and DPP9 are involved in immune responses and cancer biology^{26–28}. Here, for the first time, we report that DPP8 and DPP9 are indispensable for adipocyte differentiation. We show that DPP8 and DPP9 mRNA are significantly increased during adipocyte differentiation and DPP8/DPP9 knockdown impairs preadipocyte differentiation. We have evaluated the expression of both DPP8 and DPP9 by immunoblotting using commercially available antibodies. We succeeded to




Figure 3. Knockdown of DPP8 and DPP9 blocked adipocyte differentiation. (A) Gene expression analysis of DPP8 and DPP9 during 3T3L1 differentiation. 3T3L1 cells were cultured in 10% FBS (NC) or treated with dexamethasone, isobutylmethylxanthine and insulin (DMI) from day 0 to day 2 of differentiation, mRNA levels of DPP8 and DPP9 were measured by real time PCR. β -actin expression was used as an internal control. (B) Oil Red O staining of control-shRNA-transduced cells (shCTL) and cells with DPP8/9 shRNAs (shDPP8, shDPP9) at day 8 of differentiation. (C) Adipocyte markers, FABP4, adiponectin (AdipoQ) and leptin, were measured by real time PCR at day 8 of differentiation. β -actin expression was used as an internal control.

detect only the endogenous DPP8 protein in preadipocytes (Supplementary Fig. 4). However, after trying three DPP9 antibodies from different vendors, we have not found a suitable antibody to detect the endogenous DPP9 protein in 3T3-L1 cells. Although it is difficult to determine the natural endogenous substrate of DPP8 and DPP9 that mediate adipogenesis, there is evidence in the literature to support a potential biological role of DPP8/DPP9 in metabolism²⁹. DPP8 and DPP9 are homologous, cytoplasmic N-terminal post-proline-cleaving enzymes. To this day, more than 29 *in vivo* DPP8 and DPP9 substrate




Figure 4. Inhibition of DPP8 and DPP9 prevents PPAR γ 2 induction during 3T3-L1 differentiation. (A) PPAR γ 2, CEBP α , CEBP β and CEBP δ mRNA levels were measured in control-shRNA-transduced 3T3-L1 cells (shCTL) and cells with DPP8/9 shRNAs (shDPP8, shDPP9) at basal level. (B) PPAR γ 2 and CEBP α mRNA levels were measured in control-shRNA-transduced cells (shCTL) and cells with DPP8/9 shRNAs (shDPP8, shDPP9) after 48 hour treatment of DMI. CEBP β and CEBP δ mRNA levels were measured after 6 hour treatment. (C) PPAR γ 2 and CEBP α mRNA levels were measured after 48 hour treatment of dexamethasone, isobutylmethylxanthine and insulin (DMI) with or without the DPP4 inhibitor MK-0431 (DMI+MK-0431), the DPP8/9 inhibitor 1G244 (DMI+1G244), the FAP inhibitor 3099 (DMI+3099). CEBP β and CEBP δ mRNA levels were measured in these cells after 6 hour treatment.




Figure 5. TZDs or ectopic PPAR γ 2 rescues inhibition of DPP8/9 induced adipogenic defects in 3T3-L1 cells. (A) Representative western blot for the expression of PPAR γ in stable 3T3-L1 cells transfected with control plasmid (vehicle) or PPAR γ 2 plasmid (PPAR γ 2). The blots were cropped, and the full-length blots are presented in the supplementary information. (B) Oil Red O staining of control cells or PPAR γ 2 overexpressed 3T3-L1 cells, treated vehicle (NC), DMI (DMI), 500 μ M non-selective DPP4 family inhibitor P32/98 (DMI+P32/98) or 20 μ M DPP8/9 inhibitor 1G244 (DMI+1G244) at day 8 of differentiation. (C) Oil Red O staining of 3T3-L1 cells treated with 20 μ M DPP8/9 inhibitor 1G244 (DMI+1G244) or 1G244 plus 1 μ M rosiglitazone (DMI+1G244+Rosi) or 5 μ M troglitazone (DMI+1G244+Tro). (D) Adipocyte markers, FABP4, adiponectin (AdipoQ) and leptin, were measured in these cells by real time PCR at day 8 of differentiation. β -actin expression was used as an internal control.

candidates have been discovered using terminal amine isotopic labeling of substrates technique²⁹. Most of these substrates are involved in regulating cellular metabolism and energy homeostasis, suggesting potential roles for DPP8 and DPP9 in cellular metabolic pathways, including glycolysis, gluconeogenesis, fatty acid metabolism, and nucleotide metabolism/biosynthesis²⁹. The future goal will be to identify the substrates of DPP8/9 mediating preadipocyte differentiation.

PPAR γ is a member of the nuclear-receptor superfamily and is both necessary as well as sufficient for adipogenesis⁴. The relative roles of PPAR γ 1 and PPAR γ 2 in adipogenesis remain an open question. Both PPAR γ 1 and PPAR γ 2 are induced during adipogenesis, but PPAR γ 1 is also expressed in macrophages, liver and skeletal muscle and colonic epithelium⁴. In addition, PPAR γ 2 is highly induced during differentiation and is critical in promoting adipogenesis in preadipocytes^{30,31}. In our study, DPP8/9 inhibitor reduced both PPAR γ 1 and PPAR γ 2 induction during adipocyte differentiation (Supplementary

Figure 5). Consistent with previous reports, we have found that PPAR γ 2 is highly upregulated during 3T3-L1 differentiation. Therefore, we and others³² have selected PPAR γ 2 as a marker for preadipocyte differentiation.

PPAR γ regulates adipogenesis together with members of the CEBP family. CEBP β and CEBP δ are expressed early during preadipocyte differentiation^{33,34}, and are involved in the induction of PPAR γ and CEBP α . CEBP α and PPAR γ are induced at later stages and are also active in mature adipocytes⁴. Despite the importance of CEBPs, CEBPs are not sufficient to induce adipogenesis in the absence of PPAR γ . Some transcription factors, like KLF15, KLF5, GATA2 and GATA 3 directly target the expression of PPAR γ and CEBP α , with no effects on the expression of CEBP β and CEBP δ during the adipogenesis⁴. Similarly, here we also find that inhibition of DPP8 and DPP9 successfully attenuated the induction of PPAR γ and CEBP α , with no effects on CEBP β and CEBP δ . The activation of CEBP α and PPAR γ are partly dependent on the expression of CEBP β and CEBP δ . For example, transcription factor KLF15 induces the expression PPAR γ during adipocyte differentiation independent of the activity of CEBP β and CEBP δ ³⁵. In addition, GATA2/3 also inhibits PPAR γ expression independent of the activity of CEBP β and CEBP δ during adipogenesis^{36,37}. In contrast, KLF5 activates PPAR γ expression, induced early during adipocyte differentiation by CEBP β and CEBP δ ³⁸. Here, we find that DPP8 and DPP9 are required for 3T3-L1 differentiation by targeting the expression of CEBP α and PPAR γ . In our future directions, we plan to define whether the role of DPP8/9 on adipogenesis is dependent on CEBP β and CEBP δ .

We noticed that a relatively high dose of P32/98 inhibitor (100 μ M, 10 fold higher than IC₅₀ value) is needed in the media to block differentiation in preadipocytes. Considering that DPP8 and DPP9 are cytosolic enzymes, the inhibitors need to pass the cell membrane to block the cytosolic activity of DPP8/9. We suggest it is, at least in part, the reason we need a high concentration of P32/98 to block the preadipocyte differentiation. High dose of DPP8/9 specific inhibitor 1G244 is used to block preadipocyte differentiation in our study as well. This is consistent with previous report, in which, 1G244 ameliorated DPP8/9 activity in HEK293 cells at high extracellular concentration of 8 μ M (over 100 fold higher than IC₅₀ value)³⁹. In addition, non-inhibitory P34/98 is used as the control of P32/98. Moreover, we find that P34/98 had no effects on adipogenesis, which suggests minimal off-target effects.

A preclinical study²⁵ has shown that the DPP8/9 inhibitor, allo-isoletucyl isoindoline derivative 4, resulted in severe toxicity, including death, alopecia and thrombocytopenia. In contrast to DPP4, which has an extracellular catalytic domain, DPP8 and DPP9 are cytosolic enzymes. A recent study found that allo-isoletucyl isoindoline derivative 4 is not able to penetrate the plasma membrane of mammalian cells, even at very high concentrations³⁹. Therefore, the toxic effects observed with this inhibitor are not the result of DPP8/9 inhibition. Instead, it is most likely an “off-target” effect through an unknown mechanism outside the membrane. Moreover, a recent study found that high doses of vildagliptin, produced nearly complete inhibition of DPP8 and DPP9 enzyme activity *in vivo*, with no toxic side effects⁴⁰. These findings indicate that the report of toxicity associated with certain DPP8/9 inhibitor compounds are not related to inhibition of DPP8 and DPP9. The DPP8/9 inhibitor 1G244 has an excellent membrane penetration capacity and rats treated with high dose of 1G244 do not exhibit pathological symptoms³⁹. We observed that there was no difference dead cell counts with trypan blue staining compared to control, when 3T3-L1 cells were treated with 20 μ M 1G244.

Here we have found that DPP8 and DPP9 are required for preadipocyte differentiation. However, there is little information on the importance of DPP8 and DPP9 on *in vivo* adipogenesis. Since a knock-out mouse model either for DPP8 or DPP9 is not available, the studies on role of DPPs *in vivo* mainly depend on the DPP inhibitors. There are five DPP4 inhibitors are currently in clinical use: saxagliptin⁴¹, sitagliptin⁴² and vildagliptin⁴³, linagliptin⁴⁴ and alogliptin⁴⁵. From the literature, sitagliptin, saxagliptin, linagliptin and alogliptin have very less DPP8/9 inhibitory activity, whereas vildagliptin has quite potent activities against both DPP8 and DPP9^{13,46}. However vildagliptin is the only DPP-4 inhibitor to show reduced fasting lipolysis that is associated with a reduction in fatter liver in patient⁴⁷. Considering that DPP8 and DPP9 are cytosolic enzymes, the inhibitors need to pass the cell membrane to block the cytosolic activity of DPP8/9³⁹. In our experiment, the concentration of inhibitors is needed at least 10 fold higher than IC₅₀ value to block the adipogenesis in preadipocytes. So, the broad DPP inhibitors, like P32/98 and vildagliptin, are not perfect to study the role of DPP8/9 *in vivo*. Here we found that the DPP8/9 specific inhibitor 1G244 blocked the adipogenesis by inhibiting the expression of PPAR γ during preadipocyte differentiation. TZDs (Rosiglitazone or troglitazone) were able to rescue the adipogenic defect in DPP8/9 inactivated preadipocytes. These data suggest that DPP8/9 inhibitor 1G244 has effects opposite to those of TZDs in preadipocytes. Considering the effects of TZDs on adipogenic and antidiabetic effects⁸, we hypothesize that the DPP8/9 specific inhibitor will have important role on adipogenesis and glucose homeostasis *in vivo*.

Methods

Materials. DPP8/9 inhibitor 1G244, DPP4 inhibitor MK-0431 and FAP inhibitor 3099 were a gift from Dr. Bachovchin (Tufts University, Boston, MA). P32/98 and P34/98 were received from Probiodrug⁴⁸ (Halle, Germany). The PPAR γ antibody was purchased from Cell signaling.

Adipocyte differentiation. 3T3-L1 cells were cultured to confluence in DMEM (Dulbecco's modified Eagle's medium) and supplemented with 10% calf serum. Two days after post-confluence (designated day 0), cells were induced to differentiate with a differentiation cocktail: DMEM supplemented with 10% fetal bovine serum (FBS), 1 μ M dexamethasone (Sigma), 0.5 mM isobutylmethylxanthine (Sigma) and, 1 μ g/ml insulin (Sigma) for 48 hours. The DPP inhibitors or PPAR γ agonists were added to the differentiation cocktail as indicated. Then the media were replaced with DMEM supplemented with 10% FBS and 1 μ g/ml insulin for 48 hours. The cells were subsequently re-fed every 48 h with DMEM supplemented with 10% FBS. For 3T3-F442A differentiation: cells were induced to differentiate with 10 μ g/ml insulin (Sigma) for 4 days. The DPP inhibitors were added together with insulin as indicated.

Generation of stably transduced 3T3-L1 cells. One day prior to transfection, Phoenix Eco cells (NGVB, Indianapolis) were seeded at a density of 3×10^6 /10 cm dish. Retroviral vector was transfected into cells using TurboFectin 8.0 (Origene) according to the manufacturer's instructions. Two days later, the medium containing viruses was filtered through 0.45 μ m syringe filter and was used to infect 3T3-L1 cells. Two days after infection, cells were selected with 2.5 μ g/ml puromycin or 6 μ g/ml Blasticidine. For PPAR γ 2 overexpression experiment: pMSCVpuro-PPAR γ 2 was obtained from Addgene and control vector pMSCVpuro from Clontech. For DPP8 or DPP9 knockdown: DPP8 shRNA plasmid (Origene, TF504979) was puromycin resistance, with sequence 5' TTCCTGAGTCTGGAGAACAACACTATGAACTG-3'; DPP9 shRNA plasmid (Origene, TG515456) was puromycin resistance, with sequence 5' TGTC AAGC TGCGAGAAGGAAGTGGTACAG-3'. For DPP8 and DPP9 double-knockdown: DPP9 shRNA sequence was constructed into blasticidine resistance vector pGFP-B-RS and co-transfected with puromycin resistance DPP9 shRNA plasmid.

Reverse transcription and Real-time PCR. RNA was isolated using Roche RNA isolation kit and cDNA synthesized by iScriptcDNA synthesis kit (BioRad). Real-time PCR was performed using the CFX96 Touch™ real-time PCR detection system (BioRad) and the TaqMan PCR Reagent Kit with pre-designed primers and fluorescently labeled probes with the duplicates. The primers were from the Applied Biosystems: β actin, Mm00607939_s1; FABP4, Mm00445880_m1; Adiponectin, Mm00456425_m1; Leptin Mm00434759_m1; PPAR γ 2 Mm00440940_m1; CEBP α , Mm01265914_s1; CEBP β , Mm00843434_s1; CEBP δ Mm00786711_s1; DPP4, Mm00494538_m1; DPP8, Mm00547049_m1; DPP9, Mm00841122_m1; FAP, Mm00484254_m1. The mRNA expression levels were calculated using the comparative CT method with b-actin as the endogenous reference gene, in accordance with the Applied Biosystems' ABI PRISM, as described previously⁴⁹.

Oil red O staining. At the end of adipogenic differentiation, adipocytes were fixed in 4% paraformaldehyde (USB Corporation) for 8 min and were stained with Oil Red O for 30 min at room temperature. Subsequently, the cells were washed three times with PBS and imaged (Nikon, Melville, NY).

Western Blot. For western blotting analysis, the cells were lysed in RIPA buffer (150 mmol/l NaCl, 1% Triton X-100, 0.5% sodium deoxycholate, 0.1% SDS, 50 mmol/l Tris-HCl, pH 7.4) containing phosphatase inhibitors and a protease inhibitor cocktail (Sigma). The lysate was then subjected to SDS-PAGE, transferred to PVDF membranes and incubated with the primary antibodies, followed by IRDye secondary antibody (Li-Cor). The bound antibody was visualized by Li-Cor Odyssey CLX system.

DPP Activity. 3T3-L1 cells were lysed in 0.1% Triton X-100. DPP activity was measured by luminescence, using DPPIV-Glo™ Protease Assay (Promega) according to the manufacturer's instructions. The activity of individual DPPs was determined based on the differences between DPP activities with or without selective DPP inhibitors (10 μ M).

Statistical analysis. Every experiment has been repeated at least three times. Statistical analysis was performed using GraphPad software. Multiple comparisons were analyzed by One-way repeated measure ANOVA with post-hoc t-test using Dunnett's method, and pair-wise comparison using Student's t test. Results are presented as means \pm SD, and a p value of <0.05 was considered statistically significant.

References

1. Olshansky, S. J. *et al.* A potential decline in life expectancy in the United States in the 21st century. *N Engl J Med* **352**, 1138–1145, doi: 10.1056/NEJMSr043743 (2005).
2. Van Gaal, L. F., Mertens, I. L. & De Block, C. E. Mechanisms linking obesity with cardiovascular disease. *Nature* **444**, 875–880, doi: 10.1038/nature05487 (2006).
3. Kahn, S. E., Hull, R. L. & Utzschneider, K. M. Mechanisms linking obesity to insulin resistance and type 2 diabetes. *Nature* **444**, 840–846, doi: 10.1038/nature05482 (2006).
4. Rosen, E. D. & MacDougald, O. A. Adipocyte differentiation from the inside out. *Nature reviews. Molecular cell biology* **7**, 885–896, doi: 10.1038/nrm2066 (2006).
5. Halaas, J. L. *et al.* Weight-reducing effects of the plasma protein encoded by the obese gene. *Science* **269**, 543–546 (1995).
6. Yamauchi, T. *et al.* The fat-derived hormone adiponectin reverses insulin resistance associated with both lipoatrophy and obesity. *Nature medicine* **7**, 941–946, doi: 10.1038/90984 (2001).
7. Stepan, C. M. *et al.* The hormone resistin links obesity to diabetes. *Nature* **409**, 307–312, doi: 10.1038/35053000 (2001).

8. Ahmadian, M. *et al.* PPARgamma signaling and metabolism: the good, the bad and the future. *Nature medicine* **19**, 557–566, doi: 10.1038/nm.3159 (2013).
9. Christy, R. J. *et al.* Differentiation-induced gene expression in 3T3-L1 preadipocytes: CCAAT/enhancer binding protein interacts with and activates the promoters of two adipocyte-specific genes. *Genes & development* **3**, 1323–1335 (1989).
10. Dalen, K. T. *et al.* Adipose tissue expression of the lipid droplet-associating proteins S3-12 and perilipin is controlled by peroxisome proliferator-activated receptor-gamma. *Diabetes* **53**, 1243–1252 (2004).
11. Villanueva, C. J. *et al.* TLE3 is a dual-function transcriptional coregulator of adipogenesis. *Cell metabolism* **13**, 413–427, doi: 10.1016/j.cmet.2011.02.014 (2011).
12. Lehmann, J. M. *et al.* An antidiabetic thiazolidinedione is a high affinity ligand for peroxisome proliferator-activated receptor gamma (PPAR gamma). *The Journal of biological chemistry* **270**, 12953–12956 (1995).
13. Kirby, M., Yu, D. M., O'Connor, S. & Gorrell, M. D. Inhibitor selectivity in the clinical application of dipeptidyl peptidase-4 inhibition. *Clin Sci (Lond)* **118**, 31–41, doi: 10.1042/CS20090047 (2010).
14. Baggio, L. L. & Drucker, D. J. Harnessing the therapeutic potential of glucagon-like peptide-1: a critical review. *Treatments in endocrinology* **1**, 117–125 (2002).
15. Holst, J. J. Glucagon-like peptide-1, a gastrointestinal hormone with a pharmaceutical potential. *Current medicinal chemistry* **6**, 1005–1017 (1999).
16. Mu, J. *et al.* Chronic inhibition of dipeptidyl peptidase-4 with a sitagliptin analog preserves pancreatic beta-cell mass and function in a rodent model of type 2 diabetes. *Diabetes* **55**, 1695–1704, doi: 10.2337/db05-1602 (2006).
17. Drucker, D. J. Dipeptidyl peptidase-4 inhibition and the treatment of type 2 diabetes: preclinical biology and mechanisms of action. *Diabetes care* **30**, 1335–1343, doi: 10.2337/dc07-0228 (2007).
18. Conarello, S. L. *et al.* Mice lacking dipeptidyl peptidase IV are protected against obesity and insulin resistance. *Proceedings of the National Academy of Sciences of the United States of America* **100**, 6825–6830, doi: 10.1073/pnas.0631828100 (2003).
19. Lamont, B. J. & Drucker, D. J. Differential antidiabetic efficacy of incretin agonists versus DPP-4 inhibition in high fat fed mice. *Diabetes* **57**, 190–198, doi: 10.2337/db07-1202 (2008).
20. Ahren, B. Dipeptidyl peptidase-4 inhibitors: clinical data and clinical implications. *Diabetes care* **30**, 1344–1350, doi: 10.2337/dc07-0233 (2007).
21. Abbott, C. A., Yu, D., McCaughan, G. W. & Gorrell, M. D. Post-proline-cleaving peptidases having DP IV like enzyme activity. Post-proline peptidases. *Advances in experimental medicine and biology* **477**, 103–109, doi: 10.1007/0-306-46826-3_10 (2000).
22. Ajami, K., Abbott, C. A., McCaughan, G. W. & Gorrell, M. D. Dipeptidyl peptidase 9 has two forms, a broad tissue distribution, cytoplasmic localization and DPIV-like peptidase activity. *Biochimica et biophysica acta* **1679**, 18–28, doi: 10.1016/j.bbexp.2004.03.010 (2004).
23. Geiss-Friedlander, R. *et al.* The cytoplasmic peptidase DPP9 is rate-limiting for degradation of proline-containing peptides. *The Journal of biological chemistry* **284**, 27211–27219, doi: 10.1074/jbc.M109.041871 (2009).
24. Connolly, B. A. *et al.* Dipeptide boronic acid inhibitors of dipeptidyl peptidase IV: determinants of potency and *in vivo* efficacy and safety. *Journal of medicinal chemistry* **51**, 6005–6013, doi: 10.1021/jm800390n (2008).
25. Lankas, G. R. *et al.* Dipeptidyl peptidase IV inhibition for the treatment of type 2 diabetes: potential importance of selectivity over dipeptidyl peptidases 8 and 9. *Diabetes* **54**, 2988–2994 (2005).
26. Spagnuolo, P. A. *et al.* Inhibition of intracellular dipeptidyl peptidases 8 and 9 enhances parthenolide's anti-leukemic activity. *Leukemia* **27**, 1236–1244, doi: 10.1038/leu.2013.9 (2013).
27. Lu, C. *et al.* Dipeptidyl peptidases as survival factors in Ewing sarcoma family of tumors: implications for tumor biology and therapy. *The Journal of biological chemistry* **286**, 27494–27505, doi: 10.1074/jbc.M111.224089 (2011).
28. Schade, J. *et al.* Regulation of expression and function of dipeptidyl peptidase 4 (DP4), DP8/9, and DP10 in allergic responses of the lung in rats. *The journal of histochemistry and cytochemistry: official journal of the Histochemistry Society* **56**, 147–155, doi: 10.1369/jhc.7A7319.2007 (2008).
29. Wilson, C. H. *et al.* Identifying natural substrates for dipeptidyl peptidases 8 and 9 using terminal amine isotopic labeling of substrates (TAILS) reveals *in vivo* roles in cellular homeostasis and energy metabolism. *The Journal of biological chemistry* **288**, 13936–13949, doi: 10.1074/jbc.M112.445841 (2013).
30. Ren, D., Collingwood, T. N., Rebar, E. J., Wolffe, A. P. & Camp, H. S. PPARgamma knockdown by engineered transcription factors: exogenous PPARgamma2 but not PPARgamma1 reactivates adipogenesis. *Genes & development* **16**, 27–32, doi: 10.1101/gad.953802 (2002).
31. Mueller, E. *et al.* Genetic analysis of adipogenesis through peroxisome proliferator-activated receptor gamma isoforms. *The Journal of biological chemistry* **277**, 41925–41930, doi: 10.1074/jbc.M206950200 (2002).
32. Cristancho, A. G. *et al.* Repressor transcription factor 7-like 1 promotes adipogenic competency in precursor cells. *Proceedings of the National Academy of Sciences of the United States of America* **108**, 16271–16276, doi: 10.1073/pnas.1109409108 (2011).
33. Cao, Z., Umek, R. M. & McKnight, S. L. Regulated expression of three C/EBP isoforms during adipose conversion of 3T3-L1 cells. *Genes & development* **5**, 1538–1552 (1991).
34. Yeh, W. C., Cao, Z., Classon, M. & McKnight, S. L. Cascade regulation of terminal adipocyte differentiation by three members of the C/EBP family of leucine zipper proteins. *Genes & development* **9**, 168–181 (1995).
35. Mori, T. *et al.* Role of Kruppel-like factor 15 (KLF15) in transcriptional regulation of adipogenesis. *The Journal of biological chemistry* **280**, 12867–12875, doi: 10.1074/jbc.M410515200 (2005).
36. Tong, Q. *et al.* Function of GATA transcription factors in preadipocyte-adipocyte transition. *Science* **290**, 134–138 (2000).
37. Tong, Q., Tsai, J., Tan, G., Dalgin, G. & Hotamisligil, G. S. Interaction between GATA and the C/EBP family of transcription factors is critical in GATA-mediated suppression of adipocyte differentiation. *Molecular and cellular biology* **25**, 706–715, doi: 10.1128/MCB.25.2.706-715.2005 (2005).
38. Oishi, Y., Manabe, I. & Nagai, R. [Kruppel-like family of transcription factor 5 (KLF5). KLF5 is a key regulator of adipocyte differentiation]. *Nihon rinsho. Japanese journal of clinical medicine* **69 Suppl 1**, 264–268 (2011).
39. Wu, J. J. *et al.* Biochemistry, pharmacokinetics, and toxicology of a potent and selective DPP8/9 inhibitor. *Biochemical pharmacology* **78**, 203–210, doi: 10.1016/j.bcp.2009.03.032 (2009).
40. Burkey, B. F. *et al.* Adverse effects of dipeptidyl peptidases 8 and 9 inhibition in rodents revisited. *Diabetes, obesity & metabolism* **10**, 1057–1061, doi: 10.1111/j.1463-1326.2008.00860.x (2008).
41. Herman, G. A. *et al.* Pharmacokinetics and pharmacodynamics of sitagliptin, an inhibitor of dipeptidyl peptidase IV, in healthy subjects: results from two randomized, double-blind, placebo-controlled studies with single oral doses. *Clinical pharmacology and therapeutics* **78**, 675–688, doi: 10.1016/j.clpt.2005.09.002 (2005).
42. Augeri, D. J. *et al.* Discovery and preclinical profile of Saxagliptin (BMS-477118): a highly potent, long-acting, orally active dipeptidyl peptidase IV inhibitor for the treatment of type 2 diabetes. *Journal of medicinal chemistry* **48**, 5025–5037, doi: 10.1021/jm050261p (2005).
43. Ahren, B. *et al.* Inhibition of dipeptidyl peptidase-4 reduces glycemia, sustains insulin levels, and reduces glucagon levels in type 2 diabetes. *The Journal of clinical endocrinology and metabolism* **89**, 2078–2084, doi: 10.1210/jc.2003-031907 (2004).

44. Deacon, C. F. & Holst, J. J. Linagliptin, a xanthine-based dipeptidyl peptidase-4 inhibitor with an unusual profile for the treatment of type 2 diabetes. *Expert opinion on investigational drugs* **19**, 133–140, doi: 10.1517/13543780903463862 (2010).
45. Feng, J. *et al.* Discovery of alogliptin: a potent, selective, bioavailable, and efficacious inhibitor of dipeptidyl peptidase IV. *Journal of medicinal chemistry* **50**, 2297–2300, doi: 10.1021/jm070104l (2007).
46. Brandt, I. *et al.* Inhibition of dipeptidyl-peptidase IV catalyzed peptide truncation by Vildagliptin ((2S)-[[3-(3-hydroxyadamantan-1-yl)amino]acetyl]-pyrrolidine-2-carbonitrile). *Biochemical pharmacology* **70**, 134–143, doi: 10.1016/j.bcp.2005.04.009 (2005).
47. Macauley, M. *et al.* Effect of vildagliptin on hepatic steatosis. *The Journal of clinical endocrinology and metabolism* **100**, 1578–1585, doi: 10.1210/jc.2014-3794 (2015).
48. Frerker, N. *et al.* Neuropeptide Y (NPY) cleaving enzymes: structural and functional homologues of dipeptidyl peptidase 4. *Peptides* **28**, 257–268, doi: 10.1016/j.peptides.2006.09.027 (2007).
49. Han, R., Kitlinska, J. B., Munday, W. R., Gallicano, G. I. & Zukowska, Z. Stress hormone epinephrine enhances adipogenesis in murine embryonic stem cells by up-regulating the neuropeptide Y system. *PLoS one* **7**, e36609, doi: 10.1371/journal.pone.0036609 (2012).

Acknowledgments

The authors thank Dr. Yan Chen, Dr. Alessandro Bartolomucci, Dr. Jana Strakova, Dr. Gufa Lin, Dr. Joanna Kitlinska, Dr. Christopher T Banek and Dr. Ninitha Asirvatham Jeyaraj for their insightful comments and editing. This work was supported by U.S. National Institutes of Health grants R01HL067357 to Z.Z. and J.W.O.

Author Contributions

R.H. designed the experiment, wrote the manuscript and researched data. X.W. designed the experiment, wrote the manuscript and researched data. W.B. reviewed/edited the manuscript. Z.Z. researched data. J.W.O. reviewed/edited manuscript.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Han, R. *et al.* Inhibition of dipeptidyl peptidase 8/9 impairs preadipocyte differentiation. *Sci. Rep.* **5**, 12348; doi: 10.1038/srep12348 (2015).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>