

CASE REPORT

Migratory fishbones in the pharynx: A report of two cases

Shuyi Xu¹ | Xiaomin Li² | Yue Peng² | Jing Yang² | Qianxu Liu² | Jiefeng Guo² | Zhijian Yu² 

¹Department of Ophthalmology, Zhuhai People's Hospital (Zhuhai Hospital Affiliated with Jinan University), Zhuhai, China

²Department of Otolaryngology, Zhuhai People's Hospital (Zhuhai Hospital Affiliated with Jinan University), Zhuhai, China

Correspondence

Zhijian Yu, Department of Otolaryngology, Zhuhai People's Hospital (Zhuhai Hospital Affiliated with Jinan University), No.79 Kangning Road, Xiangzhou District, 519000 Zhuhai, Guangdong, China.
Email: yzjent@163.com

Funding information

This study was supported by grants from the Cultivation Project of Zhuhai People's Hospital (No. 2019PY-22) and Zhuhai Medical Research Fund Project (No. ZH3310200025PJL)

Abstract

The possibility of fishbone migration into the surrounding tissues, especially in cases where it cannot be identified on routine inspection. Early diagnosis of migratory fishbone and therapeutic management are essential for optimal patient survival.

KEYWORDS

fishbone, laryngoscopy, migratory, pharynx

1 | INTRODUCTION

Fishbones migrated into surrounding tissues are relatively unusual. We present two cases of patients' migratory fishbones into the retropharyngeal space and hypopharynx. All fishbones were removed without complications. Early diagnosis of migratory fishbone and therapeutic management are essential for optimal patient survival.

Foreign bodies (eg, fishbones) stuck in the pharynx are the most common emergency cases in otorhinolaryngology practice. Sharp and pointed fishbones may damage the mucosa. In a majority of cases, however, these can be easily removed. In rare cases, fishbones can migrate to the different parts of the body causing deep neck space infection,¹ large blood vessel rupture,² and gastrointestinal perforation³ if not promptly managed.

Here, we describe two cases of patients' migratory fishbones into the retropharyngeal space and hypopharynx. We

discuss our experience in treating these cases using a specific approach for each patient.

2 | CASE PRESENTATION

2.1 | Case 1: migratory fishbone in the retropharyngeal space

A 62-year-old man presented with sudden onset pharyngeal pain after eating fish at a local hospital. Fiber-optic laryngoscopy showed no abnormalities. Computed tomography (CT) revealed a foreign body lying transversely at the right piriform fossa. Painless gastroscopy was performed to remove the foreign body; however, no foreign bodies were detected. During gastroscopy, he presented with laryngeal edema, which required tracheal intubation.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2021 The Authors. *Clinical Case Reports* published by John Wiley & Sons Ltd.

Four days after admission, he was referred to our hospital for management. Cervical CT showed a foreign body in the right retropharyngeal space between the cervical vertebrae 1 and 2 (Figure 1A,B). Subsequently, he underwent surgical exploration of the retropharyngeal space under general anesthesia. During surgery, the foreign body could not be detected in the plane of the second cervical vertebra. An extended longitudinal incision was performed to find the fishbone in the surrounding tissues. Fortunately, a fishbone was found in the plane of the third cervical vertebra. A 1-week follow-up revealed the absence of pain and foreign body sensation.

2.2 | Case 2: migratory fishbone in the hypopharynx

A 52-year-old man was admitted to our department for throat pain, foreign body sensation, dysphagia, and mild sialorrhea for 3 days after eating fish. Cervical CT confirmed the presence of a linear radiopaque structure in the left laryngopharynx (Figure 2A,B). Flexible fiber-optic laryngoscopy did not identify a foreign body in the area but this revealed the absence of mucosal lesions (Figure 2C). To further ascertain the presence of a foreign body in the hypopharynx, a gastroscopy was performed showing a fishbone in the left piriform fossa (Figure 2D). The fishbone pierced from the left piriform fossa into the posterior pharyngeal wall. The object was extracted using forceps. The patient immediately felt comfortable.

3 | DISCUSSION

Here, we describe two cases of migratory fishbones in the pharynx that were not observable via flexible fiber-optic

laryngoscopy. We used a specific treatment for each patient. Fortunately, all patients had favorable outcomes.

Patients suspected of having fishbone impaction usually undergo physical and laryngoscopic examinations. A fishbone at the base of the tongue or hypopharynx usually requires flexible fiber-optic laryngoscopy for observation. However, it is difficult to determine the precise localization of fishbones in cases with an endoscopically undetectable fishbone. Sharp fishbones are more prone to cause mucosal damage. Moreover, muscle contractions after entering the pharynx may contribute to migration.⁴ Migratory fishbones are associated with an increased incidence of complications. Therefore, it is important to establish this diagnosis early and accurately. CT is considered the most sensitive (sensitivity =100%) modality for detecting ingested fishbones in soft tissues.⁵ Therefore, CT is recommended when suspecting the migration of ingested foreign bodies, as demonstrated in the present cases.

The retropharyngeal space consists of loose soft tissue between the buccopharyngeal fascia and the prevertebral fascia. Owing to this structure, fishbones can migrate freely into the retropharyngeal space. This was consistent with our observations on the first case. During the operation, it is difficult to identify the location of the fishbone in the retropharyngeal space. We suggest the use of CT to approximate the position of the fishbone, followed by an extended longitudinal incision to identify its exact location in the surrounding tissues.

The incidence of foreign bodies in the hypopharynx is common. However, complete visualization of the foreign body is usually limited by the anatomically closed hypopharynx in its resting state. Modified Killian's method can be applied to remove an endoscopically invisible fishbone.⁵ Unfortunately, the second patient had severe sore throat symptoms upon head flexion and rotation. In this case, we removed the fishbone through gastroscopy, which fully exposed the hypopharynx.





FIGURE 1 The fishbone in the retropharyngeal space. Sagittal (A) and axial (B) computed tomography (CT) images showing linear calcifications in the retropharyngeal space (red arrow)

FIGURE 2 The fishbone in the hypopharynx. Sagittal (A) and axial (B) CT images showing linear calcifications in the hypopharynx (red arrow). (C) Flexible fiberoptic laryngoscopy showing the absence of any foreign body. (D) Gastroscopic view of a fishbone in the hypopharynx (yellow arrow)



This suggests that gastroscopy may be a good alternative for the visualization and removal of fishbones in such cases. Similar to gastroscopy, laparoscopy⁶ is a good choice for visualization and removal of a fishbone that has perforated the viscus and entered the abdominal cavity.

4 | CONCLUSION

This paper highlights the possibility of fishbone migration into the surrounding tissues, especially in cases where it cannot be identified on routine inspection. Early diagnosis of migratory fishbone and therapeutic management are essential for optimal patient survival.

ACKNOWLEDGEMENTS

The authors thank the patients in this case report.

CONFLICTS OF INTERESTS

None provided.

AUTHOR CONTRIBUTION

Shuyi Xu contributed to the discussion and edited the manuscript. Xiaomin Li, Yue Peng, Jing Yang, Qianxu Liu, and

Jiefeng Guo participated in the data collection. Zhijian Yu wrote the manuscript. All authors read and approved the final manuscript.

ETHICAL APPROVAL

This study was approved with our previous project by the Institutional Review Board of Zhuhai People's Hospital, Zhuhai hospital affiliated with Jinan University (reference number KYZY-2020-11). All participants approved and signed written informed consents upon enrollment.

DATA AVAILABILITY STATEMENT

All data included in this study are available upon request by contact with the corresponding author.

ORCID

Zhijian Yu  <https://orcid.org/0000-0002-3735-6209>

REFERENCES

1. Kataria G, Saxena A, Bhagat S, Singh B, Kaur M, Kaur G. Deep neck space infections: a study of 76 cases. *Iran J Otorhinolaryngol.* 2015;27(81):293-299.
2. Jiang D, Lu Y, Zhang Y, Hu Z, Cheng H. Aortic penetration due to a fish bone: a case report. *J Cardiothorac Surg.* 2020;15:292.

3. Wu CW, Chiu YW. Unintentional fish bone ingestion causing perforation of small intestine. *Intern Emerg Med*. 2020.
4. Ho NH, Chang FC, Wang YF. Clinical approaches to migrating ingested foreign bodies in the neck. *Ear Nose Throat J*. 2020;145561320948787.
5. Kikuchi D, Ikeda M, Muroso S. Endoscopic removal of a fish bone foreign body in the hypopharynx with the modified Killian's method. *Auris Nasus Larynx*. 2020;48:1035-1037.
6. Mulita F, Papadopoulos G, Tsochatzis S, Kehagias I. Laparoscopic removal of an ingested fish bone from the head of

the pancreas: case report and review of literature. *Pan Afr Med J*. 2020;25(36):123.

How to cite this article: Xu S, Li X, Peng Y, et al. Migratory fishbones in the pharynx: A report of two cases. *Clin Case Rep*. 2021;9:e04548. <https://doi.org/10.1002/ccr3.4548>