

a national population-based serosurvey and a standardized serological test, represent a considerable burden of infections in adolescents and young adults that sharply contrast with the official surveillance reports. Also, the Bp-IgG PT ELISA commercial kit lower sensitivity than the CDC/FDA PT ELISA may lead underestimation of recent infections. In conclusion, assessing pertussis seroprevalence requires careful consideration of the right tests and epidemiological model for interpretation.

Disclosures. All authors: No reported disclosures.

671. Increased Disease Severity and Larger Household Size Among Hispanic Infants With Pertussis: Chicago, IL, 2010–2016

Enrique Ramirez, BA¹; Allison Arwady, MD, MPH²; Peter Ruestow, PhD³ and Marielle Fricchione, MD¹; Immunization Program, Chicago Department of Public Health, Chicago, Illinois, ²Chicago Department of Public Health, Chicago, Illinois, ³Communicable Diseases, Chicago Department of Public Health, Chicago, Illinois

Session: 66. Public Health: Epidemiology and Outbreaks

Thursday, October 4, 2018: 12:30 PM

Background. Infants experience higher pertussis incidence and mortality compared with other age groups and 85% of infant cases are acquired from a household member. The Chicago Department of Public Health (CDPH) noted increased pertussis incidence among Hispanics from 2014 (5.3 cases per 100,000 population) to 2016 (12.3 cases per 100,000 population) while other ethnicities remained stable.

Methods. All pertussis cases reported to CDPH through the Illinois National Electronic Disease Surveillance System with onsets from January 1, 2010 to December 31, 2016 were included in the analyses. Variables of interest were age, race/ethnicity, household size, disease severity, healthcare location of diagnosis, and hospitalization. To assess differences in demographic and care-related indicators between Hispanic and non-Hispanic White (NHW) infants, chi-square tests were performed; risk ratios and 95% confidence intervals were estimated.

Results. From 2010–2016, 1,036 pertussis cases were reported to CDPH including 276 infants aged <1 year. Among infant cases, 187(68%) were <4 months of age, 143(52%) male, 151(55%) were Hispanic and 54(20%) were NHW. Median household size for infants of all ethnicities was four members (range: 0–15). Of all infants with household member size of ≥4 persons (157), 64% were Hispanic. 185 (67%) of infants had ≥1 severe symptom (apnea, cyanosis, pneumonia, seizure, encephalopathy) and 133 (48%) were hospitalized. Diagnosis occurred in the emergency room (ER) for 199 (72%) infants. Hispanics were more likely than NHW to have household member size ≥4 (RR 1.9, $P < 0.05$), severe symptoms (RR 1.6, $P < 0.05$), hospitalization (RR 2.3, $P < 0.05$), and diagnosis in the ER (RR 2.8, $P < 0.05$) (Table 1).

Conclusion. Among Chicago infant pertussis cases, Hispanic infants have larger household sizes and more severe disease at time of diagnosis compared with NHW.

Table 1: Chicago Hispanic Infant Pertussis Case Characteristics Compared with Non-Hispanic Whites (NHW), 2010–2016

Infant Characteristics	Race/Ethnicity	N (%)	RR (95% CI)
Household size ≥4	NHW	19 (35.2)	Ref.
	Hispanic	100 (66.2)	1.9 (1.3–2.8)*
Severe symptoms	NHW	25 (46.3)	Ref.
	Hispanic†	113 (75.3)	1.6 (1.2–2.2)*
Hospitalization	NHW	14 (25.9)	Ref.
	Hispanic	90 (59.6)	2.3 (1.4–3.7)*
Diagnosis in ER	NHW	16 (29.6)	Ref.
	Hispanic	124 (82.1)	2.8 (1.8–4.2)*

* Significant at $P < 0.05$.

† population = 150, one patient missing data.

Disclosures. All authors: No reported disclosures.

672. The Impact on Humans of Treating Dogs with Amoxicillin/Clavulanate

Kirithana R. Beaulac, PharmD¹; Shira Doron, MD, MS, FIDSA²; Claire Fellman, DVM PhD³; Annie Wayne, DVM MPH¹; Laura A McDermott, MT(ASCP)² and David Snyderman, MD, FIDSA²; ¹Department of Pharmacy, Tufts Medical Center, Boston, Massachusetts, ²Division of Geographic Medicine and Infectious Diseases, Tufts Medical Center, Boston, Massachusetts, ³Cummings School of Veterinary Medicine at Tufts University, North Grafton, Massachusetts, ⁴Emergency Medicine and Critical Care, Cummings School of Veterinary Medicine at Tufts University, North Grafton, Massachusetts

Session: 66. Public Health: Epidemiology and Outbreaks

Thursday, October 4, 2018: 12:30 PM

Background. It has been shown that the use of antibiotics promotes resistance to that antibiotic, and that colonization with resistant pathogens can be transmitted from companion animals to their human owners. There is an evolving concern that use of medically important antibiotics in pets can confer colonization by resistant organisms in their human owners.

Methods. In dogs receiving amoxicillin/clavulanate for 14 days for a diagnosed infection, we screened stool both at the time of antibiotic initiation and at the end of the course for the dog taking antibiotics and their human owner. Owners had to meet a closeness score of 3 or 4, meaning living in same household with frequent contact,

with or without oral contact, and have no recent antibiotic exposure. Stool samples were quantitatively cultured for enteric flora and incubated on selective media for ampicillin/sulbactam resistance (A/S-r), ciprofloxacin resistance (CIP-r), ESBL, MRSA, and VRE.

Results. We enrolled eight dog-human pairs. All had enteric flora present at the time of antibiotic initiation (Day 1), whereas two of the seven dog samples that could be analyzed had no detectable enteric flora by the end of antibiotics (Day 14). No humans or dogs acquired MRSA or VRE. One human lost colonization with CIP-r flora, whereas two dogs acquired CIP-r during antibiotic treatment that did not transmit to their owners. One dog and one unrelated human acquired ESBL colonization by day 14 that was not present at Day 1. There were three humans colonized with A/S-r on Day 1 which persisted through the treatment period. Of their three dogs, one had no A/S-r at either time point, one newly acquired high counts of A/S-r flora, and one did not provide enough stool at Day 14 for A/S-r testing. There was one other dog that acquired A/S-r flora and one dog that had increasing counts of A/S-r, both of which had human owners with no A/S-r on Day 1 or 14.

Conclusion. In this pilot study, we identified that use of amoxicillin-clavulanate, a common antibiotic prescribed to both humans and dogs, can lead to resistant colonization in the dog taking antibiotics, and may have some impact on their close human owners who may share or transmit colonization. Further study is under way to better understand this relationship.

Disclosures. L. A. McDermott, CytisPharma Inc.: Research Contractor, Contractual agreement with Tufts Medical Center.

673. Syphilis Outbreak in Women Who Have Sex with Men in Japan: A Case-Control Study in Tokyo, 2017–2018

Shingo Nishiki, MD¹; Yuzo Arima, MPH, PhD²; Takuya Yamagishi, MD, PhD²; Takashi Hamada, MD²; Takuri Takahashi, PhD²; Tomimasa Sunagawa, MD, PhD²; Tamano Matsui, MD, PhD²; Kazunori Oishi, MD, PhD² and Makoto Ohnishi, MD, PhD⁴; ¹Field Epidemiology Training Program, National Institute of Infectious Diseases, Tokyo, Japan, ²Infectious Disease Surveillance Center, National Institute of Infectious Diseases, Tokyo, Japan, ³Shinjuku Ladies Clinic, Tokyo, Japan, ⁴Department of Bacteriology I, National Institute of Infectious Diseases, Tokyo, Japan

Session: 66. Public Health: Epidemiology and Outbreaks

Thursday, October 4, 2018: 12:30 PM

Background. In Japan, syphilis notifications have increased sharply, with Tokyo accounting for a third of all cases. Importantly, the increase in women has been remarkable, with the majority being women who have sex with men. We therefore conducted a prospective case-control study to evaluate correlates of recent syphilis infection among women in Tokyo.

Methods. We employed a test-negative design case-control study among women who sought a syphilis test at a women's health clinic in Tokyo, comparing socio-economic status, medical history, and recent sexual behaviors/partnerships, between those who were diagnosed with recent syphilis infection based on clinical signs and serological test results (cases) vs. those who were nonreactive for the nontreponemal test (controls). Participants aged ≥20 years and sexually active in the past 6 months were recruited from June 2017 to March 2018 and completed a self-administered questionnaire. We described cases and assessed for correlates of recent syphilis infection based on odds ratios (ORs) and 95% confidence intervals (CIs).

Results. A total of 524 women, composed of 60 cases and 464 controls, were enrolled. The median age was 23 years (range = 20–54) among cases and 26 years (range = 20–59) among controls. Of the 60 cases, 10 (17%) were current students and three (5%) had a history of syphilis; in the past 6 months, while 35 (58%) had engaged in commercial exchange of sexual practices, 14 (23%) had only one sex partner. Having lower educational attainment (OR = 4.1; 95% CI = 2.1–8.1), not being employed full-time (OR = 3.1; 95% CI = 1.8–5.5), and commercial exchange of sexual practices (OR = 3.4; 95% CI = 2.0–5.9) were associated with case status in univariate analysis. The association between case status and inconsistent condom use (relative to consistent condom use) during vaginal/anal sex in the past 6 months (OR = 2.0; 95% CI = 0.9–4.3) became stronger when restricted to women engaged in commercial exchange of sexual practices (OR = 4.0; 95% CI = 1.4–10.9).

Conclusion. This was the first study to evaluate potential risk factors for recent syphilis infection among women in Japan. In light of these findings, we may need a multi-pronged approach to prevent and control syphilis, for both those engaged in commercial exchange of sexual practices and the general population.

Disclosures. All authors: No reported disclosures.

674. Epidemiology of Diphtheria and Antimicrobial Resistance Among Diphtheria Cases, Bijapur District, Karnataka, India, 2012–2015

Sree Kalpana Mohankumar, MBBS, MPH, EIS¹; Suresh Paschapur, MPH² and Shankareppa Mailare, MBBS²; ¹Department of Public Health, Namakkal, India, ²Public Health, District Surveillance Unit, Bijapur, India

Session: 66. Public Health: Epidemiology and Outbreaks

Thursday, October 4, 2018: 12:30 PM

Background. India contributes to 80% of diphtheria cases. Many diphtheria outbreaks were reported from Bijapur district of Karnataka state, India. Analysis of burden and drug sensitivity pattern might help to identify risk groups and to provide guidelines for treatment of diphtheria cases. Our objectives were to describe epidemiology and drug susceptibility of diphtheria cases in Bijapur district.

Methods. We did cross-sectional study between 2012 and 2015. We defined a probable case as inflammation of upper respiratory tract with adherent membranes.