

# How Important Impact of Low Level of Hematocrit Can Be on Outcome in Patients Undergoing Off Pump Coronary Artery Bypass Surgery?

Slavenka Straus, Ilirijana Haxhibeqiri Karabdic, Sanja Grabovica, Amel Hadzimehmedagic, Muhamed Djedovic, Edin Kabil, Tarik Selimovic, Nermir Granov

Clinic for Cardiovascular Surgery, Clinical Center University of Sarajevo, Sarajevo, Bosnia and Herzegovina

Corresponding author: Slavenka Straus, MD, PhD, Clinic for Cardiovascular Surgery, Clinical Center of University of Sarajevo, Sarajevo, Bosnia and Herzegovina. Address: Bolnicka 25, Sarajevo, 71000, Bosnia and Herzegovina. Phone: +387 61 203 870. E-mail: vstraus@yahoo.com. ORCID ID: 0000-0003-4605-9334

doi: 10.5455/aim.2023.31.102-106

ACTA INFORM MED. 2023 JUN 31(2): 102-106

Received: MAY 15, 2023

Accepted: JUN 23, 2023

© 2023 Slavenka Straus, Ilirijana Haxhibeqiri Karabdic, Sanja Grabovica, Amel Hadzimehmedagic, Muhamed Djedovic, Edin Kabil, Tarik Selimovic, Nermir Granov

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

## ABSTRACT

**Background:** Cardiosurgical patients can be often anemic, and preoperative anaemia is associated with increased postoperative complications, as well as morbidity and mortality.

**Objective:** The present study was conducted aiming to determine the influence of lower preoperative hematocrit values on the early postoperative recovery of patients undergoing elective off-pump coronary artery bypass grafting (OPCABG). **Methods:** Our retrospective study included 150 patients who underwent elective OPCABG surgery from September 2021 to December 2022 at the Clinic for Cardiovascular Surgery University of Sarajevo. Patients were divided into two groups, anemic and non-anemic, with anemia defined as hemoglobin level <130g/l males and <120g/l females. Study observed comorbidities and early postoperative data. **Results:** Overall prevalence of anemia in OPCABG patients was 36,6%, more frequent among older patients. Comorbidities as chronic obstructive pulmonary disease, hypertension, ejection fraction was quite similar in both groups, but body mass index was significantly higher in the group of patients with anemia (29.9% vs 28.4%). In early recovery period, vasopressors were more used in patients with anemia (63.6% vs 42.1%) and their doses were higher. Total drainage was higher in patients with anemia (744.45±45.72 ml vs 681.58±349.06 ml). Number of transfusions was higher in the group of patients with anemia: red blood cells (0.89±1.29 vs 0.29±0.84; p<0,05), fresh frozen plasma (0.65±1.13 vs 0.41±1.15; p<0,05) and platelets (0.11±0.57 vs 0,07±0.42; p>0,05). Duration of mechanical ventilation was longer in patients with anemia compared to non-anemic patients (623-91±259.83 minutes vs 469±191.13 minutes). Atrial fibrillation occurred more often in patients with anemia (12.7 vs 7.4%). Length of stay in the intensive care unit was without significant difference. **Conclusion:** Based on our findings, patients who underwent elective OPCABG with lower than normal hematocrit, needed more blood and blood products, more vasopressor drugs in higher doses, longer mechanical ventilation, all which can prolong the patient's recovery and increase the cost of treatment.

**Keywords:** off pump coronary artery bypass graft surgery, anemia, early postoperative recovery.

## 1. BACKGROUND

Anemia can be very common among patients undergoing surgery, especially in cardiac surgery. Because of extremely limited coronary reserve, patients undergoing coronary artery bypass grafting surgery represent a group of patients potentially most sensitive to the impact of low hemoglobin level. During surgery it can be aggravated by increased use of fluid, which will cause hemodilution, and by blood loss during surgery, when it is necessary to replace

this loss with blood and blood products (1).

As we know, conventional coronary artery bypass graft surgery (CABG) can be executed with the use of cardiopulmonary bypass (CPB) so call on-pump CABG or off-pump coronary artery bypass graft surgery (OPCABG), without CPB, which has been developed to decrease perioperative complications related to the use of CPB. Recent observational and randomized studies have suggested that certain high-risk patient

subgroups are more likely to benefit from an OPCABG approach, avoiding the deleterious effects of CPB (2).

OPCAB has been found to be associated with less bleeding, less renal dysfunction, a shorter length of hospital stay, and less neurocognitive dysfunction, with beneficial effects for patients at high risk for complications. Cardiac surgical procedures are estimated to be responsible for nearly 20% of the annual blood transfusions in the United States, and blood transfusion following CABG has been reported to occur in up to 60% of cases (3).

According to the World Health Organization (WHO) definition of anemia is a hemoglobin (Hb) levels < 120 g/l in non-pregnant adult females and < 130 g/l in adult males. Using WHO definition, the general consensus for preoperative anemia prevalence is around 20–30% and increases with age and the presence of comorbidities. The most common causes of preoperative anemia in cardiac surgery patients are hospital-acquired anemia, iron-deficiency anemia, and anemia of persisting chronic disease (4).

Correcting an anemia, we use allogenic blood transfusion which has been associated with significant adverse patient effects, including the occurrence of transfusion reactions, transfusion related acute lung injury, immunosuppression and immunomodulation, infectious complications, and even reduced long term survival (3).

A significant association of anemia with increased perioperative morbidity and mortality has been established in a multitude of settings and in both cardiac and noncardiac surgery (5).

The relative contribution of preoperative anemia and red blood cells (RBC) transfusion to the adverse outcomes in cardiac surgery patients remains uncertain, however, it has been shown that the interaction of the two increases morbidity and mortality far greater than the additive effects of each exposure individually (6).

## 2. OBJECTIVE

The present study was conducted aiming to determine the influence of preoperative anemia on the early postoperative recovery of patients undergoing elective OPCABG surgery.

## 3. PATIENTS AND METHODS

Our study was retrospective, descriptive, single-center study which enrolled 150 patients in the period from September 2021 to December 2022, at the Clinic for Cardiovascular Surgery Sarajevo. All patients had undergone elective off pump coronary artery bypass surgery (OPCABG).

The inclusion criteria: adult patients (>18 years of age) for the first time on CABG surgery, elective cases, patients with silent, stable, or unstable angina.

The exclusion criteria: patients with a previous history of CABG surgery, emergency surgery, and patients with combined cardiac surgery.

We collected preoperative, intraoperative and postoperative data from our clinical database and hospital charts.

The anesthesia technique was standardized for all patients. In brief, this consisted of intravenous anesthesia with propofol or etomidate. Neuromuscular blockade was achieved with Pancuronium bromide. Anesthesia was maintained by the use of a fentanyl and propofol pump, sevoflurane was used

when needed. Intraoperative fluid administration was mainly saline solution. Heparin was given at a dose of 200 IU/kg. It was reversed with protamine sulfate in the proportion of 1:1.

Conventional median sternotomy and preparation of conduits were undertaken by using a standard technique. In all the cases, the left internal mammary artery was anastomosed to the left anterior descending artery. For additional grafts, saphenous vein was used. Cardiac exposure and stabilization was achieved by Octopus Tissue Stabilizer. An intracoronary shunt was used in all cases of to support hemodynamic stability. A cell-saving device was used routinely with frequent infusions of the salvaged blood as collected. Two chest tubes were placed in the mediastinum, and additional tubes were left in the pleura as needed. Additional protamine sulfate was used in relation to the prolonged ACT with excessive bleeding. Patients were weaned from mechanical ventilation (MV) as soon as they were fully awake, hemodynamical stable, normal body temperature, satisfied diuresis and bleeding and with laboratory tests in normal ranges.

### Statistical analysis

Results of the analysis are presented as tables and charts by number of cases, percentage, mean, standard deviation and the range. Statistical comparison was performed using the nonparametric Chi-square and Mann-Whitney test depending on the type of data. Results of the test were considered as significant at the confidence level of 95%. Analysis was performed using the statistical package IBM Statistics SPSS v 23.0.

## 4. RESULTS

The first table shows sociodemographic data that indicate that the respondents with anemia were older with an average age of 65.29±6.64 years, compared to patients without anemia and average age of 61.8.29±8.29 years (p=0,0023). According to gender there were no statistically significant differences with both groups (p>0,05).

In the group of patients without anemia more present were diabetes mellitus (41.1 vs 38.2%), stroke (6.3 vs 3.6), myo-

Variable	Anemia		p	
	Absent (N=95)	Present (N=55)		
Age – Mean±SD (Range)	61,89±8,29 (40-77)	65,29±6,64 (53-80)	0,023	
Gender – N (%)	Male	77 (81,1)	41 (74,5)	0,409
	Female	18 (18,9)	14 (25,5)	
Diabetes mellitus – N (%)	39 (41,1)	21 (38,2)	0,433	
Stroke – N (%)	6 (6,3)	2 (3,6)	0,384	
Myocardial infarction < 6 weeks N (%)	28 (29,5)	9 (16,4)	0,080	
Myocardial infarction > 6 weeks N (%)	39 (41,1)	25 (45,5)	0,612	
Chronic obstructive pulmonary disease – N (%)	4 (4,2)	4 (7,3)	0,465	
Arterial hypertension – N (%)	77 (81,1)	50 (90,9)	0,157	
Smoking – N (%)	39 (41,1)	15 (27,3)	0,113	
BMI – Mean±SD (Range)	28,48±3,89 (20,2-41,2)	29,95±3,86 (23,1-41,1)	0,040	
Ejection fraction – Mean±SD (Range)	44,41±9,32 (25-65)	44,07±9,97 (25-65)	0,834	
Percutaneous coronary intervention – N (%)	21 (22,1)	17 (30,9)	0,159	

Table 1. Sociodemographic characteristics, comorbidities and risk factors

Variable	Anemia		p
	Absent (N=95)	Present (N=55)	
Hemoglobin – Mean±SD (Range)	145,69±10,12 (124-177)	114,05±12,65 (98-177)	0,0001
Hematocrit – Mean±SD (Range)	42,91±3,52 (37-54)	34,77±2,98 (30-41)	0,0001
Platelets – Mean±SD (Range)	227,96±40,05 (127-331)	221,56±53,69 (118-327)	0,467
Creatinine – Mean±SD (Range)	81,04±20,65 (43-153)	90,69±28,51 (42-173)	0,044
Creatinine >120 – N (%)	3 (3,2)	8 (14,5)	0,019

**Table 2. Preoperative values of hemoglobin, hematocrit, platelets and serum creatinine**

Variable	Anemia		p
	Absent (N=95)	Present (N=55)	
Transfusion of RBC – Mean±SD (Range)	0,29±0,86 (0-6)	0,93±1,14 (0-3)	0,0001
Transfusion of FFP – Mean±SD (Range)	1,11±1,37 (0-3)	1,81±1,32 (0-4)	0,004
Platelet transfusion – Mean±SD (Range)	0,04±0,07 (0-3)	0,07±0,38 (0-2)	0,576

**Table 3. Intraoperative data**

cardial infarction less than 6 weeks (29.5 vs 16.4%) and more than 6 weeks before surgery (41.1 vs 45.5%).

Body mass index was statistically significantly higher in the group of patients with anemia (29.95±3.86 vs 28.48±3.89).

Ejection fraction values were higher in average in the group of patients without anemia (44.41±9.32 vs 44.07±9.97 %) and percutaneous coronary intervention (PCI) was more present in the group of patients with anemia (30.9 vs 22.1), all without statistically significant difference (p>0.05).

In the group of patients with present anemia statistically significantly higher mean values of creatinine were recorded (90.69±28.51 vs 81.04±20.62) and more patients with creatinine values above 120 (14.5 vs 3.2%) (Table 2).

Values of blood parameters were higher in the group of patients without anemia, hemoglobin (145.69±10.12 vs 114.05±12.65; p<0,05), hematocrit (42.91±3.52 vs 34.77±2.98; p<0,05) and platelets count (227.96±40.05 vs 221.56±53.69; p>0,05).

The main intraoperative data are shown in the Table 3. Patients with anemia received more transfusions during the surgery – RBC (0.93±1.14 vs 0.29±0.86; p<0.05), FFP (1.81±1.32 vs 1.11±1.37; p<0.05) and platelet transfusion (0.07±0.38 vs 0.04 vs 0.07; p>0,05).

Transfusion of RBC and FFP were significantly different between two groups.

The dose of Cyklokapron was statistically significantly higher in the group of patients with anemia (2.54±0.98 vs 2.25±0.93), unlike Protamine which was statistically significantly more often administered in patients without anemia (28.4 vs 14.5%).

Although without statistically significant difference the insulin was more used in patients with anemia (73.6 vs 28.4%), but its dose was higher in patients without anemia (1.07±4.43 vs 0.44±0.50).

Vasopressors were statistically significantly more often used in patients with anemia (63.6 vs 42.1%) and their dose was higher (0.63±0.49 vs 0.42±0.50).

Variable	Anemia		p
	Absent (N=95)	Present (N=55)	
Dose of Cyklokapron in grams – Mean±SD (Range)	2,25±0,93 (0-6)	2,54±0,98 (2-6)	0,043
Insulin – N (%)	27 (28,4)	24 (43,6)	0,298
Insulin (dose) – Mean±SD (Range)	1,07±4,43 (0-30)	0,44±0,50 (0-1)	0,193
Vasopressors – N (%)	40 (42,1)	35 (63,6)	0,017
Vasopressors – Mean±SD (Range)	0,42±0,50 (0-1)	0,63±0,49 (0-1)	0,011
Total drainage in 48 hours – Mean±SD (Range)	681,58±349,06 (258-2435)	744,45±45,72 (0-2020)	0,115
Transfusion of RBC – Mean±SD (Range)	0,29±0,84 (0-6)	0,89±1,29 (0-6)	0,001
Transfusion of FFP – Mean±SD (Range)	0,41±1,15 (0-6)	0,65±1,13 (0-3)	0,046
Transfusion of platelets – Mean±SD (Range)	0,07±0,42 (0-3)	0,11±0,57 (0-3)	0,856
Minutes of mechanical ventilation – Mean±SD (Range)	469,25±191,13 (80-1289)	623,91±259,83 (0-1200)	0,0001
Occurrence of atrial fibrillation – N (%)	7 (7,4)	7 (12,7)	0,277
Additional dose of Protamine N (%)	27 (28,4)	8 (14,5)	0,039
ICU duration of stay in days – Mean±SD (Range)	3,13±1,15 (2-11)	3,06±1,07 (2-7)	0,467

**Table 4. Early postoperative recovery - Intensive Care Unit**

Total drainage, although without statistically significant difference was higher in patients with anemia (744.45±45.72 vs 681.58±349.06).

Number of transfusions was higher in group of patients with anemia as follows: RBC (0.89±1.29 vs 0.29±0.84; p<0,05), FFP (0.65±1.13 vs 0.41±1.15; p<0,05) and platelets (0.11±0.57 vs 0,07±0.42; p>0,05).

Duration of MV was statistically significantly longer in patients with compared to patients without anemia (623-91±259.83 vs 469±191.13).

Occurrence of AF was more often in patients with anemia (12.7 vs 7.4%) but without statistically significant difference (p>0.05).

The duration of intensive care unit stay was almost identical in observed groups, longer for patients without anemia (3.13±1.15 vs 3.06±1.07 days).

There were no deaths during early postoperative recovery.

## 5. DISCUSSION

Conventional CABG with CPB is at increased risk of perioperative blood loss requiring a transfusion of blood products, CPB leads to thrombocytopenia and platelet dysfunction, dilution and consumption of coagulation factors, hyperfibrinolysis, and a residual heparin effect. So, there has been an increasing interest in OPCAB with the suggestion that this may avoid the complication of CPB and reduce postoperative blood loss, usage of blood and blood products, morbidity and mortality. All of which reduces the stay in the intensive care unit, length of hospitalization and thus the total cost of treatment. Preoperative anemia has been identified as the single-most important determinant of transfusion, and many postoperative complications (use of vasoconstrictors, specific drugs to control bleeding and use of allogeneic blood products) (7).

If patient is anemic, in order to avoid donating blood and blood products and thereby reduce postoperative complications, various preoperative blood management strategies have been studied such as iron therapy, erythropoietin administration, dietary supplements of vitamin B12 and folate, or preoperative autologous blood donation. This is particularly relevant since it makes it possible to intervene in patients whose elective cardiac surgery is scheduled (8, 9).

When assessed according to criteria, about 20% - 50% of patients undergoing cardiac surgery suffer from preoperative anemia, with especially high incidences in older patients or in patients with chronic diseases and/or multiple comorbidities. In order to correct anemia, RBC transfusion is indicated and it is another recognized risk factor for adverse events after cardiac surgery. The pathophysiologic mechanism by which transfusions might harm has not been fully elucidated, but it is known that erythrocytes undergo irreversible morphologic and biochemical changes during storage. As a result, after transfusion, they can promote a proinflammatory state, impair tissue oxygen delivery, and exacerbate tissue oxidative stress. This in turn can contribute to organ dysfunction and explain, at least in part, the increased hospital mortality and morbidity (10).

The aging of population, causes an increase of anemia in elderly people. In our study patients with anemia were older than patients without anemia with an average age of  $65.29 \pm 6.64$  years. Same results gave Cappellini MD in his study (11).

Kanic V in his study found that women suffered anemia more than men and that the prevalence of anemia increased rapidly with age in men but not in women. We found no differences between genders in anemic and non anemic group (12).

The prevalence of comorbidities in our patients with anemia is very similar to the data provided by Kulier A et al. and Hung M in their studies (13, 14).

Obesity is linked to a variety of unfavorable outcomes, including anemia, which is a serious global public health problem. The prevalence of obesity along with anemia suggests a relationship between obesity and anemia. Our anemic patients had higher BMI compared to patients without anemia which is the same date from Saad RA et al study (15).

Anand IS in his study showed that anemic patients have a better left ventricular ejection fraction (LVEF) and thus gives the same results as we found in our study (16).

Anemia is often associated with DM. Patients with diabetes need to be screened for anemia along with other risk factors and anemia should be corrected appropriately to improve overall clinical outcomes. Our study had more diabetic patients in non-anemic group, a finding that differs from numerous other studies (17).

Our study confirmed higher dose of vasopressors in anemic patients and that they are used more often. Many studies had the same data, but Pongsuthana S et al. in their study did not find differences in the use of inotropic drugs between anemic and non-anemic patients (18).

Duration of MV in our patients with anemia was significantly longer compared with patients without anemia. The same data was found in the study of Zilberberg MD et al (19).

Zochios V et al in their study found that hypoxemia and anemia on admission to ICU were the only 2 factors inde-

pendently associated with the need for escalation of respiratory support or prolonged invasive ventilation. Escalation of respiratory support or prolonged invasive ventilation is frequently seen in cardiac surgery patients and is highly associated with increased mortality and morbidity (20).

We found in our study that atrial fibrillation occurred more often in anemic patients which is the same as in the study from Lu Dai et al. and Hanna-Rivero N et al. which confirmed that anemia may be associated with the development of new-onset AF and anemia may also be more likely to experience clinical recurrence of AF (21, 22).

In our study ICU days were almost identical in both groups. Lu Dai et al. showed in their study that anemic patients experienced longer ICU (27 versus 48 hours,  $p < 0.001$ ) and postoperative hospital (6.1 versus 7.4 days,  $p < 0.001$ ) length of stay than non-anemic patients. The same results confirm study from Lorente V et al. that both ICU and hospital stay were significantly longer in patients with anemia (23).

In the investigation made by Sadeghi A it is confirmed that about one-third of the study population had anemia before surgery and these patients required blood transfusion 2.5 times more than those without anemia (24). A low preoperative Hct is an independent risk factor for the need for a transfusion, and the mortality rate among patients is directly proportional to the need for a transfusion, regardless of the type of surgery (25).

Defining the subtypes of anemia (iron-deficiency anemia, anemia of chronic disease or anemia due to both conditions) based on laboratory testing seems to be the key for specific and individualized therapy. The optimal treatment strategy remains unclear. Although therapeutic interventions shortly before surgery have been suggested, treatments should be initiated within several weeks before surgery to effectively improve RBC mass and hemoglobin values (26).

Lower pre-operative hemoglobin concentrations are independently associated with increased utilization of hospital resources after cardiac surgery. Each g/dl unit fall in pre-operative hemoglobin concentration is independently associated with increased transfusion requirements of blood products and increased post-operative length of stay in intensive care and in hospital (27).

## 6. CONSLUSION

Our study showed that patients who underwent elective OPCABG with preoperative anemia, got more units of blood and blood products, had higher doses of inotropic and vasoactive drugs, and longer mechanical ventilation. In order to reduce the use of blood and blood products, which lead to many unwanted complications, it is necessary to correct the existing preoperative anemia in elective OPCABG surgery.

- **Author's contributions:** All authors made a significant contribution to this study, whether in the conception, study design, execution, acquisition of data, analysis and interpretation, or in all these areas; the drafting, revising, or critical reviewing of the article; giving final approval of the version to be published; agreeing on the journal to which the article has been submitted; and agreeing to be accountable for all aspects of the work.
- **Conflict of interest:** The author reports no conflicts of interest or financial interest in this work.
- **Financial support and sponsorship:** This work was no funded.

REFERENCES

1. Lu Dai, Stephanie L. Mick, Keith R. McCrae, Penny L. Houghtaling, Joseph F. Sabik et al. Preoperative Anemia in Cardiac Operation: Does Hemoglobin Tell the Whole Story? *Ann Thorac Surg* 2018; 105: 100–107. doi: 10.1016/j.athoracsur.2017.06.074.
2. Paparella D, Guida P, Scrascia G, Fanelli V, Contini M, Zaccaria S, et al. On-pump versus off-pump coronary artery bypass surgery in patients with preoperative anemia. *J Thorac Cardiovasc Surg* 2015; 149: 1018-1026. doi.org/10.1016/j.jtcvs.2014.12.049.
3. LaPar DJ, Hawkins RB, McMurry TL, Isbell JM, Rich JB, Speir AM, Quader MA, Kron IL, Kern JA, Ailawadi G. Preoperative anemia versus blood transfusion: which is the culprit for worse outcomes in cardiac surgery? *J Thorac Cardiovasc Surg* 2018; 156:66-74. 10.1016/j.jtcvs.2018.03.109
4. WHO. Haemoglobin concentrations for the diagnosis of anaemia and assessment of severity. In: *Vitamin and Mineral Nutrition Information System*. Geneva: World Health Organ; 2011. <http://scholar.google.com/scholar?hl=en&btnG=Search&q=intitle:Haemoglobin+concentrations+for+the+diagnosis+of+anaemia+and+assessment+of+severity#1>.
5. Padmanabhan H, Siau K, Curtis J, Ng A, Menon S, Luckraz H, et al. Preoperative Anemia and outcomes in cardiovascular surgery: systematic review and meta-analysis. *Ann Thorac Surg* 2019;108(6):1840–1848. doi.org/10.1016/j.athoracsur.2019.04.108.
6. Nguyen Q, Meng E, Berube J, Bergstrom R, Lam W. Preoperative anemia and transfusion in cardiac surgery: a single-centre retrospective study. *Journal of Cardiothoracic Surgery*. 2021; 16: 109. doi.org/10.1186/s13019-021-01493-z
7. Ranucci M, Baryshnikova E, Castelvechio S, Pelissero G;Surgical and Clinical Outcome Research (SCORE) Group. Major bleeding, transfusions, and anemia: the deadly triad of cardiac surgery. *Ann Thorac Surg*. 2013; 96: 478-485.
8. Mueller MM, Van Remoortel H, Meybohm P, Aranko K, Aubron C, Burger R, et al. Patient blood management: recommendations from the 2018 Frankfurt consensus conference. *JAMA*. 2019; 321(10): 983–997. <https://doi.org/10.1001/jama.2019.0554>.
9. Spahn DR, Schoenrath F, Spahn GH, Seifert B, Stein P, Theusinger OM, Kaserer A, Hegemann I, Hofmann A, Maisano F, Falk V Effect of ultra-shortterm treatment of patients with iron deficiency or anaemia undergoing cardiac surgery: a prospective randomised trial. *Lancet*. 2019; 393(10187): 2201–2212. Available from: [https://doi.org/10.1016/S0140-6736\(18\)32555-8](https://doi.org/10.1016/S0140-6736(18)32555-8)
10. Kattou F, Montandrou O, Rezik M, Delentdecker P, Brini K, Zannis K, Beaussier M. Critical preoperative hemoglobin value to predict anemia-related complications after cardiac surgery. *J Cardiothorac Vasc Anesth* 2022; 36:1901-1907. doi:10.1053/jjvca.2022.01.013.
11. Cappellini MD, Motta I. Anemia in Clinical Practice-Definition and Classification: Does Hemoglobin Change With Aging? *Semin Hematol*. 2015 Oct; 52(4): 261-269. doi: 10.1053/j.seminhematol.2015.07.006.
12. Kanic V, Kompara G, Vollrath M, Suran D, Kanic Z. Age-Specific Sex-Based Differences in Anemia in Patients with Myocardial Infarction. *J Womens Health (Larchmt)*. 2019 Jul; 28(7): 1004-1010. doi: 10.1089/jwh.2018.7211.
13. Kulier A, Levin J, Moser R, Rumpold-Seitlinger G, Tudor IC, Snyder-Ramos SA, et al; Investigators of the Multicenter Study of Perioperative Ischemia Research Group; Ischemia Research and Education Foundation. Impact of preoperative anemia on outcome in patients undergoing coronary artery bypass graft surgery. *Circulation*. 2007 Jul; 116(5): 471-479.
14. Hung M, Besser M, Sharples LD, Nair SK, Klein AA. The prevalence and association with transfusion, intensive care unit stay and mortality of pre-operative anaemia in a cohort of cardiac surgery patients. *Anaesthesia*. 2011 Sep; 66(9): 812-818.
15. Saad RA, Qutob HM. The relationship between anemia and obesity. *Expert Rev Hematol*. 2022 Oct; 15(10): 911-926. doi: 10.1080/17474086.2022.2131521.
16. Anand IS, Gupta P. Anemia and Iron Deficiency in Heart Failure: Current Concepts and Emerging Therapies. *Circulation*. 2018 Jul 3; 138(1): 80-98. doi: 10.1161/CIRCULATIONAHA.118.030099.
17. Sahay M, Kalra S, Badani R, Bantwal G, Bhoraskar A, Das AK, Dhorepatil B, Ghosh S, Jeloka T, Khandelwal D, Latif ZA, Nadkar M, Pathan MF, Saboo B, Sahay R, Shimjee S, Shrestha D, Siyan A, Talukdar SH, Tiwaskar M, Unnikrishnan AG. Diabetes and Anemia: International Diabetes Federation (IDF) - Southeast Asian Region (SEAR) position statement. *Diabetes Metab Syndr*. 2017 Dec;11 Suppl 2:S685-S695. doi: 10.1016/j.dsx.2017.04.026
18. Pongsuthana S, Tanghongse V. Correlation of anemia and clinical outcomes in heart failure at Rajavithi Hospital. *J Med Assoc Thai*. 2014 Nov; 97 Suppl 11: S41-47.
19. Zilberberg MD, Stern LS, Wiederkehr DP, Doyle JJ, Shorr AF. Anemia, transfusions and hospital outcomes among critically ill patients on prolonged acute mechanical ventilation: a retrospective cohort study. *Crit Care*. 2008; 12(2): R60. doi: 10.1186/cc6885
20. Zochios V, Chandan JS, Schultz MJ, Morris AC, Parhar KK, Giménez-Milà M, Gerrard C, Vuylsteke A, Klein AA. The Effects of Escalation of Respiratory Support and Prolonged Invasive Ventilation on Outcomes of Cardiac Surgical Patients: A Retrospective Cohort Study. *J Cardiothorac Vasc Anesth*. 2020 May;34(5):1226-1234. doi: 10.1053/jjvca.2019.10.052
21. Lu Dai, Stephanie L. Mick, Keith R. McCrae, Penny L. Houghtaling, Joseph F. Sabik, III, Eugene H. Blackstone, Colleen G. Koch. Preoperative Anemia in Cardiac Operation: Does Hemoglobin Tell the Whole Story? *Ann Thorac Surg* 2018; 105: 100–107. doi: 10.1016/j.athoracsur.2017.06.074.
22. Hanna-Rivero N, Tu SJ, Elliott AD, Pitman BM, Gallagher C, Lau DH, Sanders P, Wong CX. Anemia and iron deficiency in patients with atrial fibrillation. *BMC Cardiovasc Disord*. 2022 May 4; 22(1): 204. doi.org/10.1186/s12872-022-02633-6
23. Lorente V, Aboal J, Garcia C, Sans-Roselló J, Sambola A, Andrea R, Tomás C, Bonet G, Viñas D, El Ouaddi N, Montero S, Cantalapiedra J, Pujol M, Hernández I, Pérez-Rodríguez M, Llaó I, Sánchez-Salado JC, Gual M, Ariza-Solé A. Anemia in patients with high-risk acute coronary syndromes admitted to Intensive Cardiac Care Units. *J Geriatr Cardiol*. 2020 Jan; 17(1): 35-42. doi: 10.11909/j.issn.1671-5411.2020.01.006
24. Ali Sadeghi, Rasool Ferasatkish, Avaz Heydarpour, Rasoul Azarfarin, Mohsen Ziyaeifard, Zahra Faritous, Fatemehshima Hadipourzadeh. Prevalence of Anemia in Patients Undergoing Cardiac Surgery and Need for Transfusion During Surgery Regarding Hemoglobin Levels in Rajaie Heart Center. *Iranian Heart Journal* 2020; 21(1): 94-102
25. LaPar DJ, Hawkins RB, McMurry TL, et al.: Preoperative anemia versus blood transfusion: which is the culprit for worse outcomes in cardiac surgery?. *J Thorac Cardiovasc Surg*. 2018, 156:66-74. 10.1016/j.jtcvs.2018.03.109.
26. Kloeser R, Buser A, Bolliger D. Treatment Strategies in Anemic Patients Before Cardiac Surgery. *Journal of Cardiothoracic and Vascular Anesthesia*. 2023; 37: 266-275. doi:10.1053/jjvca.2022.09.085.
27. George Hallward, Nikhail Balani, Stuart McCorkell, James Roxburgh, Victoria Corneliu. The Relationship Between Preoperative Hemoglobin Concentration, Use of Hospital Resources, and Outcomes in Cardiac Surgery. *J Cardiothorac Vasc Anesth*. 2016 Aug; 30(4): 901-908. doi: 10.1053/jjvca.2016.02.004.