

Letter to the editor

Open Access

# Chromosome-scale genome assembly of brown-spotted flathead *Platycephalus* sp.1 provides insights into demersal adaptation in flathead fish

Flatheads are valuable commercial fish species endemic to the Indo-West Pacific. Due to their economic value and unique biological traits, such as metamorphosis and camouflage, they serve as ideal marine organisms for studies on demersal adaptation and evolution. The brown-spotted flathead (*Platycephalus* sp.1) is the most widely distributed in the northwestern Pacific. Despite the lack of a valid scientific name, it has been long recognized and exploited in the marine fisheries of China, Japan, and Korea. In the current study, we applied Illumina, PacBio, and Hi-C sequencing to assemble a chromosome-scale genome for this species. The assembled genome was 660.63 Mb long with a scaffold N50 of 28.65 Mb and 100% of the contigs were anchored onto 24 chromosomes. We predicted 22 743 protein-coding genes, 94.8% of which were functionally annotated. Comparative genomic analyses suggested that *Platycephalus* sp.1 diverged from its common ancestor with *Gasterosteus aculeatus* ~88.4 million years ago. The expanded gene families were significantly enriched in immune, biosynthetic, and metabolic pathways. Furthermore, three shared Gene Ontology (GO) terms and 377 common positively selected genes were identified between flathead and flatfish species, suggesting that these genes may contribute to demersal adaptation in flatheads. The assembled genomic data provide a valuable molecular resource for further research on the biological and adaptive evolution of flathead species.

Flathead fish (family Platycephalidae) are endemic demersal carnivores distributed in the Indo-West Pacific. The genus *Platycephalus* is the most speciose in the Platycephalidae family (Imamura, 2015; Nakabo, 2002). Given their high-quality flesh and large body size, flatheads are commercially valuable and among the first fish to be targeted by fisheries in Indo-Pacific regions, including Japan, China,

and Australia (Puckridge et al., 2013). In addition, flatheads are camouflage masters and ambush predators with symmetrical body-axis metamorphosis (Hyndes et al., 1992), similar to flatfish species (Shao et al., 2017). Flatheads are hypothesized to exhibit protandrous hermaphroditism and sex-reversal (i.e., sex change from male to female) (Shinomiya et al., 2003). These biological characteristics make flatheads valuable models for investigating the molecular mechanisms underlying demersal adaptation and evolution, as well as innovations in sex determination and growth in teleosts. High-quality genome assemblies should benefit the above-mentioned research fields as well as our understanding of the adaptive evolution of flathead species.

The brown-spotted flathead, *Platycephalus* sp.1 (sensu Nakabo, 2002), is a widely distributed species in the northwestern Pacific (Nakabo, 2002). Despite the lack of a valid scientific name, *Platycephalus* sp.1 has long been recognized in Japan as “Yoshino-gochi” (Nakabo, 2002). Recently, our research team reported *Platycephalus* sp.1 as a new record species in the coastal waters of China, with several distinct morphological characters, including dark brown spots scattered on head and body, lateral line scales 83 to 99, and no yellow blotch on caudal fin (Qin et al., 2013). Considering the contributions of Japanese ichthyologists, we herein recommend “*Platycephalus yoshinoi*” as the scientific name of this species, in reference to its Japanese name. However, the lack of genomic resources and insufficient exploitation of genetic information have limited study on the evolutionary biology, fisheries management, and conservation of this species. Thus, we performed a chromosome-scale genome assembly of *Platycephalus* sp.1 using PacBio, Illumina, and Hi-C sequencing technologies.

An adult male *Platycephalus* sp.1 (Figure 1A) captured from the littoral waters of Qingdao, China (E120.275°, N36.109°) in

This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Copyright ©2021 Editorial Office of Zoological Research, Kunming Institute of Zoology, Chinese Academy of Sciences


Received: 13 August 2021; Accepted: 30 August 2021; Online: 31 August 2021

Foundation items: This study was supported by the Fundamental Research Funds for Zhejiang Provincial Universities and Research Institutes (2019JZ00001) and National Natural Science Foundation of China (41776171)

November 2019, was used for genome sequencing. Muscle tissue below the dorsal fin was collected for DNA extraction using the cetyltrimethylammonium bromide (CTAB) method. DNA quality and concentration were measured using agarose gel electrophoresis (1%), pulse field gel electrophoresis (1%), and Qubit 3.0 (Thermo Fisher Scientific, USA). To facilitate prediction of protein-coding genes, we extracted total RNA from seven flathead tissues, including muscle, gill, gonad, brain, spleen, liver, and eye, using TRIzol reagent (Invitrogen, USA) according to the manufacturer's instructions. RNA integrity and quantity were evaluated using an Agilent 2100 Bioanalyzer (Agilent, USA). Seven RNA-seq libraries were prepared using a NEBNext Ultra RNA Library Prep Kit for Illumina (New England BioLabs, USA) following the manufacturer's protocols, and were then sequenced on the Illumina Novaseq6000 sequencing platform (Illumina, USA).

Illumina short-insert (350 bp) libraries were prepared and sequenced on the Illumina Novaseq6000 platform (Illumina, USA). In total, 83.91 Gb of Illumina short-read sequencing data were generated (Supplementary Table S1). *K*-mer analysis was conducted using Jellyfish v2.2.10 (Marçais & Kingsford, 2011). The *k* value was set to 17 and genome size was estimated to be 674.96 Mb, with a heterozygosity ratio of

0.77% and repeat sequence ratio of 29.23%. One 20 kb single-molecule real-time (SMRT) sequencing bell library was constructed, and long-read sequencing was performed using the PacBio Sequel II sequencer (Pacific Biosciences, USA). The PacBio sequencing platform generated 138.12 Gb of high-quality long-read sequencing data, with a mean subread length of 20 099 bp and average read length of 31 100 bp (Supplementary Table S1). The PacBio long reads were used for *de novo* genome assembly with Wtdbg2 (Ruan & Li, 2020) and draft contigs were corrected using Arrow v2.2.1 (Chin et al., 2013) with the same PacBio dataset. Finally, the PacBio sequencing data resulted in a 660.63 Mb assembly with 60 contigs and contig N50 of 23.07 Mb. By comparison, our result is the most contiguous assembly of reported species in the order Scorpaeniformes (Supplementary Table S2). Illumina short reads from the same individual were further used to evaluate the quality of the initial genome assembly using BWA v0.7.10 (Li & Durbin, 2010) and assembly completeness was also evaluated using Benchmarking Universal Single-Copy Orthologs (BUSCO) v3.0.1 (Simão et al., 2015) and Core Eukaryotic Gene Mapping Approach (CEGMA) (Parra et al., 2007). Quality evaluation of the initial genome assembly showed that 97.64% of the Illumina data were successfully



**Figure 1 Genomic analyses of brown-spotted flathead *Platycephalus sp.1***

A: Photograph of brown-spotted flathead. B: Hi-C contact map of *Platycephalus sp.1* genome. LG 0-23 represent 24 pseudochromosomes. Color bar indicates contact density from white (low) to black (high). C: Gene family comparison between *Platycephalus sp.1* and investigated teleost species. D: Phylogenetic topology of 14 teleost species investigated in this study.

mapped to the assembled genome. BUSCO analysis against the actinopterygii\_odb9 database found that 96.4% of the conserved single-copy orthologous genes, including 95.5% complete and 0.9% fragmented genes, were found in the genome assembly, while CEGMA identified 94.76% of the core eukaryotic genes (Supplementary Table S3).

To obtain a chromosome-scale genome assembly, a Hi-C library was prepared following Hi-C library protocols (Rao et al., 2014) and sequenced using the Illumina Novaseq6000 sequencing platform (Illumina, USA). The Hi-C library generated 245.62 Gb of raw data, representing 364-fold coverage (Supplementary Table S1). Duplicate removal, sorting, and quality control were performed using HiC-Pro v2.8.0 (Servant et al., 2015). LACHESIS (Burton et al., 2013) was used to cluster, order, and orient the contigs into a chromosome-scale assembly. Using LACHESIS, 100% of the initial assembled sequences were anchored to 24 pseudochromosomes with lengths ranging from 12.68 to 35.47 Mb (Figure 1B). To the best of our knowledge, this *Platycephalus* sp.1 genome is the first marine fish genome assembly with a 100% anchoring rate. Finally, the total length of the genome assembly was 660.63 Mb, and contig N50 and scaffold N50 were 23.07 Mb and 28.65 Mb, respectively (Supplementary Table S2).

Combined homology alignment and *de novo* search were applied using our repeat annotation pipeline. A *de novo* repetitive elements database was built by LTR\_FINDER (Xu & Wang, 2007), RepeatScout (Price et al., 2005), and RepeatModeler (Hubley & Smit, 2015) with default parameters. Tandem repeats were also extracted *ab initio* using TRF v4.09 (Benson, 1999). All repeat sequences with length > 100 bp and gap “N” < 5% constituted the raw transposable element (TE) library. The homology-based predictions were searched against the Repbase database (Bao et al., 2015) using RepeatMasker v3.3.0 (Chen, 2004) and its in-house script RepeatProteinMask with default parameters. Repbase and our *de novo* TE library were processed using UCLUST (Edgar, 2010) to yield a non-redundant library and RepeatMasker was used to identify DNA-level repeats. In total, 167.49 Mb of repetitive sequences were annotated, representing 25.35% of the genome assembly (Supplementary Table S4). The repetitive sequences were dominated by long terminal repeats (LTRs, 111.04 Mb, 16.81%), DNA transposons (20.58 Mb, 3.12%), and long interspersed elements (LINEs, 9.05 Mb, 1.37%). In comparison, the dominant repetitive elements in the *Sebastes schlegelii* genome are DNA transposons rather than LTRs (He et al., 2019; Liu et al., 2019).

We employed *ab initio*, homology-based, and RNA-seq-assisted prediction to detect protein-coding genes. For homology-based prediction, the protein sequences of *Danio rerio*, *Larimichthys crocea*, *Gadus morhua*, *Oryzias latipes*, and *Gasterosteus aculeatus* were downloaded from the GenBank and Ensembl databases (Cunningham et al., 2019). The protein sequences were aligned against the *Platycephalus* sp.1 genome using TBLASTN v2.2.26 (e-value ≤ 1e-5), and matching proteins were then aligned to the homologous genome sequences for accurate spliced alignments with GeneWise v2.4.1 (Doerks et al., 2002). *Ab*

*initio* prediction was performed using Augustus v3.2.3 (Stanke & Waack, 2003), GeneID v1.4 (Blanco et al., 2007), GeneScan v1.0 (Burge & Karlin, 1997), GlimmerHMM v3.04 (Majoros et al., 2004), and SNAP v2013-11-29 (Korf, 2004) based on the repeat masked genome sequences. Transcripts of the seven tissues were assembled by Trinity and aligned to the genome using PASA (Haas et al., 2003). To optimize genome annotation, the transcriptome reads from the seven tissues were also aligned to the genome directly using HISAT v2.0.4 (Pertea et al., 2016) to identify exon regions and splice positions. The alignment results were then used as inputs in Stringtie v1.3.3 (Pertea et al., 2016) for genome-based transcript assembly, and coding regions were then predicted using TransDecoder v2.0 (<http://transdecoder.github.io>). Finally, genes predicted by the above three methods were merged into a non-redundant reference gene set with EvidenceModeler v1.1.1 (Haas et al., 2008) with identical weights. Protein-coding genes were then annotated by alignment against the SwissProt, NT, NR, GO, and Kyoto Encyclopedia of Genes and Genomes (KEGG) databases using BLAST+ v2.2.28 (McGinnis & Madden, 2004) with an e-value threshold of 1e-5. InterProScan v5.31 (Mulder & Apweiler, 2007) was used to predict protein function based on conserved domains and motifs by searching against ProDom, PRINTS, Pfam, SMRT, PANTHER, and PROSITE. Finally, 22 743 protein-coding genes were predicted and 21 555 (94.8%) were successfully annotated (Supplementary Tables S5, S6). Compared to other teleost species, including *D. rerio*, *Larimichthys crocea*, *Gadus morhua*, *O. latipes*, and *Gasterosteus aculeatus*, the predicted gene models of *Platycephalus* sp.1 showed similar distribution patterns in coding sequence (CDS) length, exon length, exon number, gene length, and intron length (Supplementary Figure S1).

For non-coding RNA (ncRNA) prediction, transfer RNAs (tRNAs) were predicted using tRNAscan-SE v1.3.1 (Lowe & Eddy, 1997); ribosomal RNAs (rRNAs) were identified by alignment to closely related rRNA sequences using BLAST; and microRNAs (miRNAs) and small nuclear RNAs (snRNAs) were identified against the Rfam database (Daub et al., 2015) using Infernal v1.1 (Nawrocki & Eddy, 2013). In total, 2 876 ncRNAs, including 1 118 miRNAs, 1 267 tRNAs, 92 rRNAs, and 399 snRNAs, were identified in the assembled genome (Supplementary Table S7).

Gene families were clustered for *Platycephalus* sp.1 and 13 other species (*D. rerio*, *Gadus morhua*, *O. latipes*, *Gasterosteus aculeatus*, *Hippocampus comes*, *Takifugu rubripes*, *Labrus bergylta*, *Cynoglossus semilaevis*, *Scophthalmus maximus*, *Lepisosteus oculatus*, *Tetraodon nigroviridis*, *Xiphophorus maculatus*, and *Gambusia affinis*) using OrthoMCL v2.0.9 (Li et al., 2003). In brief, the longest transcript of each gene was retained and all-against-all BLASTP analysis was conducted with a cutoff of 1e-5. Orthologs and paralogs were grouped using OrthoMCL v2.0.9 with an inflation index of 1.5. Comparative genomic analysis revealed that the predicted genes could be clustered into 20 743 gene families (Figure 1C). A total of 6 385 gene families were shared among all 14 species, including 1 539 orthologous gene families. Based on the single-copy orthologous genes, a species-level phylogenetic topology was generated using

PhyML v3.0 (Guindon et al., 2010) to evaluate the evolutionary relationships. The MCMCTree program implemented in the PAML package (Yang, 2007) was used to estimate divergence times among species. The divergence times of *Takifugu rubripes* and *Tetraodon nigroviridis*, *Labrus bergylta* and *Gasterosteus aculeatus*, *X. maculatus* and *Gambusia affinis*, *Gasterosteus aculeatus* and *Gadus morhua*, *D. rerio* and *O. latipes*, and *D. rerio* and *Lepisosteus oculatus* were used as calibration times and were downloaded from the TimeTree database (<http://www.timetree.org/>). The topology results indicated that *Gasterosteus aculeatus* was most closely related to *Platycephalus* sp.1, with a divergence time of 88.4 million years ago (Figure 1D). The divergence time was relatively larger than that between *Sebastes schlegelii* and *Gasterosteus aculeatus* (He et al., 2019; Liu et al., 2019), suggesting that *Platycephalus* sp.1 is a more basal species than *Sebastes schlegelii* in the order Scorpaeniformes. This suggestion is consistent with previous studies based on mitochondrial and nuclear gene sequences (Cui et al., 2019; Smith et al., 2018).

Expansion and contraction of gene families between ancestors and each selected species were determined using CAFÉ (De Bie et al., 2006). The CODEML program in the PAML package (Yang, 2007) was used to estimate the nonsynonymous to synonymous ratio (dN/dS) and two hierarchical branch-site likelihood ratio tests were applied to identify positively selected genes (PSGs). Firstly, species-specific PSGs were identified using flathead as the foreground species and *Scophthalmus maximus*, *C. semilaevis*, *Gasterosteus aculeatus*, *Takifugu rubripes*, *Labrus bergylta*, *Tetraodon nigroviridis*, *O. latipes*, *X. maculatus*, and *Gambusia affinis* as the background species. Secondly, three metamorphosed species (i.e., flathead, *Scophthalmus maximus*, and *C. semilaevis*) were selected as foreground species, and the other above-mentioned species were used as background species. For functional enrichment analyses, GO and KEGG categories were assigned to the orthologous groups according to the flathead genome reference. In *Platycephalus* sp.1, a total of 438 expanded gene families (including 1 173 genes) and 1 219 contracted gene families (including 85 genes) were detected compared to its most recent common ancestor (Supplementary Figure S2). There were 207 GO terms identified for the expanded genes, 126 of which were significantly enriched ( $P < 0.05$ ) (Supplementary Table S8), including cell-cell adhesion (GO: 0098609, 28 genes,  $P = 2.62E-20$ ), ATPase activity (GO: 0016887, 32 genes,  $P = 1.71E-14$ ), homophilic cell adhesion (GO: 0007156, 27 genes,  $P = 6.67E-13$ ), and serine-type endopeptidase activity (GO: 0004252, 22 genes,  $P = 2.26E-09$ ). Notably, three shared GO terms, i.e., ion channel activity (GO: 0005216, 34 genes,  $P = 0.01113$ ), DNA replication (GO: 0006260, eight genes,  $P = 0.02557$ ), and DNA metabolic process (GO: 0006259, 20 genes,  $P = 0.1373$ ), were detected between the flathead and flatfish species (*Paralichthys olivaceus* and *C. semilaevis*), and are known to play crucial roles in morphological and functional modifications during flatfish metamorphosis (Shao et al., 2017). These findings suggest that the expanded genes may be associated with metamorphosis and demersal adaptation in flatheads. The

expanded genes were assigned to 27 KEGG pathways, 13 of which were significantly enriched ( $P < 0.05$ ) (Supplementary Table S9). Enrichment analyses suggested that the expanded genes were highly involved in the immune, biosynthetic, and metabolic-related pathways. The contracted gene families were significantly enriched in 26 KEGG pathways (Supplementary Table S10), which were highly represented in processes related to immunity and metabolism. The annotation of expanded and contracted genes indicated that *Platycephalus* sp.1 may have an improved capacity for resilience and adaptation to the demersal environment.

For PSG identification, a total of 1 325 genes were identified in the flathead genome. There were 129 GO terms identified from the PSGs, 16 of which were significantly enriched ( $P < 0.05$ ) (Supplementary Table S11). Additionally, 377 PSGs were identified in the three metamorphosed species (i.e., flathead, *Scophthalmus maximus* and *C. semilaevis*) as foreground species (Supplementary Table S12). These PSGs were functionally enriched in RNA methylation (GO: 0001510, seven genes,  $P = 0.00376$ ), methyltransferase activity (GO: 0008168, 18 genes,  $P = 0.02356$ ), and RNA processing (GO: 0006396, 26 genes,  $P = 0.02604$ ) (Supplementary Table S13). Similarly, these detected PSGs may also play crucial roles in demersal adaptive evolution in flatheads. However, further investigations are needed to determine the functions associated with the genes in these expanded, contracted, and positively selected gene families, such as their putative roles in adaptive evolutionary processes.

In the present study, we assembled a high-quality chromosome-scale genome of *Platycephalus* sp.1, the first reference genome of flatheads in the family Platycephalidae. This study provides valuable genomic data for further studies on demersal adaptation in flathead species and the molecular mechanisms underlying their biological characteristics, such as camouflage, metamorphosis, and sex reversal.

#### DATA AVAILABILITY

Raw sequencing data for the *Platycephalus* sp.1 genome have been deposited in the NCBI Sequence Read Archive (SRA) database under accession Nos. SRP269051, SRP278978 and SRP281984 and were deposited in the Genome Sequence Archive (GSA) database under Accession No. CRA004842.

#### SUPPLEMENTARY DATA

Supplementary data to this article can be found online.

#### COMPETING INTERESTS

The authors declare that they have no competing interests.

#### AUTHORS' CONTRIBUTIONS

T.X.G. and Z.Q.H. designed the study. S.Y.X., Z.C.L., S.S.C., and T.X.G. collected the samples and extracted genomic DNA. S.Y.X., B.B.C., and D.D.H. assembled and annotated the genome and analyzed the data. S.Y.X. and Z.Q.H. wrote the draft manuscript. T.X.G., Z.Q.H., and S.Y.X. revised the manuscript. All authors read and approved the final version of

the manuscript.

Sheng-Yong Xu<sup>1</sup>, Zhi-Chuang Lu<sup>2</sup>, Bin-Bin Cao<sup>3</sup>, Dong-Dong Han<sup>3</sup>, Shan-Shan Cai<sup>1</sup>, Zhi-Qiang Han<sup>1,\*</sup>, Tian-Xiang Gao<sup>1,†</sup>

<sup>1</sup> Fishery College, Zhejiang Ocean University, Zhoushan, Zhejiang 316022, China

<sup>2</sup> Dalian Key Laboratory of Conservation Biology for Endangered Marine Mammals, Liaoning Ocean and Fisheries Science Research Institute, Dalian, Liaoning 116023, China

<sup>3</sup> Novogene Bioinformatics Institute, Beijing 100083, China

\*Corresponding authors, E-mail: d6339124@163.com; gaotianxiang0611@163.com

## REFERENCES

- Bao WD, Kojima KK, Kohany O. 2015. Repbase Update, a database of repetitive elements in eukaryotic genomes. *Mobile DNA*, **6**: 11.
- Benson G. 1999. Tandem repeats finder: a program to analyze DNA sequences. *Nucleic Acids Research*, **27**(2): 573–580.
- Blanco E, Parra G, Guigó R. 2007. Using geneid to identify genes. *Current Protocols in Bioinformatics*, **18**(1): 4.3.1–4.3.28.
- Burge C, Karlin S. 1997. Prediction of complete gene structures in human genomic DNA. *Journal of Molecular Biology*, **268**(1): 78–94.
- Burton JN, Adey A, Patwardhan RP, Qiu RL, Kitzman JO, Shendure J. 2013. Chromosome-scale scaffolding of *de novo* genome assemblies based on chromatin interactions. *Nature Biotechnology*, **31**(12): 1119–1125.
- Chen NS. 2004. Using RepeatMasker to identify repetitive elements in genomic sequences. *Current Protocols in Bioinformatics*, **5**(1): 4.10.1–4.10.14.
- Chin CS, Alexander DH, Marks P, Klammer AA, Drake J, Heiner C, et al. 2013. Nonhybrid, finished microbial genome assemblies from long-read SMRT sequencing data. *Nature Methods*, **10**(6): 563–569.
- Cui L, Cao RB, Dong YL, Gao XC, Cen JY, Lu SH. 2019. The first complete mitochondrial genome of the flathead *Cociella crocodilus* (Scorpaeniformes: Platycephalidae) and the phylogenetic relationships within Scorpaeniformes based on whole mitogenomes. *Genes*, **10**(7): 533.
- Cunningham F, Achuthan P, Akanni W, Allen J, Amode MR, Armean IM, et al. 2019. Ensembl 2019. *Nucleic Acids Research*, **47**(D1): D745–D751.
- Daub J, Eberhardt RY, Tate JG, Burge SW. 2015. Rfam: annotating families of non-coding RNA sequences. In: Picardi E. RNA Bioinformatics. New York: Humana Press, 349–363.
- De Bie T, Cristianini N, Demuth JP, Hahn MW. 2006. CAFÉ: a computational tool for the study of gene family evolution. *Bioinformatics*, **22**(10): 1269–1271.
- Doerks T, Copley RR, Schultz J, Ponting CP, Bork P. 2002. Systematic identification of novel protein domain families associated with nuclear functions. *Genome Research*, **12**(1): 47–56.
- Edgar RC. 2010. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics*, **26**(19): 2460–2461.
- Guindon S, Dufayard JF, Lefort V, Anisimova M, Hordijk W, Gascuel O. 2010. New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Systematic Biology*, **59**(3): 307–321.
- Haas BJ, Delcher AL, Mount SM, Wortman JR, Smith Jr RK, Hannick LI, et al. 2003. Improving the *Arabidopsis* genome annotation using maximal transcript alignment assemblies. *Nucleic Acids Research*, **31**(19): 5654–5666.
- Haas BJ, Salzberg SL, Zhu W, Pertea M, Allen JE, Orvis J, et al. 2008. Automated eukaryotic gene structure annotation using EvidenceModeler and the program to assemble spliced alignments. *Genome Biology*, **9**(1): R7.
- He Y, Chang Y, Bao LS, Yu MJ, Li R, Niu JJ, et al. 2019. A chromosome-level genome of black rockfish, *Sebastes schlegelii*, provides insights into the evolution of live birth. *Molecular Ecology Resources*, **19**(5): 1309–1321.
- Hubley R, Smit A. 2015. RepeatMasker Open-1.0. <http://www.repeatmasker.org/RepeatMasker.html>.
- Hyndes GA, Neira FJ, Potter IC. 1992. Reproductive biology and early life history of the marine teleost *Platycephalus speculator* Klunzinger (Platycephalidae) in a temperate Australian estuary. *Journal of Fish Biology*, **40**(6): 859–874.
- Imamura H. 2015. Taxonomic revision of the flathead fish genus *Platycephalus* Bloch, 1785 (Teleostei: Platycephalidae) from Australia, with description of a new species. *Zootaxa*, **3904**(2): 151–207.
- Korf I. 2004. Gene finding in novel genomes. *BMC Bioinformatics*, **5**: 59.
- Li H, Durbin R. 2010. Fast and accurate long-read alignment with Burrows-Wheeler transform. *Bioinformatics*, **26**(5): 589–595.
- Li L, Stoeckert Jr CJ, Roos DS. 2003. OrthoMCL: identification of ortholog groups for eukaryotic genomes. *Genome Research*, **13**(9): 2178–2189.
- Liu QH, Wang XY, Xiao YS, Zhao HX, Xu SH, Wang YF, et al. 2019. Sequencing of the black rockfish chromosomal genome provides insight into sperm storage in the female ovary. *DNA Research*, **26**(6): 453–464.
- Lowe TM, Eddy SR. 1997. tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Research*, **25**(5): 955–964.
- Majoros WH, Pertea M, Salzberg SL. 2004. TigrScan and GlimmerHMM: two open source *ab initio* eukaryotic gene-finders. *Bioinformatics*, **20**(16): 2878–2879.
- Marçais G, Kingsford C. 2011. A fast, lock-free approach for efficient parallel counting of occurrences of *k*-mers. *Bioinformatics*, **27**(6): 764–770.
- McGinnis S, Madden TL. 2004. BLAST: at the core of a powerful and diverse set of sequence analysis tools. *Nucleic Acids Research*, **32**(S2): W20–W25.
- Mulder N, Apweiler R. 2007. InterPro and InterProScan: tools for protein sequence classification and comparison. In: Bergman NH. Comparative Genomics. Totowa: Humana Press, 59–70.
- Nakabo T. 2002. Platycephalidae. In: Nakabo T. Fishes of Japan with Pictorial Keys to the Species, English Edition. Tokyo: Tokai University Press, 601–602.
- Nawrocki EP, Eddy SR. 2013. Infernal 1.1: 100-fold faster RNA homology searches. *Bioinformatics*, **29**(22): 2933–2935.
- Parra G, Bradnam K, Korf I. 2007. CEGMA: a pipeline to accurately annotate core genes in eukaryotic genomes. *Bioinformatics*, **23**(9): 1061–1067.
- Pertea M, Kim D, Pertea GM, Leek JT, Salzberg SL. 2016. Transcript-level expression analysis of RNA-seq experiments with HISAT, StringTie and Ballgown. *Nature Protocols*, **11**(9): 1650–1667.
- Price AL, Jones NC, Pevzner PA. 2005. *De novo* identification of repeat families in large genomes. *Bioinformatics*, **21**(S1): i351–358.
- Puckridge M, Andreakis N, Appleyard SA, Ward RD. 2013. Cryptic diversity in flathead fishes (Scorpaeniformes: Platycephalidae) across the Indo-West Pacific uncovered by DNA barcoding. *Molecular Ecology Resources*, **13**(1):

32–42.

- Qin Y, Song N, Zou JW, Zhang ZH, Cheng GP, Gao TX, et al. 2013. A new record of a flathead fish (Teleostei: Platycephalidae) from China based on morphological characters and DNA barcoding. *Chinese Journal of Oceanology and Limnology*, **31**(3): 617–624.
- Rao SSP, Huntley MH, Durand NC, Stamenova EK, Bochkov ID, Robinson JT, et al. 2014. A 3D map of the human genome at kilobase resolution reveals principles of chromatin looping. *Cell*, **159**: 1665–1680.
- Ruan J, Li H. 2020. Fast and accurate long-read assembly with wtdbg2. *Nature Methods*, **17**(2): 155–158.
- Servant N, Varoquaux N, Lajoie BR, Viara E, Chen CJ, Vert JP, et al. 2015. HiC-Pro: an optimized and flexible pipeline for Hi-C data processing. *Genome Biology*, **16**: 259.
- Shao CW, Bao BL, Xie ZY, Chen XY, Li B, Jia XD, et al. 2017. The genome and transcriptome of Japanese flounder provide insights into flatfish asymmetry. *Nature Genetics*, **49**(1): 119–124.
- Shinomiya A, Yamada M, Sunobe T. 2003. Mating system and protandrous sex change in the lizard flathead, *Inegocia japonica* (Platycephalidae). *Ichthyological Research*, **50**(4): 383–386.
- Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. 2015. BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics*, **31**(19): 3210–3212.
- Smith WL, Everman E, Richardson C. 2018. Phylogeny and taxonomy of flatheads, scorpionfishes, sea robins, and stonefishes (Percomorpha: Scorpaeniformes) and the evolution of the lachrymal saber. *Copeia*, **106**(1): 94–119.
- Stanke M, Waack S. 2003. Gene prediction with a hidden Markov model and a new intron submodel. *Bioinformatics*, **19**(S2): ii215–ii225.
- Xu Z, Wang H. 2007. LTR\_FINDER: an efficient tool for the prediction of full-length LTR retrotransposons. *Nucleic Acids Research*, **35**(S2): W265–W268.
- Yang ZH. 2007. PAML 4: phylogenetic analysis by maximum likelihood. *Molecular Biology and Evolution*, **24**(8): 1586–1591.