

Tracheal bronchus: A rare unforeseen anaesthetic challenge

Address for correspondence:

Dr. Mita Eunice Sarkar,
Department of
Anaesthesiology,
Bangalore Baptist Hospital,
Hebbal, Bellary Road,
Bangaluru - 560 024,
Karnataka, India.
E-mail: mitasarkar7@yahoo.
com

Mita Eunice Sarkar, Alfred Inbaraj¹, Varghese Zachariah, Sharmishta Shukla

Departments of Anaesthesiology and ¹Radiology, Bangalore Baptist Hospital, Bengaluru, Karnataka, India

ABSTRACT

The most common congenital central airway anomaly, a tracheal bronchus is of concern during airway management especially if previously undetected. If present, it can lead to inadequate ventilation both during intubation with a normal endotracheal tube and an attempted double lumen tube insertion for one lung ventilation. Meticulous preoperative assessment and use of adjuncts like bronchoscopy before lung isolation can safely and successfully assist the anaesthetic management of such cases.

Key words: Endotracheal tube, one lung ventilation, tracheal bronchus

Access this article online

Website: www.ijaweb.org

DOI: 10.4103/ija.IJA_180_18

Quick response code



INTRODUCTION

Tracheal bronchus, a rare anatomical variant of an accessory bronchus occurring due to abnormal embryogenesis more commonly on the right side, supplies a part or the entire right upper lobe.^[1] Commonly asymptomatic, a tracheal bronchus is of significance to the anaesthesiologist. We describe the anaesthetic management of a patient with an incidental chest X-ray finding of tracheal bronchus and enumerate the challenges of administering general anaesthesia and managing one-lung ventilation in such patients.

CASE REPORT

A 4 year old, 16 kg, 105 cm female patient presented with swelling on the left side of the neck since 3 years, progressively increasing in size. There was no history of fever, cough, loss of weight or appetite. She did not complain of difficulty or pain on swallowing, voice change or difficulty in breathing. Neuropsychomotor development was normal.

There was no significant past or family history.

Her heart rate was 98/min, blood pressure 82/50 mmHg and mallampati grading (MPG) II. ENT and systemic examination were normal. The American Society of Anesthesiologists physical status was I.

The left-sided neck swelling measured 5 cm × 4 cm. It was soft, cystic, mobile, non-tender and non-pulsatile. There was no bruit or lymphadenopathy. The left carotid artery was palpable. There was no airway compromise or intrathoracic extension clinically. No other swellings were seen elsewhere.

Complete blood count was normal. Chest X-ray showed no intrathoracic extension but revealed the presence of a tracheal bronchus arising about one cm above the carina [Figure 1]. Ultrasonography neck suggested a lymphangioma.

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

How to cite this article: Sarkar ME, Inbaraj A, Zachariah V, Shukla S. Tracheal bronchus: A rare unforeseen anaesthetic challenge. *Indian J Anaesth* 2018;62:621-4.



Figure 1: Chest xray showing type II tracheal bronchus indicated by arrow a and the right upper lobe bronchus by arrow b

After a written, informed anaesthetic consent from the parents and following preoperative fasting guidelines, the patient was premedicated with syrup Triclofos. In the operating room intravenous (IV) access (22G, Vasofix, BBraun) was secured on the right hand. Pulse oximetry, non invasive blood pressure, three lead electrocardiogram, temperature and anaesthetic agent analyser monitoring were used. Injection fentanyl 2 µg/kg, sevoflurane with air and oxygen 70% were used for induction. With assisted ventilation, injection Atracurium 0.5 mg/kg IV was given. The trachea was intubated with a 5.0 mm internal diameter uncuffed endotracheal tube (ETT) fixed at 14 cm at the lip. Bilateral breath sounds were confirmed.

Fibre optic bronchoscopy done after intubation confirmed a tracheal bronchus in close proximity to the carina. The tip of the ETT was neither in the tracheal bronchus nor obstructing it.

Anaesthesia was maintained with oxygen (60%), nitrous oxide and isoflurane with GE Systems, Aespire™ workstation. Incremental doses of morphine up to 0.1 mg/kg and paracetamol suppositories provided analgesia. Breath sounds were monitored intraoperatively. The cyst extending from level II to V was separated from the surrounding structures and excised. After an uneventful intraoperative period, residual neuromuscular blockade was reversed, and the trachea was extubated when the child was awake. Post-operative chest X-ray did not reveal atelectasis.

DISCUSSION

Sandifort in 1785 described an anomaly wherein the right upper lobe bronchus originated from the

trachea and termed it as a tracheal bronchus.^[2] Recent literature defines it as one originating from the trachea, usually within 2 cm, and up to 6 cm from the carina.^[3] A variant, the pig bronchus has the entire upper lobe displaced on the trachea.^[4] Right upper lobe bronchial anomalies are seven times more common than left-sided ones,^[2] with a prevalence of 0.1–2% on bronchographic examinations.^[1]

Tracheal bronchus are classified as displaced and supernumerary, the displaced variant being more common.^[1] Anomalous bronchi are usually asymptomatic, hence picked up as incidental radiological findings.^[5] Occasionally, they present with recurrent respiratory infections including pneumonia and bronchiectasis, haemoptysis, cough, stridor and acute respiratory distress.^[3,5]

The initial radiological investigation, usually a chest X-ray shows a bronchus directly arising from the tracheal wall more commonly on the right side as was seen in our case [Figure 1].^[6] To depict exact anatomy, computed tomography is the imaging modality of choice.^[6] Bronchoscopy and tracheo bronchography have been used for a definitive view.^[7] We performed a diagnostic fibre optic bronchoscopy post-intubation instead of a CT chest preoperatively owing to the cost factor and the clarity of the tracheal bronchus on chest X-ray.

Based on origin, there are three types of tracheal bronchus of significance to the anaesthesiologist. Type I has the carina arising at mid tracheal level, manifesting most severely among the three, usually in childhood with a wheeze or stridor, misdiagnosed and managed as asthma.^[7] There may be associated congenital anomalies including cyanotic heart disease.^[8] A type II tracheal bronchus arises about 1 cm above the carina. Carinal trifurcation or type III arises from the carina [Figure 2].^[7]

Intubating a type I anomaly will obstruct ventilation to the rest of the lung causing hypoxia while intubating the true trachea will bypass the upper lobe causing atelectasis.^[9,10] The usage of short tubes to prevent ETT entry into the tracheal bronchus and fibre optic confirmation of the position of the tip of the ETT post-intubation are recommendations for non-thoracic surgeries.^[10,11]

Using a standard length ETT in a type II anomaly can cause a hypoxic shunt by obstructing the tracheal



Figure 2: A schematic overview of types of tracheal bronchus of importance to the anaesthesiologist

bronchus since such offsets have been reported to be situated as much as 6 cm from the carina.^[3,7] If desaturation occurs following intubation in such anomalies, the ETT should be carefully withdrawn, monitoring for improvement in oxygen saturation while auscultating the apical zone of the lung.^[7]

Both type II and III anomalies pose problems during one-lung ventilation, especially of the right side.^[7] For right lung isolation, the opening for the right upper lobe on a right-sided double lumen tube does not correspond with the tracheal bronchus; hence, a left double lumen tube should be used. A bronchial blocker in the right main bronchus and a Fogarty catheter in the tracheal bronchus can also be considered for right lung isolation. For left lung isolation, a bronchial blocker or a left double lumen tube can be used.^[12]

Yoshimura *et al.* reported multiple unsuccessful attempts at positioning a left-sided double lumen tube to achieve right lung isolation in a previously undetected tracheal bronchus with an apically retracted left main bronchus. They managed the case with two lung ventilation using small tidal volumes and retrospectively suggested using a selective bronchial blocker or a single-lumen ETT in the left main bronchus for such patients.^[13] Apart from the planned surgical procedure, the exact location of a tracheal bronchus and the presence of intralobar micro airway communications are important to achieve successful lung isolation.^[14]

Although not practiced routinely, bronchoscopy before insertion of any lung separator can prevent inadvertent accidents in patients with undetected anomalies.^[7] Failure to identify the origin of the right

upper lobe bronchus about 1–3 cm distal to the carina on bronchoscopy should alert the anaesthesiologist to the possibility of a tracheal bronchus.^[15] Bronchoscopy can also confirm the position of the tracheal bronchus in relation to the ETT cuff.

In conclusion, recognition of a tracheal bronchus before anaesthesia can guide airway management. Although rare, preoperative chest radiographs should be observed carefully for the presence of a tracheal bronchus. The differential diagnosis of desaturation occurring post-intubation should include an undiagnosed tracheal bronchus. Furthermore, bronchoscopy before lung isolation can detect anatomical variants and guide the use of a suitable lung separator.

Declaration of patient consent

The authors certify that they have obtained all appropriate parent consent forms. In the form the parent(s) has/have given his/her/their consent for his/her/their child's images and other clinical information to be reported in the journal. The parent(s) understand that their child's names and initials will not be published and due efforts will be made to conceal their child's identity, but anonymity cannot be guaranteed.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

1. Ghaye B, Szapiro D, Fanchamps JM, Dondelinger RF. Congenital bronchial abnormalities revisited. *Radiographics* 2001;21:105-19.
2. Kubik S, Müntener M. Bronchus abnormalities: tracheal, eparterial and pre-eparterial. *Rofo* 1971;114:145-63.
3. Siegel MJ, Shackelford GD, Francis RS, McAlister WH. Tracheal bronchus. *Radiology* 1979;130:353-5.
4. Lemoine JM, Gagnon A. Main modes of division and anatomical abnormalities of the trachea and bronchi. *Bronches* 1952;2:409-21.
5. Rémy J, Smith M, Marache P, Nuyts JP. Pathogenetic left tracheal bronchus. A review of the literature in connection with four cases (author's transl). *J Radiol Electrol Med Nucl* 1977;58:621-30.
6. Manjunatha YC, Gupta AK. Tracheal bronchus (pig bronchus). *Indian J Pediatr* 2010;77:1037-8.
7. Conacher ID. Implications of a tracheal bronchus for adult anaesthetic practice. *Br J Anaesth* 2000;85:317-20.
8. McLaughlin FJ, Strieder DJ, Harris GB, Vawter GP, Eraklis AJ. Tracheal bronchus: Association with respiratory morbidity in childhood. *J Pediatr* 1985;106:751-5.
9. Pribble CG, Dean JM. An unusual cause of intraoperative hypoxemia. *J Clin Anesth* 1994;6:247-9.
10. Vredevoe LA, Brechner T, Moy P. Obstruction of anomalous

- tracheal bronchus with endotracheal intubation. *Anesthesiology* 1981;55:581-3.
11. Ikeno S, Mitsuhashi H, Saito K, Hirabayashi Y, Akazawa S, Kasuda H, *et al.* Airway management for patients with a tracheal bronchus. *Br J Anaesth* 1996;76:573-5.
 12. Lee HL, Ho AC, Cheng RK, Shyr MH. Successful one-lung ventilation in a patient with aberrant tracheal bronchus. *Anesth Analg* 2002;95:492-3.
 13. Yoshimura T, Ueda KI, Kakinuma A, Nakata Y. Difficulty in placement of a left-sided double-lumen tube due to aberrant tracheobronchial anatomy. *J Clin Anesth* 2013;25:413-6.
 14. Kawagoe I, Kohchiyama T, Hayashida M, Satoh D, Suzuki K, Inada E, *et al.* Successful one-lung ventilation with a right-sided double-lumen tube in a patient with a right upper tracheal bronchus, who underwent left pneumonectomy for left hilar lung cancer. *Masui* 2016;65:594-8.
 15. Peragallo RA, Swenson JD. Congenital tracheal bronchus: The inability to isolate the right lung with a univent bronchial blocker tube. *Anesth Analg* 2000;91:300-1.

Author Help: Online submission of the manuscripts

Articles can be submitted online from <http://www.journalonweb.com>. For online submission, the articles should be prepared in two files (first page file and article file). Images should be submitted separately.

1) **First Page File:**

Prepare the title page, covering letter, acknowledgement etc. using a word processor program. All information related to your identity should be included here. Use text/rtf/doc/pdf files. Do not zip the files.

2) **Article File:**

The main text of the article, beginning with the Abstract to References (including tables) should be in this file. Do not include any information (such as acknowledgement, your names in page headers etc.) in this file. Use text/rtf/doc/pdf files. Do not zip the files. Limit the file size to 1 MB. Do not incorporate images in the file. If file size is large, graphs can be submitted separately as images, without their being incorporated in the article file. This will reduce the size of the file.

3) **Images:**

Submit good quality color images. Each image should be less than 4096 kb (4 MB) in size. The size of the image can be reduced by decreasing the actual height and width of the images (keep up to about 6 inches and up to about 1800 x 1200 pixels). JPEG is the most suitable file format. The image quality should be good enough to judge the scientific value of the image. For the purpose of printing, always retain a good quality, high resolution image. This high resolution image should be sent to the editorial office at the time of sending a revised article.

4) **Legends:**

Legends for the figures/images should be included at the end of the article file.