



ORAL PRESENTATION

Open Access

# Assessment of suspicious oral lesions using optical coherence tomography

Zaid Hamdoon\*, Waseem Jerjes, Gordon McKenzie, Amrita Jay, Colin Hopper

From 2nd Scientific Meeting of the Head and Neck Optical Diagnostics Society  
San Francisco, CA, USA. 23-24 January 2010

## Introduction/aims

Optical biopsy systems have been investigated for various clinical applications; however the main interest remains in the diagnosis and monitoring of premalignant and malignant conditions. In this study, we compared findings of optical coherence tomography (OCT) with histopathology results of suspicious oral lesions to assess the feasibility of OCT in identifying pathological tissue.

## Material/methods

Suspicious oral lesions acquired from 120 patients were subjected to immediate ex-vivo Swept-Source Frequency-Domain OCT. Five OCT parameters were assessed (keratin, epithelial, sub-epithelial layers changes, basement membrane and microanatomical structures). Two clinicians and two pathologists, who were blind to clinical and histopathological diagnosis, examined the OCT images autonomously, provided deferential diagnosis, the most probable diagnosis and provided judgment on the need for surgical biopsy. Inter, Intra-observer differences, sensitivity and specificity was calculated.

## Results/statistics

Basic microanatomical tissue structures were identified on the mainstream of the OCT images. Recognition of the basement membrane was achieved in the majority of the lesions. Identification of changes in the parameters ruled areas of architectural changes. There was a high inter and intra-observer agreement among the two clinicians and two pathologists, who recommended a surgical biopsy when examined all the histologically proven dysplasia and cancer OCT images. Sensitivity and specificity were calculated and proved to be encouraging.

## Conclusions/clinical relevance

At this phase, OCT can definitely aide clinical examination and monitoring and can be invaluable tool for inexperienced clinicians.

## Acknowledgment

We would like to thank the BAOMS for supporting this research project.

Published: 29 October 2010

doi:10.1186/1758-3284-2-S1-O6

Cite this article as: Hamdoon *et al.*: Assessment of suspicious oral lesions using optical coherence tomography. *Head & Neck Oncology* 2010 **2**(Suppl 1):O6.

## Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)



UCL Department of Surgery, University College London Medical School, London, UK