

Hip and Elbow Surgery in Major League Baseball Pitchers: Letter to the Editor

Dear Editor:

We read with interest the article by Kantrowitz et al² on athletic hip injuries in Major League Baseball pitchers associated with ulnar collateral ligament tears. Although overlooked in their report, their data support the observation previously published demonstrating a 75% prevalence of ulnar collateral ligament surgery among Major League Baseball pitchers undergoing hip arthroscopy.¹

J.W. Thomas Byrd, MD
Nashville, Tennessee, USA

Address correspondence to J.W. Thomas Byrd, MD (email: info@nashvillehip.org).

One or more of the authors has declared the following potential conflict of interest or source of funding: J.W.T.B. receives royalties from Smith & Nephew and Springer, receives research support from Smith & Nephew, and is a paid consultant for Smith & Nephew. AOSSM checks author disclosures against the Open Payments Database (OPD). AOSSM has not conducted an independent investigation on the OPD and disclaims any liability or responsibility relating thereto.

REFERENCES

1. Byrd JWT, Jones KS. Hip arthroscopy in high-level baseball players. *Arthroscopy*. 2015;31(8):1507-1510.
2. Kantrowitz DE, Trofa DP, Ahmad CS, Lynch TS. Athletic hip injuries in Major League Baseball pitchers associated with ulnar collateral ligament tears. *Orthop J Sports Med*. 2018;6(10):2325967118800704.