



Epidemiologic Trends and Seasonality of Scabies in South Korea, 2010-2017

Jong-Hun Kim, Hae-Kwan Cheong*

Department of Social and Preventive Medicine, Sungkyunkwan University School of Medicine, Suwon 16419, Korea

Abstract: Scabies is a parasitic skin infection with intense itching. Scabies infection seriously impairs quality of life, while outbreaks in medical institutions cause financial losses. This study aimed to present the annual and seasonal trend of prevalence of scabies in the national population. Scabies cases were extracted from National Health Insurance Service database and its epidemiologic characteristics were assessed. To analyze the seasonality of scabies occurrence, temperature and humidity were included in the model as weather factors, and the per capita gross national income index was adjusted. The annual prevalence by age group was 0.56-0.69 per 1,000 persons until the age of 40 years and peaked at 3.0-4.1 per 1,000 persons in the age group over 80 years. The number of women diagnosed with scabies has been consistently higher compared to that of men since 2010. Mean number of cases diagnosed as scabies was lowest in spring, approximately 4,000 cases, when the average temperature was less than 5°C at 2 months prior, whereas more than 6,000 scabies cases occurred in autumn when temperatures exceeded 25°C at 2 months prior. This study presents the epidemiological characteristics and seasonality of all cases nationwide over 8 years and will help to establish control policies.

Key words: Scabies, seasonality, prevalence, trend, elderly population, climate

INTRODUCTION

Scabies is an infectious skin disease caused by *Sarcoptes scabiei* var. *hominis* which is accompanied by intense itching [1]. Scabies infections can severely impair patients' quality of life due to annoyance from itching and sleep disturbances. It also incurs significant financial losses for medical institutions due to ongoing infection control costs and negative publicity throughout a scabies outbreak [2,3]. In many countries, scabies has been neglected because it is not a notifiable infectious disease [4], which is the same in South Korea. It has been difficult to rapidly recognize the occurrence of scabies outbreaks. For this reason, the World Health Organization added scabies to the list of neglected tropical diseases in 2017 [5]. Scabies infections appear sporadically, making it difficult to determine the magnitude and seasonality of an outbreak in a population [4,6]. The prevalence of scabies at the national level varies greatly depending on the overcrowding of residential environ-

ments and the socioeconomic level including poverty rate [7]. In low-income countries in the tropical regions, prevalence is higher than 10%, while in developed countries, it is much lower [8,9]. Scabies is less prevalent in the temperate regions and more common in the tropical regions [9,10]. However, studies conducted in mid-latitude countries reported that scabies is more likely to occur in winter than in summer, resulting in an opposite dose-response relationship between temperature and scabies [4,11].

In developed countries, research on seasonality related to onset and reports on epidemiological characteristics such as the prevalence and tendencies of scabies in an entire population are lacking [4]. In this study, we assessed seasonal trends in relation to climate factors and estimated prevalence using scabies occurrence data in the national population.

MATERIALS AND METHODS

Extraction of scabies cases

South Korea has the National Health Insurance Service (NHIS) system that provides single-payer public health insurance for the entire population. It is run by a fee-for-service system and its payment is approved by the Health Insurance Review and Assessment (HIRA) service which reviews all claims.

•Received 28 February 2019, revised 26 May 2019, accepted 7 June 2019.

*Corresponding author (hkcheong@skku.edu)

© 2019, Korean Society for Parasitology and Tropical Medicine

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Using NHIS and HIRA data, research information can be obtained about disease diagnoses for the entire population [12]. In this study, data on subjects who were diagnosed with scabies (International Statistical Classification of Disease and Related Health Problems, Tenth Revision, code B86) were extracted from the main database of 2010 to 2017 and were used to estimate the prevalence by sex and age. Age was categorized into groups based on 10 year unit. The annual prevalence of scabies was calculated as the number of cases diagnosed per 1,000 by age group. The details of the use of medical institutions by patients diagnosed with scabies were divided into hospital and clinic. This study was approved by the Institutional Review Board of Sungkyunkwan University for its compliance with the Privacy Protection Act (SKKU 2018-01-008).

Meteorological factors

We developed a single model of total scabies cases in South Korea, meteorological factors were also included to represent the exposure level of the entire population. The national representative estimates of meteorological factors in South Korea were calculated based on data from 45 weather stations [13]. As the national average value, the Korea Meteorological Administration (KMA) has been using the average value of 45 weather stations that have been conducted stable and consecutive observations since 1973. In this study, monthly average temperature and monthly relative humidity in 2010-2017 were extracted from the above data.

Statistical analysis

Descriptive features of the prevalence stratified by demographic characteristics were presented. Non-linear regression analysis with a generalized additive model was used to determine the dose-response relationship between the monthly mean temperature and the number of monthly scabies cases. In this model, the number of monthly scabies cases was used as an independent variable, while the total annual population, average temperature, and relative humidity were used as dependent variables. To adjust for the decreasing trend of the overall number of scabies cases over time, we used the per capita gross national income (GNI) index, a socio-economic growth indicator at the national level. The per capita GNI data in South Korea were obtained from a Korean Statistical Information Service [14] as

$$Cases_{ij} = \beta_0 + \beta_1 \cdot pop_j + \beta_2 \cdot AT_{(i-2)j} + \beta_3 \cdot RH_{(i-2)j} + \beta_4 \cdot GNI_j \quad (\text{Equation 1})$$

where $Cases_{ij}$, Pop_j , and GNI_j represent the number of scabies cases in i month of j year, the total population in j year, and per capita GNI in j year, respectively. $AT_{(i-2)j}$ and $RH_{(i-2)j}$ represent the national average temperature and relative humidity in $i-2$ month of j year, respectively.

The time difference between the number of scabies cases variable and the meteorological factors variable considers the delay between the initial infection and the final diagnosis. After a review of the existing literature related to the delay to diagnosis, the time of 2 months was applied in this study.

RESULTS

According to the HIRA service data, a total of 51,331 patients in 2010 and 42,436 patients in 2017 were diagnosed with scabies (Table 1). Until 2015, the number of cases was steadily decreasing, while the number of patients has rebounded slightly since 2016. The ratio of females to males diagnosed with scabies was 0.51:0.49 in 2010, but the gap between females and males continued to increase. In 2017, the ratio of females to males was 0.57:0.43, 1.33 times higher in females than in males. The number of scabies cases diagnosed in clinics has decreased by more than 10,000 (from 43,903 to 32,510) between 2010 and 2017. However, the number of scabies cases diagnosed at hospitals fluctuated without a certain tendency, but the proportion of cases diagnosed at hospitals among all cases steadily increased from 14% in 2010 to 23% in 2017. The proportion of cases by age group in the total number of scabies cases is consistently decreasing in the population of children, adolescents, and adults under 50 years of age. However, in the population over 50 years, the proportion of cases steadily increased over time. In 2017, in terms of the number of cases, the highest frequency occurred in the 50-59 years age group, followed by the 60- and 40-year age groups.

The prevalence of scabies per 1,000 persons was similar in children, adolescents, and adults under 50 years of age, ranging from 0.56 to 1.06, but the prevalence increased from age 50 to older age (Table 2). In particular, the prevalence of 2.88-3.86 per 1,000 persons in the group over 80 years old was 4-5 times higher than that of the younger age group. From 2010 to 2017, the overall prevalence decreased in all age groups, and the prevalence of scabies in the entire population decreased from 1.03 per 1,000 persons in 2010 to 0.83 per 1,000 persons in 2017.

The occurrence of scabies in South Korea for the past 8 years

Table 1. Demographic characteristics of scabies cases in South Korea, 2010-2017

Characteristic	2010	2011	2012	2013	2014	2015	2016	2017
No. of cases	51,331	52,646	50,284	45,915	45,046	40,389	41,555	42,436
Sex								
Female	50.6	50.3	52.4	53.3	53.0	54.7	55.6	57.1
Male	49.4	49.7	47.6	46.7	47.0	45.3	44.4	42.9
Medical institution								
Clinic	85.5	84.0	81.5	79.8	79.9	79.3	78.3	76.6
Hospital	14.5	16.0	18.5	20.2	20.1	20.7	21.7	23.4
Age group (yr)								
0-9	7.7	7.6	6.8	7.7	7.2	6.4	6.4	6.1
10-19	11.6	11.6	10.8	11.2	10.9	10.3	8.0	7.1
20-29	14.2	13.6	12.5	11.2	11.2	10.6	10.4	9.9
30-39	14.0	13.7	12.3	12.1	11.5	11.0	10.6	10.2
40-49	16.4	15.7	15.1	14.9	14.8	14.2	13.7	14.0
50-59	15.3	15.6	16.7	16.7	17.1	17.8	19.1	19.2
60-69	7.7	8.4	9.2	9.5	10.2	11.5	12.9	14.5
70-79	6.7	7.1	8.5	8.4	8.4	8.5	9.1	8.8
80+	6.8	7.1	8.5	8.6	9.1	10.1	10.1	10.4

*Relative frequency, %.

Table 2. Prevalence of scabies per 1,000 persons in South Korea, 2010-2017

Age group (yr)	2010	2011	2012	2013	2014	2015	2016	2017
0-9	0.81	0.85	0.73	0.76	0.70	0.56	0.58	0.58
10-19	0.88	0.91	0.83	0.81	0.81	0.68	0.59	0.56
20-29	1.05	1.06	0.95	0.78	0.76	0.65	0.65	0.63
30-39	0.86	0.87	0.76	0.69	0.66	0.57	0.59	0.58
40-49	0.97	0.95	0.87	0.78	0.75	0.65	0.65	0.69
50-59	1.17	1.15	1.11	0.99	0.96	0.90	0.96	0.98
60-69	0.96	1.07	1.10	1.01	1.02	1.02	1.04	1.13
70-79	1.36	1.41	1.51	1.28	1.23	1.11	1.19	1.15
80+	3.86	3.84	4.13	3.54	3.41	3.37	2.94	2.88
Total	1.03	1.05	1.00	0.91	0.89	0.80	0.82	0.83




Fig. 1. No. of monthly scabies cases and meteorological factors in South Korea, 2010-2017.




Fig. 2. Effect of monthly mean temperature on scabies infection in South Korea using generalized additive model.

showed a remarkable seasonality (Fig. 1). In October, approximately 6,800 people were diagnosed with scabies, the most cases were reported throughout the year. In April, approximately 3,700 people were diagnosed, the fewest cases were observed during the year. We examined the dose-response relationship between the monthly mean temperature and the monthly scabies cases when considering 2 months as the time delay effect from the initial infection to the diagnosis. Approximately 4,000 cases occurred at an average temperature of less than 5°C in the previous 2 months (Fig. 2). As the temperature increased, the number of scabies cases gradually increased. As a result, more than 6,000 cases occurred when the average temperature was 25°C or higher at 2 months prior.

DISCUSSION

This nationwide study demonstrated the decreasing annual trend of scabies in Korea. Its prevalence was highest in the elderly population while seasonality of scabies was closely associated with climate factors.

The pattern of the age-specific prevalence of scabies in low-income countries peaks during infancy and continues to decline until adulthood [8,10]. However, in developed countries in North America and Western Europe, the prevalence was evenly low in all age groups [10,11]. In the United Kingdom (UK), the average prevalence from 1997 to 2005 was approximately 2.5 per 1,000 persons and the overall female-to-male ratio was 1.24 [11]. The prevalence of scabies by age in South Korea has an opposite pattern to that in low-income countries. In South Korea, unlike other developed countries, the prevalence of scabies increases with age, presenting the highest prevalence in elderly people over 80 years of age. The female-to-male ratio for scabies cases in South Korea was 1.02 in 2010.

However, since then, it has increased continuously and was 1.33 in 2017. From 2010 to 2017, the overall female-to-male ratio was 1.14, and more cases of women were reported, similar to that observed in the UK.

The high prevalence of scabies in the elderly can be considered from 2 aspects. One aspect is the increase in nursing homes for the elderly due to population aging and due to change toward the nuclear family [15]. The Korean government started providing long-term care insurance since 2008. As a result, the number of long-term care facilities is rapidly increasing. Also, the demand for nursing hospitals is increasing since the aging population and the number of elderly people living in a nursing home increased dramatically, from 7,864 in 2001 to 1,037,973 in 2012 [16]. In the early stage, this quantitative expansion caused problems with infection control and the quality of service was not managed properly. Since 2011, the quality level of all facilities has been evaluated and the results released to the public, thus qualitative control began to take place [17]. Long-term care facilities and hospitals have been still suspected as being the main route for scabies transmission [18-20], while patient transfers between hospitals and/or facilities have also contributed to transmitting scabies [18,21].

Our result supports the above estimations. For the last 8 years, the percentage of patients diagnosed with scabies in clinics has decreased by 9%, while those diagnosed in hospitals has increased by 9%. The other aspect is the high relative poverty rate in the elderly population, which is particularly high compared to developed countries in North America and Western Europe. However, the relative poverty rate in elderly Koreans is over 40%, which is extremely high compared to the relative poverty rate of 13.8% of the total population in Korea and the average relative poverty rate of 12.6% in the Organisation for Economic Co-operation and Development (OECD) member countries in 2015 [22-24]. Poverty in the elderly is accompanied by poor hygiene, overcrowding, and malnutrition, all of which are linked with a high prevalence of scabies [11,25].

The seasonality of scabies outbreaks has been reported in the data of a few specific age groups [4,11,26]. According to the survey on the young soldier population in Israel and on the children in the UK, more scabies cases were reported in winter than in summer. The seasonality of scabies in Europe seems to contradict our findings, however. The study in Israel reported the highest outbreak from December to February,

with average temperatures around 10-15°C [4]. The report from the South Korean multicenter study showed that the highest outbreak was reported in October and November when average temperatures around 7-15°C, suggesting similar temperature ranges in both studies [18]. In our study, the number of scabies cases showed distinct seasonality from 2010 to 2017. The number of scabies cases began to increase in late summer and reached its peak in the autumn (September to November). From the winter, the number of scabies cases began to decrease again, and the lowest point was seen in the spring (March to May). This has been a very regular seasonal pattern, with a number of cases being 1.64 times higher in autumn than in spring (Fig. 1). When scabies peaked, the average temperature was similar to that of the previous studies at about 14.5°C.

Classically, the clinical presentation of scabies is typical as a pruritic skin eruption [27]. However, the clinical features may be atypically altered among institutionalized elderly patients, so pruritus may not be a predominant symptom [28]. For elderly patients residing in long-term care facilities with skin diseases as a result of poor hygiene, or as a result of communication difficulties due to cognitive impairment, the diagnosis can be delayed or errantly made as psoriasis or eczema [29]. Patients infected with scabies usually show clinical symptoms after an incubation period of 4-6 weeks [5,30]. The older the patient, the longer it takes to diagnose [18]. Because of these characteristics, it is known that scabies originating from elderly people living in long-term care facilities are the starting point of the spread [29,31-33]. Given the time delay from the initial infection to the diagnosis through the incubation period, most of the scabies cases diagnosed in autumn in South Korea were estimated to have started in summer. An analysis of the dose-response relationship between the average temperature and the number of scabies cases, the average temperature 2 months prior explained the increase in scabies cases.

This research has some limitations. First, access to raw data was only available for information on the number of diagnosed cases per month. Because previous studies suggested that it would take 4 to 6 weeks for scabies to be diagnosed, if data on the number of diagnosed cases per week could have been obtained, the dose-response curve in the number of diagnosed cases with changes in temperature could have been more elaborate. Second, the difference in temperature between regions in South Korea at the same time point is not large, but because the national representative average temperature and

average relative humidity are used, the meteorological factors may not exactly match the spatial location of reported scabies. However, we presented the age-specific prevalence of scabies that is typically neglected disease by using insurance data for the entire population in South Korea. In addition, using meteorological data, we derived a dose-response relationship between the change in average temperature and the number of diagnosed scabies and provided the information on at what point in the year scabies cases could increase.

The main cause of the occurrence of scabies was previously considered due to a poor living environment. However, despite advances in medical technology and improvements in environmental facilities, recent outbreaks in developed countries increased in elderly care facilities due to diagnostic delay, medical staff indifference, and caregiver carelessness. Apart from socio-economic factors, when considering both the incubation period and the delayed diagnosis, the rising of average temperature in the summer season prior to about 2 months could account for the increasing of scabies in the autumn season. Population aging is rapidly progressing in South Korea, so the demand for long-term care facilities such as nursing homes is expected to increase steadily in the future. The understanding in the seasonality related to the average temperature in scabies and the identification of the prevalence according to the demographic characteristics will help to establish a strategy to control scabies.

ACKNOWLEDGMENT

This study was supported by the Government-wide R&D Fund project for infectious disease research (HG18C0088), Republic of Korea.

CONFLICT OF INTEREST

The authors declare no conflict of interest related to this study.

REFERENCES

1. Chosidow O. Scabies. *N Engl J Med* 2006; 354: 1718-1727.
2. de Beer G, Miller MA, Tremblay L, Monette J. An outbreak of scabies in a long-term care facility: the role of misdiagnosis and the costs associated with control. *Infect Control Hosp Epidemiol* 2006; 27: 517-518.
3. Mounsey KE, Murray HC, King M, Oprescu F. Retrospective

- analysis of institutional scabies outbreaks from 1984 to 2013: lessons learned and moving forward. *Epidemiol Infect* 2016; 144: 2462-2471.
4. Mimouni D, Ankol OE, Davidovitch N, Gdalevich M, Zangvil E, Grotto I. Seasonality trends of scabies in a young adult population: a 20-year follow-up. *Br J Dermatol* 2003; 149: 157-159.
 5. World Health Organization. Neglected tropical diseases [Internet]. Available from: https://www.who.int/neglected_diseases/diseases/scabies/en/.
 6. Micali G, Lacarrubba F, Verzì AE, Chosidow O, Schwartz RA, Schwartz. Scabies: Advances in Noninvasive Diagnosis. *PLoS Negl Trop Dis* 2016; 10: e0004691.
 7. Heukelbach J, Feldmeier H. Scabies. *Lancet* 2006; 367: 1767-1774.
 8. Romani L, Steer AC, Whitfeld MJ, Kaldor JM. Prevalence of scabies and impetigo worldwide: a systematic review. *Lancet Infect Dis* 2015; 15: 960-967.
 9. Fuller LC. Epidemiology of scabies. *Curr Opin Infect Dis* 2013; 26: 123-126.
 10. Karimkhani C, Colombara DV, Drucker AM, Norton SA, Hay R, Engelman D, Steer A, Whitfeld M, Naghavi M, Dellavalle RP. The global burden of scabies: a cross-sectional analysis from the Global Burden of Disease Study 2015. *Lancet Infect Dis* 2017; 17: 1247-1254.
 11. Lassa S, Campbell MJ, Bennett CE. Epidemiology of scabies prevalence in the U.K. from general practice records. *Br J Dermatol* 2011; 164: 1329-1334.
 12. Cheol Seong S, Kim YY, Khang YH, Heon Park J, Kang HJ, Lee H, Do CH, Song JS, Hyon Bang J, Ha S, Lee EJ, Ae Shin S. Data resource profile: the National Health Information Database of the national health insurance service in South Korea. *Int J Epidemiol* 2017; 46: 799-800.
 13. National Climate Data Service System. Multiple meteorological station statistics [Internet]. Available from: <https://data.kma.go.kr/cmmn/main.do>.
 14. Korean Statistical Information Service. GDP and GNI by economic activities [Internet]. Available from: <http://kosis.kr/eng/index/index.do>.
 15. Chang SJ. Lived experiences of nursing home residents in Korea. *Asian Nurs Res (Korean Soc Nurs Sci)* 2013; 7: 83-90.
 16. Ministry of Health and Welfare. 2012 statistics on the elderly care facilities [Internet]. Available from: <http://www.mw.go.kr>.
 17. Long-Term Care Insurance ACT. Article 54 (Control and Evaluation of Long-Term Care Benefits) [Internet]. Available from: https://elaw.klri.re.kr/kor_mobile/viewer.do?hseq=40189&type=sogan&key=10.
 18. Park SY, Hong JS, Roh JY, Lee JY, Kim DW, Yoon TJ, Sim WY, Lee KS, Kye YC, Lee AY, Kim MB, Cho S, Moon KC, Lee ES, Yang JM, Lee WS, Lee M, Park SD, Won YH, Shin BS, Hong C, Lee J, Yoon TY, Kim KJ, Ro YS, Kim KH, Eun HC. Epidemiological and Clinical Study of Scabies in Korea: Multicenter Prospective and Retrospective Study. *Korean J Dermatol* 2013; 51: 678-684.
 19. Makigami K, Ohtaki N, Yasumura S. A 35-month prospective study on onset of scabies in a psychiatric hospital: Discussion on patient transfer and incubation period. *J Dermatol* 2012; 39: 160-163.
 20. Choi KH, Chun SW, Song JS, Ro BI, Cho HK. Characteristics of scabies patients in Goyang city and transmission route. *Korean J Dermatol* 2013; 51: 673-677 (in Korean).
 21. Ki MR, Moon HJ, Cho H. Outbreak of scabies at geriatric long-term care facilities in Korea. *Korean J Epidemiol* 2006; 28: 100-111 (in Korean).
 22. O. Thévenon. Tackling Child Poverty in Korea. OECD Social, Employment and Migration Working Papers. OECD Publishing, Paris, French. 2018, pp 15-20.
 23. Ku I, Kim CO. Decomposition analyses of the trend in poverty among older adults: the case of South Korea. *J Gerontol B Psychol Sci Soc Sci* 2018; 1-10.
 24. Yun A, Ko K. Multidimensional Elderly Poverty Index. Policy Report. Korea Institute for Health and Social Affairs Seoul, Korea. 2018, pp 25-51 (in Korean).
 25. Vorou R, Remoudaki HD, Maltezou HC. Maltezou. Nosocomial scabies. *J Hosp Infect* 2007; 65: 9-14.
 26. Mimouni D, Gdalevich M, Mimouni FB, Haviv J, Ashkenazi I. Ashkenazi. The epidemiologic trends of scabies among Israeli soldiers: a 28-year follow-up. *Int J Dermatol* 1998; 37: 586-587.
 27. Wilson MM, Philpott CD, Breer WA. Atypical presentation of scabies among nursing home residents. *J Gerontol A Biol Sci Med Sci* 2001; 56: 424-427.
 28. Degelau J. Scabies in long-term care facilities. *Infect Control Hosp Epidemiol* 1992; 13: 421-425.
 29. Tjioe M, Vissers WHPM. Scabies outbreaks in nursing homes for the elderly: recognition, treatment options and control of reinfection. *Drugs & aging* 2008; 25: 299-306.
 30. Orion E, Matz H, Wolf R. Ectoparasitic sexually transmitted diseases: scabies and pediculosis. *Clin Dermatol* 2004; 22: 513-519.
 31. Hewitt KA, Nalabanda A, Cassell JA. Cassell. Scabies outbreaks in residential care homes: factors associated with late recognition, burden and impact. A mixed methods study in England. *Epidemiol Infect* 2015; 143: 1542-1551.
 32. Makigami K, Ohtaki N, Ishii N, Tamashiro T, Yoshida S, Yasumura S. Risk factors for recurrence of scabies: a retrospective study of scabies patients in a long-term care hospital. *J Dermatol* 2011; 38: 874-879.
 33. Lay CJ, Wang CL, Chuang HY, Chen YL, Chen HL, Tsai SJ, Tsai CC. Risk factors for delayed diagnosis of scabies in hospitalized patients from long-term care facilities. *J Clin Med Res* 2011; 3: 72-77.