

Comparison of cytokine and gene activities in tissue and blood samples of patients with celiac disease

Ensieh KhalKhal¹, Zahra Razzaghi², Hakimeh Zali³, Ayad Bahadorimonfared⁴, Majid Iranshahi⁵, Mohammad Rostami-Nejad⁵

¹*Proteomics Research Center, Faculty of Paramedical Sciences, Shahid Beheshti University of Medical Sciences, Tehran, Iran*

²*Laser Application in Medical Sciences Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran.*

³*Proteomics Research Center, School of Advanced Technologies in Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran.*

⁴*Department of Health & Community Medicine, Faculty of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran*

⁵*Gastroenterology and Liver Diseases Research Center, Research Institute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, Tehran, Iran*

ABSTRACT

Aim: The aim of this study is to explore the expression of genes associated to celiac disease (CD) in the target tissue and peripheral blood monocytes (PBMC) or serum to introduce possible potential biomarkers.

Background: Celiac disease (CD) is an autoimmune disease induced by gluten ingestion in genetically predisposed individuals. Despite technological progress, small intestine biopsy is still the gold standard for diagnosis of CD.

Methods: CD data were collected from public databases (proteomics and microarray-based techniques documents). Differentially expressed genes (DEGs) in PBMC or serum as well as small intestinal biopsies from celiac patients compared to normal were collected and analyzed to introduce common individuals. Gene ontology was done to identify the involved biological terms.

Results: Among 598 CD genes in biopsies and 260 genes in PBMC or serum, 32 common genes with a similar expression pattern in both sources were identified. A total of 48 biological terms were introduced which were involved in the CD via the determined DEGs. "Cytokine activity" was the most expanded one of the biological terms.

Conclusion: In this analysis, it was concluded that 32 potential biomarkers of CD can be assessed by complementary research to introduce effective and available biomarkers in biopsy and blood.

Keywords: Celiac disease, PBMs, Gene expression, Intestine biopsy.

(Please cite as: **KhalKhal E, Razzaghi Z, Zali H, Bahadorimonfared A, Iranshahi M, Rostami-Nejad M. Comparison of cytokine and gene activities in tissue and blood samples of patients with celiac disease. Gastroenterol Hepatol Bed Bench 2019;12(Suppl.1):S108-S116.**)

Introduction

Celiac disease (CD) with a global prevalence of 1% is a chronic autoimmune disorder in genetically susceptible individuals which is triggered by wheat gluten. It is a multifactorial disease induced by genetic

(HLA-DQ2 or HLA-DQ8) and environmental factors (gluten) (1, 2). Inflammation and villous atrophy of the small intestine are caused by undesirable immunological reactions. Villous atrophy can result in malabsorption of vitamins and nutrients (3). The CD patients may experience osteoporosis, iron deficiency, and bone disease due to nutritional deficiency (4). A permanent gluten-free diet (GFD) is the only accepted therapy for CD with most individuals responding to it (5). CD is associated with some diseases including insulin-dependent diabetes mellitus (6-8), non-alcoholic

Received: 22 September 2019 *Accepted:* 3 December 2019

Reprint or Correspondence: **Mohammad Rostami-Nejad**, PhD. *Gastroenterology and Liver Diseases Research Center, Research Institute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, Tehran, Iran.*

E-mail: m.rostami@gmail.com

ORCID ID: 0000-0003-2495-1831

fatty liver disease, autoimmune hepatitis (9), hypothyroidism (6, 8, 10), Sjögren syndrome (7), Hashimoto thyroiditis, psoriasis (7, 8), inflammatory bowel diseases, asthma, arthritis, migraine, bile stones, and menstrual abnormalities (8). CD diagnosis is based on specific serological tests and the presence of a characteristic enteropathy in an intestinal biopsy (11, 12). Genetic and non-genetic factors such as epigenetics and the environmental factors cause most human diseases which share common genetic profiles. The available information for each disease constitutes a challenge in identification of disease-specific mechanisms. Hence, the combination of information from various databases and multi-source data such as proteomics, genomics, metabolomics, and microarray-based techniques can be crucial in understanding the pathophysiology and the common biological basis among different diseases. Achievement of a deeper understanding of the molecular mechanisms of diseases can be useful for detection of many unknown molecular aspects, development of more precise treatments for diseases as well as for biomarker discovery. To this end, we have integrated data to detect the key genes involved in the CD; thus, more effective and early diagnosis could be made in patients and to identify healthy at-risk individuals (13-25).

The aim of this study is to explore the common differentially expressed genes in the intestine and

peripheral blood monocytes (PBMC) or serum of CD patients to introduce readily available biomarker candidates for CD.

Methods

All data including proteomics and microarray-based documents about CD were collected from public databases and gene expression databases. CD related data were extracted from NCBI (<https://www.ncbi.nlm.nih.gov/pubmed>) which were published by March 2019. We used “Celiac Disease”, “duodenum or biopsy”, “proteomic”, “microarray or RT-PCR”, and “PBMC” as the main keywords for searching in NCBI and Google Scholar to find the genes or proteins that are up or downregulated in CD comparison to healthy control.

The “Celiac Disease”, “celiac disease + duodenum or biopsy + proteomic”, “celiac disease + PBMC + proteomic”, “celiac disease + duodenum or biopsy + microarray or RT-PCR”, “celiac disease + PBMC + microarray or RT-PCR” parent terms were used. The common DEGs between intestine biopsy and blood samples were identified. Also, the DEGs with a similar expression pattern in both samples were enriched by Reactome application of Cytoscape software for biological processes and molecular function.

Results




Figure 1. The number of common and differentially expressed genes in both of CD blood and tissue.

S110 Comparison of cytokine and gene activities in celiac disease

Table 1. Biological processes (BP) and molecular function (MF) related to the 32 common DEGs in both CD intestine and blood samples

R	Gene name	No. of genes	Biological process and molecular function	NO. of BP and MF
1	APOA1	1	apolipoprotein A-I receptor binding, beta-amyloid binding, chemorepellent activity, high-density lipoprotein particle binding, phosphatidylcholine binding, phosphatidylcholine-sterol O-acyltransferase activator activity, phospholipid transporter activity	6
2	ARG2	1	arginase activity	1
3	CXCL10	1	cAMP-dependent protein kinase regulator activity	1
4	IL2,CHI3L1	2	carbohydrate binding	1
5	PCK1	1	carboxylic acid binding, GDP binding, manganese ion binding, phosphoenolpyruvate carboxykinase (GTP) activity	2
6	CCR9,CCR3	2	C-C chemokine receptor activity	1
7	CLU	1	chaperone binding, misfolded protein binding	1
8	CXCL8,CXCL10,CXCL11	3	chemokine activity	1
9	CCR9,CCR3	2	chemokine receptor activity	1
10	CHI3L1	1	chitinase activity, chitin binding	2
11	APOC3,APOA1	2	cholesterol binding, high-density lipoprotein particle receptor binding, lipase inhibitor activity	3
12	APOB,APOA1	2	cholesterol transporter activity	1
13	CXCL10,CXCL11	2	CXCR3 chemokine receptor binding	1
14	IL21,IL12B,IL10,IL15,IL18,IL2,IL4,IL6,IFNG,IL23A,IL17A	11	cytokine activity	1
15	TNFRSF9	1	cytokine binding,	1
16	IL12B	1	cytokine receptor activity, interleukin-12 alpha subunit binding, interleukin-12 receptor binding	3
17	IL21,IL15,IL17A	3	cytokine receptor binding	1
18	STAT1	1	double-stranded DNA binding, nuclear hormone receptor binding, RNA polymerase II core promoter sequence-specific DNA binding, transcription factor activity, RNA polymerase II core promoter sequence-specific, tumor necrosis factor receptor binding	6
19	APOC3	1	enzyme regulator activity	1
20	CHI3L1	1	extracellular matrix structural constituent	1
21	APOH	1	glycoprotein binding, lipoprotein lipase activator activity	2
22	IL2	1	glycosphingolipid binding, kappa-type opioid receptor binding, kinase activator activity	3
23	RGS1	1	G-protein alpha-subunit binding	1
24	IL12B,IL10,IL2,IL4,IL6	5	growth factor activity	1
25	RGS1,TAGAP	2	GTPase activator activity	1
26	MPO,APOH,APOB,CXCL10,CXCL11	5	heparin binding	1
27	TNFAIP3,APOH,IL12B,STAT1,APOA1	5	identical protein binding	1
28	IFNG	1	interferon-gamma receptor binding	1
29	IL10	1	interleukin-10 receptor binding	1
30	IL12B,IL23A	2	interleukin-23 receptor binding	1
31	IL21,IL2	2	interleukin-2 receptor binding	1
32	IL4	1	interleukin-4 receptor binding	1
33	IL6	1	interleukin-6 receptor binding	1
34	CXCL8	1	interleukin-8 receptor binding	1
35	TNFAIP3	1	K63-linked polyubiquitin binding, kinase binding, protein self-association, thiol-dependent ubiquitin-specific protease activity, ubiquitin binding	4
36	APOB	1	lipase binding, low-density lipoprotein particle receptor binding	2
37	MPO	1	peroxidase activity	1
38	APOH,APOB,APOC3,APOA1	4	phospholipid binding	1
39	TNFAIP3,LCN2	3	protease binding	1
40	RBP4,RBP2	2	retinol binding	1
41	REL	1	RNA polymerase II distal enhancer sequence-specific DNA binding, transcriptional activator activity, RNA polymerase II distal enhancer sequence-specific binding	3
42	SH2B3	1	signaling adaptor activity	1
43	SH2B3,STAT1	2	signal transducer activity	1
44	LCN2	1	small molecule binding	1
45	RBP2,LCN2	2	transporter activity	1
46	TNFRSF14,TNFRSF9	2	tumor necrosis factor-activated receptor activity	1
47	CLU,TNFRSF14	2	ubiquitin protein ligase binding	1
48	TNFRSF14	1	virus receptor activity	1

Table 1 presents the genes that display differential expression in PBMC or serum as well as CD small intestinal biopsies, as compared with normal small intestinal biopsies and PBMC or serum. In order to integrate data, we combine the up- and downregulated genes involved in celiac patients compared to healthy controls. There are 373 upregulated and 225 downregulated genes in the intestine tissue of celiacs

versus controls. Also, 180 upregulated and 78 downregulated genes of different gene expression profiles were identified in the peripheral blood mononuclear cells (PBMCs) or serum of celiac patients versus controls (19, 26-38).

There are 49 common differentially expressed genes including 33 genes with a similar pattern of regulation and 16 ones with an opposite outline in both CD small

intestinal biopsies and blood (Figure 1). Table 2 reports different gene expression profiles in peripheral PBMCs or serum, as well as tissue of celiac patients.

ANXA3, ARG2, CCR3, CCR6, CCR9, CLU, CXCL10, CXCL11, GUCY1B3, IFNG, IL12B, IL15, IL17A, IL2, IL21, IL23A, IL4, IL6, IL8, KRT19, LCN2, MPO, OLFM4, STAT1, REL, TAGAP, TBX21, TNFAIP3, and TNFRSF9 are upregulated while APOH, LPP, and RBP are downregulated in both samples of intestine and blood. APOA1, APOB, APOC3, CHI3L1, IL18, JAM3, KIAA1109, and PCK1 are downregulated in the intestine and upregulated in blood samples. IL10, IL6, INPP4A, RGS1, SH2B3, STAT1, and TNFRSF14 are the genes that are upregulated in the intestine biopsy and downregulated in the blood samples. The biological processes and molecular functions related to the 33 common DEGs with a similar expression change in both samples are shown in Table 1.

Discussion

In the present study, we tried to integrate CD related data (differentially expressed genes). Integration and analysis of these data is useful to uncover critical molecular mechanisms of CD and facilitate diagnostic biomarker discovery. In order to mix data, we combined up and down-regulated genes involved in celiac patients as compared to healthy controls. Amongst them, 47 CD genes had altered expression in both CD small intestinal biopsies and blood. The genes with a common expression pattern in blood and intestinal tissue were thirty-two while the other 15 were expressed in an opposite way.

As mentioned earlier, APOH, APOB, APOA1, APOA4, APOC2, and APOC3 are downregulated in the CD intestinal, while only Beta-2-glycoprotein 1 (APOH) among them is downregulated in both sources. It is shown that Intestinal apolipoproteins play a critical role in lipoprotein metabolism. Some of them are

Table 2. A list of genes that are upregulated or downregulated in the small intestinal and in PBMC or serum of celiac patients

	Up in both source	Down in both source	down in biopsy up in blood	up in biopsy down blood
1	ANXA3	APOH	APOA1	IL10
2	ARG2	LPP	APOB	IL6
3	CCR3	RBP	APOC3	INPP4A
4	CCR6		CHI3L1	RBP4
5	CCR9		IL18	RGS1
6	CLU		JAM3	SH2B3
7	CXCL10		KIAA1109	STAT1
8	CXCL11		PCK1	TNFRSF14
9	GUCY1B3			
10	IFNG			
11	IL12B			
12	IL15			
13	IL17A			
14	IL2			
15	IL21			
16	IL23A			
17	IL4			
18	IL6			
19	IL8			
20	KRT19			
21	LCN2			
22	MPO			
23	OLFM4			
24	pSTAT1			
25	REL			
26	SH2B3			
27	TAGAP			
28	TBX21			
29	TNFAIP3			
30	TNFRSF9			

S112 Comparison of cytokine and gene activities in celiac disease

synthesized with chylomicrons in the mucosa and secreted into the intestinal lymph. This seems to be important for the secretion of chylomicrons and to stay with chylomicrons during their lipolysis in plasma (39). Because of villous atrophy and apolipoprotein reduction, fat and cholesterol malabsorption in CD patients is often increased causing hypocholesterolemia. Consequently, the absorption efficiency of cholesterol is impaired (40). The APOH serum protein binds to phospholipids, heparin, and dextran sulfate and prevents intracellular coagulation cascade activation by binding to phospholipids on the surface of damaged cells. It binds to the membrane of T cell, and supports its function as a marker for membrane flip flop. In addition, it is involved in inflammatory mediated process of cell death (41) as well as activation of endothelial cells and apoptotic mechanisms (42). It is a part of the group of acute phase proteins (43) whose expression is sometimes upregulated and sometimes downregulated, reflecting its pleiotropic functions such as APOH reduction in CSF during cerebral malaria (44) and rises in several infectious diseases (45). It can be one of the major elements of the first line of the innate immune response regulating homeostasis.

The other common downregulated gene in both samples is Retinol-binding protein (RBP). It is known as an adipocyte-secreted protein which transports retinol in the blood from its storage site in the liver to the epithelial tissues. RBP is a retinol transporter protein present in the small intestinal epithelium. It plays an important role in the uptake and intracellular metabolism of vitamin A. It also modulates the rate of retinoic acid supply to the endometrial cell nuclei during the menstrual cycle (46, 47). The RBP plasma concentration is regulated by vitamin A status, where in vitamin A deficiency, RBP molecules are not secreted from the liver and decline in plasma (46). Retinol is a vitamin necessary for growth, reproduction, differentiation of epithelial tissues, and vision. Malabsorption is frequently encountered in CD-patients because of villous atrophy and crypt hyperplasia in the duodenum and jejunum; thus, vitamin and mineral deficiencies e.g. vitamin A deficiency is reported in CD-patients. Vitamin A deficiency blocks secretion of the binding protein post transnationally and results in

defective delivery and supply to the epidermal cells (48).

Lipoma-preferred partner (LPP), the third downregulated gene is involved in cell motility and cell adhesion which is crucial to maintaining the barrier integrity of the small intestine. Therefore, it might play a role in the pathogenesis of CD (49)

A total of 29 of these genes have had higher expression levels in both sources compared to controls. They are directly or indirectly involved in the inflammation process, and were significantly up-regulated in the epithelial cells of CD patients relative to controls. They have a specific role in the gluten-induced immune response (see Table 1).

Both c-REL (proto-oncogene c-REL) and TNFAIP3 (Tumor necrosis factor alpha-induced protein 3) are involved in the regulation of the nuclear factor kappa β (NF- κ B) inflammatory signaling pathway in the pathology of CD, and are the key mediators in this nuclear activating complex (50). NF- κ B is a transcription complex which plays a key role in regulating the cellular immune response to stimuli. In various inflammatory disorders, including asthma, arthritis, and inflammatory bowel disease, NF- κ B is activated whose pathway may play an independent role in the innate mechanisms of disease development (51). c-REL is a member of NF- κ B transcription factor family and has high levels of expression in B and T cells, with many target genes involved in lymphoid cell growth and survival. It plays a major role in promoting immune and inflammatory responses through producing pro-inflammatory cytokines. It has been associated with autoimmune diseases such as arthritis (52), psoriasis (37), and CD (50). REL expression increases in a variety of B and T cell malignancies as well as in other cancer types. Thus, agents that modulate REL activity may have therapeutic benefits for chronic inflammatory diseases and human cancers (53). TNFAIP3, also known as A20, has been associated with psoriasis (54), rheumatoid arthritis (55), type 1 diabetes mellitus (56), systemic lupus erythematosus (57), and CD (50). TNFAIP3 expression is up-regulated by NF- κ B activation, and acts in a negative feedback to control NF- κ B-dependent gene expression. It blocks NF- κ B activation which mediates immune responses (58); thus, TNFAIP3 plays a key role in controlling inflammation in autoimmune

diseases and is suggested as an attractive candidate for drug targeting (59). Also, TNFAIP3 is frequently inactivated in subsets of B-lineage lymphomas that are characterized by NF- κ B hyper activation. Hence, it could be suggested as a novel tumor suppressor (60).

The inflammatory process involves infiltration of macrophages, B-lymphocytes, plasma cells and mainly T lymphocytes. T-cells can be divided into Th1 (T helper 1) or Th2 (T helper 2) subsets based on their cytokine secretion; IFN- γ and IL-2 are the main cytokines secreted by Th1 cells while IL-4 is secreted by the Th2 cells. Cytokine secretion from lymphocytes is higher among celiac patients as compared to controls. Elevated IFN- γ and IL-2 secretion has been detected in mucosal tissue lymphocytes and peripheral blood T lymphocytes (61-63). TBX21 (T-box transcription factor TBX21) and pSTAT1 are important transcription factors for the development of T helper type 1 (Th1) cells and control of Th1 cytokine expression plus IFN γ . Its expression is also associated with IFN γ expression. Th1 cells and natural killer play an important role for this gene in initiating Th1 lineage development of Th-naive precursor cells (19, 64).

RGS1 (Regulator of G-protein signaling 1) controls the homing of intraepithelial lymphocytes (IELs) which are less active in CD patients than in controls while being essential for producing the gluten induced epithelial damage. IELs play an important role in preventing unwanted and over-inflammatory reactions that can damage the intestinal epithelium. It is also able to prevent the entry and spread of pathogens thus preventing the development of immunity. IELs eliminate infected and damaged cells, as well as regulatory functions that contribute to epithelium healing and repairing. A hallmark of CD is deregulated activation of IELs. It is involved in epithelial cell destruction and the subsequent development of villous atrophy.

The expression of SH2B3 (SH2B adapter protein 3) and RGS1 is lower in CD peripheral monocytes versus controls and higher in CD duodenal intestinal samples than in controls. The capacity of RGS1 to limit the egress of inflammatory and autoimmune cells could encourage the immunopathology of autoimmune disorders (65). SH2B3 is associated with many autoimmune and cardiovascular disorders, including type 1 diabetes mellitus, celiac disease, myocardial

infarction, and hypertension. It is strongly expressed in the inflamed small intestines of CD patients. This alteration leads to modulation of intermolecular interactions and inhibition of cytokine responses. Its function is found in the innate immune response. SH2B3 negatively regulates lymphopoiesis and early hematopoiesis and reduces the Th-1 cell response in the context of CD etiology. Deficiency of SH2B3 leads to enhanced production of B-cells (66).

PCK1 is an enzyme that regulates gluconeogenesis. At low glucose levels, it catalyzes the oxaloacetate (OAA) to phosphoenolpyruvate (PEP) while at high glucose levels, it catalyzes PEP to OAA. It has a metabolic function in diseases (67). KIAA1109 level is higher in CD blood monocytes and lower in CD intestinal tissue samples versus controls. KIAA1109 is expressed in many different tissues e.g. parathyroid, muscle, ear, eye, mammary gland, lymph nodes, and thymus in humans. It is also expressed in various diseases e.g. bladder carcinoma, chondrosarcoma, glioma, leukemia, lymphoma, non-neoplasia, and retinoblastoma tissues. In mammals, its function is regulating epithelial growth and differentiation, tumor development, as well as in inflammatory conditions affecting the gastrointestinal tract, type I diabetes, and rheumatoid arthritis (68-71).

Gen ontology analysis revealed more information about roles of DEGs in the onset and promotion of CD. The important biological term which appeared as a term with maximum DEGs involvement is "cytokine activity". There are 11 DEGs among the 32 introduced genes that are involved in this biological term. This finding is affected by the fact that immune system is the main dysregulated part in CD.

As an overview in CD, the genes involved in inflammatory processes are upregulated, while the genes in the cell adhesion or integrity of the intestinal barrier are downregulated (34).

Since blood is a more available source than intestinal biopsy, finding common DEGs in both samples is a useful task to introduce potential biomarkers. In this regard, about 5% of tissues DEGs were identified as blood DEGs. Most of these potential biomarkers were determined as immune system elements.

Acknowledgment

Shahid Beheshti University of Medical Sciences supported this research.

S114 Comparison of cytokine and gene activities in celiac disease

Conflict of interests

The authors declare that they have no conflict of interest.

References

1. Rostami-Nejad M, Hogg-Kollars S, Ishaq S, Rostami K. Subclinical celiac disease and gluten sensitivity. *Gastroenterol Hepatol Bed Bench* 2011;4:102-8.
2. Izadi F, Rezaei Tavirani M, Honarkar Z, Rostami-Nejad M. Celiac disease and hepatitis C relationships in transcriptional regulatory networks. *Gastroenterol Hepatol Bed Bench* 2017;10:303-10.
3. Pereyra L, Gonzalez R, Mohaidle A, Fischer C, Mella JM, Panigadi GN, et al. Risk of colorectal neoplasia in patients with celiac disease: a multicenter study. *J Crohns Colitis* 2013;7:e672-7.
4. Khoshbaten M, Rostami Nejad M, Farzady L, Sharifi N, Hashemi SH, Rostami K. Fertility disorder associated with celiac disease in males and females: fact or fiction? *J Obstet Gynaecol Res.* 2011;37:1308-12.
5. Farrell RJ, Kelly CP. Celiac sprue. *N Engl J Med* 2002;346:180-8.
6. Ch'ng CL, Jones MK, Kingham JGC. Celiac disease and autoimmune thyroid disease. *Clin Med Res* 2007;5:184-92.
7. Bibbò S, Pes GM, Usai-Satta P, Salis R, Soro S, Quarta Colosso BM, et al. Chronic autoimmune disorders are increased in coeliac disease: a case-control study. *Medicine* 2017;96:e8562-e.
8. Assa A, Frenkel-Nir Y, Tzur D, Katz LH, Shamir R. Large population study shows that adolescents with celiac disease have an increased risk of multiple autoimmune and nonautoimmune comorbidities. *Acta Paediatr* 2017;106:967-72.
9. Bakhshpour A, Kaykhaei MA, Moulaei N, Mashhadi MA. Prevalence of coeliac disease in patients with non-alcoholic fatty liver disease. *Arab J Gastroenterol* 2013;14:113-5.
10. Zubarik R, Ganguly E, Nathan M, Vecchio J. Celiac disease detection in hypothyroid patients requiring elevated thyroid supplementation: A prospective cohort study. *Eur J Int* 2015;26:825-9.
11. Green PH, Rostami K, Marsh MN. Diagnosis of coeliac disease. *Best Pract Res Clin Gastroenterol* 2005;19:389-400
12. Fasano A, Catassi C. Current approaches to diagnosis and treatment of celiac disease: an evolving spectrum. *Gastroenterology* 2001;120:636-51.
13. Golubnitschaja O, Baban B, Boniolo G, Wang W, Bubnov R, Kapalla M, et al. Medicine in the early twenty-first century: paradigm and anticipation - EPMA position paper 2016. *EPMA J* 2016;7:23.
14. Rhodes DR, Tomlins SA, Varambally S, Mahavisno V, Barrette T, Kalyana-Sundaram S, et al. Probabilistic model of the human protein-protein interaction network. *Nat Biotechnol* 2005;23:951-9.
15. Gandhi T, Zhong J, Mathivanan S, Karthick L, Chandrika K, Mohan SS, et al. Analysis of the human protein interactome and comparison with yeast, worm and fly interaction datasets. *Nat Genet* 2006;38:285-93.
16. Azodi MZ, Peyvandi H, Rostami-Nejad M, Safaei A, Rostami K, Vafae R, et al. Protein-protein interaction network of celiac disease. *Gastroenterol Hepatol Bed Bench* 2016;9:268-277.
17. Zachariou M, Minadakis G, Oulas A, Afxenti S, Spyrou GM. Integrating multi-source information on a single network to detect disease-related clusters of molecular mechanisms. *J Proteomics* 2018;188:15-29.
18. Galatola M, Auricchio R, Greco L. Gene expression profiling of celiac biopsies and peripheral blood monocytes using Taqman assays. *Methods Mol Biol* 2015;1326:105-15.
19. Frisullo G, Nociti V, Iorio R, Patanella AK, Plantone D, Bianco A, et al. T-bet and pSTAT-1 expression in PBMC from coeliac disease patients: new markers of disease activity. *Clin Exp Immunol* 2009;158:106-14.
20. Ciccocioppo R, Panelli S, Conti Bellocchi MC, Cangemi GC, Frulloni L, Capelli E, et al. The transcriptomic analysis of circulating immune cells in a celiac family unveils further insights into disease pathogenesis. *Front Med* 2018;5:182.
21. Galatola M, Cielo D, Panico C, Stellato P, Malamisura B, Carbone L, et al. Presymptomatic diagnosis of celiac disease in predisposed children: the role of gene expression profile. *J Pediatr Gastroenterol Nutr* 2017;65:314-20.
22. Brynychova I, Hoffmanova I, Dvorak M, Dankova P. Increased Expression of TLR4 and TLR7 but Not Prolactin mRNA by Peripheral Blood Monocytes in Active Celiac Disease. *Adv Clin Exp Med* 2016;25:887-93.
23. Ghasiyari H, Rostami-Nejad M, Amani D, Rostami K, Pourhoseingholi MA, Asadzadeh-Aghdai H, et al. Diverse Profiles of Toll-Like Receptors 2, 4, 7, and 9 mRNA in Peripheral Blood and Biopsy Specimens of Patients with Celiac Disease. *J Immunol Res* 2018;2018.
24. Bragde H, Jansson U, Fredrikson M, Grodzinsky E, Söderman J. Potential blood-based markers of celiac disease. *BMC Gastroenterol* 2014;14:176.
25. Broide E, Scapa E, Bloch O, Shapiro M, Kimchi N, Ben-Yehudah G, et al. Evidence for aberrant regulation of MAP kinase signal transduction pathway in peripheral blood mononuclear cells in patients with active celiac disease. *Dig Dis Sci* 2009;54:1270-5.
26. Diosdado B, Wapenaar M, Franke L, Duran K, Goerres M, Hadithi Ma, et al. A microarray screen for novel candidate genes in coeliac disease pathogenesis. *Gut* 2004;53:944-51.
27. Stulík J, Hernychová L, Porkertová S, Pozler O, Tučková L, Sánchez D, et al. Identification of new celiac disease autoantigens using proteomic analysis. *Proteomics* 2003;3:951-6.
28. Simula MP, Cannizzaro R, Canzonieri V, Pavan A, Maiero S, Toffoli G, et al. PPAR signaling pathway and cancer-related proteins are involved in celiac disease-associated tissue damage. *Mol Med* 2010;16:199-209.

29. Tavirani MR, Bashash D, Rostami FT, Tavirani SR, Nikzamir A, Tavirani MR, et al. Celiac disease microarray analysis based on system biology approach. *Gastroenterol Hepatol Bed Bench* 2018;11:216.
30. Bragde H, Jansson U, Fredrikson M, Grodzinsky E, Söderman JJC. Celiac disease biomarkers identified by transcriptome analysis of small intestinal biopsies. *Cell Mol Life Sci* 2018;75:4385-401.
31. Bragde H, Jansson U, Jarlsfelt I, Söderman JJPr. Gene expression profiling of duodenal biopsies discriminates celiac disease mucosa from normal mucosa. *Pediatr Res* 2011;69:530.
32. Sangineto M, Graziano G, D'Amore S, Salvia R, Palasciano G, Sabbà C, et al. Identification of peculiar gene expression profile in peripheral blood mononuclear cells (PBMC) of celiac patients on gluten free diet. *PLoS One* 2018;13:e0197915.
33. Pascual V, Medrano L, López-Palacios N, Bodas A, Dema B, Fernández-Arquero M, et al. Different gene expression signatures in children and adults with celiac disease. *PLoS One* 2016;11:e0146276.
34. Cielo D, Galatola M, Fernandez-Jimenez N, De Leo L, Garcia-Etxebarria K, Loganec C, et al. Combined Analysis of Methylation and Gene Expression Profiles in separate Compartments of small Bowel Mucosa Identified Celiac Disease patients' signatures. *Sci Rep* 2019;9:10020.
35. Galatola M, Izzo V, Cielo D, Morelli M, Gambino G, Zanzi D, et al. Gene expression profile of peripheral blood monocytes: a step towards the molecular diagnosis of celiac disease? *PLoS One* 2013;8:e74747.
36. Fernandez-Jimenez N, Santin I, Irastorza I, Plaza-Izurieta L, Castellanos-Rubio A, Vitoria JC, et al. Upregulation of KIR3DL1 gene expression in intestinal mucosa in active celiac disease. *Hum Immunol* 2011;72:617-20.
37. Castellanos-Rubio A, Caja S, Irastorza I, Fernandez-Jimenez N, Plaza-Izurieta L, Vitoria JC, et al. Angiogenesis-related gene expression analysis in celiac disease. *Autoimmunity* 2012;45:264-70.
38. Salvati V, Bajaj-Elliott M, Poulson R, Mazzarella G, Lundin K, Nilsen E, et al. Keratinocyte growth factor and coeliac disease. *Gut* 2001;49:176-81.
39. Green P, Glickman RMJJolr. Intestinal lipoprotein metabolism. *J Lipid Res* 1981;22:1153-73.
40. Vuoristo M, Kesäniemi Y, Gylling H, Miettinen TJM. Metabolism of cholesterol and apolipoprotein B in celiac disease. *Metabolism* 1993;42:1386-91.
41. Niessen H, Lagrand W, Rensink H, Meijer CJ, Aarden L, Hack C. Apolipoprotein H, a new mediator in the inflammatory changes ensuing in jeopardised human myocardium. *J Clin Pathol.* 2000;53:863-7.
42. Miyakis S, Giannakopoulos B, Krilis SA. Beta 2 glycoprotein I-function in health and disease. *Thromb Res* 2004;114:335-46.
43. Sellar G, Keane J, Mehdi H, Peeples M, Browne N, Whitehead AJB, et al. Characterization and Acute-Phase Modulation of Canine Apolipoprotein H (β 2-Glycoprotein 1). *Biochem Biophys Res Commun* 1993;191:1288-93.
44. Açar Ç, de Groot PG, Mörgelin M, Monk SD, van Os G, Levels JH, et al. β 2-glycoprotein I: a novel component of innate immunity. *Blood* 2011;117:6939-47.
45. Stefas I, Dubois G, Tigrett S, Lucarz E, Veas FJAappaensboh, diseases v. Apolipoprotein H, an acute phase protein, a performing tool for ultra-sensitive detection and isolation of microorganisms from different origins. In: Veas F, Editor. *Acute phase proteins as early non-specific biomarkers of human and veterinary diseases.* London: InTechOpen; 2011.
46. Kanai M, Raz A, Goodman DS. Retinol-binding protein: the transport protein for vitamin A in human plasma. *J Clin Invest* 1968;47:2025-44.
47. Rask L, Anundi H, Böhme J, Eriksson U, Fredriksson A, Nilsson S, et al. The retinol-binding protein. *Scand J Clin Lab Invest Suppl* 1980;154:45-61.
48. McGough N, Cummings JH. Coeliac disease: a diverse clinical syndrome caused by intolerance of wheat, barley and rye. *Proc Nutr Soc* 2005;64:434-50.
49. Petit MM, Meulemans SM, Van de Ven WJ. The focal adhesion and nuclear targeting capacity of the LIM containing lipoma-preferred partner (LPP) protein. *J Biol Chem* 2003;278:2157-68.
50. Trynka G, Zhernakova A, Romanos J, Franke L, Hunt K, Turner G, et al. Coeliac disease-associated risk variants in TNFAIP3 and REL implicate altered NF- κ B signalling. *Gut* 2009;58:1078-83.
51. Perkins ND. Integrating cell-signalling pathways with NF- κ B and IKK function. *Nat Rev Mol Cell Biol* 2007;8:49.
52. Gregersen PK, Amos CI, Lee AT, Lu Y, Remmers EF, Kastner DL, et al. REL, encoding a member of the NF- κ B family of transcription factors, is a newly defined risk locus for rheumatoid arthritis. *Nat Genet* 2009;41:820.
53. Gilmore TD, Kalaitzidis D, Liang MC, Starczynowski DT. The c-Rel transcription factor and B-cell proliferation: a deal with the devil. *Oncogene* 2004;23:2275.
54. Kooloos WM, de Jong DJ, Huizinga TW, Guchelaar H-JJdt. Potential role of pharmacogenetics in anti-TNF treatment of rheumatoid arthritis and Crohn's disease. *Drug Discov Today* 2007;12:125-31.
55. Plenge RM, Cotsapas C, Davies L, Price AL, De Bakker PI, Maller J, et al. Two independent alleles at 6q23 associated with risk of rheumatoid arthritis. *Nat Genet* 2007;39:1477.
56. Fung E, Smyth DJ, Howson JM, Cooper JD, Walker NM, Stevens H, et al. Analysis of 17 autoimmune disease-associated variants in type 1 diabetes identifies 6q23/TNFAIP3 as a susceptibility locus. *Genes Immun* 2009;10:188.
57. Graham RR, Cotsapas C, Davies L, Hackett R, Lessard CJ, Leon JM, et al. Genetic variants near TNFAIP3 on 6q23

S116 Comparison of cytokine and gene activities in celiac disease

- are associated with systemic lupus erythematosus. *Nat Genet* 2008;40:1059.
- 58.Wertz IE, O'rourke KM, Zhou H, Eby M, Aravind L, Seshagiri S, et al. De-ubiquitination and ubiquitin ligase domains of A20 downregulate NF- κ B signalling. *Nature* 2004;430:694.
- 59.Coornaert B, Carpentier I, Beyaert R. A20: central gatekeeper in inflammation and immunity. *J Biol Chem* 2009;284:8217-21.
- 60.Verhelst K, Carpentier I, Kreike M, Meloni L, Verstrepen L, Kensche T, et al. A20 inhibits LUBAC-mediated NF- κ B activation by binding linear polyubiquitin chains via its zinc finger 7. *EMBO J* 2012;31:3845-55.
- 61.Kontakou M, Przemioslo R, Sturgess R, Limb A, Ciclitira PJ. Expression of tumour necrosis factor- α , interleukin-6, and interleukin-2 mRNA in the jejunum of patients with coeliac disease. *Scand J Gastroenterol* 1995;30:456-63.
- 62.Romagnani S, Parronchi P, D'elios M, Romagnani P, Annunziato F, Piccinni M, et al. An update on human Th1 and Th2 cells. *Int Arch Allergy Immunol* 1997;113:153-6.
- 63.Mizrachi A, Broide E, Buchs A, Kornberg A, Aharoni D, Bistrizter T, et al. Lack of correlation between disease activity and decreased stimulated secretion of IL-10 in lymphocytes from patients with celiac disease. *Scand J Gastroenterol* 2002;37:924-30.
- 64.Fong SB. Characterization of innate T cells in response to oral bacterial infection [PhD Thesis]. Melbourne: Melbourne Dental School; 2015.
- 65.Abadie V, Discepolo V, Jabri B. Intraepithelial lymphocytes in celiac disease immunopathology. *Semin Immunopathol* 2012;34:551-66.
- 66.Takaki S. Sh2b3/Lnk family adaptor proteins in the regulation of lymphohematopoiesis. *Nihon Rinsho Meneki Gakkai Kaishi* 2008;31:440-7.
- 67.Pilz A, Willer E, Povey S, Abbott CM. The genes coding for phosphoenolpyruvate carboxykinase-1 (PCK1) and neuronal nicotinic acetylcholine receptor α 4 subunit (CHRNA4) map to human chromosome 20, extending the known region of homology with mouse chromosome 2. *Ann Hum Genet* 1992;56:289-93.
- 68.Hunt KA, Zhernakova A, Turner G, Heap GA, Franke L, Bruinenberg M, et al. Newly identified genetic risk variants for celiac disease related to the immune response. *Nat Genet* 2008;40:395.
- 69.Adamovic S, Amundsen S, Lie B, Gudjonsdottir A, Ascher H, Ek J, et al. Association study of IL2/IL21 and Fc γ RIIa: significant association with the IL2/IL21 region in Scandinavian coeliac disease families. *Genes Immun* 2008;9:364-7.
- 70.Van Heel DA, Franke L, Hunt KA, Gwilliam R, Zhernakova A, Inouye M, et al. A genome-wide association study for celiac disease identifies risk variants in the region harboring IL2 and IL21. *Nat Genet* 2007;39:827.
- 71.Bouaziz D, Fourati H, Amouri A, Marques I, Abida O, Tahri N, et al. Autoimmune diseases association study with the KIAA1109-IL2-IL21 region in a Tunisian population. *Mol Biol Rep* 2014;41:7133-9.
- 72.Adams D, Hamilton TA. Macrophages as destructive cells in host defense. In: Gallin JI, Goldstein IM, Snyderman R, editors. *Inflammation: basic principles and clinical correlates*. 2nd ed. New York: Raven Pr; 1992.
- 73.Shi C, Pamer EG. Monocyte recruitment during infection and inflammation. *Nat Rev Immunol* 2011;11:762-74.
- 74.Brown Z, Gerritsen ME, Carley WW, Strieter RM, Kunkel SL, Westwick J. Chemokine gene expression and secretion by cytokine-activated human microvascular endothelial cells. Differential regulation of monocyte chemoattractant protein-1 and interleukin-8 in response to interferon-gamma. *Am J Pathol* 1994;145:913.