

POSTER PRESENTATION

Open Access

# A phase I study of an agonist anti-CD27 human antibody (CDX-1127) in patients with advanced hematologic malignancies or solid tumors

Stephen Ansell<sup>1\*</sup>, Donald Northfelt<sup>2</sup>, Ian Flinn<sup>3</sup>, Howard Burris<sup>3</sup>, Shira Dinner<sup>4</sup>, Victor Villalobos<sup>4</sup>, Branimir Sikic<sup>4</sup>, Lana Pilja<sup>5</sup>, Michael Yellin<sup>5</sup>, Tibor Keler<sup>5</sup>, Thomas Davis<sup>5</sup>

From Society for Immunotherapy of Cancer 28th Annual Meeting  
National Harbor, MD, USA. 8-10 November 2013

CD27, a member of the tumor necrosis factor receptor superfamily, is constitutively expressed on the majority of mature T cells, memory B cells, and a subset of NK cells. We previously demonstrated anti-tumor and immunological properties of the fully human anti-CD27 mAb 1F5 (CDX-1127) in murine tumor models of hematologic and solid tumors. A Phase I dose escalation study of CDX-1127 in patients (pts) with recurrent or treatment-refractory B-cell malignancies and select solid tumors is ongoing. CDX-1127 (0.1 to 10 mg/kg IV) is administered as a single-dose with 28-day observation, followed by 4 weekly doses before restaging at Day 85. Up to 4 additional re-treatment cycles (consisting of 4 weekly doses) are permitted in pts with stable disease or tumor response. Dose-escalation in pts with solid tumors has been completed with preliminary results presented in a separate abstract. Flow cytometry and functional immune analysis of peripheral blood lymphocytes from these solid tumor pts showed an increase in expression of activation markers on T cells (HLA-DR), increases in NK cells, decrease in Tregs, and enhanced T cell response to various stimuli in *in vitro* assays. Tumor shrinkage was also seen. Dose-escalation in pts with hematologic malignancies is ongoing. To date, 13 B-cell lymphoma pts (3 Hodgkin, 3 follicular, 3 marginal zone, 4 diffuse large B-cell) have received 0.1, 0.3, or 1 mg/kg CDX-1127, with no DLT. Treatment-related toxicity, generally Grade 1-2, included fatigue, nausea, decreased appetite and anemia. One pt with Hodgkin lymphoma who had previously progressed following treatment with chemotherapy, autologous stem cell transplant, and brentuximab vedotin had a partial response (PR) with

77% shrinkage in measurable disease; treatment is ongoing. Three additional pts had stable disease (range: 4.5 to 11+ months), including a pt with follicular lymphoma who has completed 3 cycles of therapy and a pt with marginal zone lymphoma with -36% shrinkage in measurable disease. Flow cytometry and functional immune analysis of peripheral blood lymphocytes are being performed to assess the immunomodulatory activity of CDX-1127 in patients with hematological malignancies. Emerging results suggest that weekly dosing of CDX-1127 is well tolerated with promising biological and early clinical activity. In addition, these data position CDX-1127 well for combination therapy with conventional and immunotherapy approaches as demonstrated in preclinical models.

#### Authors' details

<sup>1</sup>Mayo Clinic, Rochester, MN, USA. <sup>2</sup>Mayo Clinic, Scottsdale, AZ, USA. <sup>3</sup>Sarah Cannon Research Institute, Nashville, TN, USA. <sup>4</sup>Stanford Cancer Institute, Stanford, CA, USA. <sup>5</sup>Celldex Therapeutics, Inc., Phillipsburg, NJ, USA.

Published: 7 November 2013

doi:10.1186/2051-1426-1-S1-P259

Cite this article as: Ansell *et al.*: A phase I study of an agonist anti-CD27 human antibody (CDX-1127) in patients with advanced hematologic malignancies or solid tumors. *Journal for ImmunoTherapy of Cancer* 2013 **1**(Suppl 1):P259.

<sup>1</sup>Mayo Clinic, Rochester, MN, USA

Full list of author information is available at the end of the article