

Redescription of *Pseudopoda taibaischana* (Araneae, Sparassidae), with the first description of the female

Li-Jun Gong¹, Yang Zhong¹

¹ Hubei Key Laboratory of Radiation Chemistry and Functional Materials, School of Nuclear Technology and Chemistry & Biology, Hubei University of Science and Technology, Xianning 437100, Hubei, China

Corresponding author: Yang Zhong (hubeispider@aliyun.com)

Academic editor: G. Blagoev | Received 27 July 2020 | Accepted 13 October 2020 | Published 11 November 2020

<http://zoobank.org/7312CAB8-96B1-4656-B38F-DABE9CD797F4>

Citation: Gong L-J, Zhong Y (2020) Redescription of *Pseudopoda taibaischana* (Araneae, Sparassidae), with the first description of the female. ZooKeys 991: 111–119. <https://doi.org/10.3897/zookeys.991.56969>

Abstract

Pseudopoda taibaischana Jäger, 2001 (Sparassidae) is redescribed based on new material from the type locality in Taibaishan Nation Forest Park of Shaanxi Province, China. The female is described and illustrated for the first time, and a redescription is provided for the male.

Keywords

Biodiversity, huntsman spiders, Shaanxi, taxonomy

Introduction

The genus *Pseudopoda* was established by Jäger (2000) to include nine species previously assigned to *Heteropoda*: *P. casaria* (Simon, 1897), *P. exigua* (Fox, 1938), *P. exiguoides* (Song & Zhu, 1999), *P. grahami* (Fox, 1936), *P. lushanensis* (Wang, 1990), *P. prompta* (O. Pickard-Cambridge, 1885), *P. virgata* (Fox, 1936), *P. zhangmuensis* (Hu & Li, 1987), and *P. zhejiangensis* (Zhang & Kim, 1996). Currently, *Pseudopoda* is the third largest genus of the subfamily Heteropodinae, and includes 142 species. Members of this genus are known from Bhutan, China, India, Indonesia, Japan, Laos, Myanmar, Nepal, Pakistan, Thailand, and Vietnam (World Spider Catalog 2020). From China,

63 species are known; among them, seven species are only known from females and eleven only from males (World Spider Catalog 2020). *Pseudopoda taibaischana* Jäger, 2001 was first described based on one male specimen from Taibaishan National Forest Park of Shaanxi Province, China (Jäger 2001). Recently, new material of both sexes was collected from the type locality of this species, enabling us to describe the female for the first time in this paper.

Materials and methods

Specimens were examined and measured with a Leica M205C stereomicroscope. The points arising from the tegular appendages are listed as clock-positions from the left bulb in ventral view. Male palps were examined after dissection and detachment. The epigynes were examined and illustrated after dissection: they were removed and cleared in warm lactic acid before illustration. The vulva was photographed after being embedded in Arabic gum. All photographs were taken with a Leica DFC450 digital camera attached to a Leica M205C stereomicroscope, with 10–20 photographs taken in different focal planes and combined using the image stacking software Leica LAS. Images were edited using Adobe Photoshop CC 2015.

Leg measurements are listed as: total length (femur, patella, tibia, metatarsus, tarsus). The number of spines is listed for each segment in the following order: prolateral, dorsal, retrolateral, ventral (in femora and patellae, ventral spines are absent, and the fourth digit is omitted in the spination formula).

Abbreviations used in the text and figures are given below:

SMF Senckenberg Research Institute and Museum, Frankfurt, Germany (P. Jäger);
HUST School of Nuclear Technology and Chemistry & Biology, Hubei University of Science and Technology, Xianning, Hubei, China (Y. Zhong);

ALE	anterior lateral eye;	OW	opisthosoma width;
AME	anterior median eye;	Pa	patella;
AW	anterior width of carapace;	PI	posterior incision of LL;
C	conductor;	PL	carapace length;
CO	copulatory opening;	PLE	posterior lateral eyes;
CH	clypeus height;	PME	posterior median eyes;
E	embolus;	Pp	palp;
EP	embolic projection;	PP	posterior part of spermathecae;
FD	fertilization duct;	PW	carapace width;
FE	femur;	RTA	retrolateral tibial apophysis;
FW	first winding;	S	spermathecae;
LL	lateral lobes;	T	tegulum;
Mt	metatarsus;	Ta	tarsus;
OL	opisthosoma length;	Ti	tibia. I, II, III, IV—legs I to IV.

Taxonomy

Family Sparassidae Bertkau, 1872

Subfamily Heteropodinae Thorell, 1873

Genus *Pseudopoda* Jäger, 2000

Type species. *Sarotes promptus* O. Pickard-Cambridge, 1885.

Diagnosis. See Jäger (2000) and Jiang et al. (2018).

Composition. *P. daliensis*-group (*P. anguilliformis* Zhang et al., 2017, *P. peronata* Zhang et al., 2017, *P. sicyoidea* Zhang et al., 2017, *P. daliensis* Jäger & Vedel, 2007, *P. kunmingensis* Sun & Zhang, 2012), *P. diversipunctata*-group (*P. diversipunctata* Jäger, 2001, *P. intermedia* Jäger, 2001, *P. marsupia* (Wang, 1991)), *P. latembola*-group (*P. albolineata* Jäger, 2001, *P. alta* Jäger, 2001, *P. chauki* Jäger, 2001, *P. everesta* Jäger, 2001, *P. latembola* Jäger, 2001, *P. monticola* Jäger, 2001, *P. sinopodooides* Jäger, 2001), *P. martensi*-group (*P. chulingensis* Jäger, 2001, *P. dhulensis* Jäger, 2001, *P. gogona* Jäger, 2001, *P. hyatti* Jäger, 2001, *P. kalinchoka* Jäger, 2001, *P. khimtensis* Jäger, 2001, *P. martensi* Jäger, 2001, *P. martinae* Jäger, 2001, *P. megalopora* Jäger, 2001, *P. platembola* Jäger, 2001, *P. tinjura* Jäger, 2001, *P. varia* Jäger, 2001, *P. virgata* (Fox, 1936)), *P. parvipunctata*-group (*P. biapicata* Jäger, 2001, *P. dao* Jäger, 2001, *P. jirensis* Jäger, 2001, *P. parvipunctata* Jäger, 2001, *P. schawalleri* Jäger, 2001, *P. thorelli* Jäger, 2001, *P. triapicata* Jäger, 2001, *P. lushanensis* (Wang, 1990)), *P. prompta*-group (*P. brauni* Jäger, 2001, *P. cuneata* Jäger, 2001, *P. grasshoffi* Jäger, 2001, *P. huberti* Jäger, 2001, *P. marmorea* Jäger, 2001, *P. trisuliensis* Jäger, 2001, *P. casaria* (Simon, 1897), *P. prompta* (O. Pickard-Cambridge, 1885), *P. zhangmuensis* (Hu & Li, 1987)), *P. schwendingeri*-group (*P. hirsuta* Jäger, 2001, *P. schwendingeri* Jäger, 2001), and *P. signata*-group (*P. bibulba* (Xu & Yin, 2000), *P. physematosa* Zhang et al., 2019, *P. semilunata* Zhang et al., 2019, *P. signata* Jäger, 2001, *P. wu* Jäger, Li & Krehenwinkel, 2015, *P. yinae* Jäger & Vedel, 2007, *P. yunnanensis* (Yang & Hu, 2001)) and 88 other species that have not yet been grouped.

Pseudopoda taibaischana Jäger, 2001

Figures 1–4

Pseudopoda taibaischana Jäger, 2001: 86, figs 47a–e (holotype male from Taibaishan National Forest Park of Shaanxi Province, deposited in SMF PJ1056)

Material examined. 2♂, 10♀ (HUST 0001), Shaanxi Province, Baoji City, Taibaishan National Forest Park; 34.05°N, 107.87°E; alt. 1438 m; 20.VII. 2019, Y. Zhong leg.

Diagnosis. This species resembles *Pseudopoda cangshana* Jäger & Vedel, 2007 (Jäger and Vedel 2007: figs 66–68, 70–72) in having the embolus strongly S-shaped, proximal part of embolus visible, and lateral loops of internal duct system extending laterally beyond its first winding, but can be distinguished from the latter by the




Figure 1. *Pseudopoda taibaischana* Jäger, 2001 **A–C** left male palp (**A** prolateral view **B** ventral view **C** retrolateral view) **D, E** cheliceral dentition, ventral view (**D** male **E** female) **F** schematic course of internal duct system. Abbreviations: C–conductor, E–embolus, EP–embolic projection, RTA–retrolateral tibial apophysis, T–tegulum. Scale bars: 0.5 mm.




Figure 2. *Pseudopoda taibaischana* Jäger, 2001 **A, B** Left male palpal tibia (**A** ventral view **B** retrolateral view) **C** epigyne, intact **D** epigyne, cleared **E** vulva, cleared **F** epigyne, cleared and embedded in Arabic gum **G** vulva, cleared and embedded in Arabic gum (**C, D, F** ventral view **E, G** dorsal view). Abbreviations: CO—copulatory opening, FD—fertilization duct, FW—first winding, LL—lateral lobes, PI—posterior incision of LL, S—spermathecae. Scale bars: 0.5 mm.




Figure 3. *Pseudopoda taibaischana* Jäger, 2001 **A,B** male habitus (**A** dorsal view **B** ventral view) **C,D** female habitus (**C** dorsal view **D** ventral view). Scale bars: 2 mm.

following characters: 1, male palp with laminar and rounded embolic projection (absent in *P. cangshana*); 2, tip of RTA with distinct triangular extension dorsally (absent in *P. cangshana*); 3, female epigyne with converging part of anterior margins of lateral lobes T-shaped (Y-shaped in *P. cangshana*); 4, female vulva with loops of internal duct system distinctly curved in ventral view (not curved in *P. cangshana*) (Figs 1, 2).

Description. Male. PL 3.7, PW 2.8, AW 2.0, OL 4.0, OW 3.2. Eyes and interdistances: AME 0.20, ALE 0.25, PME 0.21, PLE 0.27, AME–AME 0.18, AME–ALE 0.07, PME–PME 0.24, PME–PLE 0.30, AME–PME 0.32, ALE–PLE 0.27,



Figure 4. Collection localities of *Pseudopoda taibaischana* in Shaanxi Province, China.

CH AME 0.30, CH ALE 0.28. Spination: Palp: 131, 101, 2101; Fe: I–III 323, IV 331; Pa: I–IV 001; Ti: I–II 2026, III–IV 2126; Mt: I–II 2024, III 3024, IV 3036. Measurements of palp and legs: Palp 6.3 (2.0, 1.1, 1.2, –, 2.0), I 17.9 (4.7, 2.0, 5.4, 4.2, 1.6), II 19.4 (5.4, 2.1, 5.8, 4.3, 1.8), III 14.9 (4.4, 1.6, 4.2, 3.3, 1.4), IV 18.2 (5.2, 1.7, 4.7, 4.9, 1.7). Leg formula: 2-4-1-3. Cheliceral furrow with three anterior and four posterior teeth, each tooth with 22 denticles (Fig. 1D). Carapace yellowish brown, with fovea slightly darker and bearing more spots. Chelicerae deep reddish brown. Sternum pale yellow, with small and irregular spots. Legs yellowish brown, with medium-sized spots and slightly larger spine patches. Abdomen yellowish brown dorsally, with three pairs of dark patches laterally and an irregular pattern in posterior half; ventrally yellowish brown with small and irregular patches (Fig. 3A, B).

Palp as in diagnosis. Cymbium longer than tibia. Embolus arising from tegulum at 8 o'clock position, embolic projection making the tip of embolus look somewhat incised. Conductor curved, arising from an 11 o'clock position. Spermophor visible and slightly curved in retrolateral view. RTA arising medially from tibia, with only one apex, broad in retrolateral view (Figs 1, 2A, B).

Female. PL 3.6, PW 3.2, AW 2.3, OL 4.7, OW 3.4. Eyes and interdistances: AME 0.17, ALE 0.23, PME 0.20, PLE 0.26, AME–AME 0.16, AME–ALE 0.10, PME–PME 0.23, PME–PLE 0.30, AME–PME 0.33, ALE–PLE 0.28, CH AME 0.35, CH ALE 0.31. Spination: Palp: 131, 101, 1014, 2121; Fe: I–III 323, IV 331; Pa: I–IV 001; Ti: I 2026, III–IV 2126; Mt: I–II 2024, III 3025, IV 3036. Measurements of palp and legs: Palp 4.9 (1.6, 0.7, 1.0, –, 1.6), I 12.8 (3.8, 1.7, 3.3, 2.8, 1.2), II 14.0 (4.3, 1.3, 3.7, 3.4, 1.3), III 11.6 (3.5, 1.4, 2.8, 2.9, 1.0), IV 13.3 (4.0, 1.3, 3.3, 3.5, 1.2). Leg formula: 2-4-1-3. Cheliceral furrow with three anterior and four posterior teeth, each tooth with 32 denticles (Fig. 1E).

Epigynal field only slightly wider than long, with very short anterior bands or without such bands. Anterior margins of lateral lobes bent anteriorly at their lateral ends. Posterior incision of lateral lobe distinct, near the posterior meeting point of lateral lobes. Base of internal duct system distinctly extending laterally beyond first winding (Fig. 2C–G).

Coloration in ethanol: as in male, but generally darker, abdomen with more spots ventrally (Fig. 3C, D).

Distribution. China (Shaanxi Province) (Fig. 4).

Acknowledgements

We especially thank Dr. Peter Jäger (Senckenberg Research Institute and Museum, Germany) for helping us identify the males of this species. The manuscript benefitted greatly from comments by Drs. Cristina Rheims (Instituto Butantan, Sao Paulo, Brazil) and Alireza Zamani (Zoological Museum, Biodiversity Unit, University of Turku, Finland). This study was financially supported by the National Natural Sciences Foundation of China (NSFC-32000303), Natural Sciences Foundation of Hubei Province (2019CFB248), PhD grant from Hubei University Science and Technology (BK201811, BK202114) and Biological Applications of Nuclear Technology, Nuclear Technology Innovation team project of Hubei University of Science and Technology (H2019002).

References

- Bertkau P (1872) Über die Respirationsorgane der Araneen. *Archiv für Naturgeschichte* 38: 208–233.
- Jäger P (2000) Two new heteropodine genera from southern continental Asia (Araneae: Sparassidae). *Acta Arachnologica* 49(1): 61–71. <https://doi.org/10.2476/asjaa.49.61>

- Jäger P (2001) Diversität der Riesenkrabbenspinnen im Himalaya – die Radiation zweier Gattungen in den Schneetropen (Araneae, Sparassidae, Heteropodinae). *Courier Forschungsinstitut Senckenberg* 232: 1–136.
- Jäger P, Vedel V (2007) Sparassidae of China 4. The genus *Pseudopoda* (Araneae: Sparassidae) in Yunnan Province. *Zootaxa* 1623: 1–38. <https://doi.org/10.11646/zootaxa.1623.1.1>
- Jiang TY, Zhao QY, Li SQ (2018) Sixteen new species of the genus *Pseudopoda* Jäger, 2000 from China, Myanmar, and Thailand (Sparassidae, Heteropodinae). *ZooKeys* 791: 107–161. <https://doi.org/10.3897/zookeys.791.28137>
- Thorell T (1873) Remarks on synonyms of European spiders. Part IV. C. J. Lundström, Uppsala, 375–645.
- WSC (2020) World Spider Catalog, version 21.5. Natural History Museum, Bern. <https://doi.org/10.24436/2> [accessed 25.07.2020]
- Zhang H, Jäger P, Liu J (2017) One new *Pseudopoda* species group (Araneae: Sparassidae) from Yunnan Province, China, with description of three new species. *Zootaxa* 4318(2): 271–294. <https://doi.org/10.11646/zootaxa.4318.2.3>
- Zhang H, Jäger P, Liu J (2019) Establishing a new species group of *Pseudopoda* Jäger, 2000 with the description of two new species (Araneae, Sparassidae). *ZooKeys* 879: 91–115. <https://doi.org/10.3897/zookeys.879.35110>