



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

# Asthma in the elderly: Current understanding and future research needs—a report of a National Institute on Aging (NIA) workshop

Nicola A. Hanania, MD, MS, (Co-chair),<sup>a\*</sup> Monroe J. King, DO, (Chair),<sup>b\*</sup> Sidney S. Braman, MD,<sup>c</sup> Carol Saltoun, MD,<sup>d</sup> Robert A. Wise, MD,<sup>e</sup> Paul Enright, MD,<sup>f</sup> Ann R. Falsey, MD,<sup>g</sup> Sameer K. Mathur, MD, PhD,<sup>h</sup> Joe W. Ramsdell, MD,<sup>i</sup> Linda Rogers, MD,<sup>j</sup> David A. Stempel, MD,<sup>k</sup> John J. Lima, PharmD,<sup>l</sup> James E. Fish, MD,<sup>m</sup> Sandra R. Wilson, PhD,<sup>n</sup> Cynthia Boyd, MD, MPH,<sup>o</sup> Kushang V. Patel, PhD,<sup>o</sup> Charles G. Irvin, PhD,<sup>p</sup> Barbara P. Yawn, MD, MSc,<sup>q</sup> Ethan A. Halm, MD, MPH,<sup>r</sup> Stephen I. Wasserman, MD,<sup>i</sup> Mark F. Sands, MD,<sup>s</sup> William B. Ershler, MD,<sup>t</sup> and Dennis K. Ledford, MD,<sup>b</sup> for the Asthma in the Elderly workshop participants<sup>‡</sup> Houston and Dallas, Tex, Tampa and Jacksonville, Fla, Providence, RI, Chicago, Ill, Baltimore, Bethesda, and Gaithersburg, Md, Tucson, Ariz, New York, Rochester, and Buffalo, NY, Madison, Wis, San Diego and Palo Alto, Calif, Bellevue, Wash, Gladwyne, Pa, Burlington, Vt, and Rochester, Minn

Asthma in the elderly is underdiagnosed and undertreated, and there is a paucity of knowledge on the subject. The National Institute on Aging convened this workshop to identify what is known and what gaps in knowledge remain and suggest research directions needed to improve the understanding and care of asthma in the elderly. Asthma presenting at an advanced age often has similar clinical and physiologic consequences as seen with younger patients, but comorbid illnesses and the psychosocial effects of aging might affect the diagnosis, clinical presentation, and care of asthma in this population. At least 2 phenotypes exist among elderly patients with asthma; those with longstanding asthma have more severe airflow limitation and less complete reversibility than those with late-onset asthma. Many challenges exist in the recognition and treatment of asthma in the elderly. Furthermore, the pathophysiologic mechanisms of asthma in the elderly are likely to be different from those seen in young asthmatic patients, and these differences might influence the clinical course and outcomes of asthma in this population. (*J Allergy Clin Immunol* 2011;128:S4-24.)

**Key words:** Aging, airway, allergy, asthma, elderly, immune mechanisms, immunosenescence

The proportion of persons older than 65 years in the United States is currently about 13% but is projected to grow from about 40 million in 2005 to more than 86 million by 2050, accounting for 25% of the population. The age group with the largest growth will be those older than 85 years, which is estimated to be more than 1 million by 2050.<sup>1,2</sup> In 2004, the US prevalence of asthma for those 65 years or older was 7%, with 1,088,000 reporting an asthma attack in the previous 12 months.<sup>3</sup> Older asthmatic patients are more likely to be underdiagnosed, undertreated,<sup>4,5</sup> and hospitalized than younger asthmatic patients.<sup>3</sup> They also have the highest death rate (51.3 per million persons) of any other age group.<sup>6</sup> Older women are hospitalized more than twice as often as older men.

Asthma in older adults is superimposed on a background of aging-related changes in respiratory and immune physiology and often on multiple diseases and conditions common in older age.

\*These authors contributed equally to this work.

‡The Asthma in the Elderly workshop participants are shown in Appendix 1.

From <sup>a</sup>the Section of Pulmonary and Critical Care Medicine, Asthma Clinical Research Center, Baylor College of Medicine, Houston; <sup>b</sup>the Division of Allergy and Immunology, Department of Medicine, University of South Florida, Tampa; <sup>c</sup>Brown University, Providence; <sup>d</sup>Northwestern University, Chicago; <sup>e</sup>Johns Hopkins University, Baltimore; <sup>f</sup>the University of Arizona, Tucson; <sup>g</sup>the University of Rochester, Rochester; <sup>h</sup>the Department of Medicine, Section of Allergy, Pulmonary, and Critical Care, University of Wisconsin, Madison; <sup>i</sup>the University of California—San Diego, San Diego; <sup>j</sup>New York University School of Medicine, New York; <sup>k</sup>GlaxoSmithKline, Bellevue; <sup>l</sup>Clinical Pediatric Pharmacology, Nemours Children Clinic, Jacksonville; <sup>m</sup>Genentech, Gladwyne; <sup>n</sup>Stanford University School of Medicine, Palo Alto; <sup>o</sup>the National Institute on Aging, Bethesda; <sup>p</sup>the University of Vermont, Burlington; <sup>q</sup>the Department of Family and Community Health, University of Minnesota, Rochester; <sup>r</sup>the University of Texas Southwestern Medical Center, Dallas; <sup>s</sup>the Department of Veteran Affairs Medical Center and the University of Buffalo, State University of New York; and <sup>t</sup>the Institute for Advanced Studies in Aging, Gaithersburg.

Publication of this article was supported by the National Institute on Aging.

Disclosure of potential conflict of interest: N. A. Hanania is on the speakers' bureau for GlaxoSmithKline, AstraZeneca, and Genentech; is on advisory boards for GlaxoSmithKline, Genentech, Novartis, and Pfizer; and receives research support from GlaxoSmithKline, Genentech, Novartis, and Pfizer. M. J. King has consultant arrangements with the National Institutes of Health (NIH). R. A. Wise has consultant arrangements with AstraZeneca, Boehringer Ingelheim, GlaxoSmithKline, Novartis, Sunovion, and Merck and receives research support from GlaxoSmithKline, Boehringer Ingelheim, Forest, and Merck. A. R. Falsey has consultant arrangements with

MedImmune and Sanofi Pasteur and receives research support from Sanofi Pasteur and GlaxoSmithKline. S. K. Mathur receives research support from the NIH and is vice-chair of the American Academy of Allergy, Asthma & Immunology (AAAAI)'s Asthma and Allergic Diseases in the Elderly committee. D. A. Stempel is employed by GlaxoSmithKline. J. J. Lima receives research support from the NIH and the American Lung Association. J. E. Fish is employed by Merck & Co, Inc. S. R. Wilson receives research support from the National Heart, Lung, and Blood Institute/NIH. C. G. Irvin is on the speakers' bureau for Merck and the advisory board for MethaPharm, receives research support from the NIH and the American Lung Association, and is on the advisory board for the Vermont Department of Health. B. P. Yawn has consulted for the American Thoracic Society, receives research support from the Agency for Healthcare Research and Quality, and is the chair of the state coalition for asthma for the American Lung Association. S. I. Wasserman receives research support from the American Lung Association, has provided legal consultation or expert witness testimony in cases related to mold injury and medical malpractice, and is president of the American Board of Allergy and Immunology. The rest of the authors have declared that they have no conflict of interest.

Received for publication April 26, 2011; revised June 24, 2011; accepted for publication June 27, 2011.

Reprint requests: Monroe J. King, DO, Division of Allergy and Immunology, Department of Medicine, University of South Florida, 7674 Cumberland Ct, Largo, FL 33777. E-mail: m.james.king@me.com.

0091-6749/\$36.00

© 2011 American Academy of Allergy, Asthma & Immunology

doi:10.1016/j.jaci.2011.06.048

#### Abbreviations used

|        |  |
|--------|--|
| ADL:   | Activities of daily living                     |
| COPD:  | Chronic obstructive pulmonary disease          |
| FVC:   | Forced vital capacity                          |
| GERD:  | Gastroesophageal reflux disease                |
| hMPV:  | Human metapneumovirus                          |
| LOA:   | Late-onset asthma                              |
| LSA:   | Longstanding asthma                            |
| LTM:   | Leukotriene-modifying agent                    |
| NIA:   | National Institute on Aging                    |
| NO:    | Nitric oxide                                   |
| RSV:   | Respiratory syncytial virus                    |
| TENOR: | The Epidemiology and Natural History of Asthma |

Recognizing the paucity of research, the many challenges that exist in the recognition and treatment of asthma in older adults, and the opportunity to bridge geriatrics and the clinical specialties that focus on asthma, the National Institute on Aging (NIA) sponsored a Workshop on Asthma in the Elderly in Herndon, Virginia, on September 8 and 9, 2008. The workshop was planned by a committee of 6 physician-scientists from US academic institutions or from the Division of Geriatrics and Clinical Gerontology in the NIA. The planning committee selected speakers and participants for their expertise in asthma, pulmonology, allergy/immunology, primary care, emergency medicine, geriatrics, and/or gerontologic science (see the list of participants in Appendix 1). The immediate goals of this workshop were to summarize the current understanding of the mechanisms of asthma in older persons and to identify knowledge gaps and research opportunities leading to improved medical care and health outcomes for older persons with asthma. These research opportunities are discussed in the body of this report and summarized in Table I.<sup>7-9</sup>

In addition, the NIA, in collaboration with the National Heart, Lung, and Blood Institute and the National Institute of Allergy and Infectious Diseases, recently issued a set of program announcements inviting research proposals on asthma in older adults (<http://grants.nih.gov/grants/guide/pa-files/PA-10-263.html>, <http://grants.nih.gov/grants/guide/pa-files/PA-10-264.html>, and <http://grants.nih.gov/grants/guide/pa-files/PA-10-265.html>).

## BIOLOGY OF AGING

It is a central principle of gerontology that aging itself is not a disease.<sup>10</sup> Yet there are physiological changes within organs, tissues, and cells that result in diminished functional reserve and thereby increased susceptibility to stressors, disease, or both. A second principle is that these aging changes are highly variable and account for the great constitutional heterogeneity among older persons from very "fit" to very "frail." In fact, the concept of frailty, both its causes and consequences, has become a focus of concentrated gerontologic investigation.

### Aging at the cellular level

At the root of age-associated physiological changes are a number of genetic, epigenetic, and environmental factors.<sup>11</sup> Molecular damage accumulates over time, and the capacity for DNA repair decreases.<sup>12</sup> Cellular senescence, which is believed to be the consequence of accumulated DNA and protein damage and

reduced proliferative capacity, is becoming increasingly understood at the molecular level.<sup>13</sup> However, just how this correlates with the phenotypic changes of advanced age remains incompletely understood.<sup>14</sup>

There has been much written about cellular senescence and the events that lead to cell death.<sup>15-17</sup> After a finite number of divisions, normal somatic cells invariably enter a state of irreversibly arrested growth, a process termed *replicative senescence*.<sup>18</sup> In fact, it has been proposed that escape from the regulators of senescence is the antecedent of malignant transformation. However, the role of replicative senescence as an explanation of organismal aging remains the subject of vigorous debate. The controversy relates, in part, to the fact that certain organisms (eg, *Drosophila* species and *Caenorhabditis elegans*) undergo an aging process, yet all of their adult cells are postreplicative.

What is clear is that the loss of the proliferative capacity of human cells in culture is intrinsic to the cells and not dependent on environmental factors or even culture conditions.<sup>18</sup> Unless transformation occurs, cells age with each successive division. The number of divisions turns out to be more important than the actual amount of time passed. Thus cells held in a quiescent state for months, when allowed back into a proliferative environment, will continue approximately the same number of divisions as those that were allowed to proliferate without a quiescent period.<sup>19</sup>

The question remains whether this *in vitro* phenomenon is relevant to animal aging. One suggestive observation is that fibroblasts cultured from samples of old skin undergo fewer cycles of replication than those from young skin.<sup>20</sup> Furthermore, when various species are compared, replicative potential is directly and significantly related to lifespan.<sup>21</sup> An unusual  $\beta$ -galactosidase with activity peaks at pH 6 has proved to be a useful biomarker of *in vitro* senescence because it is expressed by senescent but not presenescent or quiescent fibroblasts.<sup>22</sup> This particular  $\beta$ -galactosidase isoform was found to have the predicted pattern of expression in skin from young and old donors, with measurably increased levels in dermal fibroblasts and epidermal keratinocytes with advancing age.<sup>22</sup> The nature of the expression of this *in vivo* biomarker of aging in other tissues will be important to discern.

### Pathways to frailty

For clinical investigators, frailty has proved hard to define primarily because of the seemingly insurmountable heterogeneity inherent in geriatric populations on the basis of these variable rates of organ system decrease and the presence or absence of 1 or more diseases.<sup>9</sup> Yet, regardless of the pathway taken to frailty, the clinical picture has common features, including a reduction in lean body mass (sarcopenia), loss of bone mass (osteopenia), cognitive impairment, functional decline, and anemia. On the basis of data derived from large cohorts of elderly patients, Fried et al<sup>23</sup> have offered an operational definition of frailty incorporating an assessment of 5 specific characteristics to ascribe a frailty index. On this 5-point scale, a score of 3 or more has been shown to be independently predictive of a range of adverse clinical outcomes, including acute illness, falls, hospitalization, nursing home placement, and early mortality.<sup>23-25</sup> Furthermore, simple performance measures, such as the assessment of walking speed, are predictive of important outcomes, including survival.<sup>26</sup>

With the phenotype better defined, attention has shifted to pathophysiology. Although frailty can occur in the absence of a diagnosable illness, the fact that some become frail and others do

**TABLE I.** Identified future research needs

|   |  |
|---|--|
| The aging lung  |  |
| Large, longitudinal, and more complete studies to determine the effects of aging on the function of the respiratory system  |  |
| Improved knowledge about lung structure-function relationships in older age using techniques of imaging and measures of lung function not requiring effort (eg, high-resolution computed tomographic scanning and forced oscillation)   |  |
| Improved assessment of lung processes underlying airflow limitation attributable to aging versus COPD or asthma, especially in asthmatic patients who smoke   |  |
| Studies to examine the effects of aging in ethnic groups and the role of gender   |  |
| Epidemiology, effect, diagnosis, and management   |  |
| Determine the true prevalence and cost of asthma in the older population  |  |
| Develop a uniform definition of asthma to be applied to health care records that will distinguish asthma from COPD and mixed asthma/COPD  |  |
| Evaluate evidence-based treatment algorithms for older asthmatic patients, such as those developed by the National Heart, Lung, and Blood Institute and Global Initiative For Asthma guidelines <sup>7</sup>  |  |
| Assess the effect of asthma treatment, including direct medical costs of care, indirect costs of care, and value of treatment in improving quality of life <sup>8,9</sup>   |  |
| Assess the effect of comorbid conditions, especially COPD and congestive heart failure, on asthma <sup>9</sup>  |  |
| Characterize phenotypes of elderly asthma with regard to responses to therapy and long-term outcomes based on age of onset, duration of disease, and environmental triggers   |  |
| Develop algorithms for electronic medical record systems that are asthma-specific   |  |
| Evaluate effects of current asthma medications in older patients compared with younger patients   |  |
| Identify pharmacogenetic determinants of response to asthma medications in older adults   |  |
| Identify simpler and safer drug delivery systems and schedules for older adults   |  |
| Develop simple methods to differentiate COPD from asthma exacerbations in older adults  |  |
| Epigenetics and environmental and microbiological triggers  |  |
| Understand how environmental or aging-related factors affect epigenetic changes in asthma in older adults   |  |
| Identify differences between older and younger asthmatic patients or between LSA and LOA with regard to inflammation, remodeling, intracellular mechanisms, responses to environmental pollutants, and allergy sensitization and their effects on the metabolism and action of asthma drugs |  |
| Identify naturally occurring age-related changes in airway cellular patterns  |  |
| Develop animal models of age-related airway inflammation  |  |
| Understand the significance of allergy sensitization associated with asthma in older adults (eg, through larger prospective studies)  |  |
| Identify the utility of allergy tests, either skin tests or serum specific IgE measurement, in reflecting allergy sensitization in older adults   |  |
| Identify the role of the microbiome in patients with LOA  |  |
| Understand the role of non-IgE mechanisms in older adults' inflammatory responses to inhaled allergens or pollutants (eg, T <sub>H</sub> 17 lymphocytes producing IL-17 or protease receptor responses to molds and dust mites)   |  |
| Determine the roles of adaptive versus innate immune mechanisms on asthma development, progression, and response to treatment in older adults   |  |
| Determine whether there are environmental pollutants peculiar to institutional settings   |  |
| Identify viruses and other microbiological agents responsible for, and the mechanisms by which they cause, asthma exacerbations in older adults, which might lead to the development of vaccine- or antiviral drug-based interventions  |  |
| Determine effects of asthma medications, viral or bacterial load, or allergy status on susceptibility to exacerbations in older patients  |  |
| Define rates of infection and specific pathogens in older asthmatic patients  |  |
| Distinguish roles of innate immunity in eosinophilic versus neutrophilic asthma   |  |

not suggests an inherent or acquired variability in homeostatic pathways. Recent evidence from observational studies has raised suspicion that dysregulated inflammatory processes are involved in, if not central to, the variable patterns of aging. Increased serum levels of certain proinflammatory cytokines, most notably IL-6, are increasingly present with advancing age and to a greater extent with frailty.<sup>27,28</sup> Furthermore, the appearance of this and other inflammatory markers has been associated with a number of adverse clinical outcomes, including decreased strength and mobility, falls, dementia, and mortality.<sup>27</sup>

### Life expectancy, lifespan, and maximum survival

From the perspective of those who study aging, there is an important distinction made between median (life expectancy) and maximum lifespan. Over the past several decades, with the advent of modern sanitation, refrigeration, and other public health measures, including vaccination and antibiotics, there has been a dramatic increase in median survival.<sup>29</sup> Early deaths have been diminished, and more patients are reaching old age. In the United

States today, life expectancy now approaches 80 years.<sup>30</sup> Median survival is what concerns public health officials and health care providers, but for those studying the biology of aging, it is maximum survival that is the focus of greatest attention. It is worthwhile to note that it has been estimated that if atherosclerosis and cancer were eliminated from the population as a cause of death, about 10 years would be added to the average lifespan, yet there would be no change in maximum lifespan.<sup>31</sup>

Although several theories have been proposed, none suffice to account for the complexities of aging. Lifespan is finite and varies generally from species to species and much less so within species. Mice live, on average, 2½ years, monkeys 30 years, and human subjects about 90 years. Among species, larger animals generally live longer than smaller animals, but within species, smaller animals are likely to live longer. It is clear that aging is not entirely explained by DNA sequence. For example, mice and bats have only 0.25% difference in their primary DNA sequence, but bats live for 25 years, 10 times longer than mice. A commonly held notion is that regulation of gene expression accounts for a longevity difference between species.

It is now clearly established that certain specific genes can alter lifespan, at least in lower animals, but whether these same genes regulate “aging” is still in question. For example, transgenic *Drosophila* species expressing increased copies of the free radical scavenging enzymes superoxide dismutase and catalase live on average a third longer than the appropriate controls.<sup>32</sup> In even lower species (eg, yeast and nematodes) the identification of specific genes that influence lifespan<sup>33,34</sup> has led to the optimistic impression that analogous genes in higher organisms will lead greater insights into the aging process. Yet the identification and functional analysis of analogous genes in human subjects remains elusive.

The oldest human being alive today is approximately 120 years old. What is intriguing is that the record has remained stable and unchanged by the public health initiatives mentioned above. In fact, there has been some recent data presented that the maximum survival is actually decreasing in the United States.<sup>35,36</sup> What is interesting is that, unlike the public health initiatives in human subjects in which median but not maximum survival has been enhanced, experimental interventions in lower species have resulted in prolongation of maximum survival. As mentioned above, transgenic *Drosophila* species producing extra copies of superoxide dismutase and catalase survived about 33% longer than controls,<sup>32</sup> and similarly, the maximum survival in C57BL/6 mice fed a calorically restricted diet enhanced by 50% or more.<sup>37,38</sup> The true mechanisms of aging might well be uncovered with a better understanding of how these interventions affect longer survival.

Future research in aging should attempt to improve our understanding of the basic biology of aging and interventions that retard the aging process. There is a need for the development and application of a standardized definition of frailty for future clinical investigation. Investigations directed at the role of comorbidities in accelerating the aging process are important. Furthermore, future research should focus on the development of cellular and animal models of typical, delayed, and accelerated aging and of large collaborative networks in which populations and resources can be shared to study aging and frailty. Leveraging on well-characterized existing cohorts, when possible, is recommended.

## THE AGING LUNG

The lungs, like other organs, age and exhibit continued loss of function as a person grows old. Lung function is traditionally assessed by means of a number of standardized methods. The most common measurement used is spirometry with the determination of FEV<sub>1</sub> and forced vital capacity (FVC). FEV<sub>1</sub> and FVC both show continuous decreases of between 25 and 30 mL with each year of life after about age 20 years.<sup>39</sup> The cause of this decrease is usually attributed to the loss of the driving forces for airflow as a result of reduced respiratory muscle performance, loss of static elastic recoil, or both.<sup>39,40</sup> The decrease in FEV<sub>1</sub> in asthmatic patients is largely a function of the decrease in FVC because of the increase in residual volume.<sup>41</sup> Stiffening of the chest wall and reduced respiratory muscle performance result in a decrease in total lung capacity and an increase in residual volume because of ever-increasing closing volume.<sup>42</sup> Accordingly, these aging processes lead to airflow limitation that might be hard to distinguish from an active disease process.

Not all older persons are able to perform spirometry, especially those with decreased cognition, coordination, and frailty. In

addition, spirometry is effort dependent, and the very old can tire quickly. Techniques of imaging and measures of lung function not requiring effort (ie, forced oscillation) should be used in future studies to extend our knowledge about lung structure-function relationships at the very end of life.

Bronchodilator responses are known to be less marked in the elderly, perhaps as a consequence of the aging effects attributed to the emphysema-like state of the senile lung<sup>43</sup>; however, this would not explain the slow temporal response to bronchodilators. Other studies do not find such age-related bronchodilator differences. Furthermore, although methacholine responsiveness has been reported to increase with aging, the exact mechanism for this is not apparent.

Increased incidence and prevalence of many lung diseases occur with age. Alterations in immune function increase the risk of many of these diseases. Studies of systemic immunity suggest that sustained antigenic stress over a lifetime leads to a decrease in naive T-cell numbers, an accumulation of memory T cells, and a decrease in T-cell repertoire and B-cell functions but a lesser decrease in innate immunity.<sup>43,44</sup> Little is known about what happens to the immune/inflammatory pathways in older asthmatic patients. The immune system changes seen with aging will be discussed in more detail in the section on pathophysiology.

## IMPACT OF ASTHMA IN THE ELDERLY

In the United States the National Health Interview Survey asks questions regarding lifetime history of asthma, current asthma prevalence, and asthma attacks in the last 12 months.<sup>3</sup> For all age groups, asthma prevalence has been steadily increasing since 1980. For the 65 years and older age group, asthma is consistently more prevalent in female than male subjects.<sup>3</sup> The National Center for Health Statistics tracks data on physician encounters for asthma. The National Ambulatory Medical Care Survey reported that those 65 years or older have the second-highest rate of outpatient office visits after those aged 0 to 4 years. Those 65 years and older did not have significantly different emergency department visits than the other adult groups. The 65 years and older age group accounts for a greater proportion of hospitalizations (23%) than the size of its population (13%) would indicate.

Not surprisingly, the elderly population is a high user of medical resources for the treatment of asthma. Hospitalizations and emergency department visits are more common for these patients than for other adult cohorts. Some of the increased costs are related to comorbid disease. For example, the presence of comorbid chronic obstructive pulmonary disease (COPD) increases the risk of an asthma-related hospitalization in Medicare patients 3.6-fold, respiratory medical costs almost 6-fold, and total medical costs 2-fold. Elderly female subjects appear at greater risk than elderly male subjects.<sup>45-48</sup>

Asthma mortality increased steadily from 1980 until it peaked in 1998. The highest mortality rates for asthma occur in the 65 years and older group. In fact, the increase in asthma mortality between 1979 and 1989 was primarily driven by the 65 years and older group. In addition, the decrease in asthma mortality between 1999 and 2005 was most evident in this age group. Elderly women with asthma tend to have higher mortality rates than elderly men with asthma.

One reason for the increasing prevalence of asthma in the elderly might be the improved longevity of the population. Also, increased office visits for asthma in the elderly might be

**TABLE II.** Possible mechanisms of asthma in the elderly

|     | Age of onset                 | Genetic role   | Infection  | Allergy  | Inflammation   | Environment  |
|-----|------------------------------|--|--|----------|--|--|
| LSA | Child or young adult (<40 y) | Likely gene-environment  | Viral: rhinovirus and RSV  | Likely   | T <sub>H</sub> 2 driven, eosinophilic  | Allergens, day care and school, workplace                  |
| LOA | Adult (>40 y)                | Likely epigenetic, including oxidative stress, shortened telomeres | Viral: RSV, influenza, and bacterial (eg, <i>Chlamydia pneumoniae</i> ), microbial superantigens | Unlikely | T <sub>H</sub> 1 or T <sub>H</sub> 2 driven, neutrophilic and/or eosinophilic, innate immunity, T <sub>H</sub> 17, proteases | Workplace, dwelling type (house, apartment, institutional) |

responsible for fewer attacks. Increasing hospital admissions might account for decreased mortality. By continuing to gather surveillance data on asthma, reasons for these trends could become clearer. In addition, surveillance data help to focus intervention efforts in areas of greatest need.

In the Cardiovascular Health Study, a large community-based cohort of subjects older than 65 years, questions were asked that were relevant to asthma and provided more insight into the prevalence and effect of asthma in this population.<sup>5,49,50</sup> *Definite asthma* was defined as a positive response to the questions indicating that the patient had current asthma and that a physician confirmed the diagnosis. *Probable asthma* was defined as a history of wheezing in the past year associated with chest tightness or breathlessness. Excluding smokers and those with a diagnosis of congestive heart failure, 4% of subjects had definite asthma, and 4% had probable asthma. Among those who smoked, 11% had definite asthma, and 14% had probable asthma. Among non-smokers, 185 subjects were identified who had definite or probable asthma; 76% were women, and 20% were older than 80 years. The age of asthma onset was spread approximately evenly among decades. Twenty-seven percent had late onset of disease after age 60 years, and 25% had onset of disease before age 20 years. As expected, respiratory symptoms in the older asthmatic subjects were more prevalent, with a 2- to 5-fold increase in cough, phlegm, wheezing, and dyspnea. Dyspnea on exertion was 1.6-fold more likely to be present in asthmatic patients than in those without the diagnosis. Lung function was reduced in those with a diagnosis of asthma. Mean FEV<sub>1</sub> was 77% of predicted value in those with definite asthma and 89% of predicted value in those with probable asthma compared with 96% in those who did not have asthma. Forty-one percent of those with a diagnosis of asthma had airflow obstruction below the fifth percentile for the age group, and peak flow lability was increased. Elderly asthmatic patients reported the most common trigger was a viral infection in 58% compared with animal allergies in 30%. Two thirds reported seasonal worsening. Asthma had a significant effect on quality of life, with 35% of patients with definite or probable asthma reporting a fair or poor health status compared with 17% of elderly patients without asthma. Sixty percent of patients with definite asthma reported seasonal allergic rhinitis compared with only 30% in the nonasthma group. Despite the high prevalence and morbidity of asthma in this population, inadequate treatment was common. Only 40% of those with definite asthma had a rescue albuterol inhaler, and only 30% had inhaled corticosteroid use.<sup>5,49,51,52</sup>

## PATHOPHYSIOLOGY AND RISK FACTORS

The pathophysiology of asthma in the older adult is poorly understood and understudied. Many questions about this issue

remain: Is asthma the same disease in older adults as it is in children and younger adults? Is late-onset asthma (LOA; asthma that starts in middle age or older) different from longstanding asthma (LSA; asthma of early onset that has persisted into older adulthood)? If LOA and LSA are the same disease, then the diagnosis and treatment should be similar. However, if LOA and LSA are different phenotypes or at least have a different cause and pathophysiology, then the diagnosis and treatment might differ (Table II).

## Epigenetics, aging, and asthma

The traditional view of disease susceptibility has been expanded to include epigenetics to account for the influence of environmental factors and aging on the genomic blueprint. *Epigenetics* is defined as heritable changes in gene expression that occur without alterations in DNA sequence. It is the process by which genotype interacts with environment to produce a phenotype and explains differences between cells, tissues, and organs despite identical genetic information. Genes function in a milieu determined by the developmental and environmental history of the cell, which constitutes the epigenotype.<sup>53,54</sup> Epigenetic changes or marks can play a major role in human disease.<sup>55,56</sup> The most common examples of epigenetic marks are DNA methylation of CpG islands by DNA methyltransferases and chromatin modification of histone proteins, particularly acetylation by histone acetyltransferases and histone deacetylases.<sup>57,58</sup> The function of epigenetic changes is to regulate gene expression. Epigenetic changes are known to contribute to cancer and autoimmune disease and are thought to contribute to common diseases, including cardiovascular disease, diabetes, and the loss of response to stress caused by aging.<sup>59</sup>

Asthma is a markedly heterogeneous disease, and recent evidence suggests that environmentally induced epigenetic changes contribute to asthma phenotypes and that airway inflammation in patients with asthma and COPD might involve epigenetic regulation.<sup>57,58,60</sup> Methylation patterns and chromatin structure change with age and are thought to contribute to the increase in the incidence of common diseases that begin in middle age.<sup>55,61</sup> The incidence of asthma in the elderly resembles the incidence of common diseases. Moreover, characteristics and asthma drug response in the elderly asthmatic patient differ from those seen in childhood asthma. Compared with younger cohorts, elderly asthmatic patients have a higher prevalence, higher rates of bronchial hyperreactivity, more severe asthma, and a lower prevalence of atopy. The symptoms of elderly asthmatic patients are more difficult to control with drug therapy, and these patients have steroid resistance and might respond better to leukotriene receptor antagonists compared with inhaled corticosteroids.<sup>62-71</sup> The contribution of epigenetics to differences

observed between elderly asthmatic patients and younger cohorts is unknown. Unlike genetic variants that contribute to disease, epigenetic changes can be reversed and therefore represent potential drug targets.<sup>72</sup>

### Role of airway inflammation

Older asthmatic patients are less responsive to albuterol treatments given in the emergency department and are more frequently admitted for hospitalization.<sup>73</sup> Thus it appears that the responsiveness to treatment is diminished and the severity of asthma exacerbations is greater. The exact reason for these disparities is not known.

Immune cell function decreases with aging, a property often termed *immunosenescence*.<sup>74</sup> One often-confusing aspect of immunosenescence is the observation that aging might be associated with opposing immunologic effects. For example, T-cell secretion of IL-2, IL-4, or IFN- $\gamma$  has been shown to be both decreased with aging and also increased with aging.<sup>75</sup> It is likely that both phenomena are correct but are dependent on the context of the immune function. Thus the effect of aging on T-cell function in the context of allergen stimulation might be different than the effects of aging on T-cell function in the context of viral infection. Given that asthma is an inflammatory disorder of the airway, it is of interest to determine whether asthmatic airway inflammation of the elderly might differ from that of younger asthmatic patients and thus represent a distinct phenotype of asthma. These changes might have implications for susceptibility to exacerbations because of viruses or other pathogens, as well as response to treatment.

The aging process has been shown to exhibit changes in airway inflammation. An examination of the cellular composition of bronchoalveolar lavage fluid from 19- to 83-year-old subjects without a history of allergies, pulmonary disease, or gastroesophageal reflux showed increased airway neutrophilia, as well as increased numbers of CD4<sup>+</sup> T cells.<sup>76,77</sup> The T cells also appeared to be more activated in the elderly, with increased expression of HLA-DR and CD69. The increase in airway neutrophils with aging has also been observed in asthmatic patients.<sup>78</sup> Because there is a phenotype of severe asthma characterized by a predominantly neutrophilic airway inflammation,<sup>79</sup> the question arises as to whether the increased presence of neutrophils contributes to greater asthma severity in the elderly.

Some of the prominent inflammatory cells recruited into the airway in asthmatic patients are eosinophils, neutrophils, and T cells, which are capable of secreting numerous inflammatory mediators, including leukotrienes and cytokines. It is not known whether immunosenescence affects the production of these mediators in elderly asthmatic patients, either at baseline or during an exacerbation of symptoms. Furthermore, it is not known whether age-related changes in their production would have any implication for the clinical presentation or management of asthma in the elderly. Peripheral blood eosinophils were isolated from younger (20-40 years old) and older (55-80 years old) subjects for *in vitro* functional assays. The eosinophil effector functions of degranulation and superoxide production were diminished in the older compared with the younger asthmatic patients.<sup>78</sup> In another study examining the expression of neutrophil mediators in younger and older asthmatic patients, there was decreased baseline expression of leukotriene B<sub>4</sub> in the sputum of older asthmatic patients despite greater numbers of neutrophils.<sup>80</sup> Whether these

findings have implications during an asthma exacerbation has yet to be determined. Nevertheless, the results demonstrate age-related changes in the function of an inflammatory cell considered pathognomonic for allergic asthma and raise the question of whether additional effects of immunosenescence are relevant to airway inflammation in asthmatic patients.

There have been several studies of animal models to address age-related changes in the airway inflammation induced by allergen challenge of sensitized aged animals (see Experimental approaches section below). These studies yielded conflicting results, and there is concern that the animal models do not accurately represent the chronic features of human asthma with seasonal allergen exposure and intermittent exacerbations. The aged animals were both sensitized and challenged at old age, which is in contrast to the typical elderly human asthmatic patient who might be exposed to allergens for several decades.

### Role of allergy

Typically, nasal and ocular symptoms on exposure to allergens diminish with age. Allergen-triggered asthma symptoms also diminish with age. The Epidemiology and Natural History of Asthma (TENOR) study examined the natural history of asthma in older (>65 years old) compared with younger patients and found that older asthmatic patients had lower total IgE levels, fewer positive skin prick test responses, and less concomitant allergic rhinitis or atopic dermatitis.<sup>81</sup> Several studies have demonstrated age-related decreases in total IgE and allergen-specific IgE levels,<sup>82-87</sup> suggesting that this might be the explanation for the decrease in allergy symptoms. There is also evidence for an age-related decrease in skin prick test responses to allergens.<sup>88</sup> However, the relationship between total IgE levels and allergic disease persists in the elderly, such that subjects with greater IgE levels remain more likely to have allergic rhinitis or asthma.<sup>89,90</sup>

Given the changes in allergic inflammation with aging, one might conclude that asthma in the elderly should be milder. However, there are several other common triggers for exacerbations of asthma, including irritants (eg, cold air) and respiratory tract infections. Estimates suggest that up to 80% of asthma exacerbations in adults are caused by viral upper respiratory tract infections.<sup>91</sup>

The role of environmental exposures and allergy in older asthmatic patients is largely unknown. In the general population, evidence regarding the effect of indoor pollution on asthma is summarized in "Clearing the air: asthma and indoor air exposure" by the Institute of Medicine for the Environmental Protection Agency published in 2000.<sup>92</sup> Evidence was reported for asthma development related to house dust mites and for asthma development associated with environmental tobacco smoke in preschool children. The report also showed evidence for causation of asthma exacerbations for house dust mite, environmental tobacco smoke (in preschool children), cat, and cockroach; an association with exacerbations was found for dogs, fungi, formaldehyde, and rhinovirus. In addition, evidence associating exacerbation of asthma-related symptoms with self-reported damp air was reported in a review of damp indoor spaces and health.<sup>93,94</sup>

There are only a few studies that have evaluated the role of atopy in elderly patients with asthma. One large national study of allergy skin tests that included older adults<sup>95</sup> and several small studies of allergy skin tests in older adults with asthma<sup>81,96-103</sup>

**TABLE III.** Studies examining allergy in patients with early-onset asthma versus those with LOA

| Location/date                        | No. | Mean age (y) | Mean age of onset (y) | Positive skin test response (%) | Positive serum specific IgE level (%) |
|--------------------------------------|-----|--------------|-----------------------|---------------------------------|---------------------------------------|
| Providence, RI, 1991 <sup>97</sup>   | 25  | >70          | <43 (n = 13)          | 62                              | ND                                    |
|                                      |     |              | >70 (n = 12)          | 0                               |                                       |
| Rochester, Minn, 1997 <sup>102</sup> | 63  | >65          | <40 (n?)              | 56                              | ND                                    |
|                                      |     |              | >41 (n?)              | 21                              | ND                                    |
|                                      |     |              | >65 (n?)              | 20                              | ND                                    |
| Boston, Mass, 1997 <sup>101</sup>    | 46  | 61           | 49 ± 15.7             | ND                              | 24 (cat)                              |
|                                      |     |              |                       |                                 | 21 (Der p 1)                          |
| Boston, Mass, 1997 <sup>101</sup>    | 33  | 61           | 61                    | ND                              | 18 (cat)                              |
|                                      |     |              |                       |                                 | 21 (Der p 1)                          |

ND, Not done.

were reviewed. Allergy skin test results were positive in 8% to 12% of all older adults. The prevalence of positive skin test results or specific IgE levels to at least 1 allergen in older adults with asthma ranged from 0% to 75%. Those whose asthma had an unknown age of onset ranged from 27% to 60%, those with onset before age 41 years ranged from 56% to 62%, and those with onset greater than age 41 years ranged from 0% to 24% (Table III).<sup>97,101,102</sup>

Although there were no studies of allergen room exposure or bronchial challenge in older adults, neither prick-puncture skin test results nor specific IgE levels predicted the nasal challenge response to dust mites.<sup>104</sup> Safety concerns for allergen challenges in older adults are unresolved. Technical limitations of allergens, environmental measurements, and age-specific norms and cutoff levels for laboratory and physiologic tests are needed.

A few epidemiologic studies suggest an association between outdoor environmental exposures and emergency department or hospital admissions in older adults.<sup>105,106</sup>

In summary, studies of the general population suggest a causation or association between indoor air pollutants and allergy exposure and asthma. There are several small studies suggesting higher levels of positive allergy test results in older adults with asthma than in the general population of older adults. When age of onset is considered, asthma with an early onset (<41 years of age) has a much higher association with positive allergy test results than late-onset asthma.

### Role of viral respiratory tract infections

Viral respiratory tract infections are common precipitants of asthma exacerbations during childhood. In approximately 80% of children with acute asthma exacerbations, a respiratory tract virus can be detected, with rhinovirus being the most frequent pathogen identified.<sup>107</sup> Although it is likely that viruses also lead to exacerbations of asthma in older adults, comprehensive studies regarding the rates and specific pathogens are lacking. Several issues have made defining the role of viruses in adult asthmatic patients problematic and include difficulty distinguishing COPD from asthma, lack of sensitive diagnostic tests, and issues with asymptomatic infection. A number of investigators have explored the incidence of viral infection in adult asthmatic patients.<sup>91,108,109</sup>

Older studies using viral culture and serology for diagnosis demonstrated infection rates of 10% to 29%.<sup>109</sup> In contrast, more recent studies, which include RT-PCR, have shown significantly higher infection rates of 44% to 55%.<sup>91,108</sup> Similar to results found in children, rhinoviruses are the most frequently detected pathogen. Few older persons were included in these studies, in which the mean ages of subjects were 30 to 39 years.

The incidence of acute respiratory tract infections decreases steadily with advancing age, and rates of viral infection in older adults are influenced by place of residence.<sup>110</sup> Among community-dwelling older adults, rates of acute respiratory tract infections are roughly 1% to 2% per year, whereas rates in senior day care centers and long-term care facilities are substantially higher at 6% to 11%.<sup>111</sup> In addition, the epidemiology of respiratory tract infections can be quite complex in these semiclosed populations, with multiple pathogens circulating simultaneously.<sup>112</sup> Influenza A, respiratory syncytial virus (RSV), and human metapneumovirus (hMPV) are the most commonly identified viruses among older persons hospitalized with acute cardiopulmonary conditions.<sup>113,114</sup> Most patients who require hospitalization during a viral respiratory tract infection have underlying heart and lung conditions. Although studies to date have largely focused on the role of viruses in COPD exacerbations, it is reasonable to extrapolate infection rates and specific pathogens from these studies to older adults with asthma. Johnston,<sup>115</sup> in Ontario, Canada, found a seasonal peak in emergency department visits for all acute respiratory tract infections, as well as exacerbations of both COPD and asthma, for persons younger and older than 50 years. All the common respiratory tract viruses have been associated with COPD exacerbations, and depending on the methodology and season of study, the specific rates of influenza, RSV, parainfluenza viruses, coronaviruses, hMPV, and rhinoviruses vary.<sup>116,117</sup> Wheezing appears to be a common symptom in older adults infected with any of the respiratory tract viruses, particularly RSV and hMPV, and 7% of adults hospitalized with RSV will have a discharge diagnosis of asthma.<sup>114,118</sup>

Because most adult infections represent reinfection, the viral load in respiratory secretions tends to be low, making detection with conventional techniques difficult. Viral culture and rapid antigen testing, which can be used successfully in children, have poor sensitivity in older adults. The use of molecular diagnostics has vastly improved the ability to detect a number of viruses, such as RSV, parainfluenza, and rhinoviruses, and allows the detection of hitherto uncultivable agents, such as hMPV and coronaviruses.

Viral infections appear to aggravate reactive airway disease through a number of different mechanisms. It has been postulated that viral infection disrupts the negative feedback loop of acetylcholine on the M2 receptor, leading to increased levels of acetylcholine and increased constriction of bronchiolar smooth muscle.<sup>119</sup> Infection of the respiratory epithelium also induces chemokines, cytokines, and immune and growth factors, which result in a proinflammatory state.<sup>120,121</sup> Immunosenescence might affect the ability of older adults to clear viruses efficiently, and thus greater and more prolonged inflammation can result.

In summary, respiratory viral tract infections are common among older adults and are likely precipitants of acute asthma exacerbations. Furthermore, viral respiratory tract infections might likely precipitate the onset of LOA, although this needs to be further examined.<sup>122</sup> Comprehensive studies regarding the rates and specific pathogens are lacking in older adults. Distinguishing COPD from asthma, lack of sensitive diagnostic tests,



and issues with asymptomatic infections make it difficult to define the role of infections in older adults.

## DIAGNOSIS OF ASTHMA IN THE ELDERLY

### Clinical assessment

Classic symptoms of asthma in the elderly are mostly similar to those seen in younger asthmatic patients.<sup>4,96</sup> Data on the clinical features of asthma in the elderly have been derived from both longitudinal community surveys and case studies.<sup>5,64,97,98,122-124</sup> Most patients complain of episodic wheezing, shortness of breath, and chest tightness. These symptoms are often worse at night and with exertion and, like those in younger asthmatic patients, are often precipitated by an upper respiratory tract infection. In fact, the majority of elderly patients who have asthma after age 65 years have their first asthmatic symptom preceded immediately by or concomitant with an upper respiratory tract infection.<sup>122</sup> Asthma can often be triggered by environmental exposures, such as aeroallergens, irritants (cigarette smoke, household aerosols, and paints), strong odors (perfumes), and inhalation of metabisulfites (found in beer, wine, and food preservatives). Asthmatic symptoms can also be triggered by medications, such as aspirin, nonsteroidal anti-inflammatory agents, angiotensin-converting enzyme inhibitors, or  $\beta$ -blockers, which are commonly used by this patient population. This emphasizes the need for the physician to perform a comprehensive review of medications taken by the older asthmatic patient.

Studies have consistently shown that elderly patients and their physicians frequently overlook symptoms caused by asthma.<sup>5,73,125</sup> Several factors contribute to the underdiagnosis and misdiagnosis of asthma. One reason, as shown in large community studies, is that most patients first have asthma in childhood or adolescence, and many physicians have had the misconception that asthma is a childhood disease. Another important reason is that the symptoms of asthma are more commonly associated with other diseases seen in this age group. The symptoms of asthma in the elderly are therefore nonspecific and might be caused by conditions that mimic asthma. The differential diagnosis of asthma in the elderly is greater than seen in younger asthmatic patients and includes congestive heart failure, emphysema and chronic bronchitis (COPD), chronic aspiration, gastroesophageal reflux disease (GERD), and tracheobronchial tumors. Comorbid illnesses and the psychosocial effects of aging might also profoundly affect the diagnosis, clinical presentation, and care of asthma in the elderly. One particular diagnosis that is often difficult to detect and frequently overlooked by the patient and physician until the condition is advanced is upper airway obstruction, including the extrathoracic and intrathoracic central airways. Common causes of upper airway obstruction include malignancy, infection, inflammatory disorders, trauma, and extrinsic compression related to enlargement of adjacent structures (eg, an enlarged thyroid gland). It appears that malignancy and benign strictures related to airway instrumentation (eg, endotracheal intubation and tracheostomy) are becoming increasingly more prevalent in the older age group.

Distinguishing chronic asthma from COPD can be very challenging, and in some patients asthma cannot be distinguished from COPD with widely available diagnostic tests. The management of these patients might have similarities to that of asthma. The distinction between LOA and COPD can be

difficult to define precisely. The Lung Health Study showed that methacholine-induced airways reactivity is present in many patients with mild-to-moderate COPD (ie, 63% of men and 87% of women). Approximately 85% of patients with tobacco-related COPD demonstrate bronchodilator reversibility at least once on repeated testing sessions. The distinction between COPD and asthma can be confounded by either the coexistence of the 2 common disease entities, the progression of common pathobiologic mechanisms induced by different environmental agents, or different disease mechanisms leading to an overlapping clinical syndrome.

It has been known for more than a century that early-morning wheezing is a prominent symptom of congestive heart failure. It has been called cardiac asthma because it can mimic the clinical picture of typical asthma. The usual symptoms of gastroesophageal reflux in the elderly, such as vomiting and heartburn, might be absent. In a study of elderly patients with esophageal reflux proved by means of intraesophageal pH monitoring, chronic cough, hoarseness, and wheezing were present in 57% of patients.<sup>126</sup> In addition to causing asthma-like symptoms, there is also evidence that GERD might be a cause of worsening asthma.

Shortness of breath is a common symptom in the elderly and is most commonly caused by heart or lung diseases. It is usually experienced during exertion. Shortness of breath at rest is not typical of heart disease or lung diseases, such as COPD or interstitial lung disease, except in advanced stages. When present, it should prompt an investigation for asthma because sudden bronchospasm can cause respiratory distress at rest or exercise. Paroxysmal nocturnal dyspnea, which is typical of congestive heart failure, is found in a smaller number of elderly patients with asthma. Many elderly patients limit their activity to avoid experiencing dyspnea, and others assume that their dyspnea results from their aging process and thus avoid seeking medical attention early in their disease process. However, aging *per se* does not cause dyspnea, and a cause needs to be always pursued in assessing an elderly patient who complains of breathlessness.

There are several other reasons why the diagnosis of asthma in the elderly might be delayed or not made at all. Elderly patients have been shown to have a reduced perception of bronchoconstriction,<sup>127</sup> and this might delay medical intervention. Many elderly patients are fearful of having an illness and dying and are reluctant to admit they are having symptoms. Underreporting of symptoms in the elderly might have many causes, including depression, cognitive impairment, social isolation, denial, and confusing symptoms with those of other comorbid illnesses.

Cough is a very prominent symptom and might occasionally be the only presenting symptom. Wheezing, on the other hand, might not be as prominent, and its presence is not very specific and does not correlate with severity of obstruction. Physical examination in elderly patients with asthma is usually nonspecific and might misguide the diagnosis: a negative examination result does not rule out asthma, and wheezing can be found in a number of conditions, such as COPD, recurrent aspiration, and "cardiac asthma" (congestive heart failure).

Two distinct clinical presentations have been described for asthma in the elderly. These are based on the onset and duration of the disease state.<sup>97,124,128</sup> Patients with LOA start having asthma symptoms for the first time when they are 65 years of age or older (some studies have suggested middle age or older). Some studies of elderly asthmatic patients have shown that, as a group, as many as 40% will have their first attack after the age of 40 years.<sup>97,98,129</sup>

Patients belonging to this group tend to have fewer atopic manifestations, higher baseline FEV<sub>1</sub>, and a more pronounced bronchodilator response than those with LSA. Patients with LSA start having asthma symptoms early in life. Patients belonging to this group tend to have a higher incidence of atopic diseases, more severe and irreversible or partially reversible airway obstruction, and more hyperinflation. The duration of the disease in this group is an important determinant of severity and of the development of irreversible airflow obstruction.<sup>128</sup>

Longitudinal studies of asthmatic populations, whether new onset or long standing, have shown that remission from asthma is uncommon in older groups, occurring in less than 20% of patients.<sup>130</sup> This contrasts with asthma in children and adolescents, in whom remission of asthma symptoms is common, especially in the second decade of life, and might be seen in as many as 60% to 70% of patients.

### Physiological assessment

Objective measures to confirm the diagnosis of asthma are uncommonly performed in primary care settings. Inhalers are prescribed for patients who are evaluated for asthma-like symptoms, and during a follow-up visit, the patient is asked whether the controller inhaler reduced the frequency of asthma symptoms or whether the albuterol inhaler quickly relieved the symptoms. Such an empiric approach might work most of the time for young patients with mild asthma but is more likely to result in an incorrect diagnosis, poorly efficacious treatment, or unnecessary medication side effects in older patients.

The onset of wheezing, shortness of breath, and cough in an elderly patient is likely to cause concern. Although the adage “all that wheezes is not asthma” is true at any age, it is especially true in the elderly. Diagnosis based on objective measures is essential. Moreover, lung function testing, even in the presence of minimal symptoms, is especially important in this age group because there is thought to be an age-related reduction in the perception of exertional dyspnea in the elderly.<sup>125</sup> An older patient with chronic, untreated, severe airway obstruction caused by asthma might reduce activity to avoid dyspnea and stoically deny impairment of activity. This might reflect either neurocognitive function or changes in lifestyle that favor sedentary activities.

There exist some barriers to lung function testing in the elderly. Spirometry might be difficult to perform in some situations because of physical or cognitive impairments. However, 80% to 90% of elderly persons are able to perform good-quality spirometry when tested by skilled technologists.<sup>129-133</sup> The Global initiative for Obstructive Lung Disease guidelines for diagnosing the airway obstruction of COPD by using a fixed FEV<sub>1</sub>/FVC ratio of less than 0.70 caused a high misclassification rate in older persons.<sup>134</sup> However, almost all computerized spirometers automatically calculate the appropriate lower limit of the normal range for FEV<sub>1</sub>/FVC ratio and for FEV<sub>1</sub> by using race-specific National Health and Nutrition Examination Survey III reference equations.

In addition, it is hard to define the lower limits of predicted normal values in this age group. Although complete reversibility of airflow obstruction is frequently seen with young asthmatic patients, most elderly asthmatic patients show incomplete reversibility despite continuous intense therapy, and many show fixed airflow obstruction as if they have COPD. However, objective measures of lung function, such as spirometric and peak flow

measurements, are generally underused in elderly patients, and this also contributes to the delay or absence of diagnosis.<sup>134,135</sup>

Lung function testing is especially important in this age group because of the age-related reduction in the perception of dyspnea seen in the elderly.<sup>127</sup> Spirometry is easily performed to determine that FEV<sub>1</sub> and FEV<sub>1</sub>/FVC ratio are demonstrated with the timed vital capacity maneuver. The flow-volume loop, which also measures inspiratory flow, is especially useful when the cause of respiratory tract symptoms is not known and an upper airway obstruction is in the differential diagnosis. Although it might be difficult to perform spirometry in the elderly in some situations because of physical and poor cognitive impairment, studies have demonstrated that between 82% and 93% of elderly patients are able to perform the test properly.<sup>131-135</sup> On the other hand, it might be more difficult to define the lower limits of predicted normal values in this age group. Traditionally, an FEV<sub>1</sub>/FVC ratio of less than 70% increases the probability of asthma in an elderly patient with asthma symptoms, but this ratio normally decreases with age because of a decrease in elastic recoil, and a ratio lower than 70% might be a normal finding.<sup>136</sup>

A brisk response to a short-acting bronchodilator might demonstrate the second cardinal feature of asthma: reversible airflow obstruction (ie, “a responder”). When airflow obstruction is found in an elderly patient, attempts should be made to demonstrate reversibility after the inhalation of a short-acting  $\beta$ -adrenergic agent, such as albuterol. Evidence of reversibility (postbronchodilator FEV<sub>1</sub> or FVC increases of >12% and 200 mL) increases the probability of a diagnosis of asthma. Elderly asthmatic patients, however, might have an impaired  $\beta$ -agonist bronchodilator response because the number of  $\beta$ -adrenergic receptors on smooth airway muscles is decreased with aging.<sup>137</sup> Although the bronchodilator response to inhaled  $\beta$ -agonists decreases with age,<sup>138</sup> this is not the case with anticholinergic agents.<sup>139</sup>

Airway obstruction might be absent at the time of testing, and further testing might be needed to facilitate the diagnosis. Bronchoprovocation testing with a methacholine challenge can be useful, and it is a safe and effective method to uncover asthma in older adults.<sup>140,141</sup> A negative test result will rule out asthma; a positive test result must be interpreted and include an assessment of pretest probability.<sup>142</sup> In addition, some studies have shown that bronchial responsiveness is heightened in older adults, and therefore aging might be an independent factor that influences airway responsiveness.<sup>143</sup> There is a relationship between the degree of bronchial hyperresponsiveness and prechallenge pulmonary function; a low FEV<sub>1</sub> predicts heightened responsiveness.<sup>144</sup> Other factors that might contribute to heightened airway responsiveness in the older population are atopy and current or previous smoking history.

Peak expiratory flow variability might be helpful in the diagnosis and follow-up of younger patients with asthma, but poor coordination and muscle weakness in some elderly patients might lead to an inaccurate reading.<sup>52,145</sup> A prospective study did not demonstrate any advantage of peak flow monitoring over symptom monitoring as an asthma management strategy for older adults with moderate-to-severe asthma when used in a comprehensive asthma management program.<sup>146</sup> Other tests, such as measuring the carbon monoxide diffusing capacity of the lung, have been advocated to distinguish between asthma and COPD because the diffusing capacity of the lung is reduced by parenchymal destruction found with emphysema. However, studies have shown that differences in lung function tests, although

statistically significant, cannot be used clinically to separate the 2 groups of subjects because of a large overlap.<sup>147</sup>

There is growing evidence that the airway function of young and middle-aged asthmatic patients decreases at a greater rate than that of healthy subjects.<sup>148-150</sup> The rate of decrease increases with increasing age and in those who smoke cigarettes.<sup>149,151</sup> In patients with LOA, there is evidence that lung function is reduced even before a diagnosis is made and decreases rapidly shortly after diagnosis.<sup>98,152</sup> Thereafter, it remains fairly stable. Although the effect on older asthmatic patients with LSA is variable, in a random survey of 1200 elderly asthmatic patients older than 65 years, only 1 in 5 patients had normal pulmonary function ( $FEV_1 >80\%$  of predicted value), whereas a similar number showed moderate-to-severe airflow obstruction ( $FEV_1 <50\%$  of predicted value) after an inhaled short-acting bronchodilator.<sup>153</sup> Because structural changes of emphysema are minimal in elderly asthmatic patients, except if they are previous smokers, airway remodeling is thought to be the main cause of fixed airflow obstruction.

### Role of nitric oxide

Nitric oxide (NO) is a gas generated by the action of NO synthase from the substrates molecular oxygen and arginine. It was originally identified as a biologically important signaling molecule with the properties of an activity previously described as endothelial-derived relaxing factor. This molecule is important in regulating vascular integrity and blood flow and is thought to be a regulator of vascular smooth muscle relaxation.

More recently, it has been found that NO can be generated by a variety of inflammatory cells, including polymorphonuclear leukocytes, mononuclear cells, and, importantly, eosinophils. This finding led to the identification of NO as a molecule present in exhaled breath. Studies of NO exhalation have found that it is increased in infection and inflammation of the airway. Although high levels of NO are found in nasally expired air, studies in pulmonary inflammation have avoided this by redirecting airflow through the oral airway. It has been found that exhaled NO reflects airways inflammation and particularly eosinophilic inflammation. Exhaled NO levels are increased during the allergy season in atopic subjects. Inhaled glucocorticoids promptly suppress exhaled NO and do so in conjunction with suppression of eosinophilic inflammatory infiltrates. Studies have demonstrated that monitoring exhaled NO might permit better regulation of asthmatic symptoms, exacerbations, and total steroid use than treatments based on guidelines or symptoms. Furthermore, increases in exhaled NO levels might predict asthma exacerbations.<sup>154</sup> It is of interest that NO levels in expired air decrease after bronchoconstrictive stimulation of asthmatic airways.

Little is known of the effects of age on NO levels in the expired air. It appears that NO production and vascular responses to NO might be diminished in the elderly, but that effect might be overcome by exercise to increase fitness. An unanswered question in airways biology is whether NO is causative of airways dysfunction, a marker for this dysfunction, or an ineffective homeostatic response to airways constriction.<sup>155-159</sup>

### Challenges in defining asthma in the elderly

There is agreement that asthma is both a common and underrecognized health problem for the elderly that leads to

impairments of lung function and quality of health and life. The first question that needs to be addressed is why we need to make such a diagnosis rather than just treat the symptoms. There are reasons that physicians must strive to assign a diagnosis to a patient with a symptom complex. The patient is given relief by letting him or her know what is wrong by giving the illness a name, which implies a cause, establishes a prognosis, and initiates a treatment plan. Moreover, advancement of the understanding of epidemiology, natural history, pathobiology, and treatment require a definable disease entity. Whether the threshold for diagnostic criteria is set at a high level of sensitivity, a high level of specificity, or a high level of accuracy depends entirely on the costs and benefits of an incorrect diagnosis versus a missed diagnosis. For example, enumeration of a disease might require a high level of accuracy, whereas diagnosis of an uncommon and difficult-to-treat disease (eg, metastatic cancer) ought to be highly specific. The diagnosis of a common and easily treatable disease (eg, vitamin deficiency) ought to be highly sensitive, even if there is a risk of overdiagnosis. Asthma tends to be one of those disorders that is relatively easy (although not inexpensive) to treat and has morbid consequences if left untreated, suggesting that the diagnostic criteria ought to be highly sensitive.

Although medical students are taught the rigorous discipline of data collection, differential diagnosis, and test confirmation, most physicians do not practice this way. In practice, physicians typically rely on a constellation of signs and symptoms along with demographic characteristics and recent experiences to establish diagnoses through the process of pattern recognition.

There are no shortages of official definitions of asthma, and modifications seem to be added every year. Most of these definitions involve the definition of a clinical syndrome (episodic cough, wheezing, and dyspnea), an underlying pathophysiology (airway hyperresponsiveness, variable, and reversible airflow obstruction), an underlying biological process (chronic eosinophilic or neutrophilic inflammation of the airways), and an associated morbid anatomy (basement membrane thickening, smooth muscle hypertrophy, and mucus cell metaplasia).

Given this, why is it so challenging to diagnose asthma in the elderly? First, the syndrome of asthma is often confused with other common diseases in the elderly, such as COPD, congestive heart failure, paroxysmal arrhythmias, pulmonary emboli, recurrent aspiration, and GERD.

Second, asthma can often coexist with these other conditions, and it can be impossible to determine which of the 2 conditions is responsible for the patient's ill health. This diagnostic confusion can be amplified by the different manifestations of asthma in the elderly. Elderly asthmatic patients can be insensitive to exertional dyspnea because of a sedentary lifestyle. They tend to be less atopic and have an incomplete response to bronchodilators. The elderly without asthma tend to show some signs suggestive of asthma: slower emptying of the lung during forced expiration, decreased lung elastic recoil, and a higher prevalence of nonspecific airways reactivity.

The hope that formal testing of airways reactivity would prove useful in diagnosing asthma has led to disappointment. In young adults a history of asthma, wheeze, or treatment for asthma plus a positive methacholine challenge test result is highly specific for asthma (99%) but misses about half of the asthmatic population.<sup>160</sup>

In epidemiologic studies that have examined various criteria for diagnosing asthma, it turns out that the solution is relatively

simple. Patients who answer yes to the question “Have you ever had asthma?” have nearly 100% specificity and 48% to 100% sensitivity when compared with those receiving an independent expert’s diagnosis.<sup>161,162</sup>

The problem of diagnosing asthma in the elderly is more complicated because of the overlap with COPD. Asthma is typically considered a disease of onset in youth driven by atopy and eosinophilic inflammation causing reversible airflow limitation. COPD, in contrast, is considered to be a disease of onset in middle age driven by cigarette smoking and neutrophilic inflammation and leading to irreversible airflow limitation. As evidence presented in this workshop has shown, asthma in the elderly displays many of the features of COPD. The disease can have its symptomatic onset late in life, often is only partially reversible, and is associated with neutrophilic inflammation. Moreover, the current cohort of elderly patients has a high prevalence of past smoking, reflecting the health habits in the United States in the 1940s and 1950s.

The failure to deal with the population of elderly patients who have overlapping signs of asthma and COPD is not just a matter of classification of disease. It has significant health consequences in that such patients are systematically eliminated from clinical trials and are not covered by treatment guidelines. Little is known about how best to treat the elderly patient with asthma who smokes or the elderly patient with COPD who has reversible airflow limitation.

This confusion is manifest by diagnostic coding in older Medicaid patients. Of those who were hospitalized with an initial diagnosis of COPD, 43% had an asthma diagnosis within 3 years. Of those who had an initial hospital diagnosis of asthma, 46% had a diagnosis of COPD within 3 years. Price et al<sup>161</sup> attempted to develop a discriminant function using clinical and demographic information that would separate patients with COPD from those with asthma by using strict physiologic criteria. Although several discriminating characteristics were found, the best diagnostic criteria were only 78% sensitive and only 75% specific.

We need to ask whether it really is important to make the distinction between asthma and COPD in the elderly in terms of prognosis or treatment. One study by Hansen et al<sup>136</sup> suggests that regardless of whether a person is given a diagnosis of asthma or COPD, the prognosis is mostly determined by the impairment in FEV<sub>1</sub>.

## MONITORING ASTHMA IN THE ELDERLY

### Standardized monitoring tools

There are a number of ways to measure the effect of asthma in both young and elderly patients. Assessments of symptoms, functional limitations, quality-of-life measures, and risk of adverse events are several that have been suggested by current asthma guidelines.<sup>163</sup> In addition, measuring a patient’s satisfaction with his or her asthma symptom control and overall asthma care has been advocated.

The use of objective measures of asthma control and satisfaction can be especially important in the elderly because the perception of symptoms might be impaired with advancing age. In addition, many elderly patients unconsciously accommodate to their symptoms or assume that the symptoms are a function of the aging process itself. Because the number of unscheduled ambulatory visits, emergency department visits, and hospitalizations are high in elderly asthmatic patients,<sup>164,165</sup> and quality-of-life

scores are low in elderly patients with persistent asthma when compared with those with mild asthma or no asthma at all,<sup>5</sup> careful assessment of asthma control is essential in this age group. Despite severe symptoms and physiologic impairment, most elderly patients with asthma can lead active productive lives if their asthma is appropriately managed. In fact, when elderly patients with severe or difficult-to-treat asthma have been identified by a physician’s assessment, they appear to do better than younger patients. In the TENOR study, despite lower lung function, older asthmatic patients (mean age, 72 years) had lower rates of unscheduled office visits, emergency department visits, and corticosteroid bursts.<sup>81</sup> Patients reported in the TENOR study received more aggressive care than younger adults, including higher use of inhaled and oral corticosteroids, and this undoubtedly had an effect on outcomes.

The tools to measure asthma outcomes include questionnaires and other self-report tools, such as diaries and standardized medical history forms. Standardized questionnaires that assess asthma impairment include the Asthma Control Test,<sup>166,167</sup> the Asthma Control Questionnaire,<sup>168,169</sup> the Asthma Therapy Assessment Questionnaire,<sup>170,171</sup> and others.<sup>172-175</sup>

There are many tools available to clinicians to assess the quality of life of asthmatic patients.<sup>176-183</sup> Unfortunately, these psychometric instruments that claim to measure the same outcomes might produce disparate results, and none have been targeted for the elderly. In general, results that measure several domains are more accurate when a composite score is derived rather than when subscores of specific domains are compared. Medical, administrative, and pharmacy records have also been used, especially to study larger asthmatic populations; these have proved useful for the assessment of a patient’s change over time and to measure group differences.

Clinical trials of asthma therapy and educational, self-management, and health services interventions have used psychometric instruments to assess elderly patients with asthma. In most of these studies, however, the majority of subjects are younger. There are no studies that have specifically determined the reliability and validity of these instruments in elderly persons. This is true of patient-satisfaction measures that have been used to assess asthma care.<sup>184-186</sup> This is much needed because using lung function testing to measure outcomes has potential limitations in this age group. There are difficulties in defining normal predicted values at a very advanced age, and many patients with physical or cognitive impairment cannot reliably perform these tests. It is hopeful that newer biomarkers of lung inflammation have a particular role to play in the assessment of asthma control in the elderly.

### Tools assessing physical function: Self-reported and objective

A major goal of geriatric and gerontologic research is to reduce the decrease in cognitive and physical function and prevent disability among older adults. Accordingly, many functional status measures have been developed and used to understand the disabling process, as well as to evaluate interventions to prevent functional decline. It is useful to identify instruments that measure functional limitations and disability to investigate the functional consequences of asthma in older adults and to understand the pathway from asthma to disability. Functional limitations are restrictions in performing basic physical and mental actions at the whole-person level (eg, walking or climbing stairs),

whereas disability refers to limitations or difficulty in performing socially defined roles or tasks of everyday living in a given environmental context (eg, grocery shopping or bathing).<sup>187,188</sup> Both self-reported and objective measures can be used to measure these different stages of disablement.<sup>189</sup>

Self-reported measures can provide an indication of how well a patient is functioning in daily life and provide an assessment of care needs. These measures incorporate self-perception of function and can assess adaptations made to compensate for decrements in function.<sup>190</sup> For disability assessment, self-reported difficulty or inability to perform basic activities of daily living (ADL) is commonly used. For example, a composite score of 8 ADL items has been used as an outcome to evaluate the efficacy of a program to prevent functional decline in frail older adults.<sup>191</sup> Other composite scores assessing difficulty in ambulation, stair climbing, transferring, upper extremity function, and basic and instrumental ADLs have been developed.<sup>192</sup> These comprehensive instruments of function and disability are amenable to computer adaptive testing.<sup>193</sup>

Several objective measures of physical performance are used in studies of older adults and in disease-specific patient populations. Tests of physical performance eliminate subjective attitudinal differences in the patient's reporting of physical function limitations. They have the advantage of providing an objective measure for comparisons across populations.<sup>194</sup> These tests are sensitive to change over time and can detect decrements in function that might not be observed with self-reported instruments. Many studies in older adults have used physical performance tests as predictors of adverse health events, as well as outcomes.

For example, the Short Physical Performance Battery, which consists of timed balance, walking, and chair-rise tasks, is a powerful predictor of disability, nursing home admission, and mortality.<sup>195,196</sup> The Short Physical Performance Battery was also used as a screening instrument to identify functionally limited older adults and as an outcome in a randomized controlled trial of exercise.<sup>197</sup> Increasingly, objective measures of physical function are used to summarize the effect of total disease burden, including subclinical conditions and impairments, and to identify physiologic reserve that might help some older adults cope with disease burden. Clinically meaningful differences have been established for commonly used performance measures.<sup>198</sup>

## MANAGEMENT OF ASTHMA IN THE ELDERLY

The goals of asthma therapy in elderly patients are not different from those for younger asthmatic patients.<sup>96</sup> They are to treat acute symptoms, prevent chronic symptoms, decrease emergency department visits and hospitalizations, preserve normal activity level, and optimize pulmonary function with a minimal adverse effect from medications.<sup>163,199,200</sup> Optimal management should also focus on improving health status (quality of life) in these patients, which is often complicated by depressive symptoms and side effects from the drugs commonly used for asthma.<sup>201</sup> Unlike many younger adults who might require no medication or just as-needed  $\beta$ -agonist therapy for occasional symptoms, most older asthmatic patients need continuous treatment programs to control their disease. At a time when memory loss is common and financial resources are often limited, many older patients require complicated and frequent dosing with multiple expensive drugs. Unfortunately, this has led to a significant rate of noncompliance among the elderly population in general.<sup>45</sup> Sex, socioeconomic

factors, educational level, marital status, and severity of disease do not seem to be good predictors of compliance in elderly asthmatic patients. In summary, there are many challenges in the treatment of asthma in the elderly, which include a greater propensity to experience adverse events from medication use, as well as potential drug interactions with medications used for the treatment of comorbidities,<sup>4,96</sup> and thus it is particularly important to treat any disease in the elderly, including asthma, with a minimum of therapy while attaining maximum efficacy.

A thorough understanding is required regarding which medications will be most effective in the treatment of asthma in the elderly to achieve this balance. Because many current therapeutic options and those in development for asthma focus on specific inflammatory cells and mediators, any age-related changes in the airway inflammatory milieu will likely affect their therapeutic efficacy. Therefore a rigorous characterization of age-related changes in airway inflammation will facilitate the management of asthma in the elderly.

## Pharmacologic interventions

The therapeutic approach to asthma in elderly patients does not differ from what is recommended for young patients. Statements on the standard of care for treating asthma have been published by the National Institutes of Health and are widely used as guidelines.<sup>163,199</sup> Treatment protocols use step-care pharmacologic therapy based on the intensity of asthma symptoms and the clinical response to these interventions. As symptoms and lung function worsen, step-up or add-on therapy is given. As symptoms improve, therapy can be stepped down. In this age group special attention should also be given to the potential adverse effects of commonly used medications.

Corticosteroids are capable of reducing airway inflammation, thereby improving lung function, decreasing bronchial hyperreactivity, reducing symptoms, and improving overall quality of life. Oral corticosteroids should be avoided if possible because they place the patient at risk for bone fracture and increased likelihood of cataracts, muscle weakness, back pain, bruising, and oral candidiasis.<sup>202</sup> Many studies have shown that inhaled corticosteroids are safe and effective treatment for persistent asthma, but none have specifically targeted the elderly population. Long-term use of inhaled corticosteroids has been associated with a good safety profile, but higher doses of inhaled steroids (eg, >1000  $\mu\text{g}/\text{d}$ ) are capable of causing hypothalamic-pituitary-adrenal axis suppression. Local adverse effects, such as hoarseness, dysphonia, cough, and oral candidiasis, do occur but can usually be avoided by the use of a spacer or holding chamber with the metered-dose inhaler and by rinsing the mouth after each use. Despite the pivotal role of inhaled corticosteroids in asthma, many elderly patients are undertreated with this group of medications.<sup>5,203</sup>

Leukotriene-modifying agents (LTMs) are also asthma controllers. These agents have been shown to be effective in preventing allergen-induced asthma, exercise-induced asthma, and aspirin-induced bronchospasm. Studies on their use in the elderly are limited. When compared with LTMs, low-dose inhaled corticosteroids have favored the latter. The LTMs might also reduce asthma exacerbation rates and the need for steroid bursts. The LTMs are generally very safe.<sup>66,204</sup>

$\beta$ -Adrenergic agents are important medications in the acute and chronic management of asthma. Elderly patients with asthma

might be less responsive to certain bronchodilators compared with younger patients.<sup>73,138</sup>

Inhaled short-acting  $\beta_2$ -adrenergic agonists are the treatment of choice for the acute exacerbation of asthma symptoms. Despite the minimal systemic absorption seen with these agents, slight tachycardia might be observed. This is presumably because of vasodilatation, which results from the stimulation of  $\beta_2$ -receptors in vascular smooth muscle. Tremor can also occur and is especially troublesome in the geriatric patient. Tremor is thought to be caused by stimulation of  $\beta_2$ -receptors in skeletal muscle. In general, they have been proven to be safe and effective in all age groups.<sup>205</sup> However,  $\beta$ -agonists can cause (1) a dose-dependent decrease in serum potassium levels and (2) a dose-dependent increase in the QT interval on electrocardiography. Because sudden death from ventricular arrhythmia can be caused by both of these mechanisms, as well as being a complication of ischemic heart disease, the use of  $\beta$ -agonists in the elderly should be closely monitored. Short-acting  $\beta_2$ -agonists should be used for rescue of symptoms, whereas long-acting agents should be used as maintenance medications only as an add-on to inhaled corticosteroids and never as stand-alone therapy.

Anticholinergics, such as inhaled ipratropium, a short-acting bronchodilator, and tiotropium, a bronchodilator with 24-hour action, have an excellent safety profile in the elderly. They should be considered when additional bronchodilator therapy is necessary; however, their role in long-term maintenance of asthma in the elderly has not been established.

Theophylline is an effective bronchodilator and has some anti-inflammatory properties. However, its use has been greatly reduced over the past decade because of safety concerns, especially in the elderly. The narrow therapeutic range of theophylline, the frequency of concomitant illnesses that alter theophylline kinetics, and many drug interactions that affect the clearance of theophylline make it essential to closely monitor blood theophylline levels in older asthmatic patients. Theophylline toxicity can cause seizures and cardiac arrhythmias, such as atrial fibrillation, supraventricular tachycardia, ventricular ectopy, and ventricular tachycardia. The most common cause for theophylline toxicity is a self-administered increase in medication.

## Nonpharmacologic interventions

**Controlling triggers.** Measures should be taken to avoid triggers that can cause worsening of symptoms. As with asthma at any age, education concerning avoidance of aggravating factors that can lead to severe bronchospasm is very useful. Although aeroallergens are less important in provoking symptoms in the elderly than in young patients, a program implementing environmental control measures, such as avoiding or minimizing aeroallergen exposure, should be instituted in patients with documented sensitivity to specific allergens. However, such programs might not be successful in all cases, especially because lifestyle changes in the elderly population might be difficult.

The most important provocative factors include viral respiratory tract infections and irritants, such as cigarette smoke, paints, varnish, and household aerosols. Pharmacologic agents that are often prescribed for concomitant illnesses (ischemic heart disease and hypertension), such as  $\beta$ -adrenoreceptor antagonists ( $\beta$ -blockers), can also provoke bronchospasm.<sup>206</sup> This includes both noncardioselective agents (propranolol, pindolol, and timolol) and, to a lesser extent, cardioselective agents (metoprolol and

acebutolol). Topical  $\beta$ -blockers are also widely used in the elderly to reduce intraocular pressure in wide-angle glaucoma. With such treatment, sufficient systemic absorption might cause fatal status asthmaticus.<sup>207</sup> The severity of  $\beta$ -blocker-induced bronchoconstriction correlates with the severity of underlying airflow obstruction and the degree of bronchial reactivity and might be reduced by the use of a cardioselective topical  $\beta$ -blocking agent, such as betaxolol.<sup>208</sup> Aspirin and nonsteroidal anti-inflammatory agents might precipitate acute bronchospasm in certain asthmatic patients, and angiotensin-converting enzyme inhibitors might cause dry cough in some, worsening the symptoms of asthma. GERD should also be considered a cause of worsening asthma symptoms.

**Asthma education.** The complexity of the prescription regimen (number and frequency of medications), coupled with the memory loss and cognitive dysfunction that might be present in this group of patients, contribute partially to poor compliance with therapy.<sup>4,96</sup>

Patient education is an effective tool and should be an integral part in the management of asthma.<sup>209</sup> Active participation by a patient and family members in monitoring lung function, avoidance of provocative agents, and decisions regarding medications provide asthma management skills that give that patient the confidence to control his or her own disease. Mastering the technique of an inhaled medication delivery device is a challenging problem in elderly patients, and the great majority of elderly patients are unable to properly use the metered-dose inhaler, even after proper instruction.<sup>210-213</sup> Use of dry powder devices, although simpler, requires the generation of an adequate inspiratory flow that might be suboptimal in frail patients and those with severe airway obstruction. In such situations the use of spacer devices or nebulizers might be beneficial. Patients should recognize the rationale behind using the different medications, the correct way to use them, and their side effects, and polypharmacy should also be avoided. Asthma in the elderly can be effectively managed, and despite severe symptoms and physiologic impairment, most patients can lead active and productive lives.

A demographic study of 380 low-income elderly persons in Chicago found that 26 (10%) without a previous diagnosis of asthma or emphysema had symptoms compatible with those of obstructive lung disease.<sup>214</sup> Of patients with a previous diagnosis, only 18% were compliant with medications, and this was largely due to the cost of medications. In addition, health care use was high in this population. Telephone intervention offers a simple option in the management of elderly patients with asthma. It has been shown that asthma care by means of telephone triage of adult asthmatic patients can lead to a higher percentage of asthmatic patients being reviewed at less cost per patient and without loss of asthma control when compared with usual routine care in the outpatient clinic.<sup>215</sup> However, it has not been determined whether such an intervention could improve asthma care specifically in persons aged 65 years or older. The following study was designed to evaluate this question.

Fifty-two elderly asthmatic patients who used their rescue inhalers more than twice a week and had at least 1 emergency department or urgent care visit in the previous year were randomized to an intervention or control group.<sup>216</sup> All patients received 2 telephone calls over a 12-month period. The intervention group received an asthma-specific questionnaire, and the control group received a general health questionnaire. Medication use and health care use were evaluated at the beginning and end of

a 12-month period. The study was completed by 23 control and 25 intervention subjects. Baseline data were similar in both groups. After 12 months, 72% (n = 18) of the intervention group was taking an inhaled corticosteroid compared with 40% (n = 10) of the control group. The intervention group had fewer emergency department visits when compared with the control group. Sixty-four percent (n = 16) of the intervention group had an asthma action plan compared with 26% (n = 6) of the control group. This study provides evidence that using a simple telephone questionnaire can successfully improve asthma care in the elderly. By empowering the elderly with the appropriate knowledge regarding their asthma, an appropriate discussion about their asthma care can be initiated with their primary care physicians.

**Pulmonary rehabilitation.** Although pulmonary rehabilitation is recommended as the standard of care for patients with COPD, there are only a few studies that evaluate the benefit of rehabilitation for asthmatic patients, and none of these consider elderly asthmatic patients. One study looked at the effects of a 10-week outpatient rehabilitation program for 58 asthmatic patients after 3 years.<sup>217-219</sup> They found that 39 of 58 subjects continued to exercise regularly all 3 years; there was a decreased number of emergency department visits and a decrease in asthma symptoms. Further studies are needed to assess empowerment strategies for elderly patients with asthma, as well as the potential benefits of pulmonary rehabilitation on morbidity and mortality.

## AREAS OF RESEARCH

### Experimental approaches

Asthma pathogenesis is complex and incompletely understood. Research into the pathophysiologic mechanisms is made more difficult by multiple factors, including the heterogeneity of the disease itself, variable presentations in different stages of life, and the lack of highly relevant animal models.<sup>220-224</sup> In the last decade, increasing interest in asthma in the elderly has triggered more intensive investigation in both human and animal systems by using ever more sophisticated immunologic methodologies.

Early investigation with rats revealed a lack of total and allergen-specific IgE in response to ovalbumin.<sup>225</sup> This was born out by several later *in vivo* studies.<sup>226-228</sup> IgG subset analysis (IgG<sub>1</sub> vs IgG<sub>2</sub>) provided further support for this phenomenon. IgG<sub>1</sub>, correlating in the mouse to a T<sub>H</sub>2 response (vs IgG<sub>2</sub> [T<sub>H</sub>1]) was shown to follow a similar pattern.<sup>227,228</sup> Recent studies<sup>226-228</sup> of cytokine profiles in aged rodents compared with young control animals enhanced the paradigm that age resulted in less robust T<sub>H</sub>2 cytokines, particularly IL-4, IL-5, and IL-13, in favor of T<sub>H</sub>1 gene and protein expression.<sup>227,228</sup> This pattern was not fully supported in a recent chronic murine asthma model<sup>229</sup> wherein IL-5 was greater in aged sensitized mice, making the picture more complex. IFN- $\gamma$ , a key T<sub>H</sub>1 cytokine, has been consistently overexpressed in aged versus young rodents.<sup>226-229</sup>

Eosinophilia, which is considered a key component of (allergic) asthma, was more pronounced in younger versus older animals (bronchoalveolar lavage fluid, lung tissue, or both) after most,<sup>226-228,230</sup> although not all,<sup>229</sup> sensitization paradigms.

Molecular genetics and T-cell subset analysis has allowed further insight into possible mechanisms underlying the waning T<sub>H</sub>2 response observed in most models.<sup>227-229</sup> Specifically, elderly mice appear to have more memory T cells, less activated

CD4<sup>+</sup> T<sub>H</sub>2 cells, and less activated monocytes.<sup>227,228</sup> Resident goblet cells also appear to express upregulation of mucin and mucin gene expression.<sup>229</sup> A key to the impaired T<sub>H</sub>2 response was recently found in the GATA3 pathway.<sup>228</sup> Elderly mice do not phosphorylate components of the extracellular signal-regulated protein kinase/mitogen-activated protein kinase pathway, resulting in lack of downstream signaling with GATA3, with subsequent impairment of promoter regions for key T<sub>H</sub>2 cytokines, including IL-4. This could be an overarching explanation for many findings in the elderly asthmatic patient, including less IgE (IL-4 and IL-13 are needed for opening switch regions for IgE production); IL-4 and IL-13 are highly associated with airway hyperreactivity, and IL-5 is associated with eosinophil activation, survival, and, to a lesser extent, trafficking.

Finally, airway hyperreactivity has been universally found to be greater in young versus aged animals.<sup>226-229,231</sup> The mechanisms might be complex, including both an altered key cytokine milieu and alterations in muscle function at the muscarinic receptor level.<sup>231-233</sup>

### Clinical and translational research

Research into the pathogenesis of asthma in recent years has led to the discovery of a number of novel, potentially important targets for the development of new treatment options. Much of this research has focused on T<sub>H</sub>2 lymphocyte-driven processes underlying allergic asthma and its characteristic eosinophilic airway inflammation. Abundant information supporting this research has been derived from bronchial biopsy and bronchoalveolar lavage studies largely carried out in a young adult population. It is recognized, however, that the role of allergy and allergic triggers in asthma diminishes with age.<sup>69,234</sup> In addition, LOA is often less reversible, more severe, and frequently occurs in response to a viral respiratory tract infection.<sup>153</sup> A distinct asthma phenotype characterized by normal airway eosinophil numbers has been described.<sup>235</sup> Moreover, normal airway eosinophilia might also be associated with abnormal sputum neutrophilia.<sup>236,237</sup> Recent studies have shown that neutrophilic asthma might be associated with activation of innate immune pathways in contrast to the adaptive immune response associated with T<sub>H</sub>2-mediated allergic asthma.<sup>238</sup> Thus alternative immune pathways involving natural killer T or T<sub>H</sub>17 lymphocyte subtypes have been hypothesized as being potentially important in the pathogenesis of asthma, particularly in adult-onset asthma.<sup>239,240</sup> Just as the discovery of T<sub>H</sub>2-related pathways has led to important leads in drug discovery for allergic asthma, further clinical research into these alternative pathways should be carried out with the goal of identifying new and exciting targets for future drug discovery. This research should focus not only on the discovery of new molecular targets but also on the identification of noninvasive biomarkers that will help predict the success of any new therapy in an individual patient.

### SUMMARY AND CONCLUSIONS

Asthma is an important disease in the older adult, affecting 7% of the population older than 65 years, which is understudied and frequently underdiagnosed. There are data to suggest that asthma in older adults is phenotypically different from that in young patients, with a potential effect on the diagnosis, assessment, and

management in this population. This workshop brought together many disciplines to further our current understanding, resolve gaps in knowledge, and explore future areas of research and education. Table I lists specific areas in need of research and study.

We thank the NIA for recognizing the need for this workshop, especially Susan Nayfield, MD, who convened the workshop and provided tremendous support in moving this research field forward; Evan Hadley, MD, the Director of the Division of Geriatrics and Clinical Gerontology; and Basil Eldadah, MD, who continued the work of Dr Nayfield and contributed valuable advice and encouragement to the authors in completing these proceedings.

## REFERENCES

- Lutz W, Sanderson W, Scherbov S. The coming acceleration of global population ageing. *Nature* 2008;451:716-9.
- US Census Bureau, Population Division. The Census Bureau on prospects for US population growth in the Twenty-First Century. Population and development review. 2000;62:197-200. doi: 10.1111/j.1728-4457.2000.00197.x.
- Moorman JE, Rudd RA, Johnson CA, King M, Minor P, Bailey C, et al. National surveillance for asthma—United States, 1980-2004. *MMWR Surveill Summ* 2007;56:1-54.
- King MJ, Hanania NA. Asthma in the elderly: current knowledge and future directions. *Curr Opin Pulm Med* 2010;16:55-9.
- Enright PL, McClelland RL, Newman AB, Gottlieb DJ, Lebowitz MD. Underdiagnosis and undertreatment of asthma in the elderly. *Cardiovascular Health Study Research Group*. *Chest* 1999;116:603-13.
- Moorman JE, Mannino DM. Increasing U.S. asthma mortality rates: who is really dying? *J Asthma* 2001;38:65-71.
- Craig BM, Kraus CK, Chewing BA, Davis JE. Quality of care for older adults with chronic obstructive pulmonary disease and asthma based on comparisons to practice guidelines and smoking status. *BMC Health Serv Res* 2008;8:144.
- Blanchette CM, Gutierrez B, Ory C, Chang E, Akazawa M. Economic burden in direct costs of concomitant chronic obstructive pulmonary disease and asthma in a Medicare Advantage population. *J Manag Care Pharm* 2008;14:176-85.
- Yancik R, Ershler W, Satariano W, Hazzard W, Cohen HJ, Ferrucci L. Report of the National Institute on Aging task force on comorbidity. *J Gerontol A Biol Sci Med Sci* 2007;62:275-80.
- Ling SM, Simonsick EM, Ferrucci L. A painful interface between normal aging and disease. *J Gerontol A Biol Sci Med Sci* 2007;62:613-5.
- Ferrucci L, Giallauria F, Guralnik JM. Epidemiology of aging. *Radiol Clin North Am* 2008;46:643-52, v.
- Hart RW, Setlow RB. Correlation between deoxyribonucleic acid excision-repair and life-span in a number of mammalian species. *Proc Natl Acad Sci U S A* 1974;71:2169-73.
- Narita M, Lowe SW. Executing cell senescence. *Cell Cycle* 2004;3:244-6.
- Kirkwood TB. A systematic look at an old problem. *Nature* 2008;451:644-7.
- Cristofalo VJ. Overview of biological mechanism of aging. *Annu Rev Gerontol Geriatr* 1990;10:1-22.
- Cristofalo VJ, Gerhard GS, Pignolo RJ. Molecular biology of aging. *Surg Clin North Am* 1994;74:1-21.
- Cristofalo VJ, Lorenzini A, Allen RG, Torres C, Tresini M. Replicative senescence: a critical review. *Mech Ageing Dev* 2004;125:827-48.
- Hayflick L. The limited in vitro lifetime of human diploid cell strains. *Exp Cell Res* 1965;37:614-36.
- Cristofalo VJ. Cell culture aging: insights for cell aging in vivo? *Aging (Milano)* 1999;11:1-3.
- Schneider EL, Mitsui Y. The relationship between in vitro cellular aging and in vivo human age. *Proc Natl Acad Sci U S A* 1976;73:3584-8.
- Rohme D. Evidence for a relationship between longevity of mammalian species and life spans of normal fibroblasts in vitro and erythrocytes in vivo. *Proc Natl Acad Sci U S A* 1981;78:5009-13.
- Dimri GP, Lee X, Basile G, Acosta M, Scott G, Roskelley C, et al. A biomarker that identifies senescent human cells in culture and in aging skin in vivo. *Proc Natl Acad Sci U S A* 1995;92:9363-7.
- Fried LP, Tangen CM, Walston J, Newman AB, Hirsch C, Gottdiener J, et al. Frailty in older adults: evidence for a phenotype. *J Gerontol A Biol Sci Med Sci* 2001;56:M146-56.
- Bandein-Roche K, Xue QL, Ferrucci L, Walston J, Guralnik JM, Chaves P, et al. Phenotype of frailty: characterization in the women's health and aging studies. *J Gerontol A Biol Sci Med Sci* 2006;61:262-6.
- Fried LP, Hadley EC, Walston JD, Newman AB, Guralnik JM, Studenski S, et al. From bedside to bench: research agenda for frailty. *Sci Aging Knowledge Environ* 2005;2005:e24.
- Studenski S, Perera S, Patel K, Rosano C, Faulkner K, Inzitari M, et al. Gait speed and survival in older adults. *JAMA* 2011;305:50-8.
- Ershler WB, Keller ET. Age-associated increased interleukin-6 gene expression, late-life diseases, and frailty. *Annu Rev Med* 2000;51:245-70.
- Leng SX, Yang H, Walston JD. Decreased cell proliferation and altered cytokine production in frail older adults. *Aging Clin Exp Res* 2004;16:249-52.
- Christensen K, Vaupel JW. Determinants of longevity: genetic, environmental and medical factors. *J Intern Med* 1996;240:333-41.
- Arias E. United States life tables, 2004. *Natl Vital Stat Rep* 2007;56:1-39.
- Mackenbach JP, Kunst AE, Laatenbach H, Oei YB, Bijlsma F. Gains in life expectancy after elimination of major causes of death: revised estimates taking into account the effect of competing causes. *J Epidemiol Community Health* 1999;53:32-7.
- Orr WC, Sohal RS. Extension of life-span by overexpression of superoxide dismutase and catalase in *Drosophila melanogaster*. *Science* 1994;263:1128-30.
- Murakami S, Johnson TE. A genetic pathway conferring life extension and resistance to UV stress in *Caenorhabditis elegans*. *Genetics* 1996;143:1207-18.
- Sun J, Kale SP, Childress AM, Pinswasdi C, Jazwinski SM. Divergent roles of RAS1 and RAS2 in yeast longevity. *J Biol Chem* 1994;269:18638-45.
- Hirsch HR. Can an improved environment cause maximum lifespan to decrease? Comments on lifespan criteria and longitudinal Gompertzian analysis. *Exp Gerontol* 1994;29:119-37.
- Riggs JE. Longitudinal Gompertzian analysis of stroke mortality in the U.S., 1951-1986: declining stroke mortality is the natural consequence of competitive deterministic mortality dynamics. *Mech Ageing Dev* 1990;55:235-43.
- Roth GS, Ingram DK, Joseph JA. Nutritional interventions in aging and age-associated diseases. *Ann N Y Acad Sci* 2007;1114:369-71.
- Weindruch R. Caloric restriction and aging. *Sci Am* 1996;274:46-52.
- Janssens JP, Pache JC, Nicod LP. Physiological changes in respiratory function associated with ageing. *Eur Respir J* 1999;13:197-205.
- Gibson GJ, Pride NB, O'Cain C, Quagliato R. Sex and age differences in pulmonary mechanics in normal nonsmoking subjects. *J Appl Physiol* 1976;41:20-5.
- Turner JM, Mead J, Wohl ME. Elasticity of human lungs in relation to age. *J Appl Physiol* 1968;25:664-71.
- Begin R, Renzetti AD Jr, Bigler AH, Watanabe S. Flow and age dependence of airway closure and dynamic compliance. *J Appl Physiol* 1975;38:199-207.
- Franceschi C, Bonafe M, Valensin S. Human immunosenescence: the prevailing of innate immunity, the failing of clonotypic immunity, and the filling of immunological space. *Vaccine* 2000;18:1717-20.
- Franceschi C, Bonafe M, Valensin S, Olivieri F, De Luca M, Ottaviani E, et al. Inflamm-aging. An evolutionary perspective on immunosenescence. *Ann N Y Acad Sci* 2000;908:244-54.
- Barr RG, Somers SC, Speizer FE, Camargo CA Jr. Patient factors and medication guideline adherence among older women with asthma. *Arch Intern Med* 2002;162:1761-8.
- Kringsman K, Moen J, Nilsson JL, Ring L. Refill adherence by the elderly for asthma/chronic obstructive pulmonary disease drugs dispensed over a 10-year period. *J Clin Pharm Ther* 2007;32:603-11.
- Kung HC, Hoyert DL, Xu J, Murphy SL. Deaths: final data for 2005. *Natl Vital Stat Rep* 2008;56:1-120.
- Zeiger RS, Hay JW, Contreras R, Chen W, Quinn VP, Seal B, et al. Asthma costs and utilization in a managed care organization. *J Allergy Clin Immunol* 2008;121:885-92.
- Enright PL, Ward BJ, Tracy RP, Lasser EC. Asthma and its association with cardiovascular disease in the elderly. The Cardiovascular Health Study Research Group. *J Asthma* 1996;33:45-53.
- Fried LP, Borhani NO, Enright P, Furberg CD, Gardin JM, Kronmal RA, et al. The Cardiovascular Health Study: design and rationale. *Ann Epidemiol* 1991;1:263-76.
- Enright PL, Kronmal RA, Higgins MW, Schenker MB, Haponik EF. Prevalence and correlates of respiratory symptoms and disease in the elderly. *Cardiovascular Health Study*. *Chest* 1994;106:827-34.
- Enright PL, McClelland RL, Buist AS, Lebowitz MD. Correlates of peak expiratory flow lability in elderly persons. *Chest* 2001;120:1861-8.
- Bjornsson HT, Sigurdsson MI, Fallin MD, Irizarry RA, Aspelund T, Cui H, et al. Intra-individual change over time in DNA methylation with familial clustering. *JAMA* 2008;299:2877-83.
- Tang WY, Ho SM. Epigenetic reprogramming and imprinting in origins of disease. *Rev Endocr Metab Disord* 2007;8:173-82.
- Sedivy JM, Banumathy G, Adams PD. Aging by epigenetics—a consequence of chromatin damage? *Exp Cell Res* 2008;314:1909-17.



56. Feinberg AP. Epigenetics at the epicenter of modern medicine. *JAMA* 2008;299:1345-50.
57. Bhavsar P, Ahmad T, Adcock IM. The role of histone deacetylases in asthma and allergic diseases. *J Allergy Clin Immunol* 2008;121:580-4.
58. Miller RL, Ho SM. Environmental epigenetics and asthma: current concepts and call for studies. *Am J Respir Crit Care Med* 2008;177:567-73.
59. Lu Q, Qiu X, Hu N, Wen H, Su Y, Richardson BC. Epigenetics, disease, and therapeutic interventions. *Ageing Res Rev* 2006;5:449-67.
60. Adcock IM, Tsaprouni L, Bhavsar P, Ito K. Epigenetic regulation of airway inflammation. *Curr Opin Immunol* 2007;19:694-700.
61. Fraga MF, Ballestar E, Paz MF, Ropero S, Setien F, Ballestar ML, et al. Epigenetic differences arise during the lifetime of monozygotic twins. *Proc Natl Acad Sci U S A* 2005;102:10604-9.
62. Griffith KA, Sherrill DL, Siegel EM, Manolio TA, Bonekat HW, Enright PL. Predictors of loss of lung function in the elderly: the Cardiovascular Health Study. *Am J Respir Crit Care Med* 2001;163:61-8.
63. Barua P, O'Mahony MS. Overcoming gaps in the management of asthma in older patients: new insights. *Drugs Aging* 2005;22:1029-59.
64. Busse PJ. Allergic respiratory disease in the elderly. *Am J Med* 2007;120:498-502.
65. Enright PL. The diagnosis and management of asthma is much tougher in older patients. *Curr Opin Allergy Clin Immunol* 2002;2:175-81.
66. Korenblat PE, Kemp JP, Scherger JE, Minkwitz MC, Mezzanotte W. Effect of age on response to zafirlucast in patients with asthma in the Accolate Clinical Experience and Pharmacoeconomics Trial (ACCEPT). *Ann Allergy Asthma Immunol* 2000;84:217-25.
67. Quadrelli SA, Roncoroni AJ. Is asthma in the elderly really different? *Respiration* 1998;65:347-53.
68. Zureik M, Orehek J. Diagnosis and severity of asthma in the elderly: results of a large survey in 1,485 asthmatics recruited by lung specialists. *Respiration* 2002;69:223-8.
69. Braman SS. Asthma in the elderly. *Clin Geriatr Med* 2003;19:57-75.
70. Newnham DM. Asthma medications and their potential adverse effects in the elderly: recommendations for prescribing. *Drug Saf* 2001;24:1065-80.
71. Kornman KS. Interleukin 1 genetics, inflammatory mechanisms, and nutrigenetic opportunities to modulate diseases of aging. *Am J Clin Nutr* 2006;83(suppl):475S-83S.
72. Choi JH, Oh SW, Kang MS, Kwon HJ, Oh GT, Kim DY. Trichostatin A attenuates airway inflammation in mouse asthma model. *Clin Exp Allergy* 2005;35:89-96.
73. Banerji A, Clark S, Afilalo M, Blanda MP, Cydulka RK, Camargo CA Jr. Prospective multicenter study of acute asthma in younger versus older adults presenting to the emergency department. *J Am Geriatr Soc* 2006;54:48-55.
74. Aw D, Silva AB, Palmer DB. Immunosenescence: emerging challenges for an ageing population. *Immunology* 2007;120:435-46.
75. Pawelec G, Ouyang Q, Wagner W, Biol D, Wikby A. Pathways to a robust immune response in the elderly. *Immunol Allergy Clin North Am* 2003;23:1-13.
76. Meyer KC, Soergel P. Variation of bronchoalveolar lymphocyte phenotypes with age in the physiologically normal human lung. *Thorax* 1999;54:697-700.
77. Meyer KC, Rosenthal NS, Soergel P, Peterson K. Neutrophils and low-grade inflammation in the seemingly normal aging human lung. *Mech Ageing Dev* 1998;104:169-81.
78. Mathur SK, Schwantes EA, Jarjour NN, Busse WW. Age-related changes in eosinophil function in human subjects. *Chest* 2008;133:412-9.
79. Wenzel SE. Asthma: defining of the persistent adult phenotypes. *Lancet* 2006;368:804-13.
80. Nyenhuis SM, Schwantes EA, Mathur SK. Characterization of leukotrienes in a pilot study of older asthma subjects. *Immun Ageing* 2010;7:8.
81. Slavin RG, Haselkorn T, Lee JH, Zheng B, Deniz Y, Wenzel SE. Asthma in older adults: observations from the epidemiology and natural history of asthma: outcomes and treatment regimens (TENOR) study. *Ann Allergy Asthma Immunol* 2006;96:406-14.
82. Delespesse G, de Maubeuge J, Kennes B, Nicaise R, Govaerts A. IgE mediated hypersensitivity in ageing. *Clin Allergy* 1977;7:155-60.
83. Hanneuse Y, Delespesse G, Hudson D, de Halleux F, Jacques JM. Influence of ageing on IgE-mediated reactions in allergic patients. *Clin Allergy* 1978;8:165-74.
84. Jarvis D, Luczynska C, Chinn S, Burney P. The association of age, gender and smoking with total IgE and specific IgE. *Clin Exp Allergy* 1995;25:1083-91.
85. Omenaas E, Bakke P, Elsayed S, Hanoa R, Gulsvik A. Total and specific serum IgE levels in adults: relationship to sex, age and environmental factors. *Clin Exp Allergy* 1994;24:530-9.
86. Stoy PJ, Roitman-Johnson B, Walsh G, Gleich GJ, Mendell N, Yunis E, et al. Ageing and serum immunoglobulin E levels, immediate skin tests, RAST. *J Allergy Clin Immunol* 1981;68:421-6.
87. Wittig HJ, Belloit J, De Fillipi I, Royal G. Age-related serum immunoglobulin E levels in healthy subjects and in patients with allergic disease. *J Allergy Clin Immunol* 1980;66:305-13.
88. Barbee RA, Kaltenborn W, Lebowitz MD, Burrows B. Longitudinal changes in allergen skin test reactivity in a community population sample. *J Allergy Clin Immunol* 1987;79:16-24.
89. Annema JT, Sparrow D, O'Connor GT, Rijcken B, Koeter GH, Postma DS, et al. Chronic respiratory symptoms and airway responsiveness to methacholine are associated with eosinophilia in older men: the Normative Aging Study. *Eur Respir J* 1995;8:62-9.
90. King MJ, Bukantz SC, Phillips S, Mohapatra SS, Tamulis T, Lockey RF. Serum total IgE and specific IgE to *Dermatophagoides pteronyssinus*, but not eosinophil cationic protein, are more likely to be elevated in elderly asthmatic patients. *Allergy Asthma Proc* 2004;25:321-5.
91. Nicholson KG, Kent J, Ireland DC. Respiratory viruses and exacerbations of asthma in adults. *BMJ* 1993;307:982-6.
92. Institute of Medicine. Clearing the air: asthma and indoor air exposures. Executive summary. Available at: <http://www.nap.edu/books/0309064961/html/>. Accessed July 22, 2011.
93. Fisk WJ, Lei-Gomez Q, Mendell MJ. Meta-analyses of the associations of respiratory health effects with dampness and mold in homes. *Indoor Air* 2007;17:284-96.
94. Institute of Medicine. Damp indoor spaces. Available at: <http://www.iom.edu/~media/Files/Report%20Files/2004/Damp-Indoor-Spaces-and-Health/dampin-door2paperforPDF.pdf>. Accessed July 22, 2011.
95. Gergen PJ, Turkeltaub PC, Sempos CT. Is allergen skin test reactivity a predictor of mortality? Findings from a national cohort. *Clin Exp Allergy* 2000;30:1717-23.
96. Braman SS, Hanania NA. Asthma in older adults. *Clin Chest Med* 2007;28:685-702, v.
97. Braman SS, Kaemmerlen JT, Davis SM. Asthma in the elderly. A comparison between patients with recently acquired and long-standing disease. *Am Rev Respir Dis* 1991;143:336-40.
98. Burrows B, Barbee RA, Cline MG, Knudson RJ, Lebowitz MD. Characteristics of asthma among elderly adults in a sample of the general population. *Chest* 1991;100:935-42.
99. Huss K, Naumann PL, Mason PJ, Nanda JP, Huss RW, Smith CM, et al. Asthma severity, atopic status, allergen exposure, and quality of life in elderly persons. *Ann Allergy Asthma Immunol* 2001;86:524-30.
100. Lee HY, Stretton TB. Asthma in the elderly. *BMJ* 1972;4:93-5.
101. Litonjua AA, Sparrow D, Weiss ST, O'Connor GT, Long AA, Ohman JL Jr. Sensitization to cat allergen is associated with asthma in older men and predicts new-onset airway hyperresponsiveness. The Normative Aging Study. *Am J Respir Crit Care Med* 1997;156:23-7.
102. Reed CE. The role of allergy and airway inflammation. In: Barbee RA, editor. *Asthma in the elderly*. New York: Marcel Dekker; 1997. p. 33-52.
103. Rogers L, Cassino C, Berger KI, Goldring RM, Norman RG, Klugh T, et al. Asthma in the elderly: cockroach sensitization and severity of airway obstruction in elderly nonsmokers. *Chest* 2002;122:1580-6.
104. King MJ, Tamulis T, Lockey RF. Prick puncture skin tests and serum specific IgE as predictors of nasal challenge response to *Dermatophagoides pteronyssinus* in older adults. *Ann Allergy Asthma Immunol* 2008;101:12-7.
105. Anderson HR, Atkinson RW, Bremner SA, Marston L. Particulate air pollution and hospital admissions for cardiorespiratory diseases: are the elderly at greater risk? *Eur Respir J Suppl* 2003;40:398-468.
106. Meng YY, Wilhelm M, Rull RP, English P, Ritz B. Traffic and outdoor air pollution levels near residences and poorly controlled asthma in adults. *Ann Allergy Asthma Immunol* 2007;98:455-63.
107. Johnston SL, Pattermore PK, Sanderson G, Smith S, Lampe F, Josephs L, et al. Community study of role of viral infections in exacerbations of asthma in 9-11 year old children. *BMJ* 1995;310:1225-9.
108. Atmar RL, Guy E, Guntupalli KK, Zimmerman JL, Bandi VD, Baxter BD, et al. Respiratory tract viral infections in inner-city asthmatic adults. *Arch Intern Med* 1998;158:2453-9.
109. Teichtahl H, Buckmaster N, Pertnikovs E. The incidence of respiratory tract infection in adults requiring hospitalization for asthma. *Chest* 1997;112:591-6.
110. Monto AS, Ullman BM. Acute respiratory illness in an American community. The Tecumseh study. *JAMA* 1974;227:164-9.
111. Falsey AR, McCann RM, Hall WJ, Tanner MA, Criddle MM, Formica MA, et al. Acute respiratory tract infection in daycare centers for older persons. *J Am Geriatr Soc* 1995;43:30-6.
112. Falsey AR, Dallah GE, Formica MA, Andolina GG, Hamer DH, Leka LL, et al. Long-term care facilities: a cornucopia of viral pathogens. *J Am Geriatr Soc* 2008;56:1281-5.

113. Falsey AR, Hennessey PA, Formica MA, Cox C, Walsh EE. Respiratory syncytial virus infection in elderly and high-risk adults. *N Engl J Med* 2005;352:1749-59.
114. Walsh EE, Peterson DR, Falsey AR. Human metapneumovirus infections in adults: another piece of the puzzle. *Arch Intern Med* 2008;168:2489-96.
115. Johnston NW. The similarities and differences of epidemic cycles of chronic obstructive pulmonary disease and asthma exacerbations. *Proc Am Thorac Soc* 2007;4:591-6.
116. Hutchinson AF, Ghimire AK, Thompson MA, Black JF, Brand CA, Lowe AJ, et al. A community-based, time-matched, case-control study of respiratory viruses and exacerbations of COPD. *Respir Med* 2007;101:2472-81.
117. Varkey JB, Varkey B. Viral infections in patients with chronic obstructive pulmonary disease. *Curr Opin Pulm Med* 2008;14:89-94.
118. Falsey AR, Walsh EE. Respiratory syncytial virus infection in elderly adults. *Drugs Aging* 2005;22:577-87.
119. Jacoby DB. Virus-induced asthma attacks. *J Aerosol Med* 2004;17:169-73.
120. Mallia P, Johnston SL. How viral infections cause exacerbation of airway diseases. *Chest* 2006;130:1203-10.
121. Proud D, Chow CW. Role of viral infections in asthma and chronic obstructive pulmonary disease. *Am J Respir Cell Mol Biol* 2006;35:513-8.
122. Bauer BA, Reed CE, Yunginger JW, Wollan PC, Silverstein MD. Incidence and outcomes of asthma in the elderly. A population-based study in Rochester, Minnesota. *Chest* 1997;111:303-10.
123. Braman SS, Corrao WM, Kaemmerlen JT. The clinical outcome of asthma in the elderly: a 7-year follow-up study. *Ann N Y Acad Sci* 1991;629:449-50.
124. Weiner P, Magadle R, Waizman Y, Weiner M, Rabner M, Zamir D. Characteristics of asthma in the elderly. *Eur Respir J* 1998;12:564-8.
125. Parameswaran K, Hildreth AJ, Chadha D, Keaney NP, Taylor IK, Bansal SK. Asthma in the elderly: underperceived, underdiagnosed and undertreated; a community survey. *Respir Med* 1998;92:573-7.
126. Raiha I, Impivaara O, Seppala M, Knuts LR, Sourander L. Determinants of symptoms suggestive of gastroesophageal reflux disease in the elderly. *Scand J Gastroenterol* 1993;28:1011-4.
127. Connolly MJ, Crowley JJ, Charan NB, Nielson CP, Vestal RE. Reduced subjective awareness of bronchoconstriction provoked by methacholine in elderly asthmatic and normal subjects as measured on a simple awareness scale. *Thorax* 1992;47:410-3.
128. Cassino C, Berger KI, Goldring RM, Norman RG, Kammerman S, Ciotoli C, et al. Duration of asthma and physiologic outcomes in elderly nonsmokers. *Am J Respir Crit Care Med* 2000;162:1423-8.
129. Braman SS. Asthma in the elderly. *Exp Lung Res* 2005;31(suppl 1):6-7.
130. Quadrelli SA, Roncoroni A. Features of asthma in the elderly. *J Asthma* 2001;38:377-89.
131. Lehmann S, Vollset SE, Nygaard HA, Gulsvik A. Factors determining performance of bronchodilator reversibility tests in middle-aged and elderly. *Respir Med* 2004;98:1071-9.
132. Pezzoli L, Giardini G, Consonni S, Dallera I, Bilotta C, Ferrario G, et al. Quality of spirometric performance in older people. *Age Ageing* 2003;32:43-6.
133. De Fillipi F, Tana F, Vanzati S, Balzarini B, Galetti G. Study of respiratory function in the elderly with different nutritional and cognitive status and functional ability assessed by plethysmographic and spirometric parameters. *Arch Gerontol Geriatr* 2003;37:33-43.
134. Bellia V, Pistelli R, Catalano F, Antonelli-Incalzi R, Grassi V, Melillo G, et al. Quality control of spirometry in the elderly. The S.A.R.A. study. *SAIute Respirazione nell'Anziano = Respiratory Health in the Elderly*. *Am J Respir Crit Care Med* 2000;161:1094-100.
135. Bellia V, Catalano F, Pistelli R, Antonelli-Incalzi R. Aging on quality of spirometry. *Am J Respir Crit Care Med* 2004;170:100-1.
136. Hansen EF, Phanareth K, Laursen LC, Kok-Jensen A, Dirksen A. Reversible and irreversible airflow obstruction as predictor of overall mortality in asthma and chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 1999;159:1267-71.
137. Pfeifer MA, Weinberg CR, Cook D, Best JD, Reenan A, Halter JB. Differential changes of autonomic nervous system function with age in man. *Am J Med* 1983;75:249-58.
138. Connolly MJ, Crowley JJ, Charan NB, Nielson CP, Vestal RE. Impaired bronchodilator response to albuterol in healthy elderly men and women. *Chest* 1995;108:401-6.
139. Ullah MI, Newman GB, Saunders KB. Influence of age on response to ipratropium and salbutamol in asthma. *Thorax* 1981;36:523-9.
140. Braman SS, Corrao WM. Bronchoprovocation testing. *Clin Chest Med* 1989;10:165-76.
141. Connolly MJ, Kelly C, Walters EH, Hendrick DJ. An assessment of methacholine inhalation tests in elderly asthmatics. *Age Ageing* 1988;17:123-8.
142. Crapo RO, Casaburi R, Coates AL, Enright PL, Hankinson JL, Irvin CG, et al. Guidelines for methacholine and exercise challenge testing—1999. This official statement of the American Thoracic Society was adopted by the ATS Board of Directors, July 1999. *Am J Respir Crit Care Med* 2000;161:309-29.
143. Scichilone N, Messina M, Battaglia S, Catalano F, Bellia V. Airway hyperresponsiveness in the elderly: prevalence and clinical implications. *Eur Respir J* 2005;25:364-75.
144. O'Connor GT, Sparrow D, Weiss ST. Normal range of methacholine responsiveness in relation to prechallenge pulmonary function. The Normative Aging Study. *Chest* 1994;105:661-6.
145. Enright PL, Burchette RJ, Peters JA, Lebowitz MD, McDonnell WF, Abbey DE. Peak flow lability: association with asthma and spirometry in an older cohort. *Chest* 1997;112:895-901.
146. Buist AS, Vollmer WM, Wilson SR, Frazier EA, Hayward AD. A randomized clinical trial of peak flow versus symptom monitoring in older adults with asthma. *Am J Respir Crit Care Med* 2006;174:1077-87.
147. Fabbri LM, Romagnoli M, Corbetta L, Casoni G, Busljetic K, Turato G, et al. Differences in airway inflammation in patients with fixed airflow obstruction due to asthma or chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2003;167:418-24.
148. Jedrychowski W, Krzyzanowski M, Wysocki M. Are chronic wheezing and asthma-like attacks related to FEV1 decline? The Cracow Study. *Eur J Epidemiol* 1988;4:335-42.
149. Peat JK, Woolcock AJ, Cullen K. Rate of decline of lung function in subjects with asthma. *Eur J Respir Dis* 1987;70:171-9.
150. Ulrik CS, Lange P. Decline of lung function in adults with bronchial asthma. *Am J Respir Crit Care Med* 1994;150:629-34.
151. Jedrychowski W, Maugeri U, Gomola K, Tobiasz-Adamczyk B, Bianchi I. Effects of domestic gas cooking and passive smoking on chronic respiratory symptoms and asthma in elderly women. *Int J Occup Environ Health* 1995;1:16-20.
152. Burrows B, Lebowitz MD, Barbee RA, Cline MG. Findings before diagnoses of asthma among the elderly in a longitudinal study of a general population sample. *J Allergy Clin Immunol* 1991;88:870-7.
153. Reed CE. The natural history of asthma in adults: the problem of irreversibility. *J Allergy Clin Immunol* 1999;103:539-47.
154. Taddei S, Galetta F, Viridis A, Ghiadoni L, Salvetti G, Franzoni F, et al. Physical activity prevents age-related impairment in nitric oxide availability in elderly athletes. *Circulation* 2000;101:2896-901.
155. Baraldi E, Carra S, Dario C, Azzolin N, Ongaro R, Marcer G, et al. Effect of natural grass pollen exposure on exhaled nitric oxide in asthmatic children. *Am J Respir Crit Care Med* 1999;159:262-6.
156. de Gouw HW, Hendriks J, Woltman AM, Twiss IM, Sterk PJ. Exhaled nitric oxide (NO) is reduced shortly after bronchoconstriction to direct and indirect stimuli in asthma. *Am J Respir Crit Care Med* 1998;158:315-9.
157. Olin AC, Rosengren A, Thelle DS, Lissner L, Bake B, Toren K. Height, age, and atopy are associated with fraction of exhaled nitric oxide in a large adult general population sample. *Chest* 2006;130:1319-25.
158. Smith AD, Cowan JO, Brassett KP, Herbison GP, Taylor DR. Use of exhaled nitric oxide measurements to guide treatment in chronic asthma. *N Engl J Med* 2005;352:2163-73.
159. Tsang KW, Ip SK, Leung R, Tipoe GL, Chan SL, Shum IH, et al. Exhaled nitric oxide: the effects of age, gender and body size. *Lung* 2001;179:83-91.
160. Pekkanen J, Pearce N. Defining asthma in epidemiological studies. *Eur Respir J* 1999;14:951-7.
161. Price DB, Tinkelman DG, Halbert RJ, Nordyke RJ, Isonaka S, Nonikov D, et al. Symptom-based questionnaire for identifying COPD in smokers. *Respiration* 2006;73:285-95.
162. Toren K, Brisman J, Jarvholm B. Asthma and asthma-like symptoms in adults assessed by questionnaires. A literature review. *Chest* 1993;104:600-8.
163. NIH NHLBI. National Heart, Lung and Blood Institute National Institute of Health. National Asthma Education and Prevention Program: Expert Panel Report 2—guidelines for the diagnosis and management of asthma. Bethesda: National Heart, Lung, and Blood Institute; 2007.
164. Diette GB, Krishnan JA, Dominici F, Haponik E, Skinner EA, Steinwachs D, et al. Asthma in older patients: factors associated with hospitalization. *Arch Intern Med* 2002;162:1123-32.
165. Griswold SK, Nordstrom CR, Clark S, Gaeta TJ, Price ML, Camargo CA Jr. Asthma exacerbations in North American adults: who are the "frequent fliers" in the emergency department? *Chest* 2005;127:1579-86.
166. Nathan RA, Sorkness CA, Kosinski M, Schatz M, Li JT, Marcus P, et al. Development of the asthma control test: a survey for assessing asthma control. *J Allergy Clin Immunol* 2004;113:59-65.
167. Schatz M, Mosen DM, Kosinski M, Vollmer WM, Magid DJ, O'Connor E, et al. Validity of the Asthma Control Test completed at home. *Am J Manag Care* 2007;13:661-7.

168. Juniper EF, O'Byrne PM, Guyatt GH, Ferrie PJ, King DR. Development and validation of a questionnaire to measure asthma control. *Eur Respir J* 1999;14:902-7.
169. Juniper EF, Bousquet J, Abetz L, Bateman ED. Identifying 'well-controlled' and 'not well-controlled' asthma using the Asthma Control Questionnaire. *Respir Med* 2006;100:616-21.
170. Vollmer WM, Markson LE, O'Connor E, Sanocki LL, Fitterman L, Berger M, et al. Association of asthma control with health care utilization and quality of life. *Am J Respir Crit Care Med* 1999;160:1647-52.
171. Vollmer WM, Markson LE, O'Connor E, Frazier EA, Berger M, Buist AS. Association of asthma control with health care utilization: a prospective evaluation. *Am J Respir Crit Care Med* 2002;165:195-9.
172. Bayliss MS, Espindle DM, Buchner D, Blaiss MS, Ware JE. A new tool for monitoring asthma outcomes: the ITG Asthma Short Form. *Qual Life Res* 2000;9:451-66.
173. Katz PP, Yelin EH, Smith S, Blanc PD. Perceived control of asthma: development and validation of a questionnaire. *Am J Respir Crit Care Med* 1997;155:577-82.
174. Chang CH, Sharp LK, Kimmel LG, Grammer LC, Kee R, Shannon JJ. A 6-item brief measure for assessing perceived control of asthma in culturally diverse patients. *Ann Allergy Asthma Immunol* 2007;99:130-5.
175. Boulet LP, Boulet V, Milot J. How should we quantify asthma control? A proposal. *Chest* 2002;122:2217-23.
176. Gupchup GV, Wolfgang AP, Thomas J III. Reliability and validity of the Asthma Quality of Life Questionnaire—marks in a sample of adult asthmatic patients in the United States. *Clin Ther* 1997;19:1116-25.
177. Katz PP, Eisner MD, Henke J, Shiboski S, Yelin EH, Blanc PD. The Marks Asthma Quality of Life Questionnaire: further validation and examination of responsiveness to change. *J Clin Epidemiol* 1999;52:667-75.
178. Marks GB, Dunn SM, Woolcock AJ. An evaluation of an asthma quality of life questionnaire as a measure of change in adults with asthma. *J Clin Epidemiol* 1993;46:1103-11.
179. Marks GB, Dunn SM, Woolcock AJ. A scale for the measurement of quality of life in adults with asthma. *J Clin Epidemiol* 1992;45:461-72.
180. Juniper EF, Buist AS, Cox FM, Ferrie PJ, King DR. Validation of a standardized version of the Asthma Quality of Life Questionnaire. *Chest* 1999;115:1265-70.
181. Juniper EF, Guyatt GH, Cox FM, Ferrie PJ, King DR. Development and validation of the Mini Asthma Quality of Life Questionnaire. *Eur Respir J* 1999;14:32-8.
182. Barr JT, Schumacher GE, Freeman S, LeMoine M, Bakst AW, Jones PW. American translation, modification, and validation of the St. George's Respiratory Questionnaire. *Clin Ther* 2000;22:1121-45.
183. Jones PW, Quirk FH, Baveystock CM, Littlejohns P. A self-complete measure of health status for chronic airflow limitation. The St. George's Respiratory Questionnaire. *Am Rev Respir Dis* 1992;145:1321-7.
184. Richards JM Jr, Dolce JJ, Windsor RA, Bailey WC, Brooks CM, Soong S. Patient characteristics relevant to effective self-management: scales for assessing attitudes of adults toward asthma. *J Asthma* 1989;26:99-108.
185. Juncal S, Vazquez MI, Seoane G, Blanco-Aparicio M, Vereza H. Applicability of the Asthma Opinion Survey in the Spanish population: distribution and relationship with sociodemographic and clinical variables. *J Asthma* 2000;37:469-79.
186. Mathias SD, Warren EH, Colwell HH, Sung JC. A new treatment satisfaction measure for asthmatics: a validation study. *Qual Life Res* 2000;9:873-82.
187. Nagi SZ. An epidemiology of disability among adults in the United States. *Milbank Mem Fund Q Health Soc* 1976;54:439-67.
188. Verbrugge LM, Jette AM. The disablement process. *Soc Sci Med* 1994;38:1-14.
189. Guralnik JM, Ferrucci L. Assessing the building blocks of function: utilizing measures of functional limitation. *Am J Prev Med* 2003;25(suppl 2):112-21.
190. Fried LP, Bandeen-Roche K, Chaves PH, Johnson BA. Preclinical mobility disability predicts incident mobility disability in older women. *J Gerontol A Biol Sci Med Sci* 2000;55:M43-52.
191. Gill TM, Baker DI, Gottschalk M, Peduzzi PN, Allore H, Byers A. A program to prevent functional decline in physically frail, elderly persons who live at home. *N Engl J Med* 2002;347:1068-74.
192. Ettinger WH Jr, Burns R, Messier SP, Applegate W, Rejeski WJ, Morgan T, et al. A randomized trial comparing aerobic exercise and resistance exercise with a health education program in older adults with knee osteoarthritis. The Fitness Arthritis and Seniors Trial (FAST). *JAMA* 1997;277:25-31.
193. Jette AM, Haley SM, Ni P, Olarsch S, Moed R. Creating a computer adaptive test version of the late-life function and disability instrument. *J Gerontol A Biol Sci Med Sci* 2008;63:1246-56.
194. Melzer D, Lan TY, Tom BD, Deeg DJ, Guralnik JM. Variation in thresholds for reporting mobility disability between national population subgroups and studies. *J Gerontol A Biol Sci Med Sci* 2004;59:1295-303.
195. Guralnik JM, Simonsick EM, Ferrucci L, Glynn RJ, Berkman LF, Blazer DG, et al. A short physical performance battery assessing lower extremity function: association with self-reported disability and prediction of mortality and nursing home admission. *J Gerontol* 1994;49:M85-94.
196. Guralnik JM, Ferrucci L, Simonsick EM, Salive ME, Wallace RB. Lower-extremity function in persons over the age of 70 years as a predictor of subsequent disability. *N Engl J Med* 1995;332:556-61.
197. Pahor M, Blair SN, Espeland M, Fielding R, Gill TM, Guralnik JM, et al. Effects of a physical activity intervention on measures of physical performance: results of the lifestyle interventions and independence for Elders Pilot (LIFE-P) study. *J Gerontol A Biol Sci Med Sci* 2006;61:1157-65.
198. Perera S, Mody SH, Woodman RC, Studenski SA. Meaningful change and responsiveness in common physical performance measures in older adults. *J Am Geriatr Soc* 2006;54:743-9.
199. NIH/NAEPP. NAEPP working group: Consideration for diagnosing and managing asthma in the elderly. Bethesda: National Institutes of Health; 1996.
200. Smyrnios NA. Asthma: a six-part strategy for managing older patients. *Geriatrics* 1997;52:36-40.
201. Braman SS. Drug treatment of asthma in the elderly. *Drugs* 1996;51:415-23.
202. Walsh LJ, Wong CA, Osborne J, Cooper S, Lewis SA, Pringle M, et al. Adverse effects of oral corticosteroids in relation to dose in patients with lung disease. *Thorax* 2001;56:279-84.
203. Sin DD, Tu JV. Underuse of inhaled steroid therapy in elderly patients with asthma. *Chest* 2001;119:720-5.
204. Creticos P, Knobil K, Edwards LD, Rickard KA, Dorinsky P. Loss of response to treatment with leukotriene receptor antagonists but not inhaled corticosteroids in patients over 50 years of age. *Ann Allergy Asthma Immunol* 2002;88:401-9.
205. Devoy MA, Fuller RW, Palmer JB. Are there any detrimental effects of the use of inhaled long-acting beta 2-agonists in the treatment of asthma? *Chest* 1995;107:1116-24.
206. Tafreshi MJ, Weinacker AB. Beta-adrenergic-blocking agents in bronchospastic diseases: a therapeutic dilemma. *Pharmacotherapy* 1999;19:974-8.
207. Prince DS, Carliner NH. Respiratory arrest following first dose of timolol ophthalmic solution. *Chest* 1983;84:640-1.
208. Dunn TL, Gerber MJ, Shen AS, Fernandez E, Iseman MD, Cherniack RM. The effect of topical ophthalmic instillation of timolol and betaxolol on lung function in asthmatic subjects. *Am Rev Respir Dis* 1986;133:264-8.
209. Anderson CJ, Bardana EJ Jr. Asthma in the elderly: the importance of patient education. *Compr Ther* 1996;22:375-83.
210. Chapman KR, Love L, Brubaker H. A comparison of breath-actuated and conventional metered-dose inhaler inhalation techniques in elderly subjects. *Chest* 1993;104:1332-7.
211. Allen SC, Jain M, Ragab S, Malik N. Acquisition and short-term retention of inhaler techniques require intact executive function in elderly subjects. *Age Ageing* 2003;32:299-302.
212. Allen SC, Prior A. What determines whether an elderly patient can use a metered dose inhaler correctly? *Br J Dis Chest* 1986;80:45-9.
213. Hayden ML. Asthma in the elderly. A diagnostic and management challenge. *Adv Nurse Pract* 2000;8:30-5, 83.
214. Malik A, Saltoun CA, Yarnold PR, Grammer LC. Prevalence of obstructive airways disease in the disadvantaged elderly of Chicago. *Allergy Asthma Proc* 2004;25:169-73.
215. Gruffydd-Jones K, Hollinghurst S, Ward S, Taylor G. Targeted routine asthma care in general practice using telephone triage. *Br J Gen Pract* 2005;55:918-23.
216. Patel RR, Saltoun CA, Grammer LC. Improving asthma care for the elderly: a randomized controlled trial using a simple telephone intervention. *J Asthma* 2009;46:30-5.
217. Emtner M, Hedin A, Stalenheim G. Asthmatic patients' views of a comprehensive asthma rehabilitation programme: a three-year follow-up. *Physiother Res Int* 1998;3:175-93.
218. Emtner M, Finne M, Stalenheim G. A 3-year follow-up of asthmatic patients participating in a 10-week rehabilitation program with emphasis on physical training. *Arch Phys Med Rehabil* 1998;79:539-44.
219. Emtner M, Herala M, Stalenheim G. High-intensity physical training in adults with asthma. A 10-week rehabilitation program. *Chest* 1996;109:323-30.
220. Zosky GR, Sly PD. Animal models of asthma. *Clin Exp Allergy* 2007;37:973-88.
221. Finkelman FD, Wills-Karp M. Usefulness and optimization of mouse models of allergic airway disease. *J Allergy Clin Immunol* 2008;121:603-6.
222. Kips JC, Anderson GP, Fredberg JJ, Herz U, Inman MD, Jordana M, et al. Murine models of asthma. *Eur Respir J* 2003;22:374-82.
223. Kumar RK, Foster PS. Modeling allergic asthma in mice: pitfalls and opportunities. *Am J Respir Cell Mol Biol* 2002;27:267-72.
224. Corry DB, Irvin CG. Promise and pitfalls in animal-based asthma research: building a better mousetrap. *Immunol Res* 2006;35:279-94.
225. Pauwels R, Bazin H, Platteau B, Van der Straeten M. The effect of age on IgE production in rats. *Immunology* 1979;36:145-9.

226. Ide K, Hayakawa H, Yagi T, Sato A, Koide Y, Yoshida A, et al. Decreased expression of Th2 type cytokine mRNA contributes to the lack of allergic bronchial inflammation in aged rats. *J Immunol* 1999;163:396-402.
227. Gelfand EW, Joetham A, Cui ZH, Balhorn A, Takeda K, Taube C, et al. Induction and maintenance of airway responsiveness to allergen challenge are determined at the age of initial sensitization. *J Immunol* 2004;173:1298-306.
228. Hasegawa A, Miki T, Hosokawa H, Hossain MB, Shimizu C, Hashimoto K, et al. Impaired GATA3-dependent chromatin remodeling and Th2 cell differentiation leading to attenuated allergic airway inflammation in aging mice. *J Immunol* 2006;176:2546-54.
229. Busse PJ, Zhang TF, Srivastava K, Schofield B, Li XM. Effect of ageing on pulmonary inflammation, airway hyperresponsiveness and T and B cell responses in antigen-sensitized and -challenged mice. *Clin Exp Allergy* 2007;37:1392-403.
230. Yagi T, Sato A, Hayakawa H, Ide K. Failure of aged rats to accumulate eosinophils in allergic inflammation of the airway. *J Allergy Clin Immunol* 1997;99:38-47.
231. Lee HK, Lim MY, Bok SM, Cho ES, Lee EM, Kim SW, et al. Age differences in cholinergic airway responsiveness in relation with muscarinic receptor subtypes. *Life Sci* 2007;81:204-9.
232. Takayanagi I, Kurata R, Kubo H, Kizawa Y. Effect of ageing on nicotine-induced contraction of guinea-pig bronchial preparation. *Gen Pharmacol* 1990;21:299-301.
233. Wills-Karp M. Effects of age on muscarinic agonist-induced contraction and IP accumulation in airway smooth muscle. *Life Sci* 1991;49:1039-45.
234. Sunyer J, Anto JM, Castellsague J, Soriano JB, Roca J. Total serum IgE is associated with asthma independently of specific IgE levels. The Spanish Group of the European Study of Asthma. *Eur Respir J* 1996;9:1880-4.
235. Green RH, Brightling CE, Woltmann G, Parker D, Wardlaw AJ, Pavord ID. Analysis of induced sputum in adults with asthma: identification of subgroup with isolated sputum neutrophilia and poor response to inhaled corticosteroids. *Thorax* 2002;57:875-9.
236. Wenzel SE, Schwartz LB, Langmack EL, Halliday JL, Trudeau JB, Gibbs RL, et al. Evidence that severe asthma can be divided pathologically into two inflammatory subtypes with distinct physiologic and clinical characteristics. *Am J Respir Crit Care Med* 1999;160:1001-8.
237. Gibson PG, Simpson JL, Saltos N. Heterogeneity of airway inflammation in persistent asthma: evidence of neutrophilic inflammation and increased sputum interleukin-8. *Chest* 2001;119:1329-36.
238. Simpson JL, Grissell TV, Douwes J, Scott RJ, Boyle MJ, Gibson PG. Innate immune activation in neutrophilic asthma and bronchiectasis. *Thorax* 2007;62:211-8.
239. Kim EY, Battaile JT, Patel AC, You Y, Agapov E, Grayson MH, et al. Persistent activation of an innate immune response translates respiratory viral infection into chronic lung disease. *Nat Med* 2008;14:633-40.
240. Molet S, Hamid Q, Davoine F, Nutku E, Taha R, Page N, et al. IL-17 is increased in asthmatic airways and induces human bronchial fibroblasts to produce cytokines. *J Allergy Clin Immunol* 2001;108:430-8.

**APPENDIX 1.**

**Planning Committee**

**Chair:**

Monroe James King, DO  
Asthma, Allergy and Immunology CRU  
University of South Florida  
Largo, Fla

**Co-chair:**

Nicola A. Hanania, MD, MS  
Section of Pulmonary and Critical Care Medicine  
Asthma Clinical Research Center  
Baylor College of Medicine  
Houston, Tex

**Members**

Sidney S. Braman, MD  
Warren Alpert Medical School, Brown University  
Division of Pulmonary, Critical Care, and Sleep Medicine  
Rhode Island Hospital  
Providence, RI

Carol Saltoun, MD  
Northwestern Feinberg School of Medicine  
Chicago, Ill

Susan G. Nayfield, MD, MSc  
Geriatrics and Clinical Gerontology Program  
National Institute on Aging  
Bethesda, Md

Robert A. Wise, MD  
Johns Hopkins University  
Baltimore, Md

**Participants**

Alexander Auais, MD  
Genentech, Inc  
Gladwyne, Pa

Jerome A. Dempsey, PhD  
University of Wisconsin School of Medicine  
Medical Sciences Building  
Madison, Wis

Cynthia M. Boyd, MD, MPH  
Division of Geriatric Medicine and Gerontology  
Johns Hopkins Department of Medicine  
Baltimore, Md

Paul Enright, MD  
The University of Arizona  
Tucson, Ariz

Ellen Brown  
American Lung Association  
Washington, DC

William B. Ershler, MD  
Harbor Hospital  
Baltimore, Md

Carlos A. Camargo, MD, DrPH  
Massachusetts General Hospital  
Boston, Mass

Ann Regina Falsey, MD  
University of Rochester School of Medicine  
Rochester General Hospital  
Rochester, NY

James E. Fish, MD  
Immunology, Tissue Growth and Repair  
Genentech  
Gladwyne, Pa

Carlos A. Vaz Fragoso, MD (OAIC)  
Clinical Epidemiology Research Center  
VA Connecticut Healthcare System  
West Haven, Conn

Paul Garbe, DVM, MPH  
Air Pollution and Respiratory Health Branch  
Division of Environmental Hazards and Health Effects  
National Center for Environmental Health, CDC  
Atlanta, Ga

Janet Holbrook, PhD, MPH  
Center for Clinical Trials  
Johns Hopkins University  
Baltimore, Md

Lydia Gilbert-McClain, MD  
Division of Pulmonary and Allergy Products  
US Food and Drug Administration  
Silver Spring, Md

Charles G. Irvin, PhD  
Vermont Lung Center  
University of Vermont  
Burlington, Vt

Ethan A. Halm, MD, MPH  
University of Texas Southwestern Medical Center  
Dallas, Tex

Dennis K. Ledford, MD  
University of South Florida College of Medicine  
James A. Haley VA Hospital  
Tampa, Fla

Robert G. Hamilton, PhD  
Johns Hopkins University School of Medicine  
Johns Hopkins Asthma and Allergy Center  
Baltimore, Md

Charlene Levine  
American Lung Association  
Washington, DC

Sameer K. Mathur, MD, PhD  
Department of Medicine  
Section of Allergy, Pulmonary and Critical Care  
University of Wisconsin School of Medicine  
Madison, Wis

John J. Lima, PharmD  
Clinical Pediatric Pharmacology  
Nemours Children's Clinic  
Jacksonville, Fla

Jeanne Moorman, MS  
Centers for Disease Control  
Atlanta, Ga

Katherine Pruitt  
American Lung Association  
Washington, DC

(continued)

**APPENDIX 1.** (Continued)

|   |  |
|---|--|
| Enid R. Neptune, MD<br>Division of Pulmonary and Critical Care Medicine<br>Institute of Genetic Medicine<br>Johns Hopkins School of Medicine<br>Baltimore, Md | Joe W. Ramsdell, MD<br>Division of General Internal Medicine<br>University of California, San Diego<br>San Diego, Calif  |
| Amy Pastva, PT, PhD<br>Department of Cell Biology<br>Duke University<br>Durham, NC  | Charles Reed, MD (keynote speaker)<br>Department of Medicine<br>Mayo Medical School<br>Rochester, Minn   |
| Kushang Patel, PhD<br>Laboratory of Epidemiology, Demography, and Biometry<br>National Institute on Aging<br>Bethesda, Md                                     | Theodore Reiss, MD<br>Merck & Co, Inc<br>Rahway, NJ  |
| Sergei Romashkan, MD<br>Clinical Trials Branch<br>Geriatrics and Clinical Gerontology Program<br>National Institute on Aging<br>Bethesda, Md                  | Linda Rogers, MD<br>NYU School of Medicine<br>Bellevue Hospital Asthma Clinic<br>New York, NY  |
| Mark F. Sands, MD<br>University at Buffalo, State University of New York<br>Division of Allergy/Immunology<br>Buffalo General Hospital<br>Buffalo, NY         | Michael L. Terrin, MD, CM, MPH<br>University of Maryland School of Medicine<br>Baltimore, Md   |
| David A. Stempel, MD<br>GlaxoSmithKline<br>Bellevue, Wash   | Sandra R. Wilson, PhD<br>Stanford University School of Medicine<br>Department of Health Services Research<br>Palo Alto Medical Foundation Research Institute<br>Palo Alto, Calif |
| Stephen I. Wasserman, MD<br>University of California—San Diego<br>San Diego, Calif  | Barbara P. Yawn, MD, MSc<br>Department of Family and Community Health<br>University of Minnesota<br>Rochester, Minn  |
| <b>NIH representatives</b>  |  |
| <b>National Institute on Aging (NIA)</b>  |  |
| Basil Eldadah, MD<br>Geriatrics Branch<br>Geriatrics and Clinical Gerontology Program<br>National Institute on Aging<br>Bethesda, Md                          | Ying Tian, MD, PhD<br>Geriatrics Branch<br>Geriatrics and Clinical Gerontology Program<br>National Institute on Aging<br>Bethesda, Md  |
| Evan Hadley, MD<br>Geriatrics and Clinical Gerontology Program<br>National Institute on Aging<br>Bethesda, Md   |  |
| <b>National Institute of Allergy and Infectious Diseases (NIAID)</b>  |  |
| Gang Dong, MD, PhD<br>Division of Allergy, Immunology and Transplantation<br>NIAID/NIH<br>Bethesda, Md  | Peter J. Gergen, MD, MPH<br>NIH/DAIT/AAIB<br>Bethesda, Md  |
| <b>National Heart Lung and Blood Institute (NHLBI)</b>  |  |
| James P. Kiley, MD<br>Division of Lung Diseases<br>National Heart Lung and Blood Institute<br>Bethesda, Md  | Virginia Taggart, MPH<br>Division of Lung Diseases<br>National Heart Lung and Blood Institute<br>Rockledge II, Room 10042<br>Bethesda, Md  |
| <b>National Institute of Nursing Research (NINR)</b>  |  |
| Karen Huss, PhD<br>NINR/NIH<br>Bethesda, Md   |  |