

1 **Convergent Antibody Responses to SARS-CoV-2 Infection in Convalescent Individuals**

2

3 Davide F. Robbiani^{1*,#}, Christian Gaebler^{1*}, Frauke Muecksch^{2*}, Julio C. C. Lorenzi^{1*}, Zijun
4 Wang^{1*}, Alice Cho^{1*}, Marianna Agudelo^{1*}, Christopher O. Barnes^{6*}, Anna Gazumyan^{1*}, Shlomo
5 Finkin^{1*}, Thomas Hagglof^{1*}, Thiago Y. Oliveira^{1*}, Charlotte Viant^{1*}, Arlene Hurley⁴, Hans-
6 Heinrich Hoffmann³, Katrina G. Millard¹, Rhonda G. Kost⁵, Melissa Cipolla¹, Kristie Gordon¹,
7 Filippo Bianchini¹, Spencer T. Chen¹, Victor Ramos¹, Roshni Patel¹, Juan Dizon¹, Irina
8 Shimeliovich¹, Pilar Mendoza¹, Harald Hartweg¹, Lilian Nogueira¹, Maggi Pack¹, Jill Horowitz¹,
9 Fabian Schmidt², Yiska Weisblum², Eleftherios Michailidis³, Alison W. Ashbrook³, Eric Waltari⁷,
10 John E. Pak⁷, Kathryn E. Huey-Tubman⁶, Nicholas Koranda⁶, Pauline R. Hoffman⁶, Anthony P.
11 West, Jr.⁶, Charles M. Rice³, Theodora Hatziioannou², Pamela J. Bjorkman⁶, Paul D. Bieniasz^{2,8,#},
12 Marina Caskey^{1,#}, Michel C. Nussenzweig^{1,8,#}

13

14 ¹Laboratory of Molecular Immunology, The Rockefeller University, New York, NY 10065, USA

15 ²Laboratory of Retrovirology, The Rockefeller University, New York, NY 10065, USA

16 ³Laboratory of Virology and Infectious Disease, The Rockefeller University, New York, NY
17 10065, USA

18 ⁴Hospital Program Direction, The Rockefeller University, New York, NY 10065, USA

19 ⁵Hospital Clinical Research Office, The Rockefeller University, New York, NY 10065, USA

20 ⁶Division of Biology and Biological Engineering, California Institute of Technology, Pasadena,
21 CA 91125, USA

22 ⁷Chan Zuckerberg Biohub, 499 Illinois Street, San Francisco, CA 94158, USA

23 ⁸Howard Hughes Medical Institute

24 *Equal contribution

25 #Send correspondence to Paul D. Bieniasz: pbieniasz@rockefeller.edu, Marina Caskey:

26 mcaskey@rockefeller.edu, Michel C. Nussenzweig: nussen@rockefeller.edu, or Davide F.

27 Robbiani: dobbiani@rockefeller.edu

28 **Abstract**

29 **During the COVID-19 pandemic, SARS-CoV-2 infected millions of people and claimed**
30 **hundreds of thousands of lives. Virus entry into cells depends on the receptor binding domain**
31 **(RBD) of the SARS-CoV-2 spike protein (S). Although there is no vaccine, it is likely that**
32 **antibodies will be essential for protection. However, little is known about the human**
33 **antibody response to SARS-CoV-2¹⁻⁵. Here we report on 149 COVID-19 convalescent**
34 **individuals. Plasmas collected an average of 39 days after the onset of symptoms had variable**
35 **half-maximal neutralizing titers ranging from undetectable in 33% to below 1:1000 in 79%,**
36 **while only 1% showed titers >1:5000. Antibody cloning revealed expanded clones of RBD-**
37 **specific memory B cells expressing closely related antibodies in different individuals. Despite**
38 **low plasma titers, antibodies to three distinct epitopes on RBD neutralized at half-maximal**
39 **inhibitory concentrations (IC₅₀s) as low as single digit ng/mL. Thus, most convalescent**
40 **plasmas obtained from individuals who recover from COVID-19 do not contain high levels**
41 **of neutralizing activity. Nevertheless, rare but recurring RBD-specific antibodies with potent**
42 **antiviral activity were found in all individuals tested, suggesting that a vaccine designed to**
43 **elicit such antibodies could be broadly effective.**

44 Between April 1 and May 8, 2020, 157 eligible participants enrolled in the study. Of these, 111
45 (70.7%) were individuals diagnosed with SARS-CoV-2 infection by RT-PCR (cases), and 46
46 (29.3%) were close contacts of individuals diagnosed with SARS-CoV-2 infection (contacts).
47 While inclusion criteria allowed for enrollment of asymptomatic participants, 8 contacts that did
48 not develop symptoms were excluded from further analyses. The 149 cases and contacts were free
49 of symptoms suggestive of COVID-19 for at least 14 days at the time of sample collection.
50 Participant demographics and clinical characteristics are shown in Table 1 and Extended Data
51 Tables 1 and 2. Only one individual who tested positive for SARS-CoV-2 infection by RT-PCR
52 remained asymptomatic. The other 148 participants reported symptoms suggestive of COVID-19
53 with an average onset of approximately 39 days (range 17 to 67 days) before sample collection. In
54 this cohort, symptoms lasted for an average of 12 days (0-35 days), and 11 (7%) of the participants
55 were hospitalized. The most common symptoms were fever (83.9%), fatigue (71.1%), cough
56 (62.4%) and myalgia (61.7%) while baseline comorbidities were infrequent (10.7%) (Table 1 and
57 Extended Data Tables 1 and 2). There were no significant differences in duration or severity (see
58 Methods) of symptoms, or in time from onset of symptoms to sample collection between genders
59 or between cases and contacts. There was no age difference between females and males in our
60 cohort (Extended Data Fig. 1).

61
62 Plasma samples were tested for binding to the SARS-CoV-2 RBD and trimeric spike (S) proteins
63 by ELISA using anti-IgG or -IgM secondary antibodies for detection (Fig. 1, Extended Data Table
64 1 and Extended Data Figs. 2 and 3). Eight independent negative controls and the plasma sample
65 from participant 21 (COV21) were included for normalization of the area under the curve (AUC).
66 Overall, 78% and 70% of the plasma samples tested showed anti-RBD and anti-S IgG AUCs that

67 were at least 2 standard deviations above the control (Fig. 1 a, b). In contrast, only 15% and 34%
68 of the plasma samples showed IgM responses to anti-RBD and anti-S that were at least 2 standard
69 deviations above control, respectively (Fig. 1 c, d). There was no positive correlation between anti-
70 RBD or -S IgG or IgM levels and duration of symptoms or the timing of sample collection relative
71 to onset of symptoms (Fig. 1e, and Extended Data Figs. 3 a-c and 3 g-j). On the contrary, as might
72 be expected, anti-RBD IgM titers were negatively correlated with duration of symptoms and the
73 timing of sample collection (Fig. 1e and Extended Data Fig. 3h). Anti-RBD IgG levels were
74 modestly correlated to age, and the severity of symptoms including hospitalization (Fig. 1 f, g and
75 Extended Data Fig. 3k). Interestingly, females had lower anti-RBD and -S IgG titers than males
76 (Fig. 1h, Extended Data Fig. 2f).

77
78 To measure the neutralizing activity in convalescent plasmas we used HIV-1-based virions
79 carrying a nanoluciferase reporter that were pseudotyped with the SARS-CoV-2 spike (SARS-
80 CoV-2 pseudovirus, see Methods, Fig. 2 and Extended Data Fig. 4). The overall level of
81 neutralizing activity in the cohort, as measured by the half-maximal neutralizing titer (NT_{50}) was
82 generally low, with 33% undetectable and 79% below 1,000 (Fig. 2 a, b). The geometric mean
83 NT_{50} was 121 (arithmetic mean = 714), and only 2 individuals reached NT_{50} s above 5,000 (Fig. 2
84 a, b and Extended Data Table 1).

85
86 Notably, levels of anti-RBD- and -S IgG antibodies correlated strongly with NT_{50} (Fig. 2 c, d).
87 Neutralizing activity also correlated with age, duration of symptoms and symptom severity
88 (Extended Data Fig. 5). Consistent with this observation, hospitalized individuals with longer
89 symptom duration showed slightly higher average levels of neutralizing activity than non-

90 hospitalized individuals ($p=0.0495$, Fig. 2e). Finally, we observed a significant difference in
91 neutralizing activity between males and females ($p=0.0031$, Fig. 2f). The difference between males
92 and females was consistent with higher anti-RBD and -S IgG titers in males, and could not be
93 attributed to age, severity, timing of sample collection relative to onset of symptoms or duration
94 of symptoms (Fig. 1h, Extended Data Fig. 1 a-d and 2f).

95
96 To determine the nature of the antibodies elicited by SARS-CoV-2 infection we used flow
97 cytometry to isolate individual B lymphocytes with receptors that bound to RBD from the blood
98 of 6 selected individuals including the 2 top and 4 high to intermediate neutralizers (Fig. 3). The
99 frequency of antigen-specific B cells, identified by their ability to bind to both Phycoerythrin (PE)-
100 and AF647-labeled RBD, ranged from 0.07 to 0.005% of all circulating B cells in COVID-19
101 convalescents but they were undetectable in pre-COVID-19 controls (Fig. 3a and Extended Data
102 Fig. 6). We obtained 534 paired IgG heavy and light chain (IGH and IGL) sequences by reverse
103 transcription and subsequent PCR from individual RBD-binding B cells from the 6 convalescent
104 individuals (see Methods and Extended Data Table 3). When compared to the human antibody
105 repertoire, several IGHV and IGLV genes were significantly over-represented (Extended Data Fig.
106 7). The average number of V genes nucleotide mutations for IGH and IGL was 4.2 and 2.8,
107 respectively (Extended Data Fig. 8), which is lower than in antibodies cloned from individuals
108 suffering from chronic infections such as Hepatitis B or HIV-1, and similar to antibodies derived
109 from primary malaria infection or non-antigen-enriched circulating IgG memory cells⁶⁻⁸ (Wang et
110 al, in press, <https://www.biorxiv.org/content/10.1101/2020.03.04.976159v1.full>). Among other
111 antibody features, IGH CDR3 length was indistinguishable from the reported norm and
112 hydrophobicity was below average (Extended Data Fig. 8)⁹.

113

114 As is the case with other human pathogens, there were expanded clones of viral antigen binding B
115 cells in all COVID-19 individuals tested (see Methods and Fig. 3 b,c). Overall, 32.2% of the
116 recovered IGH and IGL sequences were from clonally expanded B cells (range 21.8-57.4% across
117 individuals, Fig. 3b). Antibodies that shared specific combinations of IGHV and IGLV genes in
118 different individuals comprised 14% of all the clonal sequences (colored pie slices in Fig. 3 b,c).
119 Remarkably, the amino acid sequences of some antibodies found in different individuals were
120 nearly identical (Fig. 3 e,d). For example, antibodies expressed by clonally expanded B cells with
121 IGHV1-58/IGKV3-20 and IGHV3-30-3/IGKV1-39 found repeatedly in different individuals had
122 amino acid sequence identities of up to 99% and 92%, respectively (Fig. 3d and Extended Data
123 Table 4). We conclude that the IgG memory response to the SARS-CoV-2 RBD is rich in recurrent
124 and clonally expanded antibody sequences.

125

126 To examine the binding properties of anti-SARS-CoV-2 antibodies, we expressed 84
127 representative antibodies, 56 from clones and 28 from singlets (Extended Data Table 5). ELISA
128 assays showed that 94% (79 out of 84) of the antibodies tested including clonal and unique
129 sequences bound to the SARS-CoV-2 RBD with an average half-maximal effective concentration
130 (EC_{50}) of 6.1 ng/mL (Fig. 4a and Extended Data Fig. 9a). A fraction of those (7 out of 79, or 9%)
131 cross-reacted with the RBD of SARS-CoV with a mean EC_{50} of 120.1 ng/mL (Extended Data Fig.
132 9b and c). No significant cross-reactivity was noted to the RBDs of MERS, HCoV-OC43, HCoV-
133 229E or HCoV-NL63.

134

135 To determine whether the monoclonal antibodies have neutralizing activity, we tested them against
136 the SARS-CoV-2 pseudovirus (Fig. 4 and Extended Data Table 6). Among 79 RBD binding
137 antibodies tested, we found 40 that neutralized SARS-CoV-2 pseudovirus with nanogram per
138 milliliter half-maximal inhibitory concentrations (IC_{50} s) ranging from 3 to 709 (Fig. 4 b,c and e,
139 Extended Data Table 6). A subset of the most potent of these antibodies were also tested against
140 authentic SARS-CoV-2 and neutralized with IC_{50} s of less than 5 ng/ml (Fig. 4 d,e).

141
142 Potent neutralizing antibodies were found in individuals irrespective of their plasma NT_{50} s. For
143 example, C121, C144, and C135 with IC_{50} s of 1.64, 2.55 and 2.98 ng/mL against authentic SARS-
144 CoV-2, respectively, were obtained from individuals COV107, COV47, and COV72 whose
145 plasma NT_{50} values were of 297, 10,433 and 3,138, respectively (Figs. 2b and 4). Finally, clones
146 of antibodies with shared IGHV and IGLV genes were among the best neutralizers, e.g., antibody
147 C002 composed of IGHV3-30/IGKV1-39 is shared by the 2 donors with the best plasma
148 neutralizing activity (red pie slice in Fig. 3b and Fig. 4). We conclude that even individuals with
149 modest plasma neutralizing activity harbor rare IgG memory B cells that produce potent SARS-
150 CoV-2 neutralizing antibodies.

151
152 To determine whether human anti-SARS-CoV-2 monoclonal antibodies with neutralizing activity
153 can bind to distinct domains on the RBD, we performed bilayer interferometry experiments in
154 which a preformed antibody-RBD immune complex was exposed to a second monoclonal. The
155 antibodies tested comprised 3 groups, all of which differ in their binding properties from CR3022,
156 an antibody that neutralizes SARS-CoV and binds to, but does not neutralize SARS-CoV-2^{10,11}.
157 Representatives of each of the 3 groups include: C144 and C101 in Group 1; C121 and C009 in

158 Group 2; C135 in Group 3. All of these antibodies can bind after CR3022. Groups 1 and 2 also
159 bind after Group 3, and Groups 1 and 2 differ in that Group 1 can bind after Group 2 but not vice
160 versa (Fig. 4 f-n). We conclude that similar to SARS-CoV, there are multiple distinct neutralizing
161 epitopes on the RBD of SARS-CoV-2.

162

163 To further define the binding characteristics of Groups 1 and 2 antibodies, we imaged SARS-CoV-
164 2 S–Fab complexes by negative stain electron microscopy (nsEM) using C002 (Group 1, an
165 IGHV3-30/IGKV1-39 antibody, which is clonally expanded in 2 donors), C119 and C121 (both in
166 Group 2) Fabs (Fig. 4 f-r and Extended Data Fig. 10). Consistent with the conformational
167 flexibility of the RBD, 2D class averages showed heterogeneity in both occupancy and
168 conformation of bound Fabs for both groups (Fig. 4o-q). The 3D reconstructions of S-Fab
169 complexes revealed both a fully-occupied RBD and partial density for a second bound Fab,
170 suggesting that Fabs from both groups are able to recognize “up” and “down” states of the RBD
171 as previously described for some of the antibodies targeting this epitope^{12,13}. The 3D
172 reconstructions are also consistent with BLI measurements indicating that Groups 1 and 2
173 antibodies bind a RBD epitope distinct from antibody CR3022, which is only accessible on the
174 RBD “up” conformational state¹¹, and bind the RBD with different angles of approach, with Group
175 1 antibodies most similar to the approach angle of the SARS-CoV antibody S230 (Fig. 4r)¹⁴.

176

177 Human monoclonal antibodies with neutralizing activity against pathogens ranging from viruses
178 to parasites have been obtained from naturally infected individuals by single cell antibody cloning.
179 Several have been shown to be effective in protection and therapy in model organisms and in early
180 phase clinical studies, but only one antiviral monoclonal is currently in clinical use¹⁵. Antibodies

181 are relatively expensive and more difficult to produce than small molecule drugs. However, they
182 differ from drugs in that they can engage the host immune system through their constant domains
183 that bind to Fc gamma receptors on host immune cells¹⁶. These interactions can enhance immunity
184 and help clear the pathogen or infected cells, but they can also lead to disease enhancement during
185 Dengue¹⁷ and possibly coronavirus infections¹⁸. This problem has impeded Dengue vaccine
186 development but would not interfere with the clinical use of potent neutralizing antibodies that can
187 be modified to prevent Fc gamma receptor interactions and remain protective against viral
188 pathogens¹⁹.

189
190 Antibodies are essential elements of most vaccines and will likely be crucial component of an
191 effective vaccine against SARS-CoV-2²⁰(PMID:32434945; PMID:32434946). Recurrent
192 antibodies have been observed in other infectious diseases and vaccinal responses²¹⁻²⁴(Wang et
193 al, in press, <https://www.biorxiv.org/content/10.1101/2020.03.04.976159v1.full>). The observation
194 that plasma neutralizing activity is low in most convalescent individuals, but that recurrent anti-
195 SARS-CoV-2 RBD antibodies with potent neutralizing activity can be found in individuals with
196 unexceptional plasma neutralizing activity suggests that humans are intrinsically capable of
197 generating anti-RBD antibodies that potently neutralize SARS-CoV-2. Thus, vaccines that
198 selectively and efficiently induce antibodies targeting the SARS-CoV-2 RBD may be especially
199 effective.

200 **Table**


Table 1. Cohort characteristics

Gender	n	Average age	Case/Contact	Average duration		Average Sx Severity (0-10)	ELISA binding (AUC)				Neutralization (NT50)
				Sx total	Sx onset to visit		RBD		S		
						IgG	IgM	IgG	IgM		
Male	83	45 (19-76)	65/18	12 (0-31)	39 (21-63)	5.8 (0-10)	2.44	1.61	4.65	1.62	867
Female	66	42 (19-75)	46/20	12 (1-35)	38 (17-67)	5.4 (1-9)	1.99	1.58	4.36	1.86	522

Sx=symptoms

201


202 **Figures**



203

204 **Figure 1. Plasma antibodies against SARS-CoV-2.** a-d, Graphs show results of ELISAs
205 measuring plasma reactivity to RBD (a, b) and S protein (c, d). Left shows optical density units at
206 450 nm (OD, Y axis) and reciprocal plasma dilutions (X axis). Negative controls in black;
207 individuals 21, and 47 in blue and red lines and arrowheads, respectively. Right shows normalized
208 area under the curve (AUC) for controls and each of 149 individuals in the cohort. e, Symptom


209 (Sx) onset to time of sample collection in days (X axis) plotted against normalized AUC for IgM
210 binding to RBD (Y axis) $r=0.5517$ and $p<0.0001$. **f**, Participant age in years (X axis) plotted
211 against normalized AUC for IgG binding to RBD (Y axis) $r=0.1827$ and $p=0.0258$. The r and p
212 values for the correlations in **e** and **f** were determined by two-tailed Spearman's. **g**, IgG anti-RBD
213 normalized AUC for outpatients and hospitalized individuals $p=0.0178$. **h**, IgG anti-RBD
214 normalized AUC for males and females $p=0.0063$. For **g** and **h** horizontal bars indicate median
215 values. Statistical significance was determined using two-tailed Mann-Whitney U test.



216

217 **Figure 2. Neutralization of SARS-CoV-2 pseudovirus by plasma.** a, Graph shows normalized
 218 relative luminescence values (RLU, Y axis) in cell lysates of 293T_{ACE2} cells 48 hours after
 219 infection with nanoluc-expressing SARS-CoV-2 pseudovirus in the presence of increasing
 220 concentrations of plasma (X axis) derived from 149 participants (grey, except individuals 47 and
 221 21 in red, and blue lines, bars and arrowheads, respectively) and 3 negative controls (black lines).

222 Standard deviations of duplicates of one representative experiment are shown. **b**, Ranked average
223 half-maximal inhibitory plasma neutralizing titer (NT_{50}) for the 59 of 149 individuals with NT_{50} s
224 >500 and individual 107. See also Extended Data Table 1. Asterisks indicate donors from which
225 antibody sequences were derived. **c**, AUC for anti-RBD IgG ELISA (X axis) plotted against NT_{50}
226 (Y axis) $r=0.6432$, $p<0.0001$. **d**, AUC for anti-S IgG ELISA (X axis) plotted against NT_{50} (Y
227 axis) $r=0.6721$, $p<0.0001$. **e**, NT_{50} for outpatients and hospitalized individuals $p=0.0495$. **f**, NT_{50}
228 for all males and females in the cohort $p=0.0031$. Dotted line in c to f ($NT_{50}=5$) represents lower
229 limit of detection (LLOD). Samples with undetectable neutralizing titers were plotted at LLOD.
230 Correlations in **c** and **d** were determined by two-tailed Spearman's. Statistical significance in **e** and
231 **f** was determined using two-tailed Mann-Whitney U test. Horizontal bars indicate median values.



232

233 **Figure 3. Anti-SARS-CoV-2 RBD antibodies.** **a.** Representative flow cytometry plots showing

234 dual AF647- and PE-RBD binding B cells in control and 6 study individuals (for gating strategy

235 see Extended Data Fig. 6). Percentages of antigen specific B cells are indicated. Control is a

236 healthy control sample obtained pre-COVID-19. **b,** Pie charts depicting the distribution of


237 antibody sequences from 6 individuals. The number in the inner circle indicates the number of

238 sequences analyzed for the individual denoted above the circle. White indicates sequences isolated

239 only once, and grey or colored pie slices are proportional to the number of clonally related

240 sequences. Red, blue, orange and yellow pie slices indicate clones that share the same IGHV and

241 IGLV genes. **c**, Circos plot shows sequences from all 6 individuals with clonal relationships
242 depicted as in **b**. Interconnecting lines indicate the relationship between antibodies that share V
243 and J gene segment sequences at both IGH and IGL. Purple, green and gray lines connect related
244 clones, clones and singles, and singles to each other, respectively. **d**, Sample sequence alignment
245 for antibodies originating from different individuals that display highly similar IGH V(D)J and
246 IGL VJ sequences including CDR3s. Amino acid differences in CDR3s to the bolded reference
247 sequence above are indicated in red and dots represent identities.



248

249 **Figure 4. Anti-SARS-CoV-2 RBD antibody reactivity.** **a**, Graph show results of ELISA assays
 250 measuring monoclonal antibody reactivity to RBD. Optical density units at 450 nm (OD, Y axis)
 251 vs. antibody concentrations (X axis). C121, C135 C144 and isotype control in red, green, purple,
 252 and black respectively, in all panels. **b**, Graph shows normalized relative luminescence values
 253 (RLU, Y axis) in cell lysates of 293T_{ACE2} cells 48 hours after infection with SARS-CoV-2
 254 pseudovirus in the presence of increasing concentrations of monoclonal antibodies (X axis). **c**,
 255 RLU for SARS-CoV-2 pseudovirus assay (Y axis) vs. titration of monoclonal antibodies C121,
 256 C135 and C144 in one of two independent experiments (see Extended Data Table 6). **d**, SARS-

257 CoV-2 real virus neutralization assay. Infected cells (Y axis) vs. titration of monoclonal antibodies
258 C121, C135 and C144 in two independent experiments. For a and b panels, isotype control
259 antibody in black. **e**, IC50s for antibodies assayed in **b** and **d**. **f**, Diagrammatic representation of
260 biolayer interferometry experiment. **g**, Graph shows binding of C144, C101, C121, C009, C135,
261 and CR3022^{10,11} to RBD. **h-m**, Secondary antibody binding to preformed IgG-RBD complexes
262 (Ab1). The table displays the shift in nanometers after second antibody (Ab2) binding to the
263 antigen in the presence of the first antibody (Ab1). Values are normalized by the subtraction of the
264 autologous antibody control. **o-q**, Representative 2D-class averages and 3D reconstructed
265 volumes for SARS-CoV-S 2P trimers complexed with C002, C119, and C121 Fabs. 2D-class
266 averages with observable Fab density are boxed. **r**, Overlay of S-Fab complexes with fully-
267 occupied C002 (blue), C121 (magenta) and C119 (orange) Fabs aligned on the RBD “up”
268 conformational state. The SARS-CoV-2 S model with 1 “up” RBD state (PDB 6VYB) was fit into
269 the density and the SARS-CoV antibody S230 (PDB 6NB6) shown as reference (green ribbon).
270
271

272 **Methods**

273 **Study participants.** Study participants were recruited at the Rockefeller University Hospital in
274 New York from April 1 through May 8, 2020. Eligible participants were adults aged 18-76 years
275 who were either diagnosed with SARS-CoV-2 infection by RT-PCR and were free of symptoms
276 of COVID-19 for at least 14 days (cases), or who were close contacts (e.g., household, co-workers,
277 members of same religious community) with someone who had been diagnosed with SARS-CoV-
278 2 infection by RT-PCR and were free of symptoms suggestive of COVID-19 for at least 14 days
279 (contacts). Exclusion criteria included presence of symptoms suggestive of active SARS-CoV-2
280 infection, or hemoglobin < 12 g/dL for males and < 11 g/dL for females.

281 Most study participants were residents of the Greater New York City tri-state region and were
282 enrolled sequentially according to eligibility criteria. Participants were first interviewed by phone
283 to collect information on their clinical presentation, and subsequently presented to the Rockefeller
284 University Hospital for a single blood sample collection. Participants were asked to rate the
285 highest severity of their symptoms on a numeric rating scale ranging from 0 to 10. The score was
286 adapted from the pain scale chart, where 0 was the lack of symptoms, 4 were distressing symptoms
287 (e.g. fatigue, myalgia, fever, cough, shortness of breath) that interfered with daily living activities,
288 7 were disabling symptoms that prevented the performance of daily living activities, and 10 was
289 unimaginable/unspeakable discomfort (in this case, distress due to shortness of breath). All
290 participants provided written informed consent before participation in the study and the study was
291 conducted in accordance with Good Clinical Practice.

292

293 **Blood samples processing and storage.** Peripheral Blood Mononuclear Cells (PBMCs) were
294 obtained by gradient centrifugation and stored in liquid nitrogen in the presence of FCS and

295 DMSO. Heparinized plasma and serum samples were aliquoted and stored at -20°C or less. Prior
296 to experiments, aliquots of plasma samples were heat-inactivated (56°C for 1 hour) and then stored
297 at 4°C.

298

299 **Cloning, expression and purification of recombinant coronavirus proteins.** Codon-optimized

300 nucleotide sequences encoding the SARS-CoV-2 S ectodomain (residues 16-1206) and receptor

301 binding domain (RBD; residues 331-524) were synthesized and subcloned into the mammalian

302 expression pTwist-CMV BetaGlobin vector by Twist Bioscience Technologies based on an early

303 SARS-CoV-2 sequence isolate (GenBank MN985325.1). The SARS-CoV-2 RBD construct

304 included an N-terminal human IL-2 signal peptide and dual C-terminal tags ((GGGGS)₂-

305 HHHHHHHH (octa-histidine), and GLNDIFEAQKIEWHE (AviTag)). In addition, the

306 corresponding S1^B or receptor binding domains for SARS-CoV S (residues 318-510; GenBank

307 AAP13441.1), MERS-CoV S (residues 367-588; GenBank JX869059.2), HCoV-NL63 (residues

308 481-614; GenBank AAS58177.1), HCoV-OC43 (residues 324-632; GenBank AAT84362.1), and

309 HCoV-229E (residues 286-434; GenBank AAK32191.1) were synthesized with the same N- and

310 C-terminal extensions as the SARS-CoV-2 RBD construct and subcloned into the mammalian

311 expression pTwist-CMV BetaGlobin vector (Twist Bioscience Technologies). The SARS-CoV-2

312 S ectodomain was modified as previously described⁴. Briefly, the S ectodomain construct included

313 an N-terminal mu-phosphatase signal peptide, 2P stabilizing mutations (K986P and V987P),

314 mutations to remove the S1/S2 furin cleavage site (₆₈₂RRAR₆₈₅ to GSAS), a C-terminal extension

315 (IKGSG-RENLYFQG (TEV protease site), GGGSG-YIPEAPRDGQAYVRKDGWVLLSTFL

316 (foldon trimerization motif), G-HHHHHHHH (octa-histidine tag), and GLNDIFEAQKIEWHE

317 (AviTag)). The SARS-CoV-2 S 2P ectodomain and RBD constructs were produced by transient

318 transfection of 500 mL of Expi293 cells (Thermo Fisher) and purified from clarified transfected
319 cell supernatants four days post-transfection using Ni²⁺-NTA affinity chromatography (GE Life
320 Sciences). Affinity-purified proteins were concentrated and further purified by size-exclusion
321 chromatography (SEC) using a Superdex200 16/60 column (GE Life Sciences) running in 1x TBS
322 (20 mM Tris-HCl pH 8.0, 150 mM NaCl, and 0.02% NaN₃). Peak fractions were analyzed by SDS-
323 PAGE, and fractions corresponding to soluble S 2P trimers or monomeric RBD proteins were
324 pooled and stored at 4°C.

325
326 **ELISAs.** ELISAs to evaluate antibodies binding to SARS-CoV-2 RBD and trimeric spike proteins
327 were performed by coating of high binding 96 half well plates (Corning #3690) with 50 µL per
328 well of a 1µg/mL protein solution in PBS overnight at 4°C. Plates were washed 6 times with
329 washing buffer (1xPBS with 0.05% Tween 20 (Sigma-Aldrich)) and incubated with 170 µL per
330 well blocking buffer (1xPBS with 2% BSA and 0.05% Tween20 (Sigma)) for 1 hour at room
331 temperature (RT). Immediately after blocking, monoclonal antibodies or plasma samples were
332 added in PBS and incubated for 1 hr at RT. Plasma samples were assayed at a 1:200 starting
333 dilution and seven additional 3-fold serial dilutions. Monoclonal antibodies were tested at 10µg/ml
334 starting concentration and 10 additional 4-fold serial dilutions. Plates were washed 6 times with
335 washing buffer and then incubated with anti-human IgG or IgM secondary antibody conjugated to
336 horseradish peroxidase (HRP) (Jackson Immuno Research 109-036-088 and 109-035-129) in
337 blocking buffer at a 1:5000 dilution. Plates were developed by addition of the HRP substrate, TMB
338 (ThermoFisher) for 10 minutes, then the developing reaction was stopped by adding 50µl 1M
339 H₂SO₄ and absorbance was measured at 450nm with an ELISA microplate reader (FluoStar
340 Omega, BMG Labtech). For plasma samples, a positive control (plasma from patient COV21,

341 diluted 200-fold in PBS) was added in duplicate to every assay plate. The average of its signal was
342 used for normalization of all the other values on the same plate with Excel software prior to
343 calculating the area under the curve using Prism 8 (GraphPad). For monoclonal antibodies, the
344 half-maximal effective concentration (EC_{50}) was determined using 4-parameter nonlinear
345 regression (GraphPad Prism).

346
347 **293T_{ACE2} cells.** For constitutive expression of ACE2 in 293T cells, a cDNA encoding ACE2,
348 carrying two inactivating mutations in the catalytic site (H374N & H378N), was inserted into CSIB
349 3' to the SFFV promoter²⁵. 293T_{ACE2} cells were generated by transduction with CSIB based virus
350 followed by selection with 5 μ g/ml Blasticidin.

351
352 **SARS-CoV-2 and SARS-CoV pseudotyped reporter viruses.** A plasmid expressing a C-
353 terminally truncated SARS-CoV-2 S protein (pSARS-CoV2-S_{trunc}) was generated by insertion of
354 a human-codon optimized cDNA encoding SARS-CoV-2 S lacking the C-terminal 19 codons
355 (Geneart) into pCR3.1. The S ORF was taken from “Wuhan seafood market pneumonia virus
356 isolate Wuhan-Hu-1” (NC_045512). For expression of full-length SARS-CoV S protein, “Human
357 SARS coronavirus Spike glycoprotein Gene ORF cDNA clone expression plasmid (Codon
358 Optimized)” (here referred to as pSARS-CoV-S) was obtained from SinoBiological (Cat:
359 VG40150-G-N). An *env*-inactivated HIV-1 reporter construct (pNL4-3 Δ Env-nanoluc) was
360 generated from pNL4-3²⁶ by introducing a 940 bp deletion 3' to the *vpu* stop-codon, resulting in a
361 frameshift in *env*. The human codon-optimized nanoluc Luciferase reporter gene (*Nluc*, Promega)
362 was inserted in place of nucleotides 1-100 of the *nef*-gene. To generate pseudotyped viral stocks,
363 293T cells were transfected with pNL4-3 Δ Env-nanoluc and pSARS-CoV2-S_{trunc} or pSARS-CoV-

364 S using polyethyleneimine. Co-transfection of pNL4-3ΔEnv-nanoluc and S-expression plasmids
365 leads to production of HIV-1-based virions carrying either the SARS-CoV-2 or SARS-CoV spike
366 protein on the surface. Eight hours after transfection, cells were washed twice with PBS and fresh
367 media was added. Supernatants containing virions were harvested 48 hours post transfection,
368 filtered and stored at -80°C. Infectivity of virions was determined by titration on 293T_{ACE2} cells.

369
370 **Pseudotyped virus neutralization assay.** Five-fold serially diluted plasma from COVID-19
371 convalescent individuals and healthy donors or four-fold serially diluted monoclonal antibodies
372 were incubated with the SARS-CoV-2 or SARS-CoV pseudotyped virus for 1 hour at 37°C
373 degrees. The mixture was subsequently incubated with 293T_{ACE2} cells for 48 hours after which
374 cells were washed twice with PBS and lysed with Luciferase Cell Culture Lysis 5x reagent
375 (Promega). Nanoluc Luciferase activity in lysates was measured using the Nano-Glo Luciferase
376 Assay System (Promega). Relative luminescence units obtained were normalized to those derived
377 from cells infected with SARS-CoV-2 or SARS-CoV pseudotyped virus in the absence of plasma
378 or monoclonal antibodies. The half-maximal inhibitory concentration for plasma (NT₅₀) or
379 monoclonal antibodies (IC₅₀) was determined using 4-parameter nonlinear regression (GraphPad
380 Prism).

381 **Cell lines, virus and virus titration.** VeroE6 kidney epithelial cells (*Chlorocebus sabaesus*) and
382 Huh-7.5 hepatoma cells (*H. sapiens*) were cultured in Dulbecco's Modified Eagle Medium
383 (DMEM) supplemented with 1% nonessential amino acids (NEAA) and 10% fetal bovine serum
384 (FBS) at 37°C and 5% CO₂. All cell lines have been tested negative for contamination with
385 mycoplasma and were obtained from the ATCC (with the exception for Huh-7.5).

386 SARS-CoV-2, strain USA-WA1/2020, was obtained from BEI Resources and amplified in VeroE6
387 cells at 33°C. Viral titers were measured on Huh-7.5 cells by standard plaque assay (PA). Briefly,
388 500 µL of serial 10-fold virus dilutions in Opti-MEM were used to infect 400,000 cells seeded the
389 day prior in a 6-well plate format. After 90 min adsorption, the virus inoculum was removed, and
390 cells were overlaid with DMEM containing 10% FBS with 1.2% microcrystalline cellulose
391 (Avicel). Cells were incubated for five days at 33°C, followed by fixation with 3.5% formaldehyde
392 and crystal violet staining for plaque enumeration. All experiments were performed in a biosafety
393 level 3 laboratory.

394 **Microscopy-based neutralization assay of authentic SARS-CoV-2.** The day prior to infection
395 VeroE6 cells were seeded at 12,500 cells/well into 96-well plates. Antibodies were serially diluted
396 in BA-1, mixed with a constant amount of SARS-CoV-2 (grown in VeroE6) and incubated for 60
397 min at 37°C. The antibody-virus-mix was then directly applied to VeroE6 cells (MOI of ~0.1
398 PFU/cell). Cells were fixed 18 hours post infection by adding an equal volume of 7% formaldehyde
399 to the wells, followed by permeabilization with 0.1% Triton X-100 for 10 min. After extensive
400 washing, cells were incubated for 1 hour at room temperature with blocking solution of 5% goat
401 serum in PBS (catalog no. 005–000-121; Jackson ImmunoResearch). A rabbit polyclonal anti-
402 SARS-CoV-2 nucleocapsid antibody (catalog no. GTX135357; GeneTex) was added to the cells
403 at 1:500 dilution in blocking solution and incubated at 4°C overnight. A goat anti-rabbit
404 AlexaFluor 594 (catalog no. A-11012; Life Technologies) at a dilution of 1:2,000 was used as a
405 secondary antibody. Nuclei were stained with Hoechst 33342 (catalog no. 62249; Thermo
406 Scientific) at a 1:1,000 dilution. Images were acquired with a fluorescence microscope and
407 analyzed using ImageXpress Micro XLS (Molecular Devices, Sunnyvale, CA). All statistical
408 analyses were done using Prism 8 software (GraphPad).

409 **Biotinylation of viral protein for use in flow cytometry.** Purified and Avi-tagged SARS-CoV-2
410 RBD was biotinylated using the Biotin-Protein Ligase-BIRA kit according to manufacturer's
411 instructions (Avidity). Ovalbumin (Sigma, A5503-1G) was biotinylated using the EZ-Link Sulfo-
412 NHS-LC-Biotinylation kit according to the manufacturer's instructions (Thermo Scientific).
413 Biotinylated Ovalbumin was conjugated to streptavidin-BV711 (BD biosciences, 563262) and
414 RBD to streptavidin-PE (BD biosciences, 554061) and streptavidin-Alexa Fluor 647 (AF647,
415 Biologend, 405237) respectively²⁷.

416
417 **Single cell sorting by flow cytometry.** PBMCs were enriched for B cells by negative selection
418 using a pan B cell isolation kit according to the manufacturer's instructions (Miltenyi Biotec, 130-
419 101-638). The enriched B cells were incubated in FACS buffer (1 X Phosphate-buffered Saline
420 (PBS), 2% calf serum, 1 mM EDTA) with the following anti-human antibodies: anti-CD20-PECy7
421 (BD Biosciences, 335793), anti-CD3-APC-eFluro 780 (Invitrogen, 47-0037-41), anti-CD8-APC-
422 eFluro 780 (Invitrogen, 47-0086-42), anti-CD16-APC-eFluro 780 (Invitrogen, 47-0168-41), anti-
423 CD14-APC-eFluro 780 (Invitrogen, 47-0149-42), as well as Zombie NIR (BioLegend, 423105),
424 and fluorophore-labeled RBD and Ovalbumin for 30 minutes on ice²⁷. Single CD3⁻CD8⁻CD16⁻
425 CD20⁺Ova⁻RBD-PE⁺RBD-AF647⁺ B cells were sorted into individual wells of a 96-well plates
426 containing 4 µl of lysis buffer (0.5 X PBS, 10mM DTT, 3000 units/mL RNasin Ribonuclease
427 Inhibitors (Promega, N2615) per well using a FACS Aria III (Becton Dickinson). The sorted cells
428 were frozen on dry ice, and then stored at -80°C or immediately used for subsequent RNA reverse
429 transcription. Although cells were not stained for IgG expression, they are memory B cells based
430 on the fact that they are CD20⁺ (a marker absent in plasmablasts) and they express IgG (since
431 antibodies were amplified from these cells using IgG-specific primers).

432

433 **Antibody sequencing, cloning and expression.** Antibodies were identified and sequenced as
434 described previously^{22,28,29}. Briefly, RNA from single cells was reverse-transcribed (SuperScript
435 III Reverse Transcriptase, Invitrogen, 18080-044) and the cDNA stored at -20°C or used for
436 subsequent amplification of the variable IGH, IGL and IGK genes by nested PCR and Sanger
437 sequencing²⁸. Amplicons from the first PCR reaction were used as templates for Sequence- and
438 Ligation-Independent Cloning (SLIC) into antibody expression vectors. Recombinant monoclonal
439 antibodies and Fabs were produced and purified as previously described^{30,31}.

440

441 **Biolayer interferometry.**

442 BLI assays were performed on the Octet Red instrument (ForteBio) at 30°C with shaking at 1,000
443 r.p.m. Epitope binding assays were performed with protein A biosensor (ForteBio 18-5010),
444 following the manufacture protocol “classical sandwich assay”. (1) Sensor check: sensors
445 immersed 30 sec in buffer alone (buffer ForteBio 18-1105). (2) Capture 1st Ab: sensors immersed
446 10min with Ab1 at 40 µg/mL. (3) Baseline: sensors immersed 30sec in buffer alone. (4) Blocking:
447 sensors immersed 5 min with IgG isotype control at 50 µg/mL. (6) Antigen association: sensors
448 immersed 5 min with RBD at 100 µg/mL. (7) Baseline: sensors immersed 30 sec in buffer alone.
449 (8) Association Ab2: sensors immersed 5min with Ab2 at 40 µg/mL. Curve fitting was performed
450 using the Data analysis software (ForteBio).

451

452 **Computational analyses of antibody sequences.** Antibody sequences were trimmed based on
453 quality and annotated using Igblastn v1.14.0³² with IMGT domain delineation system. Annotation
454 was performed systematically using Change-O toolkit v.0.4.5³³. Heavy and light chains derived

455 from the same cell were paired, and clonotypes were assigned based on their V and J genes using
456 in-house R and Perl scripts (Fig. 3 b,c). All scripts and the data used to process antibody sequences
457 are publicly available on GitHub (<https://github.com/stratust/igpipeline>).

458 The frequency distributions of human V genes in anti-SARS-CoV-2 antibodies from this study
459 was compared to Sequence Read Archive SRP010970³⁴. The V(D)J assignments were done using
460 IMGT/High V-Quest and the frequencies of heavy and light chain V genes were calculated for 14
461 and 13 individuals, respectively, using sequences with unique CDR3s. The two-tailed t test with
462 unequal variances was used to determine statistical significance (Extended Data Fig. 7).

463 Nucleotide somatic hypermutation and CDR3 length were determined using in-house R and Perl
464 scripts. For somatic hypermutations, IGHV and IGLV nucleotide sequences were aligned against
465 their closest germlines using IgBlast and the number of differences were considered nucleotide
466 mutations. The average mutations for V genes was calculated by dividing the sum of all nucleotide
467 mutations across all patients by the number of sequences used for the analysis. To calculate the
468 GRAVY scores of hydrophobicity³⁵ we used Guy H.R. Hydrophobicity scale based on free energy
469 of transfer (kcal/mole)³⁶ implemented by the R package Peptides available in the Comprehensive
470 R Archive Network repository (<https://journal.r-project.org/archive/2015/RJ-2015-001/RJ-2015-001.pdf>). We used 534 heavy chain CDR3 amino acid sequences from this study and 22,654,256
471 IGH CDR3 sequences from the public database of memory B-cell receptor sequences³⁷. The
472 Shapiro-Wilk test was used to determine whether the GRAVY scores are normally distributed.
473 The GRAVY scores from all 533 IGH CDR3 amino acid sequences from this study
474 (sequence COV047_P4_IgG_51-P1369 lacks CDR3 amino acid sequence) were used to perform
475 the test and 5000 GRAVY scores of the sequences from the public database were randomly
476 selected. The Shapiro-Wilk p-values were 6.896×10^{-3} and 2.217×10^{-6} for sequences from this

478 study and the public database, respectively, indicating the data are not normally distributed.
479 Therefore, we used the Wilcoxon non-parametric test to compare the samples, which indicated a
480 difference in hydrophobicity distribution ($p = 5 \times 10^{-6}$; Extended Data Fig. 8).

481

482 **Negative-stain EM Data Collection and Processing**

483 Purified Fabs (C002, C119, and C121) were complexed with SARS-CoV-2 S trimer at a 2-fold
484 molar excess for 1 min and diluted to 40 $\mu\text{g}/\text{mL}$ in TBS immediately before adding 3 μL to a
485 freshly-glow discharged ultrathin, 400 mesh carbon-coated copper grid (Ted Pella, Inc.). Samples
486 were blotted after a 1 min incubation period and stained with 1% uranyl formate for an additional
487 minute before imaging. Micrographs were recorded on a Thermo Fisher Talos Arctica transmission
488 electron microscope operating at 200 keV using a K3 direct electron detector (Gatan, Inc) and
489 SerialEM automated image acquisition software³⁸. Images were acquired at a nominal
490 magnification of 28,000x (1.44 $\text{\AA}/\text{pixel}$ size) and a -1.5 to -2.0 μm defocus range. Images were
491 processed in cryoSPARC v2.14, and reference-free particle picking was completed using a
492 gaussian blob picker³⁹. Reference-free 2D class averages and *ab initio* volumes were generated in
493 cryoSPARC, and subsequently 3D-classified to identify classes of S-Fab complexes, that were
494 then homogenously refined. Figures were prepared using UCSF Chimera⁴⁰.

495

496 **Acknowledgements:** We thank all study participants who devoted time to our research; Drs. Barry
497 Coller and Sarah Schlesinger, the Rockefeller University Hospital Clinical Research Support
498 Office and nursing staff. Dr. Joseph L. DeRisi for facilitating interactions with the Chan
499 Zuckerberg BioHub. All members of the M.C.N. laboratory for helpful discussions, Drs. Amelia
500 Escolano, Gaëlle Breton and Bernardo Reis, and Maša Jankovic for laboratory support. This work
501 was supported by NIH grant P01-AI138398-S1 (M.C.N., C.M.R., P.J.B.) and 2U19AI111825
502 (M.C.N. and C.M.R.); the Caltech Merkin Institute for Translational Research and P50 AI150464
503 (P.J.B.), George Mason University Fast Grant and the European ATAC consortium (EC
504 101003650) to D.F.R.; 3 R01-AI091707-10S1 to C.M.R.; R37-AI64003 to P.D.B.; R01AI78788
505 to T.H.; The G. Harold and Leila Y. Mathers Charitable Foundation to C.M.R.. Electron
506 microscopy was performed in the Caltech Beckman Institute Resource Center for Transmission
507 Electron Microscopy (Drs. Songye Chen and Andrey Malyutin, Directors). C.G. was supported by
508 the Robert S. Wennett Post-Doctoral Fellowship, in part by the National Center for Advancing
509 Translational Sciences (National Institutes of Health Clinical and Translational Science Award
510 program, grant UL1 TR001866), and by the Shapiro-Silverberg Fund for the Advancement of
511 Translational Research. P.D.B. and M.C.N. are Howard Hughes Medical Institute Investigators.

512
513 **Contributions:** D.F.R., P.D.B., P.J.B., T.H., C.M.R. and M.C.N. conceived, designed and
514 analyzed the experiments. D.F.R., M.C. and C.G. designed clinical protocols. F.M., J.C.C.L.,
515 Z.W., A.C., M.A., C.O.B., S.F., T.H., C.V., K.G., F.B., S.T.C., P.M., H.H., L.N., F.S., Y.W., H.-
516 H.H., E.M., A.W.A., K.E.H.T., N.K. and P.R.H. carried out all experiments. A.G. and M.C.
517 produced antibodies. C.O.B., J.P. and E.W. produced SARS-CoV-2 proteins. A.H., R.K., J.H.,
518 K.G.M., C.G. and M.C. recruited participants and executed clinical protocols. R.P., J.D., M.P. and

519 I.S. processed clinical samples. T.Y.O., A.P.W. and V.R. performed bioinformatic analysis.

520 D.F.R., P.D.B., P.J.B., T.H., C.M.R. and M.C.N. wrote the manuscript with input from all co-

521 authors.

522

523 **Declaration of conflict:** In connection with this work The Rockefeller University has filed a

524 provisional patent application on which D.F.R. and M.C.N. are inventors.

525 **Extended Data Tables**

Extended Data Table 1. Individual participant demographics and clinical characteristics

ID	Age	Gender	Race	Ethnicity	Case/Contact	Duration (days)		Sx Severity (0-10)	ELISA binding (AUC)				Neutralization (NT50)
						Sx total	Sx onset to visit		RBD		S		
								IgG	IgM	IgG	IgM		
5	43	M	White	Non-Hispanic	Case	9	41	9	2.52	2.35	5.51	2.43	5.0
7	40	M	White	Non-Hispanic	Case	11	30	6	2.92	2.54	7.39	2.37	2730.4
8	37	M	White	Non-Hispanic	Case	3	57	5	2.11	0.81	4.46	0.88	5.0
9	35	F	White	Non-Hispanic	Case	12	54	5	2.90	1.07	4.44	2.80	5.0
12	27	F	White	Non-Hispanic	Contact	7	24	3	1.47	1.71	4.31	1.23	5.0
13	28	M	White	Non-Hispanic	Case	5	25	3	1.97	4.10	3.95	1.38	173.2
18	55	M	White	Non-Hispanic	Case	16	28	6	2.57	1.75	3.92	1.31	410.3
20	26	F	White	Non-Hispanic	Case	2	17	5	1.80	2.00	4.49	1.68	5.0
21	54	M	White	Hispanic	Contact	11	27	7	5.60	2.88	7.60	2.47	5052.7
24	34	M	White	Non-Hispanic	Case	15	30	4	2.29	1.97	4.36	1.45	280.7
26	66	M	White	Non-Hispanic	Case	2	35	3	1.67	1.86	4.00	1.58	276.1
27	28	M	White	Non-Hispanic	Case	9	32	4	1.79	1.64	4.41	1.30	739.3
28	26	M	White	Non-Hispanic	Case	7	21	4	2.39	2.25	4.93	1.66	888.9
29	26	F	White	Non-Hispanic	Contact	5	35	4	1.69	1.54	3.93	1.97	5.0
30	30	M	White	Non-Hispanic	Contact	3	35	4	3.07	2.55	4.86	1.34	5.0
31	51	M	White	Non-Hispanic	Case	9	33	3	1.29	1.60	4.69	1.20	192.3
32	46	F	White	Non-Hispanic	Case	8	32	3	1.30	2.18	3.79	0.83	47.4
37	27	M	White	Non-Hispanic	Case	6	28	6	2.13	1.52	4.11	1.04	286.3
38	57	F	White	Non-Hispanic	Case	10	38	4	1.87	2.02	4.30	1.12	518.9
40	44	M	White	Non-Hispanic	Case	7	23	5	1.72	1.50	3.81	1.47	42.1
41	35	M	White	Non-Hispanic	Contact	10	29	8	2.13	1.67	4.15	0.98	302.5
42	40	M	White	Non-Hispanic	Contact	20	36	8	2.37	1.50	5.82	1.43	627.1
45	55	F	White	Non-Hispanic	Case	7	48	6	1.95	0.97	3.53	0.75	5.0
46	39	M	White	Non-Hispanic	Case	8	30	2	2.68	1.45	4.14	1.29	59.2
47	43	F	White	Non-Hispanic	Case	11	33	5	3.02	2.30	6.23	2.66	10433.3
48	37	F	White	Non-Hispanic	Case	7	21	5	1.51	1.85	3.47	1.51	173.4
50	27	F	White	Non-Hispanic	Contact	7	28	4	1.64	2.32	5.42	1.45	924.7
51	21	M	White	Non-Hispanic	Contact	8	31	5	1.76	1.69	3.94	1.03	1499.2
54	40	F	White	Non-Hispanic	Contact	3	24	3	1.80	2.10	4.99	1.42	5.0
55	36	M	White	Non-Hispanic	Case	3	49	2	1.43	1.23	3.90	2.09	5.0
56	75	F	White	Non-Hispanic	Case	22	40	3	1.79	1.81	5.89	1.47	1388.4
57	66	M	White	Non-Hispanic	Case	6	21	5	1.54	2.10	4.33	1.00	2048.9
58	64	F	White	Non-Hispanic	Contact	1	32	2	1.20	1.71	3.95	1.21	5.0
64	28	F	White	Non-Hispanic	Contact	11	32	6	2.36	2.06	4.48	1.66	776.7
67	19	F	N/A	Hispanic	Case	5	29	6	2.56	1.98	5.48	1.32	2052.9
71	45	F	White	Non-Hispanic	Case	12	48	7	1.55	2.04	3.86	1.53	33.3
72	42	M	White	Non-Hispanic	Case	16	35	8	4.05	3.58	6.05	2.59	3138.2
75	46	F	White	Non-Hispanic	Case	10	36	4	1.64	2.37	3.98	1.20	271.5
76	49	F	White	Non-Hispanic	Case	28	34	4	1.88	1.65	5.17	0.97	219.8
77	37	M	White	Non-Hispanic	Contact	6	33	4	1.33	1.83	3.56	1.16	5.0
81	44	F	White	Non-Hispanic	Contact	3	35	2	1.58	1.73	3.82	1.10	5.0
82	46	M	N/A	Non-Hispanic	Case	0		0	2.19	1.92	5.41	1.81	130.7
88	41	M	White	Non-Hispanic	Case	7	23	4	1.82	3.32	4.97	1.37	424.7
95	44	M	White	Non-Hispanic	Case	9	36	6	2.61	1.62	6.03	1.62	961.9
96	48	F	White	Non-Hispanic	Case	9	30	3	3.93	1.93	6.25	2.26	927.7
97	39	M	White	Non-Hispanic	Case	9	31	3	1.58	1.61	4.03	2.33	202.7
98	35	F	White	Non-Hispanic	Case	2	24	4	1.78	1.47	5.38	1.22	249.0
99	36	F	White	Non-Hispanic	Case	13	29	5	2.50	3.27	4.38	2.49	1127.6
107	53	F	White	Non-Hispanic	Contact	10	29	4	1.74	1.41	4.66	0.90	297.5
108	75	M	White	Non-Hispanic	Case	16	41	7	1.37	0.88	3.47	1.36	557.5
110	27	M	White	Non-Hispanic	Case	1	25	1	1.30	1.60	4.00	0.93	5.0
114	30	F	White	Non-Hispanic	Case	15	36	7	1.65	1.92	3.53	1.55	110.9
115	65	F	White	Non-Hispanic	Contact	20	41	6	2.10	3.28	4.32	3.27	1127.7
119	56	M	White	Non-Hispanic	Case	13	48	3	1.36	1.26	4.41	1.59	650.3
120	56	F	White	Non-Hispanic	Case	26	48	6	0.99	0.97	3.65	1.21	100.6
121	19	M	White	Non-Hispanic	Contact	3	42	2	0.85	0.87	3.56	1.26	5.0
122	21	F	White	Non-Hispanic	Contact	3	38	1	1.44	0.94	3.39	1.22	5.0
123	26	M	White	Non-Hispanic	Contact	12	34	6	0.94	0.95	3.39	1.40	5.0
124	63	F	Asian	Non-Hispanic	Contact	4	37	3	1.58	2.06	3.49	1.32	5.0
125	51	F	White	Non-Hispanic	Case	10	26	3	1.92	3.49	3.86	1.24	126.5
127	24	F	White	Non-Hispanic	Case	10	43	6	1.80	2.50	4.37	2.41	883.5
130	39	M	White	Non-Hispanic	Contact	7	28	5	1.24	1.72	3.91	1.34	5.0
131	39	M	White	Non-Hispanic	Case	5	25	4	1.46	1.38	4.44	1.03	7.8
132	36	M	White	Non-Hispanic	Contact	10	50	6	2.15	1.76	4.84	1.97	5.0
134	27	F	White	Non-Hispanic	Contact	16	22	5	2.51	2.18	6.84	1.94	2700.6
135	62	F	White	Non-Hispanic	Case	8	31	6	2.20	2.02	3.80	1.13	350.0
140	63	F	White	Non-Hispanic	Case	28	47	1	1.05	1.24	3.58	1.28	52.4
149	41	M	White	Non-Hispanic	Contact	17	28	6	1.68	2.02	3.67	1.09	494.9
150	50	F	White	Non-Hispanic	Contact	12	45	7	1.15	0.43	3.18	1.82	5.0
154	68	M	Asian	Non-Hispanic	Case	16	30	9	3.19	2.19	4.85	1.29	928.2
157	50	M	White	Non-Hispanic	Case	10	32	8	2.40	2.86	3.90	2.06	741.7
166	28	F	White	Non-Hispanic	Case	13	45	2	1.27	0.66	3.45	0.94	5.0
167	50	F	White	Non-Hispanic	Contact	11	41	6	1.43	3.71	3.93	8.74	5.0
172	38	F	White	Non-Hispanic	Case	8	22	9	1.71	2.58	4.29	1.20	301.1
173	47	M	White	Non-Hispanic	Case	5	47	7	2.57	4.14	4.78	4.48	646.9
178	26	F	White	Non-Hispanic	Case	6	24	4	1.54	1.59	3.66	1.02	5.0
179	39	M	White	Non-Hispanic	Contact	10	37	3	1.89	2.25	3.83	1.73	370.1
182	44	F	White	Non-Hispanic	Contact	10	38	6	3.80	1.77	5.36	8.05	1503.7
183	43	F	White	Non-Hispanic	Case	13	44	8	1.56	1.02	3.88	1.55	240.1
185	54	M	White	Non-Hispanic	Case	11	44	8	3.51	1.39	5.55	2.11	1806.8
186	38	F	N/A	N/A	Case	8	26	2	1.73	2.47	4.37	1.23	296.9

190	54	F	White	Non-Hispanic	Case	18*	63	9	3.24	1.24	7.38	1.42	598.1
195	24	M	White	Non-Hispanic	Case	18	42	5	2.74	2.55	5.01	2.33	1315.1
200	60	F	White	Non-Hispanic	Case	17	39	7	2.40	1.00	4.38	1.51	1014.4
201	50	M	White	Non-Hispanic	Contact	15	33	6	4.37	2.57	6.15	1.57	3897.4
202	57	M	White	Non-Hispanic	Case	21	34	7	2.10	2.08	5.07	1.17	257.9
205	64	M	White	Non-Hispanic	Case	7	36	4	4.51	0.70	6.12	1.69	924.4
222	28	M	Asian	Non-Hispanic	Case	11	29	7	1.28	0.69	3.94	3.46	5.0
229	45	M	White	Non-Hispanic	Case	10	63	4	2.92	1.42	4.90	1.58	1272.9
230	50	M	White	Non-Hispanic	Case	18	33	7	3.80	0.47	3.48	0.88	5.0
232	38	F	White	Non-Hispanic	Case	13	43	7	1.57	0.70	4.24	5.70	94.3
233	55	M	White	Non-Hispanic	Case	20	41	3	2.07	2.11	4.51	1.07	173.2
241	36	M	White	Non-Hispanic	Case	12	30	7	2.27	2.66	4.54	1.46	923.1
242	59	M	White	Non-Hispanic	Case	10	42	6	4.91	1.94	4.81	2.16	1353.0
243	30	F	Asian	Non-Hispanic	Case	6	26	5	2.92	2.57	5.06	1.14	1300.2
246	44	F	White	Non-Hispanic	Case	10	38	7	2.05	2.79	6.09	1.32	566.0
255	33	M	White	Non-Hispanic	Case	14	44	6	2.14	0.70	4.20	1.24	172.5
256	63	F	White	Non-Hispanic	Case	27	42	6	1.72	1.96	4.26	7.79	141.6
258	52	M	White	Non-Hispanic	Contact	14	48	6	2.64	1.20	4.52	1.85	4145.9
279	41	M	White	Non-Hispanic	Case	7	38	8	1.68	2.13	3.77	1.90	308.9
280	59	M	White	Non-Hispanic	Case	6	32	7	2.53	3.07	4.61	1.19	1072.1
302	47	F	White	Non-Hispanic	Case	35*	49	7	1.48	0.97	4.06	1.26	5.0
310	34	F	White	Non-Hispanic	Case	17	35	5	3.95	1.24	9.44	3.07	485.5
314	46	M	White	Non-Hispanic	Case	11	38	7	2.12	0.88	4.56	1.51	667.1
315	29	F	White	Non-Hispanic	Case	15	42	8	3.02	0.69	3.98	1.03	376.5
319	50	M	White	Non-Hispanic	Case	5	38	6	3.71	2.28	3.79	1.05	5.0
323	39	F	White	Non-Hispanic	Case	7	45	7	1.05	1.03	3.53	1.42	5.0
325	52	M	White	Non-Hispanic	Case	16	38	8	2.25	1.47	4.83	2.28	1603.3
343	21	F	White	Non-Hispanic	Case	16	49	5	1.63	0.94	3.37	1.70	5.0
352	44	M	White	Non-Hispanic	Case	16	43	4	3.54	0.92	4.50	1.07	519.2
353	60	M	White	Non-Hispanic	Case	14	49	6	5.38	1.12	5.69	1.05	855.5
356	22	F	White	Non-Hispanic	Contact	16	38	3	1.37	0.61	2.98	1.09	5.0
357	27	F	White	Non-Hispanic	Contact	34	56	5	7.49	1.10	2.77	1.13	5.0
364	29	M	White	Non-Hispanic	Contact	14	49	6	0.97	0.58	2.90	0.89	5.0
366	41	F	White	Non-Hispanic	Contact	9	34	7	0.98	0.52	3.51	1.19	5.0
373	35	F	White	Non-Hispanic	Case	12	51	7	1.69	1.26	5.14	1.69	5.0
388	47	F	White	Non-Hispanic	Contact	14	41	9	1.57	1.06	3.61	1.69	5.0
393	69	M	White	Non-Hispanic	Case	23*	54	9	1.28	1.74	3.81	2.65	715.4
394	48	F	Multiple	Hispanic	Case	7	67	4	2.05	0.87	4.34	2.02	1281.5
397	52	M	White	Non-Hispanic	Case	22	45	8	3.32	0.59	5.01	0.87	1516.9
403	52	M	Asian	Non-Hispanic	Case	18*	39	10	5.36	1.09	10.01	1.36	3887.8
406	65	M	White	Non-Hispanic	Case	20	56	8	4.69	0.90	7.51	1.15	1288.7
410	34	M	White	Non-Hispanic	Case	12	46	8	1.06	0.56	3.95	0.76	5.0
421	62	F	White	Non-Hispanic	Contact	12	43	9	0.95	1.07	3.34	1.35	5.0
426	65	M	White	Non-Hispanic	Case	18	51	6	2.07	0.55	3.95	1.49	804.8
437	43	F	Asian	Non-Hispanic	Case	14	34	7	2.54	0.47	4.30	1.44	698.8
460	36	M	White	Non-Hispanic	Case	11	39	6	2.94	3.18	5.51	2.80	1906.7
461	49	M	White	Non-Hispanic	Case	7	39	5	3.38	0.94	4.67	2.02	1076.6
462	28	F	White	Non-Hispanic	Case	16	45	5	1.36	0.38	3.07	1.11	5.0
470	28	F	White	Non-Hispanic	Case	17	51	4	1.26	0.86	3.97	1.50	5.0
478	31	M	White	Non-Hispanic	Case	16	52	4	1.43	0.93	3.70	1.97	263.2
481	28	F	Asian	Non-Hispanic	Case	15	43	8	1.70	0.39	3.46	1.24	5.0
486	64	F	White	Non-Hispanic	Case	11	41	10	1.70	1.00	3.68	1.29	5.0
500	46	M	White	Non-Hispanic	Case	12	53	5	1.10	0.82	3.49	1.34	5.0
501	32	M	Asian	Non-Hispanic	Case	18*	53	10	2.62	0.65	4.51	1.21	718.8
502	52	M	White	Non-Hispanic	Case	16*	53	9	5.10	0.61	5.10	1.62	2171.8
506	46	M	White	Non-Hispanic	Case	12	59	9	0.84	0.81	3.13	1.21	5.0
507	39	M	White	Non-Hispanic	Case	15	60	8	1.92	0.96	4.56	1.52	5.0
509	36	M	White	Non-Hispanic	Case	11	50	5	1.99	1.01	3.99	1.45	5.0
526	49	M	Asian	Non-Hispanic	Case	11	34	7	3.36	1.45	5.88	1.57	4193.3
537	52	M	White	Non-Hispanic	Case	15	45	6	1.47	0.95	3.65	1.58	923.3
539	73	F	White	Non-Hispanic	Case	19*	54	10	2.82	0.63	4.46	1.45	487.9
547	59	M	White	Non-Hispanic	Case	15*	36	9	2.97	1.53	5.08	2.59	2900.6
587	54	M	PI	N/A	Case	17*	51	8	3.22	0.60	4.01	1.49	473.1
632	38	M	White	Non-Hispanic	Contact	10	43	6	2.49	0.86	4.50	1.63	572.3
633	39	M	White	Non-Hispanic	Contact	8	57	4	1.25	1.04	3.38	1.73	5.0
652	76	M	White	Non-Hispanic	Case	18*	56	10	4.75	1.46	8.96	3.80	2324.0
664	45	F	White	Non-Hispanic	Case	17*	42	10	1.68	0.43	3.93	1.32	5.0
675	47	M	White	Non-Hispanic	Contact	31	47	5	0.79	0.93	2.94	1.38	5.0

*hospitalized, Sx=symptoms

Extended Data Table 2. Frequency of symptoms and comorbidities reported by participants

Symptom	Participants (n=149)	%
Fever	125	83.9
Fatigue	106	71.1
Cough	93	62.4
Myalgia	92	61.7
Shortness of breath	66	44.3
Headache	63	42.3
Loss of smell/taste	50	33.3
Sore throat	38	25.3
Diarrhea	32	21.3
Presence of comorbidities (HTN, CAD, DM, COPD, asthma, cancer)	16	10.7

528

HTN (hypertension), CAD (coronary artery disease), DM (diabetes mellitus), COPD (chronic obstructive pulmonary disease)

529 Extended Data Tables 3 and 4 are provided as separate Excel files.

Extended data table 5. Sequences of cloned recombinant antibodies

Antibody ID	IGH VDJ (nt)	IGH VDJ (aa)	IGL VJ (nt)	IGL VJ (aa)
C002	GAGGTGCAGC ^c EVQLVESGGGV		GACATCCAG1DIQLTQSPSSL ^ε	
C003	GAGGTGCAGC ^c EVQLVESGGGLI		GAAATTGTGT EIVLTQSPGTL ^ε	
C004	CAGGTGCAGC ^c QVQLVQSGAEV		GCCATCCGG AIRMTQSPSSL ^ε	
C005	CAGGTGCAGC ^c QVQLVQSGPEV		GAAATTGTGT EIVLTQSPGTL ^ε	
C006	CAGGTGCAGC ^c QVQLVESGGGL		CAGTCTGTGC QSVLTQPPSAS	
C008	GAGGTGCAGC ^c EVQLVESGGGV		GACATCCAG/DIQMTQSPSTL ^ε	
C009	CAGGTGCAGC ^c QVQLVQSGAEV		CAGTCTGCC(QSALTQPPSAS	
C010	GAGGTGCAGC ^c EVQLVESGGGV		GACATCCAG1DIQLTQSPSSL ^ε	
C013	CAGGTGCAGC ^c QVQLVQSGAEV		GAAATTGTGT EIVLTQSPATLS	
C016	GAGGTGCAGC ^c EVQLVESGGGV		GCCATCCGG AIRMTQSPSSL ^ε	
C017	GAAGTGCAGC1EVQLVESGGGL ^l		GAAATTGTGT EIVLTQSPATLS	
C018	GAGGTGCAGC ^c EVQLVESGGGV		GACATCCAG1DIQLTQSPSSL ^ε	
C019	CAGGTGCAGC ^c QVQLVQSGAEV		TCCTATGAGC SYELTQPPSVS	
C021	CAGGTGCAGC ^c QVQLQESGPGL ^l		GATATTGTGA DIVMTQSPSLI	
C022	CAGGTGCAGT1QVQLQESGPGL ^l		GACATCCAG/DIQMTQSPSTL ^ε	
C027	GAGGTGCAGC ^c EVQLVESGGGV		GACATCCAG/DIQMTQSPSTL ^ε	
C029	CAGGTGCAGC ^c QVQLQESGPGL ^l		GATATTGTGA DIVMTQSPSLI	
C030	GAGGTGCAGC ^c EVQLVESGGGV		GACATCCAG/DIQMTQSPSTL ^ε	
C031	GAGGTGCAGC ^c EVQLVESGGGL ^l		GACATCCAG/DIQMTQSPSSL ^ε	
C101	CAGGTGCAGC ^c QVQLVESGGGL		GAAATTGTGT EIVLTQSPGTL ^ε	
C102	CAGGTGCAGC ^c QVQLVESGGGL		GAAATTGTGT EIVLTQSPGTL ^ε	
C103	CAGGTGCAGC ^c QVQLQWVWAGL		GAAATTGTGT EIVLTQSPGTL ^ε	
C104	CAGGTGCAGC ^c QVQLQWVWAGL		GAAATTGTGT EIVLTQSPGTV ^ε	
C105	CAGGTGCAGC ^c QVQLVESGGGL		CAGTCTGCC(QSALTQPPSAS	
C106	CAGTGCAGC ^c QLQLQESGPGL ^l		TCCTATGAGC SYELTQPPSVS	
C107	CAGGTGCAGC ^c QVQLVQSGAEV		CAGTCTGTGC QSVLTQPPSAS	
C108	CAGGTGCAGC ^c QVQLQESGPGL ^l		CAGTCTGCC(QSALTQPPASVS	
C109	GAAGTGCAGC1EVQLVESGGGL ^l		CAGTCTGCC(QSALTQPPSAS	
C110	CAGGTACAGC1QVQLQSGAEV		GACATCCAG/DIQMTQSPSTL ^ε	
C111	CAGTGCAGC ^c QLQLQESGPGL ^l		CAGTCTGTGC QSVLTQPPSVS	
C112	CAGGTGCAGC ^c QVQLVESGGGV		CAGTCTGCC(QSALTQPPASVS	
C113	CAGGTGCAGC ^c QVQLVESGGGV		GACATCCAG/DIQMTQSPSTL ^ε	
C114	CAGGTGCAGC ^c QVQLVESGGGL		CAGTCTGTGC QSVLTQPPSVS	
C115	CAGGTGCAGC ^c QVQLVESGGGL		GATATTGTGA DIVMTQSPSL ^l	
C116	CAGGTGCAGC ^c QVQLVESGGGV		AATTTTATGC ^c NFMLTQPHSV ^ε	
C117	CAGGTGCAGC ^c QVQLVESGGGV		CAGTCTGTGC QSVLTQPPSVS	
C118	CAGGTGCAGC ^c QVQLVESGGGV		CAGCCTGTG(QPVLQSPASVS	
C119	CAGGTCCAGC ^c QVQLVQSGAEV		CAGTCTGCC(QSALTQPPASVS	
C120	GAGGTGCAGC ^c EVQLVESGGGLI		GCCATCCGG AIRMTQSPSSL ^ε	
C121	CAGGTGCAGC ^c QVQLVQSGAEV		CAGTCTGCC(QSALTQPPASVS	
C122	GAGGTGCAGC ^c EVQLVESGGGLI		GACATCCAG1DIQLTQSPSFL ^ε	
C123	GAGGTGCAGC ^c EVQLVESGGGLI		GACATCCAG1DIQLTQSPSFL ^ε	
C124	GAGGTGCAGC ^c EVQLVESGGGL ^l		GAAATTGTGT EIVLTQSPATLS	
C125	CAGGTGCAGC ^c QVQLVQSGPEV		GAAATTGTGT EIVLTQSPGTL ^ε	
C126	CAGGTGCAGC ^c QVQLQESGPGL ^l		AATTTTATGC ^c NFMLTQPHSV ^ε	
C127	CAGGTGCAGC ^c QVQLVQSGAEV		CAGTCTGTGC QSVLTQPPSAS	
C128	GAGGTGCAGC ^c EVQLVESGGGL ^l		GAAATTGTGT EIVLTQSPGTL ^ε	
C130	CAGGTGCAGC ^c QVQLVQSGAEV		TCCTATGAGC SYELTQPPSVS	
C131	CAGGTGCAGC ^c QVQLVQSGSEV		GAAATAGTGA EIVMTQSPATL ^ε	
C132	CAGGTGCAGC ^c QVQLQESGPGL ^l		CAGTCTGCC(QSALTQPPASVS	
C133	CAGGTGCAGC ^c QVQLVESGGGV		GACATCCAG/DIQMTQSPSSL ^ε	
C134	GAGGTGCAGC ^c EVQLVESGGGL ^l		TCCTATGAGC SYELTQPPSVS	
C135	CAGGTGCAGC ^c QVQLVESGGGV		GACATCCAG/DIQMTQSPSTL ^ε	
C138	GAGGTGCAGC ^c EVQLVESGGGL ^l		AATTTTATGC ^c NFMLTQPHSV ^ε	
C139	GAGGTGCAGC ^c EVQLVESGGGV		GACATCCAG/DIQMTQSPSSL ^ε	
C140	GAGGTGCAGC ^c EVQLVESGGGL ^l		GACATCCAG1DIQLTQSPSFL ^ε	
C141	GAGGTGCAGC ^c EVQLVESGGGV		AATTTTATGC ^c NFMLTQPHSV ^ε	
C143	GAGGTGCAGC ^c EVQLVESGGGL ^l		CAGTCTGCC(QSALTQPPASVS	
C144	GAGGTGCAGC ^c EVQLVESGGGLI		CAGTCTGCC(QSALTQPPASVS	
C145	GAGGTGCAGC ^c EVQLVESGGGLI		CAGTCTGCC(QSALTQPPASVS	
C146	GAGGTGCAGC ^c EVQLVESGGGL ^l		CAGTCTGCC(QSALTQPPASVS	
C147	GAGGTGCAGC ^c EVQLVQSGAEV		CAGGCCTGTG(QAVVTQEPSTL ^ε	
C148	GAGGTGCAGC ^c EVQLVESGGGL ^l		GAAATAGTGA EIVMTQSPATL ^ε	
C150	GAGGTGCAGC ^c EVQLVESGGGL ^l		CAGTCTGCC(QSALTQPPASVS	
C151	GAGGTGCAGC ^c EVQLVESGGGL ^l		AATTTTATGC ^c NFMLTQPHSV ^ε	
C153	GAGGTGCAGC ^c EVQLVESGGGLI		CAGTCTGCC(QSALTQPPASVS	
C154	CAGGTGCAGC ^c QVQLVESGGGV		GACATCCAG/DIQMTQSPSSL ^ε	
C155	GAGGTGCAGC ^c EVQLVESGGGLI		GAAATAGTGA EIVMTQSPATL ^ε	
C156	CAGGTGCAGC ^c QVQLVESGGGV		TCCTATGAGC SYELTQPPSVS	
C164	GAGGTGCAGC ^c EVQLVESGGGL ^l		CAGTCTGCC(QSALTQPPASVS	
C165	CAGGTGCAGC ^c QVQLVQSGAEV		GAAATTGTGT EIVLTQSPGTL ^ε	
C201	GAAGTGCAGC1EVQLVESGGGL ^l		CATCCGGAT(IRMTQSPSSVS	
C202	GAGGTGCAGC ^c EVQLVESGGGL ^l		GACATCCAG1DIQLTQSPSSL ^ε	
C204	GAGGTGCAGC ^c EVQLVESGGGL ^l		GACATCCAG1DIQLTQSPSSL ^ε	
C205	CAGGTGCAGC ^c QVQLVQSGAEV		GAAATTGTGT EIVLTQSPGTL ^ε	
C207	GAGGTGCAGC ^c EVQLVESGGGL ^l		GAAATTGTGT EIVLTQSPATLS	
C208	GAGGTGCAGC ^c EVQLVQSGAEV		GAAATTGTGT EIVLTQSPGTL ^ε	
C209	CAGGTGCAGC ^c QVQLVQSGAEV		TCCTATGAGC SYELTQPPSVS	
C210	GAGGTGCAGC ^c EVQLVESGGGLI		GACATCCAG1DIQLTQSPSFL ^ε	
C211	GAGGTGCAGC ^c EVQLVESGGGL ^l		GAAATAGTGA EIVMTQSPATL ^ε	
C212	CAGGTGCAGC ^c QVQLVQSGAEV		CTGACTCAG(LTQPPASVSGSP	
C214	CAGGTGCAGC ^c QVQLVESGGGV		GACATCCAG1DIQLTQSPSSL ^ε	
C215	GAGGTGCAGC ^c EVQLVESGGGL ^l		GACATCCAG1DIQLTQSPSSL ^ε	
C216	GAGGTGCAGC ^c EVQLVESGGGL ^l		GACATCCAG1DIQLTQSPSSL ^ε	


530

531

Extended Data Table 6. Effective and inhibitory concentrations of the monoclonal antibodies

Participant ID	Antibody ID	SARS-CoV-2	SARS-CoV-2	SARS-CoV-2	SARS-CoV-2 RBD	SARS-CoV RBD
		IC50 ng/ml	IC80 ng/ml	IC90 ng/ml	EC50 ng/ml	EC50 ng/ml
COV21	C002	8.88	21.95	37.61	3.14	>1000
COV21	C003	313.79	992.62	>1000	6.37	>1000
COV21	C004	10.67	41.08	91.71	2.39	>1000
COV21	C005	60.49	130.65	205.20	4.41	>1000
COV21	C006	321.51	>1000	>1000	1.81	>1000
COV21	C008	625.46	>1000	>1000	4.63	>1000
COV21	C009	4.82	14.54	29.34	1.80	>1000
COV21	C010	>1000	>1000	>1000	5.44	>1000
COV21	C013	42.48	360.59	>1000	2.59	>1000
COV21	C016	>1000	>1000	>1000	7.41	>1000
COV21	C017	72.67	256.18	543.87	1.63	>1000
COV21	C018	>1000	>1000	>1000	1.53	>1000
COV21	C019	>1000	>1000	>1000	11.85	>1000
COV21	C021	>1000	>1000	>1000	1.25	>1000
COV21	C022	73.57	314.71	736.87	2.40	5.99
COV21	C027	>1000	>1000	>1000	2.65	696.05
COV21	C029	>1000	>1000	>1000	4.13	>1000
COV21	C030	>1000	>1000	>1000	2.89	>1000
COV21	C031	>1000	>1000	>1000	22.69	>1000
COV107	C101	8.20	30.15	65.30	1.51	>1000
COV107	C102	34.03	84.21	143.23	4.54	>1000
COV107	C103	4.38	12.58	23.59	3.77	>1000
COV107	C104	23.31	72.12	140.28	8.31	>1000
COV107	C105	26.09	72.24	133.70	5.20	>1000
COV107	C106	>1000	>1000	>1000	19.03	106.75
COV107	C107	>1000	>1000	>1000	11.55	>1000
COV107	C108	480.69	>1000	>1000	5.32	>1000
COV107	C110	18.44	45.11	77.28	7.29	>1000
COV107	C112	111.79	701.99	>1000	3.38	>1000
COV107	C113	>1000	>1000	>1000	6.93	>1000
COV107	C114	>1000	>1000	>1000	9.51	>1000
COV107	C115	198.33	958.18	>1000	3.40	>1000
COV107	C116	>1000	>1000	>1000	37.56	>1000
COV107	C117	348.00	>1000	>1000	5.38	>1000
COV107	C118	103.69	417.76	>1000	3.45	3.817
COV107	C119	9.12	39.45	97.78	3.57	>1000
COV107	C120	13.26	26.73	40.30	1.41	>1000
COV107	C121	6.73	14.31	22.33	2.85	>1000
COV107	C122	22.80	57.77	100.12	2.67	>1000
COV107	C123	149.22	355.47	595.51	1.92	>1000
COV107	C124	341.82	937.26	>1000	2.23	>1000
COV107	C125	43.32	92.54	144.26	1.87	>1000
COV107	C126	>1000	>1000	>1000	4.78	>1000
COV107	C127	68.74	190.96	347.31	2.62	>1000
COV072	C128	101.22	263.35	460.73	1.95	>1000
COV072	C130	>1000	>1000	>1000	2.05	>1000
COV072	C131	30.52	178.90	759.11	1.67	>1000
COV072	C132	708.67	>1000	>1000	2.92	>1000
COV072	C133	>1000	>1000	>1000	1.98	>1000
COV072	C134	>1000	>1000	>1000	1.57	>1000
COV072	C135	16.61	32.81	48.90	1.80	>1000
COV072	C136	>1000	>1000	>1000	33.61	>1000
COV072	C137	>1000	>1000	>1000	25.22	>1000
COV072	C138	>1000	>1000	>1000	2.94	>1000
COV072	C139	>1000	>1000	>1000	1.89	>1000
COV072	C140	23.88	66.24	120.69	2.19	>1000
COV072	C141	>1000	>1000	>1000	1.71	>1000
COV047	C143	>1000	>1000	>1000	3.66	>1000
COV047	C144	6.91	17.28	29.66	3.24	>1000
COV047	C145	3.04	14.51	36.79	3.86	>1000
COV047	C146	>1000	>1000	>1000	>1000	>1000
COV047	C147	>1000	>1000	>1000	>1000	>1000
COV047	C148	>1000	>1000	>1000	64.69	>1000
COV047	C150	>1000	>1000	>1000	8.11	>1000
COV047	C151	31.79	363.97	>1000	4.30	>1000
COV047	C153	70.71	490.08	>1000	3.17	>1000
COV047	C154	435.50	>1000	>1000	2.92	10.65
COV047	C155	11.00	35.75	77.01	3.30	>1000
COV047	C156	>1000	>1000	>1000	3.32	>1000
COV047	C164	239.15	865.40	>1000	2.06	>1000
COV072	C165	40.81	138.66	297.38	4.25	>1000
COV96	C201	>1000	>1000	>1000	2.98	>1000
COV96	C202	>1000	>1000	>1000	3.40	>1000
COV96	C204	>1000	>1000	>1000	3.73	9.41
COV96	C205	>1000	>1000	>1000	>1000	>1000
COV96	C207	158.52	960.39	>1000	1.87	>1000
COV96	C208	>1000	>1000	>1000	>1000	>1000
COV96	C209	>1000	>1000	>1000	3.79	>1000
COV96	C210	50.73	155.24	298.90	2.83	>1000
COV96	C211	12.79	34.87	62.89	2.82	>1000
COV96	C212	>1000	>1000	>1000	>1000	>1000
COV96	C214	>1000	>1000	>1000	5.75	>1000
COV96	C215	>1000	>1000	>1000	5.33	17.944
COV96	C216	>1000	>1000	>1000	9.53	>1000


533 **Extended Data Figures**



534

535 **Extended Data Figure 1. Clinical correlates.** **a**, Age distribution (Y axis) for all males and
536 females in the cohort $p=0.2074$. **b**, Duration of symptoms in days (Y axis) for all males and females
537 in the cohort $p=0.8704$. **c**, Time between symptom onset and plasma collection (Y axis) for all
538 males and females in the cohort $p=0.5514$. **d**, Subjective symptom severity on a scale of 0-10 (Y
539 axis) for all males and females in the cohort $p=0.1888$. **e**, Age distribution (Y axis) for all cases
540 and contacts in the cohort $p=0.0305$. **f**, Duration of symptoms in days (Y axis) for all cases and
541 contacts in the cohort $p=0.1241$. **g**, Time between symptom onset and plasma collection in days
542 (Y axis) for all cases and contacts in the cohort $p=0.1589$. **h**, Symptom severity (Y axis) for all
543 cases and contacts in the cohort $p=0.0550$. **i**, Age distribution (Y axis) for all outpatient and
544 hospitalized participants $p=0.0024$. **j**, Duration of symptoms in days (Y axis) for all outpatient and
545 hospitalized participants in the cohort $p<0.0001$. **k**, Time between symptom onset and plasma
546 collection in days (Y axis) for all outpatient and hospitalized participants in the cohort $p=0.0001$.

547 **I**, Symptom severity (Y axis) for all outpatient and hospitalized participants in the cohort
548 $p < 0.0001$. Horizontal bars indicate median values. Statistical significance was determined using
549 two-tailed Mann-Whitney U test.



550

551 **Extended Data Figure 2. Clinical correlates of plasma antibody titers. a**, AUC for IgG anti-

552 RBD (Y axis) for all cases and contacts in the cohort p=0.0107. **b**, AUC for IgM anti-RBD (Y

553 axis) for all cases and contacts in the cohort p=0.5371. **c**, AUC for IgG anti-S (Y axis) for all cases

554 and contacts in the cohort p=0.0135. **d**, AUC for IgM anti-S (Y axis) for all cases and contacts in

555 the cohort p=0.7838. **e**, AUC for IgM anti-RBD (Y axis) for all males and females in the cohort

556 p=0.9597. **f**, AUC for IgG anti-S (Y axis) for all males and females in the cohort p=0.0275. **g**,

557 AUC for IgM anti-S (Y axis) for all males and females in the cohort p=0.5363. **h**, AUC for IgM


558 anti-RBD (Y axis) for all outpatient and hospitalized participants in the cohort p=0.0059. **i**, AUC

559 for IgG anti-S (Y axis) for all outpatient and hospitalized participants in the cohort p=0.0623. **j**,

560 AUC for IgM anti-S (Y axis) for all outpatient and hospitalized participants in the cohort p=0.2976.

561 Horizontal bars indicate median values. Statistical significance was determined using two-tailed

562 Mann-Whitney U test.



563

564 **Extended Data Figure 3. Additional clinical correlates of plasma antibody titers. a,** Time

565 between symptom onset and plasma collection in days (X axis) plotted against AUC for IgG anti-

566 RBD (Y axis) $r=-0.0261$ $p=0.7533$. **b,** Time between symptom onset and plasma collection in days

567 (X axis) plotted against AUC for IgG anti-S (Y axis) $r=-0.1495$ $p=0.0697$. **c,** Time between

568 symptom onset and plasma collection in days (X axis) plotted against AUC for IgM anti-S (Y axis)

569 $r=0.1496$ $p=0.0695$. **d,** Age (X axis) plotted against AUC for IgM anti-RBD (Y axis) $r=0.0172$


570 $p=0.8355$. **e,** Age (X axis) plotted against AUC for IgG anti-S (Y axis) $r=0.1523$ $p=0.0638$. **f,** Age

571 (X axis) plotted against AUC for IgM anti-S (Y axis) $r=0.0565$ $p=0.4934$. **g,** Duration of symptoms

572 in days (X axis) plotted against AUC for IgG anti-RBD (Y axis) $r=0.1525$, $p=0.0633$. **h,** Duration

573 of symptoms in days (X axis) plotted against AUC for IgM anti-RBD (Y axis) $r=-0.3187$,

574 $p < 0.0001$. **i**, Duration of symptoms in days (X axis) plotted against AUC for IgG anti-S (Y axis)
575 $r = 0.0329$, $p = 0.6904$. **j**, Duration of symptoms in days (X axis) plotted against AUC for IgM anti-
576 S (Y axis) $r = 0.0824$, $p = 0.3177$. **k**, Severity of symptoms (X axis) plotted against AUC for IgG
577 anti-RBD (Y axis) $r = 0.2679$ $p = 0.0010$. **l**, Severity of symptoms (X axis) plotted against AUC for
578 IgM anti-RBD (Y axis) $r = -0.1943$ $p = 0.0176$. **m**, Severity of symptoms (X axis) plotted against
579 AUC for IgG anti-S (Y axis) $r = 0.1187$ $p = 0.1492$. **n**, Severity of symptoms (X axis) plotted against
580 AUC for IgM anti-S (Y axis) $r = 0.1597$ $p = 0.0517$. All correlations were analyzed by two-tailed
581 Spearman's.



582

583 **Extended Data Figure 4. Diagrammatic representation of the SARS-CoV2 pseudovirus**

584 **luciferase assay. a**, Co-transfection of pNL4-3ΔEnv-nanoluc and pSARS-CoV-2 spike vectors

585 into 293T cells leads to production of SARS-CoV-2 Spike-pseudotyped HIV-1 particles (SARS-


586 CoV-2 pseudovirus) carrying the *Nanoluc* gene. **b**, SARS-CoV-2 pseudovirus is incubated for 1 h

587 at 37°C with plasma or monoclonal antibody dilutions. The virus-antibody mixture is used to infect

588 ACE2-expressing 293T cells, which will express nanoluc Luciferase upon infection. **c**, Relative

589 luminescence units (RLU) reads from lysates of ACE2-expressing 293T cells infected with


590 increasing amounts of SARS-CoV-2 pseudovirus.



591
592 **Extended Data Figure 5. Clinical correlates of neutralization.** **a**, Anti-RBD IgM AUC (X axis)
593 plotted against NT₅₀ (Y axis) $r=0.3119$, $p=0.0001$. **b**, Anti-S IgM AUC (X axis) plotted against
594 NT₅₀ (Y axis) $r=0.3211$, $p=<0.0001$. **c**, Duration of symptoms in days (X axis) plotted against NT₅₀
595 (Y axis) $r=0.1997$, $p=0.0146$. **d**, Time between symptom onset and plasma collection in days (X
596 axis) plotted against NT₅₀ (Y axis) $r=-0.1344$, $p=0.1033$. **e**, Symptom severity (X axis) plotted


597 against NT_{50} (Y axis) $r=0.2234$, $p=0.0062$. **f**, Age (X axis) plotted against NT_{50} (Y axis) $r=0.3005$,
598 $p=0.0002$. All correlations were analyzed by two-tailed Spearman's. Dotted line ($NT_{50}=5$)
599 represents lower limit of detection (LLOD) of pseudovirus neutralization assay. Samples with
600 undetectable neutralizing titers were plotted at LLOD.

601



602

603 **Extended Data Figure 6. Flow cytometry.** Gating strategy used for cell sorting. Gating was on
604 singlets that were CD20⁺ and CD3⁻CD8⁻CD16⁻Ova⁻. Sorted cells were RBD-PE⁺ and RBD-
605 AF647⁺.




606

607 **Extended Data Figure 7. Frequency distributions of human V genes.** The two-tailed t test with

608 unequal variance was used to compare the frequency distributions of human V genes of anti-

609 SARS-CoV-2 antibodies from this study to Sequence Read Archive SRP010970³⁴.



610

611 **Extended Data Figure 8. Analysis of antibody somatic hypermutation and CDR3 length. a,**


612 For each individual, the number of somatic nucleotide mutations (Y axis) at the IGVH and IGVL

613 are shown on the left panel, and the amino acid length of the CDR3s (Y axis) are shown on the

614 right panel. The horizontal bar indicated the mean. **b**, same as in a but for all antibodies combined.

615 **c**, Distribution of the hydrophobicity GRAVY scores at the IGH CDR3 in antibody sequences

616 from this study compared to a public database (see Methods).




617

618 **Extended Data Figure 9. Binding of the monoclonal antibodies to the RBD of SARS-CoV-2**

619 **and SARS-CoV. a, EC₅₀ values for binding to the RBD of SARS-CoV-2. b and c, Binding curves**

620 **and EC₅₀ values for binding to the RBD of SARS-CoV.**

621



622

623 **Extended Data Figure 10. Biolayer interferometry experiment.** showing binding of antibodies

624 C144, C101, C002, C121, C009, C019 (see also main text Fig. 4). Graphs show secondary

625 antibody binding to preformed C121 IgG-RBD complexes. The table displays the shift in

626 nanometers after second antibody (Ab2) binding to the antigen in the presence of the first antibody

627 (Ab1). Values are normalized by the subtraction of the autologous antibody control.

628

629 **References**

- 630 1 Graham, R. L., Donaldson, E. F. & Baric, R. S. A decade after SARS: strategies for
631 controlling emerging coronaviruses. *Nat Rev Microbiol* **11**, 836-848,
632 doi:10.1038/nrmicro3143 (2013).
- 633 2 Gralinski, L. E. & Baric, R. S. Molecular pathology of emerging coronavirus infections. *J*
634 *Pathol* **235**, 185-195, doi:10.1002/path.4454 (2015).
- 635 3 Hoffmann, M. *et al.* SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is
636 Blocked by a Clinically Proven Protease Inhibitor. *Cell* **181**, 271-280 e278,
637 doi:10.1016/j.cell.2020.02.052 (2020).
- 638 4 Walls, A. C. *et al.* Structure, Function, and Antigenicity of the SARS-CoV-2 Spike
639 Glycoprotein. *Cell* **181**, 281-292 e286, doi:10.1016/j.cell.2020.02.058 (2020).
- 640 5 Jiang, S., Hillyer, C. & Du, L. Neutralizing Antibodies against SARS-CoV-2 and Other
641 Human Coronaviruses. *Trends Immunol*, doi:10.1016/j.it.2020.03.007 (2020).
- 642 6 Scheid, J. F. *et al.* Broad diversity of neutralizing antibodies isolated from memory B
643 cells in HIV-infected individuals. *Nature* **458**, 636-640, doi:10.1038/nature07930 (2009).
- 644 7 Tiller, T. *et al.* Autoreactivity in human IgG+ memory B cells. *Immunity* **26**, 205-213,
645 doi:10.1016/j.immuni.2007.01.009 (2007).
- 646 8 Murugan, R. *et al.* Clonal selection drives protective memory B cell responses in
647 controlled human malaria infection. *Sci Immunol* **3**, doi:10.1126/sciimmunol.aap8029
648 (2018).
- 649 9 Briney, B., Inderbitzin, A., Joyce, C. & Burton, D. R. Commonality despite exceptional
650 diversity in the baseline human antibody repertoire. *Nature* **566**, 393-397,
651 doi:10.1038/s41586-019-0879-y (2019).
- 652 10 ter Meulen, J. *et al.* Human monoclonal antibody combination against SARS coronavirus:
653 synergy and coverage of escape mutants. *PLoS Med* **3**, e237,
654 doi:10.1371/journal.pmed.0030237 (2006).
- 655 11 Yuan, M. *et al.* A highly conserved cryptic epitope in the receptor binding domains of
656 SARS-CoV-2 and SARS-CoV. *Science* **368**, 630-633, doi:10.1126/science.abb7269
657 (2020).
- 658 12 Walls, A. C. *et al.* Unexpected Receptor Functional Mimicry Elucidates Activation of
659 Coronavirus Fusion. *Cell* **176**, 1026-1039 e1015, doi:10.1016/j.cell.2018.12.028 (2019).
- 660 13 Pinto, D. *et al.* Cross-neutralization of SARS-CoV-2 by a human monoclonal SARS-CoV
661 antibody. *Nature*, doi:10.1038/s41586-020-2349-y (2020).
- 662 14 Zhu, Z. *et al.* Potent cross-reactive neutralization of SARS coronavirus isolates by human
663 monoclonal antibodies. *Proc Natl Acad Sci U S A* **104**, 12123-12128,
664 doi:10.1073/pnas.0701000104 (2007).
- 665 15 Salazar, G., Zhang, N., Fu, T. M. & An, Z. Antibody therapies for the prevention and
666 treatment of viral infections. *NPJ Vaccines* **2**, 19, doi:10.1038/s41541-017-0019-3
667 (2017).
- 668 16 Bournazos, S. & Ravetch, J. V. Anti-retroviral antibody Fcγ-mediated effector
669 functions. *Immunol Rev* **275**, 285-295, doi:10.1111/imr.12482 (2017).
- 670 17 Feinberg, M. B. & Ahmed, R. Advancing dengue vaccine development. *Science* **358**,
671 865-866, doi:10.1126/science.aaq0215 (2017).
- 672 18 Iwasaki, A. & Yang, Y. The potential danger of suboptimal antibody responses in
673 COVID-19. *Nat Rev Immunol*, doi:10.1038/s41577-020-0321-6 (2020).

- 674 19 Van Rompay, K. K. A. *et al.* A combination of two human monoclonal antibodies limits
675 fetal damage by Zika virus in macaques. *Proc Natl Acad Sci U S A* **117**, 7981-7989,
676 doi:10.1073/pnas.2000414117 (2020).
- 677 20 Plotkin, S. A. Correlates of protection induced by vaccination. *Clin Vaccine Immunol* **17**,
678 1055-1065, doi:10.1128/CVI.00131-10 (2010).
- 679 21 Scheid, J. F. *et al.* Sequence and structural convergence of broad and potent HIV
680 antibodies that mimic CD4 binding. *Science* **333**, 1633-1637,
681 doi:10.1126/science.1207227 (2011).
- 682 22 Robbiani, D. F. *et al.* Recurrent Potent Human Neutralizing Antibodies to Zika Virus in
683 Brazil and Mexico. *Cell* **169**, 597-609 e511, doi:10.1016/j.cell.2017.04.024 (2017).
- 684 23 Ehrhardt, S. A. *et al.* Polyclonal and convergent antibody response to Ebola virus vaccine
685 rVSV-ZEBOV. *Nat Med* **25**, 1589-1600, doi:10.1038/s41591-019-0602-4 (2019).
- 686 24 Pappas, L. *et al.* Rapid development of broadly influenza neutralizing antibodies through
687 redundant mutations. *Nature* **516**, 418-422, doi:10.1038/nature13764 (2014).
- 688 25 Kane, M. *et al.* Identification of Interferon-Stimulated Genes with Antiretroviral Activity.
689 *Cell Host Microbe* **20**, 392-405, doi:10.1016/j.chom.2016.08.005 (2016).
- 690 26 Adachi, A. *et al.* Production of acquired immunodeficiency syndrome-associated
691 retrovirus in human and nonhuman cells transfected with an infectious molecular clone. *J*
692 *Virol* **59**, 284-291 (1986).
- 693 27 Wang, Z. *et al.* Isolation of single HIV-1 Envelope specific B cells and antibody cloning
694 from immunized rhesus macaques. *J Immunol Methods* **478**, 112734,
695 doi:10.1016/j.jim.2019.112734 (2020).
- 696 28 Tiller, T. *et al.* Efficient generation of monoclonal antibodies from single human B cells
697 by single cell RT-PCR and expression vector cloning. *J Immunol Methods* **329**, 112-124,
698 doi:10.1016/j.jim.2007.09.017 (2008).
- 699 29 von Boehmer, L. *et al.* Sequencing and cloning of antigen-specific antibodies from
700 mouse memory B cells. *Nat Protoc* **11**, 1908-1923, doi:10.1038/nprot.2016.102 (2016).
- 701 30 Klein, F. *et al.* Enhanced HIV-1 immunotherapy by commonly arising antibodies that
702 target virus escape variants. *J Exp Med* **211**, 2361-2372, doi:10.1084/jem.20141050
703 (2014).
- 704 31 Schoofs, T. *et al.* Broad and Potent Neutralizing Antibodies Recognize the Silent Face of
705 the HIV Envelope. *Immunity* **50**, 1513-1529 e1519, doi:10.1016/j.immuni.2019.04.014
706 (2019).
- 707 32 Ye, J., Ma, N., Madden, T. L. & Ostell, J. M. IgBLAST: an immunoglobulin variable
708 domain sequence analysis tool. *Nucleic Acids Res* **41**, W34-40, doi:10.1093/nar/gkt382
709 (2013).
- 710 33 Gupta, N. T. *et al.* Change-O: a toolkit for analyzing large-scale B cell immunoglobulin
711 repertoire sequencing data. *Bioinformatics* **31**, 3356-3358,
712 doi:10.1093/bioinformatics/btv359 (2015).
- 713 34 Rubelt, F. *et al.* Onset of immune senescence defined by unbiased pyrosequencing of
714 human immunoglobulin mRNA repertoires. *PLoS One* **7**, e49774,
715 doi:10.1371/journal.pone.0049774 (2012).
- 716 35 Kyte, J. & Doolittle, R. F. A simple method for displaying the hydropathic character of a
717 protein. *J Mol Biol* **157**, 105-132, doi:10.1016/0022-2836(82)90515-0 (1982).
- 718 36 Guy, H. R. Amino acid side-chain partition energies and distribution of residues in
719 soluble proteins. *Biophys J* **47**, 61-70, doi:10.1016/S0006-3495(85)83877-7 (1985).

- 720 37 DeWitt, W. S. *et al.* A Public Database of Memory and Naive B-Cell Receptor
721 Sequences. *PLoS One* **11**, e0160853, doi:10.1371/journal.pone.0160853 (2016).
722 38 Mastronarde, D. N. Automated electron microscope tomography using robust prediction
723 of specimen movements. *J Struct Biol* **152**, 36-51, doi:10.1016/j.jsb.2005.07.007 (2005).
724 39 Punjani, A., Rubinstein, J. L., Fleet, D. J. & Brubaker, M. A. cryoSPARC: algorithms for
725 rapid unsupervised cryo-EM structure determination. *Nat Methods* **14**, 290-296,
726 doi:10.1038/nmeth.4169 (2017).
727 40 Goddard, T. D., Huang, C. C. & Ferrin, T. E. Visualizing density maps with UCSF
728 Chimera. *J Struct Biol* **157**, 281-287, doi:10.1016/j.jsb.2006.06.010 (2007).
729