

Article

## Study on the Synthesis, Biological Activity and Spectroscopy of Naphthalimide-Diamine Conjugates

Zhi-Yong Tian <sup>1,\*</sup>, Jing-Hua Li <sup>2</sup>, Qian Li <sup>2</sup>, Feng-Lei Zang <sup>1</sup>, Zhong-Hua Zhao <sup>1</sup>  
and Chao-Jie Wang <sup>1,2,\*</sup>

<sup>1</sup> Institute of Chemical Biology, Henan University, Kaifeng 475004, China;

E-Mails: Zangfenglei@sohu.com (F.-L.Z.); Zhaozhonghua@sohu.com (Z.-H.Z.)

<sup>2</sup> Key Laboratory of Natural Medicine and Immuno-Engineering, Henan University,

Kaifeng 475004, China; E-Mails: Lijinghua@henu.edu.cn (J.-H.L.); Liqian426@163.com (Q.L.)

\* Authors to whom correspondence should be addressed; E-Mail: tzynew@sohu.com (Z.-Y.T.);

wcjsxq@henu.edu.cn (C.-J.W.); Tel./Fax: +86-0371-2388-0680 (Z.-Y.T.);

Fax: +86-0371-2286-4665 (C.-J.W.).

Received: 20 February 2014; in revised form: 23 May 2014 / Accepted: 27 May 2014 /

Published: 10 June 2014

---

**Abstract:** Eleven novel naphthalimide-diamine conjugates were synthesized and their structures were confirmed by elemental analysis, <sup>1</sup>H-NMR, <sup>13</sup>C-NMR and MS. Their *in vitro* antitumor activities were assessed using MTT assays on two cancerous cell lines K562, HCT116, and one normal hepatoma cell line QSG 7701. Compound **7f** exhibited potent antitumor activity on HCT116 cells and favorable cell selectivity toward QSG 7701 compared with the positive control, amonafide. Moreover, **7f** could block HeG2 cells in the G2/M phase and induce HeG2 cells apoptosis. The interaction of compound **7f** with herring sperm DNA was studied by UV/vis absorption and fluorescence spectroscopy under physiological conditions (pH = 7.4). The observed spectral quenching of compound **7f** by DNA and the displacement of EB from DNA-EB complex by compound **7f** indicated that compound **7f** could intercalate into DNA base pairs, which was also corroborated by the effect of KI on compound-DNA interaction. Further calorimetric fluorescent tests revealed that the quenching mechanism was a static type. Meanwhile, the binding constants, thermodynamic parameters and the effect of NaCl on compound-DNA interaction showed that the type of interaction force was mainly hydrogen bonds and the binding process was driven by hydrogen and van der Waals bonding.

**Keywords:** naphthalimide; diamine conjugates; synthesis; antitumor activity; spectroscopy


## 1. Introduction

As a common chemical motif, naphthalimides are among a growing class of compounds with desirable anticancer activity. Numerous mono- or bis-naphthalimide derivatives displayed potent antitumor properties against a variety of murine and human tumor cells [1–4], and some of them such as mitonafide [5], amonafide [6], azonafide [7], DMP-840 [8,9] and Lu-79553 [10] have been tested in clinic trials for the treatment of solid tumors. However, most clinical trials have failed because of a poor therapeutic index, poor water-solubility or dose-limiting bone marrow toxicity such as in the case of mitonafide [11]. Subsequent efforts to improve therapeutic properties of naphthalimides have been made by modifying the naphthalimide skeleton [12–25].

Previous studies revealed that polyamines are a kind of promising carriers to transport cytotoxic agents into cancer cells [26]. Polyamines are important for tumor cell growth and function, the biosynthetic pathway of native polyamines (putrescine, spermidine and spermine) has been a popular target for therapeutic intervention during the last decades [27].

Our previous work proved that conjugates **1–3** (Figure 1), composed of 1,8-naphthalimide units covalently attached to a polyamine such as spermidine or homospermidine, possessed remarkable cell selectivity through to human hepatoma Bel-7402 and human normal hepatocyte QSG-7701 trials [28,29]. In addition, naphthalimide-polyamine conjugates have been proved to induce cancer cell apoptosis *via* different pathways [28–30]. These results encouraged us to screen more substituted naphthalimide-polyamine conjugates in order to assess their antitumor activity.

**Figure 1.** The structures of compounds **1–3**.



DNA as carrier of genetic information is a major target for drug interaction due to its abilities to interfere with transcription (gene expression and protein synthesis) and DNA replication, a major step in cell growth and division. Generally, a variety of small molecules interact reversibly with DNA in three primary ways, including intercalation of planar or approximately planar aromatic ring systems between base-pairs [31], groove binding in which the small molecules bound on nucleic acids are located in the major or minor groove [18,19] and binding along the exterior of DNA helix through interactions which are generally nonspecific and are primarily electrostatic. The 1,8-naphthalimide derivatives are the DNA intercalating agents because they consist of a flat, generally  $p$ - $\pi$  deficient aromatic system of which binds to DNA by insertion between the base pairs of the double helix. However, there are rare reports on the interaction mechanism of naphthalimide-polyamine conjugates and DNA. In this work, naphthalimide-diamine conjugates were synthesized and their antitumor activity


assessed *in vitro*. The interactions between a representative compound **7f** and herring sperm DNA were first studied by UV and fluorescence spectroscopy. The binding constants and main sorts of binding force were also investigated. Moreover, mechanism of how the novel conjugate **7f** killed HeG2 cells **7f** was reported.

## 2. Results and Discussion

### 2.1. Chemistry

The synthetic route to the naphthalimide-diamine conjugates **7a–k** is shown in Scheme 1. N-(3-bromopropyl)-1,8-naphthalimide (**4a**), N-(4-bromobutyl)-1,8-naphthalimide (**4b**) and N-(5-bromopentyl)-1,8-naphthalimide (**4c**) were prepared routinely from 1,8-naphthalimide and 1,4-dibromobutane (or 1,3-dibromopropane, 1,5-dibromopentane) in the presence of potassium carbonate and potassium iodide. The Boc-protected diamines **5** were prepared according to a modified procedure reported previously [32]. The N-alkylation reactions of **5** with **4** in the presence of potassium carbonate in dry acetonitrile at 45 °C generated intermediates which were difficult to purify because of their instability [33]. Without separation, N-Boc protection with Boc<sub>2</sub>O led to the formation of stable intermediates **6**. Subsequently the Boc groups were removed with 4M HCl at room temperature to provide target compounds **7** as dihydrochloride salts in yields of 50%–80%. The structures of target compounds **7** were confirmed by <sup>1</sup>H-NMR, <sup>13</sup>C-NMR, ESI-MS, and elemental analysis.

**Scheme 1.** Synthesis of naphthalimide-diamine conjugates.



### 2.2. Cytotoxic Effects

The cytotoxicities of the novel conjugates were assessed *in vitro* by the MTT assay in the presence of aminoguanidine (an inhibitor of amine oxidase) against three cell lines, human leukemia K562, cancer of colon HCT116 and human normal hepatocyte QSG-7701 [34]. As is shown in Table 1, most of the naphthalimide-diamine conjugates **7a–k** showed good antitumor activities against K562,


HCT116 cells, in which compound **7f** seemed to have equipotent antitumor activity against HCT116 and cell selectivity toward QSG-7701 compared to the control, amonafide. At the same time, compounds **7f–h**, which have putrescine backbones, exhibited good antitumor activities against K562, HCT116 cells. In addition, compounds **7j–k**, which have longer side chains than the others, also exhibited good antitumor activities. Moreover, The compound **7f** with a primary terminal amino group in the polyamine motif exhibited better biological properties than the corresponding compounds with a terminal tertiary amino group [35]. However, the dose-responsive curves shown in Figure 2 indicated that compound **7f** displayed much better selectivity between HCT116 and QSG-7701 than amonafide, especially at the concentrations around the  $IC_{50}$  values (5–10  $\mu\text{M}$ ). Thus, compound **7f** was selected for further investigation.

**Table 1.** *In vitro* activity of compound naphthalimide-diamine conjugates (**7a–k**).

Compd.	$IC_{50}$ ( $\mu\text{M}$ )		
	K562	HCT116	7701
<b>7a</b>	48.76 $\pm$ 3.93	19.33 $\pm$ 4.39	>50
<b>7b</b>	25.60 $\pm$ 3.74	13.93 $\pm$ 2.68	33.89 $\pm$ 4.32
<b>7c</b>	38.75 $\pm$ 1.76	9.89 $\pm$ 1.11	>50
<b>7d</b>	19.71 $\pm$ 7.23	8.95 $\pm$ 3.87	8.04 $\pm$ 3.35
<b>7e</b>	22.36 $\pm$ 1.21	9.52 $\pm$ 3.31	22.16 $\pm$ 0.67
<b>7f</b>	18.87 $\pm$ 1.97	5.45 $\pm$ 1.71	19.69 $\pm$ 3.42
<b>7g</b>	14.03 $\pm$ 3.32	13.88 $\pm$ 2.54	11.97 $\pm$ 4.07
<b>7h</b>	15.89 $\pm$ 1.53	16.21 $\pm$ 4.28	18.78 $\pm$ 4.94
<b>7i</b>	41.35 $\pm$ 1.76	10.97 $\pm$ 1.71	>50
<b>7j</b>	12.56 $\pm$ 0.81	16.44 $\pm$ 1.75	13.13 $\pm$ 3.94
<b>7k</b>	15.89 $\pm$ 1.83	9.95 $\pm$ 1.81	14.70 $\pm$ 3.91
<b>Amonafide</b>	10.10 $\pm$ 2.59	6.86 $\pm$ 1.89	20.29 $\pm$ 2.43

All data are expressed as means  $\pm$  SD from three separate determinations.  $IC_{50}$  values were given only if they were less than 50  $\mu\text{M}$ , which was the maximum concentration tested.


**Figure 2.** The proliferation inhibition of **7f** and amonafide assayed by MTT.



### 2.3. Cell Cycle Analysis and Apoptosis


To reveal the cytotoxic mechanism of compound **7f**, we first examined its effects on the cell cycle perturbation. The DNA content analysis by High Content Screening (HCS) confirmed that **7f** could induce HepG2 cell cycle perturbation (Figure 3). The exposure of HepG2 cells to **7f** resulted in the dose-dependent accumulation of cells in G2/M phase from 29.7% of the control to 49.8% of **7f**. Meanwhile, the corresponding reduction of G0/G1 phase from 52.6% of the control to 29.7% of **7f** was also observed, accompanying by the little change in the S phase. The results showed that compound **7f** could block HepG2 cells in the G2/M phase.

**Figure 3.** Arrest of cell cycle progress in HepG2 cells treated with **7f** for 48 h. Cells were fixed with ethanol and stained with PI. Cell cycle distribution was analyzed by HCS. (A) Control; (B) Compound **7f**: 10  $\mu$ M; (C) Compound **7f**: 20  $\mu$ M.



Apoptosis, or namely programmed cell death, can be triggered by several stimuli. Both naphthalimides and polyamine derivatives could trigger cell apoptosis [36]. In order to determine whether the antitumor-proliferative effect of compound **7f** was associated with cell apoptosis, the HepG2 cell apoptosis was detected by the staining of DNA with Hoechst 33342, and the apoptotic cells were counted by selecting 200 cells randomly (Figure 4). The apoptotic bodies were observed clearly in **7f**-treated groups by laser scanning confocal microscope. **7f** could trigger apoptosis in a dose-dependent manner with the 47.0% apoptotic ratio of **7f** at 20  $\mu$ M compared with the less than five percent in the control group.


**Figure 4.** **7f** induced HepG2 cell apoptosis. (A) Control; (B) Compound **7f**: 10  $\mu$ M; (C) Compound **7f**: 20  $\mu$ M.



## 2.4. UV Spectroscopy

As is shown in Figure 5, the UV spectrum of compound **7f** in the absence and presence of herring sperm DNA was measured by an ultraviolet visible range spectrophotometer. It was observed that a continuous decrease in the absorbance of compound **7f** followed with the increasing concentration of DNA, implying compound **7f** could insert into the base pairs of DNA. The spectral effects have been rationalized that the empty  $\pi^*$ -orbital of the small molecule couples with the  $\pi^*$ -orbital of the DNA base pairs, which causes an energy decrease and a decrease of the  $\pi$ - $\pi^*$  transition energy [37–39]. At the same time, the empty  $\pi^*$ -orbital is partially filled with electrons to reduce the transition probability. Therefore, the absorption of small molecules should exhibit hypochromism.

**Figure 5.** UV absorption spectra of compound **7f** with herring sperm DNA. Numbers 1–13 indicated the DNA concentration:  $0.0$ ,  $4.56 \times 10^{-6}$ ,  $9.13 \times 10^{-6}$ ,  $13.69 \times 10^{-6}$ ,  $27.4 \times 10^{-6}$ ,  $41.08 \times 10^{-6}$ ,  $54.77 \times 10^{-6}$ ,  $68.46 \times 10^{-6}$ ,  $82.15 \times 10^{-6}$ ,  $95.84 \times 10^{-6}$ ,  $109.54 \times 10^{-6}$ ,  $123.23 \times 10^{-6}$  and  $136.92 \times 10^{-6} \text{ mol}\cdot\text{L}^{-1}$ , respectively. The concentration of compound **7f** applied were  $80 \times 10^{-6} \text{ mol}\cdot\text{L}^{-1}$ .




Utilizing the absorption spectrum obtained by UV, we could also calculate compound **7f**'s apparent binding constant according to the following formula [40]:

$$\frac{A_0}{A - A_0} = \frac{\epsilon_G}{\epsilon_{H-G} - \epsilon_G} + \frac{\epsilon_G}{\epsilon_{H-G} - \epsilon_G} \frac{1}{K [DNA]} \quad (1)$$

where  $A_0$  and  $A$  denote the absorbance in the absence and presence of DNA, respectively and where  $\epsilon_G$  and  $\epsilon_{H-G}$  denote the molar absorption coefficient of compound and its formed complex with DNA. The value of apparent binding constant could be measured from the intercept and slope by plotting  $A_0/(A - A_0)$  against  $c_{DNA}$ , and the corresponding value of  $K$  was  $9.806 \times 10^3$  (Figure 6).

**Figure 6.** Plot of  $A_0/(A - A_0)$  versus  $1/c_{DNA}$  of the interaction between compound **7f** and herring sperm DNA ( $10^{-6}$  mol·L $^{-1}$ ).




## 2.5. Fluorescence Spectroscopy

### 2.5.1. Fluorescence Quenching

To evaluate the DNA binding properties of naphthalimide homospermidine conjugate, the inherent fluorescence of compound **7f** allowed us to investigate its interaction with herring sperm DNA by fluorescence spectrometry.


**Figure 7.** Fluorescence spectroscopy of compound **7f** and herring sperm DNA. Numbers 1–13 indicated the DNA concentration:  $0.0$ ,  $4.56 \times 10^{-6}$ ,  $9.13 \times 10^{-6}$ ,  $13.69 \times 10^{-6}$ ,  $27.4 \times 10^{-6}$ ,  $41.08 \times 10^{-6}$ ,  $54.77 \times 10^{-6}$ ,  $68.46 \times 10^{-6}$ ,  $82.15 \times 10^{-6}$ ,  $95.84 \times 10^{-6}$ ,  $109.54 \times 10^{-6}$ ,  $123.23 \times 10^{-6}$  and  $136.92 \times 10^{-6}$  mol·L $^{-1}$ , respectively. Compound **7f** applied was  $80 \times 10^{-6}$  mol·L $^{-1}$ . Scan condition of compound **7f**: EX = 345nm, EM = 355~600 nm; Slits of both EX and EM were 5 nm and 2.5 nm, respectively.



As is shown in Figure 7, the fluorescence of compound **7f** was quenched upon the addition of DNA. This indicated that DNA is one potential target of compound **7f** as expected.

Ethidium bromide (EB) is a well known DNA intercalator, which is often used as a spectral probe to establish the mode of binding of small molecules to double-helical DNA [41]. The fluorescence of EB increases after binding with DNA due to intercalation. Like EB, if naphthalimides intercalate into the helix of DNA, it would compete with EB for its intercalation sites in DNA, and the displacement of EB from the DNA-EB complex leads to a significant decrease in the fluorescence intensity of the DNA-EB complex [42–44]. Therefore, herring sperm DNA-EB complex in the presence of increasing concentrations of naphthalimide-diamine conjugate **7f** was also measured. As is shown in Figure 8, the fluorescence intensity of DNA-EB complex was decreased by gradually growing concentrations of compound **7f**, suggesting that compound **7f** could intercalate into DNA and a new complex was possibly formed between compound **7f** and DNA [45].

**Figure 8.** Fluorescence spectroscopy of compound **7f** with herring sperm DNA-EB. Numbers 1–13 indicated the compound **7f** concentration:  $0.0$ ,  $4 \times 10^{-6}$ ,  $8 \times 10^{-6}$ ,  $12 \times 10^{-6}$ ,  $24 \times 10^{-6}$ ,  $36 \times 10^{-6}$ ,  $48 \times 10^{-6}$ ,  $60 \times 10^{-6}$ ,  $72 \times 10^{-6}$ ,  $84 \times 10^{-6}$ ,  $96 \times 10^{-6}$ ,  $108 \times 10^{-6}$  and  $120 \times 10^{-6}$  mol·L<sup>-1</sup>, respectively. DNA and EB applied was  $13.7 \times 10^{-6}$  and  $15.7 \times 10^{-6}$  mol·L<sup>-1</sup>, respectively. Scan condition: EX = 510 nm, EM = 520~800 nm; Slits of both EX and EM were 5 nm and 10 nm, respectively.



### 2.5.2. Fluorescence Quenching Mechanism

Fluorescence quenching can occur by different mechanisms, which are usually classified as dynamic quenching and static quenching. Dynamic quenching refers to a process whereby the fluorophore and the quencher come into contact during the transient existence of the excited state, so the bimolecular quenching constants would be larger at higher temperatures. Static quenching, however, results from the formation of a ground state complex between the fluorophore and the quencher, which decreases [46]. To elucidate the quenching mechanism of the interaction between naphthalimide-diamine conjugate and DNA (or DNA-EB), fluorescence quenching tests were also performed at 298, 303 and 310 K, which could be described by Stern-Volmer equation [47–49].




The Stern–Volmer equation is the following:


$$F_0/F = 1 + K_{SV} c = 1 + K \tau_0 c \quad (2)$$

where  $F_0$  and  $F$  are the fluorescence intensities in the absence and presence of quencher (DNA for compound **7f** or compound **7f** for DNA-EB, respectively),  $K_{SV}$  is the Stern–Volmer quenching constant,  $[c]$  is the concentration of DNA (or compound **7f**),  $K_q$  is the biomolecule quenching rate constant and  $K_q = K_{SV}/\tau_0$ .  $\tau_0$  is the average lifetime of the molecule without any quencher and the fluorescence lifetime of the biopolymer is  $10^{-8}$  s [50]. The Stern–Volmer plots of  $F_0/F$  versus  $[c]$  at the three temperatures were shown in Figures 9 and 10, and the calculated  $K_{SV}$  and  $K_q$  values were presented in Tables 2 and 3.

**Figure 9.** Stern–Volmer plot of fluorescence quenching of compound **7f** by herring sperm DNA at different temperatures.



**Figure 10.** Stern–Volmer plot of fluorescence quenching of herring sperm DNA-EB by compound **7f** at different temperatures.



**Table 2.** Quenching constants of the interaction between compound **7f** and herring sperm DNA at different temperatures.

<i>T/K</i>	$K_{SV}/(\text{L}\cdot\text{mol}^{-1})$	$K_q/(\text{L}\cdot\text{mol}^{-1})$	<i>R</i>
298	$5.567 \times 10^3$	$5.567 \times 10^{11}$	0.9956
303	$4.789 \times 10^3$	$4.789 \times 10^{11}$	0.9987
310	$3.651 \times 10^3$	$3.651 \times 10^{11}$	0.9945

**Table 3.** Quenching constants of the interaction between compound **7f** and herring sperm DNA-EB at different temperatures.

<i>T/K</i>	$K_{SV}/(\text{L}\cdot\text{mol}^{-1})$	$K_q/(\text{L}\cdot\text{mol}^{-1})$	<i>R</i>
298	$5.028 \times 10^3$	$5.028 \times 10^{11}$	0.9907
303	$4.564 \times 10^3$	$4.654 \times 10^{11}$	0.9922
310	$4.176 \times 10^3$	$4.176 \times 10^{11}$	0.9809


The values of the quenching constant  $K_{SV}$  decreased with increasing temperature and the values of  $K_q$  were much greater than that of the maximum scattering collision quenching constant ( $2.000 \times 10^{10} \text{ L}\cdot\text{mol}^{-1}$ ), indicating that the fluorescence quenching of compound **7f** initiated by DNA or DNA-EB complex initiated by compound **7f** was static quenching [51].

### 2.5.3. Interaction Mode between Compounds and DNA


When small molecules bind independently to a set of equivalent sites in a macromolecule, the binding constant ( $K_b$ ) can be determined by the following equation [40,52]:

$$\log[1/c] = \log[F/(F_0 - F)] + \log K_b \quad (3)$$

where  $K_b$  denotes the binding constant for interaction of naphthalimide–DNA, and  $F_0$ ,  $F$ ,  $[c]$  have the same meanings as in Equation (2).

**Figure 11.** Plot of  $\log [1/c_{DNA}]$  versus  $\log [F/(F_0 - F)]$  of the interaction between compound **7f** and herring sperm DNA at different temperatures.

**Figure 12.** Plot of  $\log [1/c_{7f}]$  versus  $\log [F/(F_0 - F)]$  of the interaction between compound **7f** and herring sperm DNA-EB at different temperatures.



**Table 4.** Binding constants and thermodynamic parameters of the interaction between compound **7f** and herring sperm DNA at different temperatures.

$T/K$	$K_b/(\text{kJ}\cdot\text{mol}^{-1})$	$\Delta G^\circ/(\text{kJ}\cdot\text{mol}^{-1})$	$\Delta H^\circ/(\text{kJ}\cdot\text{mol}^{-1})$	$\Delta S^\circ/(\text{kJ}\cdot\text{mol}^{-1})$	$R$
298	$6.059 \times 10^3$	-21.578	-32.983	-0.0383	0.9897
303	$4.864 \times 10^3$	-21.387	-32.983	-0.0383	0.9965
310	$3.462 \times 10^3$	-21.004	-32.983	-0.0383	0.9923

**Table 5.** Binding constants and thermodynamic parameters of the interaction between compound **7f** and herring sperm DNA-EB at different temperatures.

$T/K$	$K_b/(\text{kJ}\cdot\text{mol}^{-1})$	$\Delta G^\circ/(\text{kJ}\cdot\text{mol}^{-1})$	$\Delta H^\circ/(\text{kJ}\cdot\text{mol}^{-1})$	$\Delta S^\circ/(\text{kJ}\cdot\text{mol}^{-1})$	$R$
298	$5.482 \times 10^3$	-21.330	-59.229	-0.127	0.9884
303	$3.695 \times 10^3$	-21.172	-59.229	-0.127	0.9982
310	$3.611 \times 10^3$	-21.112	-59.229	-0.127	0.9975

The values of  $K_b$  could be measured from the intercept and slope by plotting  $\log [1/c]$  against  $\log [F/(F_0 - F)]$  (intercept =  $\log K_b$ ) (Figures 11 and 12), and the corresponding values of  $K_b$  were listed in Tables 4 and 5. The down-regulated trend of  $K_b$  with increasing temperature was in accordance with  $K_{SV}$ 's dependence on temperature as mentioned above, implying that the binding between naphthalimide and DNA was moderate, and a reversible naphthalimide-DNA complex might be formed [53].

There are several acting forces between a small organic molecule and biomacromolecules, such as hydrophobic force, hydrogen bond, van der Waals force, electrostatic interactions, *etc.* It is assumed that the interaction enthalpy change ( $\Delta H^\circ$ ) does not vary significantly over the limited temperature range studied, thus the thermodynamic parameters can be calculated from the van't Hoff equation:

$$\ln (K_2/K_1) = (1/T_1 - 1/T_2) \Delta H^\circ/R \quad (4)$$

$$\Delta G^\circ = -RT \ln K = \Delta H^\circ - T\Delta S^\circ \quad (5)$$


In Equations (4) and (5),  $K$  is analogous to the binding constant at the corresponding temperature and  $R$  is gas constant. The enthalpy change ( $\Delta H^\circ$ ) and entropy change ( $\Delta S^\circ$ ) were calculated from the Equations (4) and (5), and the corresponding results were listed in Tables 4 and 5. From Tables 4 and 5, it can be seen that the negative  $\Delta H^\circ$  and negative  $\Delta S^\circ$  values showed that the hydrogen bond and weak van der Waals played a dominant role in the interactions between compound **7f** and DNA [54].

Kenaka [55] found that hydrogen bonding was the main feature of DNA intercalating agents, and the present evidences indicated that compound **7f** was a DNA intercalator. Therefore, it was inferred that the process of interaction between compound **7f** and DNA was driven by hydrogen bonding and van der Waals forces. It was also inferred that planar structure of naphthalene ring intercalated into the DNA base pairs when compound **7f** bound to DNA between the double helix and that the hydrogen bond and weak van der Waals played a dominant role in the interactions between compound **7f** and DNA.

#### 2.5.4. Iodide Quenching Studies

A highly negatively charged quencher is expected to be repelled by the negatively charged phosphate backbone of DNA, therefore an intercalative bound drug molecules should be protected from being quenched by anionic quencher, but the free aqueous complexes or groove binding drugs should be quenched readily by anionic quenchers. At the same time, whether the quencher accesses to fluorophore also plays a role in free and bound one [56].

**Figure 13.** Fluorescence quenching plots of compound **7f** with KI in the absence and presence of DNA.  $c(\mathbf{7f}) = 20 \times 10^{-6} \text{ mol}\cdot\text{L}^{-1}$ ;  $c(\text{DNA}) = 22.84 \times 10^{-6} \text{ mol}\cdot\text{L}^{-1}$ ; the KI concentration was  $4\sim 120 \times 10^{-3} \text{ mol}\cdot\text{L}$ .




Negatively charged  $\Gamma^-$  was selected for this purpose. The quenching constants ( $K_{sv}$ ) were obtained from the Stern–Volmer equation. The values of  $K_{sv}$  of compound **7f** with  $\Gamma^-$  ion in the absence and presence of DNA were 38.178 and 31.575 ( $\text{L}\cdot\text{mol}^{-1}$ ), respectively (shown in Figure 13). It was apparent that iodide quenching effect was decreased when compound **7f** was bound to DNA, which suggested that the compound **7f** is likely intercalated into the base pairs of DNA.

### 2.5.5. Effect of Ionic Intensity on the Compound **7f** and DNA Interaction

DNA is an anionic polyelectrolyte with phosphate groups and monitoring the spectral change with different ionic strength is an efficient method to distinguish the binding modes between molecules and DNA. NaCl is used to control the ionic strength of the solutions. The addition of Na<sup>+</sup> would attenuate the electrostatic interaction between molecules and DNA because of its competition for phosphate groups in DNA [57]. Hence the effect of NaCl on the fluorescence of DNA–compound **7f** system was studied. As is seen from Figure 14, the fluorescence intensity of compound **7f**-DNA complex was basically unchanged with increasing concentration of NaCl. The results revealed that interaction between compound **7f** and DNA could exclude the electrostatic interaction mode and was prompted by hydrogen bond and van der Waals force.

**Figure 14.** Effects of NaCl on the fluorescence intensity of compound **7f**-DNA system.



## 3. Experimental Section

### 3.1. General Information

All chemicals (reagent grade) used were commercially available. All the <sup>1</sup>H-NMR and <sup>13</sup>C-NMR spectra were recorded on a Bruker AV-400 model spectrometer in D<sub>2</sub>O. Chemical shifts ( $\delta$ ) for <sup>1</sup>H-NMR spectra were reported in parts per million to residual solvent protons. ESI-MS spectra were recorded on an ESQUIRE-LC Mass spectrometer. Elemental analyses were performed on a Gmbe VarioEL Elemental instrument and were within 0.4% of the theoretical values.

### 3.2. Synthesis of Naphthalimide-Diamine Conjugates

The Boc-protected amine **5** (3 mmol) was dissolved in anhydrous acetonitrile (30 mL), and solid K<sub>2</sub>CO<sub>3</sub> (0.69 g, 5 mmol) was added. After the mixture was stirred for 15 minutes at ambient temperature, *N*-(3-bromopropyl)-1,8-naphthalimide (**4a**) (or *N*-(4-bromobutyl)-1,8-naphthalimide (**4b**) and *N*-(5-bromopentyl)-1,8-naphthalimide (**4c**), 2 mmol) in acetonitrile (10 mL) was added dropwise

with constant stirring at 45 °C, then the reaction mixture was stirred overnight. The volatiles were removed under vacuum to give a residue which was redissolved in CHCl<sub>3</sub> (30 mL) and washed with aqueous Na<sub>2</sub>CO<sub>3</sub> (10%, w/v). The organic phase was separated, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated. The above residue was dissolved in methanol (30 mL), a solution of di-*tert*-butyl dicarbonate (3 mmol) in methanol (10 mL) was added dropwise and stirred at ambient temperature overnight. The solvents were removed under reduced pressure to give a residue, which was redissolved in CHCl<sub>3</sub> (30 mL) and washed with water. The organic phase was separated, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated. The residue was subjected to flash chromatography (20% Petroleum /EtOAc, v/v, R<sub>f</sub> = 0.25) to obtain the Boc protected intermediates **6**.

The respective N-Boc-protected amine (1.2 mmol) was dissolved in EtOH (20 mL) and stirred at 0 °C for 10 min. 4 M HCl (15 mL) was added dropwise at 0 °C. The reaction mixture was stirred at room-temperature overnight. The solution typically gave a bright white solid as a precipitate. The solid was filtered, washed several times with absolute ethanol and dried under vacuum to give the pure target compounds **7**.

*2-[3-(2-Aminoethylamino)propyl]1H-benz-[de]isoquinoline-1,3(2H)-dione dihydrochloride (7a)*. Yield: 76.3%; m.p.: 233.4–235.1 °C; <sup>1</sup>H-NMR (D<sub>2</sub>O) δ: 7.37 (d, 2H, *J* = 8.0 Hz), 7.33 (d, 2H, *J* = 7.2 Hz), 7.0 (t, 2H, *J* = 7.6 Hz), 3.55 (t, 2H, *J* = 7.2 Hz), 3.46 (s, 4H), 3.15 (t, 2H, *J* = 8.0 Hz), 1.83 (t, 2H, *J* = 6.8 Hz); <sup>13</sup>C-NMR (D<sub>2</sub>O) δ: 163.75, 134.73, 130.76, 129.54, 126.52, 125.10, 118.75, 45.79, 44.35, 37.16, 35.50, 24.11; ESI-MS *m/z*: 298.2 (M+H-2HCl)<sup>+</sup>; Anal. calcd. for C<sub>17</sub>H<sub>21</sub>Cl<sub>2</sub>N<sub>3</sub>O<sub>2</sub> C, 55.14; H, 5.72; N, 11.35; found: C, 54.83; H, 5.48; N, 10.94.

*2-[4-(2-Aminoethylamino)butyl]1H-benz-[de]isoquinoline-1,3(2H)-dione dihydrochloride (7b)*. Yield: 58.1%; m.p.: 243.1–244.1 °C; <sup>1</sup>H-NMR (D<sub>2</sub>O) δ: 7.39 (d, 2H, *J* = 8.0 Hz), 7.00 (d, 2H, *J* = 7.2 Hz), 7.12 (t, 2H, *J* = 7.6 Hz), 3.43–3.47 (m, 6H), 3.18 (t, 2H, *J* = 8.0 Hz), 1.71–1.74 (m, 2H), 1.45–1.46 (m, 2H); <sup>13</sup>C-NMR (D<sub>2</sub>O) δ: 163.84, 134.60, 130.69, 129.59, 126.48, 125.17, 118.95, 47.60, 44.13, 39.47, 35.49, 23.99, 23.18; ESI-MS *m/z*: 312.2 (M+H-2HCl)<sup>+</sup>; Anal. calcd. for C<sub>18</sub>H<sub>23</sub>Cl<sub>2</sub>N<sub>3</sub>O<sub>2</sub>·0.5H<sub>2</sub>O: C, 56.26; H, 6.03; N, 10.93; found: C, 56.18; H, 5.76; N, 10.66.

*2-[5-(2-Aminoethylamino)pentyl]1H-benz-[de]isoquinoline-1,3(2H)-dione dihydrochloride (7c)*. Yield: 63.7%; m.p.: 233.8–235.1 °C; <sup>1</sup>H-NMR (D<sub>2</sub>O) δ: 7.42 (d, 2H, *J* = 8.0 Hz), 7.35 (d, 2H, *J* = 7.2 Hz), 7.02 (t, 2H, *J* = 7.6 Hz), 3.40–3.45 (m, 6H), 3.15 (t, 2H, *J* = 8.0 Hz), 1.73–1.77 (m, 2H), 1.36–1.38 (m, 4H); <sup>13</sup>C-NMR (D<sub>2</sub>O) δ: 163.89, 134.54, 130.67, 129.64, 126.46, 125.23, 119.06, 48.08, 44.12, 39.86, 35.48, 26.40, 25.10, 23.21; ESI-MS *m/z*: 326.2 (M+H-2HCl)<sup>+</sup>; Anal. calcd. for C<sub>19</sub>H<sub>25</sub>Cl<sub>2</sub>N<sub>3</sub>O<sub>2</sub>·0.3H<sub>2</sub>O: C, 56.52; H, 6.39; N, 10.41; N, 10.09; found: C, 56.70; H, 6.10; N, 10.13.

*2-[3-(3-Aminopropylamino)propyl]1H-benz-[de]isoquinoline-1,3(2H)-dione dihydrochloride (7d)*. Yield: 68.8%; m.p.: 233.1–234.2 °C; <sup>1</sup>H-NMR (D<sub>2</sub>O) δ: 7.60 (d, 2H, *J* = 8.0 Hz), 7.57 (d, 2H, *J* = 7.2 Hz), 7.18 (t, 2H, *J* = 7.6 Hz), 3.69 (t, 2H, *J* = 7.2 Hz), 3.11–3.24 (m, 6H), 2.14–2.16 (m, 2H), 1.89–1.93 (m, 2H); <sup>13</sup>C-NMR (D<sub>2</sub>O) δ: 164.20, 134.93, 130.98, 129.91, 126.68, 125.55, 119.18, 45.43, 44.72, 37.27, 36.61, 24.16, 23.79; ESI-MS *m/z*: 312.2 (M+H-2HCl)<sup>+</sup>; Anal. calcd. For C<sub>18</sub>H<sub>23</sub>Cl<sub>2</sub>N<sub>3</sub>O<sub>2</sub>·0.5H<sub>2</sub>O: C, 56.26; H, 6.03; N, 10.93; found: C, 56.29; H, 5.88; N, 10.69.

2-[5-(3-Aminopropylamino)pentyl]1H-benz-[de]isoquinoline-1,3(2H)-dione dihydrochloride (**7e**). Yield: 72.5%; m.p.: 238.6–239.8 °C;  $^1\text{H-NMR}$  ( $\text{D}_2\text{O}$ )  $\delta$ : 7.36 (d, 2H,  $J = 8.0$  Hz), 7.29 (d, 2H,  $J = 7.2$  Hz), 6.97 (t, 2H,  $J = 7.6$  Hz), 3.36 (t, 2H,  $J = 7.2$  Hz), 3.06–3.20 (m, 6H), 2.11–2.15 (m, 2H), 1.70–1.72 (m, 2H), 1.30–1.34 (m, 4H);  $^{13}\text{C-NMR}$  ( $\text{D}_2\text{O}$ )  $\delta$ : 163.78, 134.49, 130.60, 129.55, 126.41, 125.12, 118.95, 47.69, 44.51, 39.85, 36.61, 26.40, 25.07, 23.79, 23.26; ESI-MS  $m/z$ : 340.3 ( $\text{M}+\text{H}-2\text{HCl}$ ) $^+$ ; Anal. calcd. for  $\text{C}_{20}\text{H}_{27}\text{Cl}_2\text{N}_3\text{O}_2 \cdot 0.3\text{H}_2\text{O}$ : C, 57.50; H, 6.66; N, 10.06; found: C, 57.52; H, 6.30; N, 9.78.

2-[3-(4-Aminobutylamino)propyl]1H-benz-[de]isoquinoline-1,3(2H)-dione dihydrochloride (**7f**). Yield: 69.7%; m.p.: 235.1–235.9 °C;  $^1\text{H-NMR}$  ( $\text{D}_2\text{O}$ )  $\delta$ : 7.59 (d, 2H,  $J = 8.0$  Hz), 7.56 (d, 2H,  $J = 7.2$  Hz), 7.17 (t, 2H,  $J = 7.6$  Hz), 3.66 (t, 2H,  $J = 7.2$  Hz), 3.01–3.11 (m, 6H), 1.76–1.88 (m, 6H);  $^{13}\text{C-NMR}$  ( $\text{D}_2\text{O}$ )  $\delta$ : 164.00, 134.70, 130.75, 129.69, 126.44, 125.35, 119.97, 46.83, 44.99, 39.58, 37.03, 23.91, 23.72, 22.56; ESI-MS  $m/z$ : 326.2 ( $\text{M}+\text{H}-2\text{HCl}$ ) $^+$ ; Anal. calcd. for  $\text{C}_{19}\text{H}_{25}\text{Cl}_2\text{N}_3\text{O}_2 \cdot 0.5\text{H}_2\text{O}$ : C, 56.02; H, 6.43; N, 10.32; found: C, 56.29; H, 6.26; N, 10.69.

2-[4-(4-Aminobutylamino)butyl]1H-benz-[de]isoquinoline-1,3(2H)-dione dihydrochloride (**7g**). Yield: 78.6%; m.p.: 226.7–228.3 °C;  $^1\text{H-NMR}$  ( $\text{D}_2\text{O}$ )  $\delta$ : 7.75 (d, 2H,  $J = 8.0$  Hz), 7.72 (d, 2H,  $J = 7.2$  Hz), 7.31 (t, 2H,  $J = 7.6$  Hz), 3.70 (t, 2H,  $J = 7.2$  Hz), 3.11–3.20 (m, 6H), 1.78–1.86 (m, 6H), 1.60–1.63 (m, 2H);  $^{13}\text{C-NMR}$  ( $\text{D}_2\text{O}$ )  $\delta$ : 163.92, 134.66, 130.74, 129.68, 126.52, 125.27, 119.04, 47.08, 46.92, 39.52, 38.86, 24.10, 23.97, 23.19, 22.83; ESI-MS  $m/z$ : 340.3 ( $\text{M}+\text{H}-2\text{HCl}$ ) $^+$ ; Anal. calcd. for  $\text{C}_{20}\text{H}_{27}\text{Cl}_2\text{N}_3\text{O}_2$ : C, 58.25; H, 6.60; N, 10.19; found: C, 56.14, H, 5.88, N, 10.69.

2-[5-(4-Aminobutylamino)pentyl]1H-benz-[de]isoquinoline-1,3(2H)-dione dihydrochloride (**7h**). Yield: 59.6%; m.p.: 236.6–237.9 °C;  $^1\text{H-NMR}$  ( $\text{D}_2\text{O}$ )  $\delta$ : 7.29 (d, 2H,  $J = 8.0$  Hz), 7.21 (d, 2H,  $J = 7.2$  Hz), 6.90 (t, 2H,  $J = 7.6$  Hz), 3.30 (t, 2H,  $J = 7.2$  Hz), 3.03–3.13 (m, 6H), 1.79–1.81 (m, 4H), 1.68–1.71 (m, 2H), 1.27–1.32 (m, 4H);  $^{13}\text{C-NMR}$  ( $\text{D}_2\text{O}$ )  $\delta$ : 163.66, 134.44, 130.54, 129.48, 126.38, 125.02, 118.87, 47.54, 46.92, 39.84, 38.88, 26.41, 25.08, 24.00, 23.31, 22.83; ESI-MS  $m/z$ : 354.3 ( $\text{M}+\text{H}-2\text{HCl}$ ) $^+$ ; Anal. calcd. for  $\text{C}_{21}\text{H}_{29}\text{Cl}_2\text{N}_3\text{O}_2 \cdot 0.7\text{H}_2\text{O}$ : C, 57.46; H, 6.98; N, 9.57; found: C, 57.39; H, 6.92; N, 9.37.

2-[3-(6-Aminohexylamino)propyl]1H-benz-[de]isoquinoline-1,3(2H)-dione dihydrochloride (**7i**). Yield: 60.7%; m.p.: 230.6–231.8 °C;  $^1\text{H-NMR}$  ( $\text{D}_2\text{O}$ )  $\delta$ : 7.62 (d, 2H,  $J = 8.0$  Hz), 7.58 (d, 2H,  $J = 7.2$  Hz), 7.19 (t, 2H,  $J = 7.6$  Hz), 3.68 (t, 2H,  $J = 7.2$  Hz), 3.00–3.10 (m, 6H), 1.88–1.91 (m, 2H), 1.69–1.76 (m, 4H), 1.43–1.47 (m, 4H);  $^{13}\text{C-NMR}$  ( $\text{D}_2\text{O}$ )  $\delta$ : 164.22, 134.94, 130.98, 129.94, 126.70, 125.50, 119.22, 47.65, 45.08, 39.38, 37.31, 26.50, 25.36, 25.27, 25.16, 24.16; ESI-MS  $m/z$ : 354.3 ( $\text{M}+\text{H}-2\text{HCl}$ ) $^+$ ; Anal. calcd. for  $\text{C}_{21}\text{H}_{29}\text{Cl}_2\text{N}_3\text{O}_2 \cdot 1.0\text{H}_2\text{O}$ : 56.76; H, 7.03; N, 9.46; found: C, 56.85; H, 6.71; N, 9.85.

2-[4-(6-Aminohexylamino)butyl]1H-benz-[de]isoquinoline-1,3(2H)-dione dihydrochloride (**7j**). Yield: 64.5%; m.p.: 235.9–237.2 °C;  $^1\text{H-NMR}$  ( $\text{D}_2\text{O}$ )  $\delta$ : 7.57 (d, 2H,  $J = 8.0$  Hz), 7.52 (d, 2H,  $J = 7.2$  Hz), 7.15 (t, 2H,  $J = 7.6$  Hz), 3.54 (t, 2H,  $J = 7.2$  Hz), 2.98–3.09 (m, 6H), 1.66–1.71 (m, 6H), 1.41–1.48 (m, 6H);  $^{13}\text{C-NMR}$  ( $\text{D}_2\text{O}$ )  $\delta$ : 164.11, 134.76, 130.86, 129.87, 126.60, 125.51, 119.27, 47.48, 46.96, 39.53, 39.33, 26.48, 25.37, 25.25, 25.14, 24.15, 23.16; ESI-MS  $m/z$ : 368.3 ( $\text{M}+\text{H}-2\text{HCl}$ ) $^+$ ; Anal. calcd. for  $\text{C}_{22}\text{H}_{31}\text{Cl}_2\text{N}_3\text{O}_2 \cdot 1.0\text{H}_2\text{O}$ : C, 57.64; H, 7.26; N, 9.17; found: C, 57.36; H, 7.13; N, 8.85.

2-[5-(6-Aminohexylamino)pentyl]1H-benz-[de]isoquinoline-1,3(2H)-dione dihydrochloride (**7k**). Yield: 77.2%; m.p.: 229.1–230.3 °C; <sup>1</sup>H-NMR (D<sub>2</sub>O) δ: 7.34 (d, 2H, *J* = 8.0 Hz), 7.30 (d, 2H, *J* = 7.2 Hz), 6.96 (t, 2H, *J* = 7.6 Hz), 3.33 (t, 2H, *J* = 7.2 Hz), 2.91–2.99 (m, 6H), 1.61–1.63 (m, 6H), 1.27–1.37c (m, 8H); <sup>13</sup>C-NMR (D<sub>2</sub>O) δ: 163.96, 134.59, 130.71, 129.75, 126.52, 125.35, 119.19, 47.51, 47.39, 39.92, 39.40, 26.52, 26.49, 25.38, 25.31, 25.17, 25.11, 23.34; ESI-MS *m/z*: 382.3 (M+H-2HCl)<sup>+</sup>; Anal. calcd. for C<sub>23</sub>H<sub>33</sub>Cl<sub>2</sub>N<sub>3</sub>O<sub>2</sub>·0.9H<sub>2</sub>O: C, 58.69; H, 7.45; N, 8.93; found: C, 58.36; H, 7.06; N, 8.60.

### 3.3. Biological Materials and Methods

All chemicals used in bioassay were purchased from Sigma (Beijing, China), unless otherwise indicated. RPMI1640 and fetal calf serum (FCS) were purchased from Gibco (Shanghai, China). Stock solution (10 mM) was prepared in DMSO and diluted to various concentrations with serum-free culture medium.

#### 3.3.1. Cell Culture

Cell lines, K562, 7721 and HCT116 were obtained from American Type Culture Collection (ATCC, Shanghai, China). Cells were cultured in RPMI1640, supplemented with 10% heat-inactivated fetal calf serum (FCS), antibiotics (penicillin, 100 units/mL; streptomycin sulfate, 100 µg/mL) at 37 °C, in an atmosphere of 95% air and 5% CO<sub>2</sub> under humidified conditions. Aminoguanidine (1 mM) was added as an inhibitor of amine oxidase derived from FCS and had no effect on the various parameters of the cell measured in this study.

#### 3.3.2. MTT Assay

Chemosensitivity was assessed using the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay. Briefly, exponentially growing K562 cells were seeded into 96-well plates at 4000 cells/well and treated with indicated concentrations of samples for 48 h, and then 10 µL of MTT (10 mg/mL) was added. After incubation for 4 h at 37 °C, the purple formazan crystals (*i.e.*, a reduced form of MTT) generated from viable cells were dissolved by adding 100 µL 10% sodium dodecyl sulphate (SDS) in each well. The absorbance of each well was then read at 570 nm.

In addition, exponentially growing HCT116 or 7721 cells were seeded into 96-well plates at 5000 cells/well and allowed to attach overnight, and then 100 µL of MTT (1 mg/mL) was added. After incubation for 4 h at 37 °C, the MTT solution was removed and the remaining formazan crystals were dissolved with 150 µL DMSO in each well. The absorbance of each well was then read at 570 nm.

### 3.4. Cell Cycle Analysis and Apoptosis

#### 3.4.1. Cell Cycle Analysis

Exponentially growing HepG2 cells were seeded in 96 well plates (4 × 10<sup>5</sup> cells/well), cultured for 24 h and then treated with different concentrations of compound **7f** for 48 h. After incubation for 48 h, cells were washed twice with ice-cold 10% PBS, fixed and permeabilized with ice-cold 70% ethanol at −20 °C overnight. The cells were treated with 50 µg/mL RNase A at room temperature



for 30 min after washed with ice-cold PBS, and finally stained with 50 µg/mL propidium iodide (PI) in the dark at 4 °C for 30 min. The distribution of cell cycle phases with different DNA contents was read in image analysis system of high content screening living cells.

### 3.4.2. Apoptosis

Exponentially growing HepG2 cells were seeded in 96 well plates ( $6 \times 10^4$  cells/well) and treated with indicated concentrations of **7f**. After incubated for 48 h, cells were washed three times with PBS, and finally stained with with 10 µM Hoechst 33342 in the dark at 37 °C for 20 min. They were detected by laser scanning confocal microscope and the apoptotic cells were count by selecting 200 cells randomly.

## 3.5. Spectroscopy Measurement

### 3.5.1. Apparatus

UV–vis absorption spectra were measured on a Unicam UV 500 spectrophotometer (Beijing, China) using a 1.0 cm cell. Fluorescence measurements were performed with a Cary Eclipse spectrofluorimeter (Shanghai, China).

### 3.5.2. UV–Vis Measurements

2 mL solution of compound **7f** ( $2.00 \times 10^{-4}$  mol·L<sup>-1</sup> in Tris-HCl (pH = 7.4) was mixed with 0.0, 0.10, 0.20, 0.30, 0.60, 0.90 1.20, 1.50, 1.80, 2.10, 2.40, 2.70 and 3.0 mL of herring sperm DNA ( $2.284 \times 10^{-4}$  mol·L<sup>-1</sup>) respectively. The mixture was diluted to 5 mL with Tris-HCl (pH = 7.4). Thus, samples were prepared in the concentration of DNA at 0.0, 4.56, 9.13, 13.69, 27.4, 41.08, 54.77, 68.46, 82.15, 95.84, 109.54, 123.23 and  $136.92 \times 10^{-6}$  mol·L<sup>-1</sup>. One contained only compound **7f** ( $80 \times 10^{-6}$  mol·L<sup>-1</sup>) as control, the others contained different concentration of DNA but had the same concentration of compound **7f**. All the above solution was shaken for 30 min. at room temperature.

### 3.5.3. Fluorescence Measurement

#### 3.5.3.1. Interaction of Compound **7f** with DNA

Preparation of sample is the same as that of UV-Vis measurement. Fluorescence wavelengths and intensity areas of samples were measured at following conditions: EX = 345 nm, EM = 355~600 nm temperature: 298, 303 and 310 K.

#### 3.5.3.2. Interaction of Compound **7f** with DNA-EB Complex

Solution of herring sperm DNA (0.3 mL,  $2.284 \times 10^{-5}$  mol·L<sup>-1</sup> in Tris-HCl (pH = 7.4) and 0.5 mL EB ( $1.57 \times 10^{-5}$  mol·L<sup>-1</sup>) was mixed with 0.0, 0.10, 0.20, 0.30, 0.60, 0.90 1.20, 1.50, 1.80, 2.10, 2.40, 2.70 and 3.00 mL of compound **7f** ( $2.0 \times 10^{-4}$  mol·L<sup>-1</sup>) respectively. The mixture was diluted to 5 mL with Tris-HCl (pH = 7.4). Thus, samples were prepared in the concentration of compound **7f** at 0.0, 4.0, 8.0, 12.0, 24.0, 36.0, 48.0, 60.0, 72.0, 84.0, 96.0 and 108.0 and 120.0  $\times 10^{-6}$  mol·L<sup>-1</sup>. One contained

only DNA ( $13.7 \times 10^{-6} \text{ mol}\cdot\text{L}^{-1}$ ) and EB ( $15.7 \times 10^{-6} \text{ mol}\cdot\text{L}^{-1}$ ) as control, the others contained different concentration of compound **7f** but had the same concentration of DNA and EB. All the above solution was shaken for 30 min. at room temperature. Fluorescence wavelengths and intensity areas of samples were measured at following conditions: EX = 510 nm, EM = 520~800 nm; temperature: 298, 303 and 310 K.

### 3.5.3.3. Iodide Quenching

Solution of compound **7f** (0.5 mL,  $2.00 \times 10^{-4} \text{ mol/L}$  and herring sperm DNA 0.5 mL ( $22.84 \times 10^{-4} \text{ mol/L}$ ) in Tris-HCl (pH = 7.4) were mixed with 0.0, 0.10, 0.20, 0.30, 0.60, 0.90 1.20, 1.50, 1.80, 2.10, 2.40, 2.70 and 3.00 of KI ( $2.0 \times 10^{-2} \text{ mol}\cdot\text{L}^{-1}$ ) respectively. Meanwhile, 0.5 mL solution of compound **7f** ( $2.00 \times 10^{-4} \text{ mol/L}$ ) was only mixed with 0.0, 0.10, 0.20, 0.30, 0.60, 0.90 1.20, 1.50, 1.80, 2.10, 2.40, 2.70 and 3.00 mL of KI ( $2.0 \times 10^{-2} \text{ mol}\cdot\text{L}^{-1}$ ) respectively. The mixture was diluted to 5 mL with Tris-HCl (pH = 7.4). Thus, samples were prepared in the concentration of KI at 0.0, 400, 800, 1,200, 2,400, 3,600, 4,800, 6,000, 7,200, 8,400, 9,600, 10,800, 12,000  $\times 10^{-6} \text{ mol}\cdot\text{L}^{-1}$ . One kind of samples contained compound **7f** ( $20 \times 10^{-6} \text{ mol}\cdot\text{L}^{-1}$ ), DNA ( $22.82 \times 10^{-6} \text{ mol}\cdot\text{L}^{-1}$ ) and different concentrations of KI. The other groups of samples contained only different concentration of KI and the same concentration of compound **7f** as the control. All the above solution was shaken for 30 min. at room temperature. Fluorescence wavelengths and intensity areas of samples were measured at following conditions: EX = 345 nm, EM = 355~600 nm.

### 3.5.3.4. Effect of Ionic Intensity on The Interaction between Compound **7f** and DNA

Solution of compound **7f** (1.0 mL,  $2.00 \times 10^{-4} \text{ mol}\cdot\text{L}^{-1}$ ) and herring sperm DNA 1.0 mL ( $2.284 \times 10^{-4} \text{ mol}\cdot\text{L}^{-1}$ ) in Tris-HCl (pH = 7.4) were mixed with 0.0, 0.10, 0.20, 0.30, 0.60, 0.90 1.20, 1.50, 1.80, 2.10, 2.40, 2.70 and 3.00 mL of NaCl ( $4.0 \times 10^{-2} \text{ mol}\cdot\text{L}^{-1}$ ) respectively. The mixture was diluted to 5 mL with Tris-HCl (pH = 7.4). Thus, samples were prepared with the concentration of NaCl at 0.0, 800, 1,600, 2,400, 4,800, 7,200, 9,600, 12,000, 14,400, 16,800, 19,200, 21,600 and 24,000  $\times 10^{-6} \text{ mol}\cdot\text{L}^{-1}$ . The above mixtures were divided into two groups, in which one contained only compound **7f** ( $40 \times 10^{-6} \text{ mol}\cdot\text{L}^{-1}$ ) and DNA ( $45.68 \times 10^{-6} \text{ mol}\cdot\text{L}^{-1}$ ) as control, the others contained different concentration of NaCl with the same concentration of compound **7f** and DNA. All the above solution was shaken for 30 min. at room temperature. Fluorescence wavelengths and intensity areas of samples were measured at following conditions: EX = 345 nm, EM = 355~600 nm.

## 4. Conclusions

A series of naphthalimide-diamine conjugates were synthesized and their *in vitro* antitumor activities were evaluated. Compound **7f** was found to have potent antitumor activity and good cell selectivity with amonafide as a control. Moreover, compound **7f** could arrest HepG2 cells in the G2/M phase and induce HepG2 cells apoptosis. The interaction of compound **7f** with DNA was first studied by spectroscopic methods. The binding of compound **7f** to DNA resulted in a series of changes in the spectral characteristics. The absorption spectra of compound **7f** with added DNA showed a hypochromic effect and the fluorescence emission of DNA-EB was efficiently quenched by compound

**7f.** These observed spectral data and the iodide quenching effect suggested that compound **7f** interacts with DNA through an intercalative mode. Further fluorescent assays at different temperatures disclosed that the quenching mechanism of both compound **7f** with DNA and compound **7f** with DNA-EB was a static type. Meanwhile, the binding constant, thermodynamic parameters obtained from the same calorimetric fluorescent tests and the effect of NaCl on compound **7f**-DNA interaction suggested that the binding process was driven by hydrogen bonding and van der Waals forces.

### Acknowledgments

This work was supported by the China Postdoctoral Science Foundation Funded Project (No. 20110490991), the Henan Natural Science Foundation (Nos. 112300410181, 134200510009), the Henan Programs for Science and Technology Development (No. 132102310026) and the Natural Science Foundation of Education (No. 2011B3500001, 14A350004).

### Author Contributions

Z.-Y.T. and C.-J.W. designed the research; all authors performed the experimental work; Z.-Y.T., Q.L. and C.-J.W. wrote the manuscript; All authors discussed, edited and approved the final version.

### Conflicts of Interest

The authors declare no conflict of interest.

### References

1. Braña, M.F.; Sanz, A.M.; Castellano, J.M.; Roldán, C.M.; Roldán, C. Synthesis and cytostatic activity of benz[de]isoquinoline-1,3-diones, structure-activity relationships. *Eur. J. Med. Chem. Chim. Ther.* **1981**, *16*, 207–212.
2. Sami, S.M.; Dorr, R.T.; Alberts, D.S.; Remers, W.A. 2-Substituted 1,2-dihydro-3H-dibenz[de,h]isoquinoline-1,3-diones. A new class of antitumor agent. *J. Med. Chem.* **1993**, *36*, 765–770.
3. Baily, C.; Braña, M.F.; Waring, M.J. Sequence-selective intercalation of antitumor bisnaphthalimides into DNA. Evident for an approach via the major groove. *Eur. J. Biochem.* **1996**, *240*, 195–208.
4. Braña, M.F.; Castellano, J.M.; Morán, M.; Pérez de Vega, M.J.; Perron, D.; Conlon, D.; Bousquet, P.F.; Romerdahl, C.A.; Robinson, S.P. Bis naphthalimides 3: Synthesis and antitumor activity of N,N'-bis[2-(1,8-naphthalimido)-ethyl]alkanedimines. *Anticancer Drug Des.* **1996**, *11*, 297–309.
5. Rosell, R.; Carles, J.; Abad, A.; Ribelles, N.; Barnadas, A.; Benavides, A.; Martin, M. Phase I study of mitonafide in 120 h continuous infusion in non-small cell lung cancer. *Investig. New Drugs* **1992**, *10*, 171–175.
6. Costanza, M.E.; Berry, D.; Henderson, I.C.; Ratain, M.J.; Wu, K.; Shapiro, C.; Duggan, D.; Kalra, J.; Berkowitz, I.; Lyss, A.P. Amonafide: An active agent in the treatment of previously untreated advanced breast cancer—a cancer and leukemia group B study (CALGB 8642). *Clin. Cancer Res.* **1995**, *1*, 699–704.

7. Sami, S.M.; Dorr, R.T.; Sólyom, A.M.; Alberts, D.S.; Remers, W.A. Amino-substituted 2-[2'-(dimethyl-amino)ethyl 1,2-dihydro-3H-dibenz[de,h] isoquinoline-1,3-diones. Synthesis, antitumor activity and quantitative structure- activity relation-ship. *J. Med. Chem.* **1995**, *38*, 983–993.
8. Thompson, J.; Pratt, C.B.; Stewart, C.F.; Avery, L.; Bowman, L.; Zamboni, W.C.; Pappo, A. Phase I study of DMP 840 in pediatric patients with refractory solid tumors. *Investig. New Drugs* **1998**, *16*, 45–49.
9. Nitiss, J.L.; Zhou, J.; Rose, A.; Hsiung, Y.; Gale, K.C.; Osheroff, N. The bis(naphthalimide) DMP-840 causes cytotoxicity by its action against eukaryotic topoisomerase II. *Biochemistry* **1998**, *37*, 3078–3085.
10. Bousquet, P.F.; Braña, M.F.; Conlon, D.; Fitzgerald, K.M.; Perron, D.; Cocchiaro, C.; Miller, R.; Moran, M.; George, J.; Qian, X.D.; *et al.* Preclinical evaluation of LU 79553: A novel bisnaphthalimide with potent antitumor activity. *Cancer Res.* **1995**, *55*, 1176–1180.
11. Diaz-Rubio, E.; Martin, M.; López-Vega, J.M.; Casado, A.; Benavides, A. Phase I study of mitonafide with a 3-day administration schedule: Early interruption due to severe central nervous system toxicity. *Investig. New Drugs* **1994**, *12*, 277–281.
12. Braña, M.F.; Domínguez, G.; Sáez, B.; Romerdahl, C.; Robinson, S.; Barlozzari, T. Synthesis and antitumour activity of new dendritic polyamines-(imide-DNA-intercalator) conjugates: Potent Lck inhibitors. *Eur. J. Med. Chem.* **2002**, *37*, 541–551.
13. Braña, M.F.; Cacho, M.; Ramos, A.; Dominguez, M.T.; Pozuelo, J.M.; Abradelo, C.; Rey-Stolle, M.F.; Yuste, M.; Carrasco, C.; Bailly, C. Synthesis, biological evaluation and DNA binding properties of novel mono and bisnaphthalimides. *Org. Biomol. Chem.* **2003**, *1*, 648–654.
14. Van Quaquebeke, E.; Mahieu, T.; Dumont, P.; Dewelle, J.; Ribaucour, F.; Simon, G.; Sauvage, S.; Gaussin, J.F.; Tuti, J.; El Yazidi, M.; *et al.* 2,2,2-Trichloro-N-({2-[2-(dimethylamino)ethyl]-1,3-dioxo-2,3-dihydro-1H-benzo[de] isoquinolin-5-yl} car-bamoyl) acetamide (UN BS3157), a novel nonhematotoxic naphthalimide derivative with potent antitumor activity. *J. Med. Chem.* **2007**, *50*, 4122–4134.
15. Chen, Z.; Liang, X.; Zhang, H.Y.; Xie, H.; Liu, J.W.; Xu, Y.F.; Zhu, W.P.; Wang, Y.; Wang, X.; Tan, S.Y.; *et al.* A new class of naphthalimide-based antitumor agents that inhibit topoisomerase II and induce lysosomal membrane permeabilization and apoptosis. *J. Med. Chem.* **2010**, *53*, 2589–2600.
16. Sk, U.H.; Prakasha Gowda, A.S.; Crampsie, M.A.; Yun, J.K.; Spratt, T.E.; Amin, S.; Sharma, A.K. Development of novel naphthalimide derivatives and their evaluation as potential melanoma therapeutics. *Eur. J. Med. Chem.* **2011**, *46*, 3331–3338.
17. Roy, S.S.; Ghosh, P.; Sk, U.H.; Chakraborty, P.; Biswas, J.; Mandal, S.; Bhattacharjee, A.; Bhattacharya, S. Naphthalimide based novel organo-selenocyanates: Finding less toxic forms of selenium that would retain protective efficacy. *Bioorg. Med. Chem. Lett.* **2010**, *21*, 6951–6955.
18. Filosa, R.; Peduto, A.; Micco, S.D.; Caprariis, P.D.; Festa, M.; Petrella, A.; Capranico, G.; Bifulco, G. Molecular modeling studies, synthesis and biological activity of a series of novel bisnaphthalimides and their development as new DNA topoisomerase II inhibitors. *Bioorg. Med. Chem.* **2009**, *17*, 13–24.

19. Kamal, A.; Ramu, R.; Tekumalla, V.; Khanna, G.B.; Barkume, M.S.; Juvekar, A.S.; Zingde, S.M. Remarkable DNA binding affinity and potential anticancer activity of pyrrolo[2,1-c][1,4] benzodiazepine-naphthalimide conjugates linked through piperazine side-armed alkane spacers. *Bioorg. Med. Chem.* **2008**, *16*, 7218–7224.
20. Machado, K.E.; Oliveira, K.N.; Santos-Bubniak, L.; Licínio, M.A.; Nunes, R.J.; Santos-Silva, M.C. Evaluation of apoptotic effect of cyclic imide derivatives on murine B16F10 melanoma cells. *Bioorg. Med. Chem.* **2011**, *19*, 6285–6291.
21. Minarini, A.; Milelli, A.; Tumiatti, V.; Ferruzzi, L.; Marton, M.R.; Turrini, E.; Hrelia, P.; Fimognari, C. Design, synthesis and biological evaluation of new naphthalene diimides bearing isothiocyanate functionality. *Eur. J. Med. Chem.* **2012**, *48*, 124–131.
22. Barron, G.A.; Bermano, G.; Gordon, A.; Kong Thoo Lin, P. Synthesis, cytotoxicity and DNA-binding of novel bisnaphthalimidopropyl derivatives in breast cancer MDA-MB-231 cells. *Eur. J. Med. Chem.* **2010**, *45*, 1430–1437.
23. Banerjee, S.; Kitchen, J.A.; Bright, S.A.; O'Brien, J.E.; Williams, D.C.; Kelly, J.M.; Gunnlaugsson, T. Synthesis, spectroscopic and biological studies of a fluorescent Pt(II) (terpy) based 1,8-naphthalimide conjugate as a DNA targeting agent. *Chem. Commun.* **2013**, *49*, 8522–8524.
24. Banerjee, S.; Kitchen, J.A.; Gunnlaugsson, T.; Kelly, J.M. The effect of the 4-amino functionality on the photophysical and DNA binding properties of alkyl-pyridinium derived 1,8-naphthalimides. *Org. Biomol. Chem.* **2013**, *11*, 5642–5655.
25. Banerjee, S.; Kitchen, J.A.; Gunnlaugsson, T.; Kelly, J.M. Synthesis and photophysical evaluation of a pyridinium 4-amino-1,8-naphthalimide derivative that upon intercalation displays preference for AT-rich double-stranded DNA. *Org. Biomol. Chem.* **2012**, *10*, 3033–3043.
26. Wang, C.J.; Delcros, J.G.; Cannon, L.; Konate, F.; Carias, H.; Bigger staff, J.; Gardner, R.A.; Phanstiel, O., IV. Defining the molecular requirements for the selective delivery of polyamine conjugates into cells containing active polyamine transporters. *J. Med. Chem.* **2003**, *46*, 5129–5138.
27. Casero, R.A., Jr.; Marton, L.J. Targeting polyamine metabolism and function in cancer and other hyperproliferative diseases. *Nat. Rev. Drug Discov.* **2007**, *6*, 373–390.
28. Tian, Z.Y.; Xie, S.Q.; Du, Y.W.; Ma, Y.F.; Zhao, J.; Gao, W.Y.; Wang, C.J. Synthesis, cytotoxicity and apoptosis of naphthalimide polyamine conjugates as antitumor agents. *Eur. J. Med. Chem.* **2009**, *44*, 393–399.
29. Tian, Z.Y.; Xie, S.Q.; Mei, Z.H.; Zhao, J.; Gao, W.Y.; Wang, C.J. Conjugation of substituted naphthalimides to polyamines as cytotoxic agents targeting the Akt/mTOR signal pathway. *Org. Biomol. Chem.* **2009**, *7*, 4551–4560.
30. Yang, L.H.; Li, W.; Tian, Z.Y.; Zhao, J.; Wang, C.J. Mononaphthalimide spermidine conjugate induces cell proliferation inhibition and apoptosis in HeLa cells. *Toxicol. In Vitro* **2011**, *52*, 882–889.
31. Pasternack, R.F.; Gibbs, E.J.; Villafranca, J.J. Interactions of water soluble porphyrins with Z-poly (dG-dC). *Biochemistry* **1983**, *22*, 2406–2414.
32. Xie, S.Q.; Cheng, P.F.; Liu, G.C.; Ma, Y.F.; Zhao, J.; Chehtane, M.; Khaled, A.R.; Phanstiel, O., IV; Wang, C.J. Synthesis and bioevaluation of N-(arylalkyl)-homospermidine conjugates. *Bioorg. Med. Chem. Lett.* **2007**, *17*, 4471–4475.

33. Wang, C.J.; Delcros, J.G.; Bigger staff, J.; Phanstiel, O., IV. Molecular requirements for targeting the polyamine transport system. Synthesis and biological evaluation of polyamine-anthracene conjugates. *J. Med. Chem.* **2003**, *46*, 2672–2682.
34. Mosmann, T. Rapid colorimetric assay for cellular growth and survival: Application to proliferation and cytotoxicity assays. *J. Immunol. Meth.* **1983**, *65*, 55–63.
35. Tian, Z.Y.; Su, L.P.; Xie, S.Q.; Zhao, J.; Wang, C.J. Study on synthesis, biological activity and fluorescence spectroscopy of naphthalimide-polyamine conjugates. *Chin. J. Org. Chem.* **2013**, *33*, 1514–1521.
36. Zhu, H.; Huang, M.; Yang, F.; Chen, Y.; Miao, Z.H.; Qian, X.H.; Xu, Y.F.; Qin, Y.X.; Luo, H.B.; Shen, X.; *et al.* Ganoderic acid DM, a natural triterpenoid, induces DNA damage, G1 cell cycle arrest and apoptosis in human breast cancer cells. *Mol. Cancer Ther.* **2007**, *6*, 484–495.
37. Long, E.C.; Barton, J.K. On demonstrating DNA intercalation. *Acc. Chem. Res.* **1990**, *23*, 271–273.
38. Tysoe, S.A.; Morgan, R.J.; Baker, A.D.; Streckas, T.C. Spectroscopic investigation of differential binding modes of  $\Delta$ - and  $\Lambda$ -Ru(bpy)<sub>2</sub>(ppz)<sup>2+</sup> with calf thymus DNA. *J. Phys. Chem.* **1993**, *97*, 1707–1711.
39. Li, N.; Ma, Y.; Yang, C.; Gong, L.P.; Yang, X.H. Interaction of anticancer drug mitoxantrone with DNA analyzed by electrochemical and spectroscopic methods. *Biophys. Chem.* **2005**, *116*, 199–205.
40. Sirajuddin, M.; Ali, S.; Badshah, A. Drug-DNA interactions and their study by UV-Visible, fluorescence spectroscopies and cyclic voltametry. *J. Photoch. Photobio. B: Biol.* **2013**, *124*, 1–19.
41. Sahoo, B.K.; Ghosh, K.S.; Bera, R.; Dasgupta, S. Studies on the interaction of diacetylcurcumin with calf thymus-DNA. *Chem. Phys.* **2008**, *351*, 163–169.
42. Ghaderi, M.; Bathaie, S.Z.; Saboury, A.A.; Sharghi, H.; Tangestaninejad, S. Interaction of an Fe derivative of TMAP (Fe (TMAP) OAc) with DNA in comparison with free-base TMAP. *Int. J. Biol. Macromol.* **2007**, *41*, 173–179.
43. Hassanien, M.M.; Abou-EI-Sherbini, K.S. Synthesis and characterisation of morin-functionalised silica gel for the enrichment of some precious metal ions. *Talanta* **2006**, *68*, 1550–1559.
44. Selim, M.; Chowdhury, S.R.; Mukherjea, K.K. DNA binding and nuclease activity of a one-dimensional heterometallic nitrosyl complex. *Int. J. Biol. Macromol.* **2007**, *41*, 579–583.
45. Lin, H.; Lan, J.F.; Guan, M.; Sheng, F.L.; Zhang, H.X. Spectroscopic investigation of interaction between mangiferin and bovine serum albumin. *Spectrochim. Acta. A. Mol. Biomol. Spectrosc.* **2009**, *73*, 936–941.
46. Hu, Y.J.; Liu, Y.; Zhao, R.M.; Dong, J.X.; Qu, S.S. Spectroscopic studies on the interaction between methylene blue and bovine serum albumin. *J. Photochem. Photobiol. A: Chem.* **2006**, *179*, 324–329.
47. Gelamo, E.L.; Silva, C.H.; Imasato, H.; Tabak, M. Interaction of bovine (BSA) and human (HSA) serum albumins with ionic surfactants: Spectroscopy and modeling. *Biochim. Biophys. Acta* **2002**, *1594*, 84–99.

48. Sun, Y.; Wei, S.; Yin, C.; Liu, L.S.; Hu, C.M.; Zhao, Y.Y.; Ye, Y.X.; Hu, X.Y.; Fan, J. Synthesis and spectroscopic characterization of 4-butoxyethoxy-N-octadecyl-1,8-naphthalimide as a new fluorescent probe for the determination of proteins. *Bioorg. Med. Chem. Lett.* **2011**, *21*, 3798–3804.
49. Lakowicz, J.R. *Principles of Fluorescence Spectroscopy*; Plenum Press: New York, NY, USA, 1999; pp. 237–265.
50. Lakowicz, J.R. *Principles of Fluorescence Spectroscopy*; Springer: New York, NY, USA, 2006; pp. 97–155.
51. Hu, Y.J.; Liu, Y.; Zhang, L.X.; Zhao, R.M.; Qu, S.S. Study of interaction between colchicines and bovine serum albumin by fluorescence quenching method. *J. Mol. Struct.* **2005**, *750*, 174–178.
52. Feng, Q.; Li, N.Q.; Jiang, Y.Y. Electrochemical studies of porphyrin interacting with DNA and determination of DNA. *Anal. Chim. Acta* **1997**, *344*, 97–104.
53. Hu, Y.J.; Liu, Y.; Xiao, X.H. Investigation of the interaction between Berberine and human serum albumin. *Biomacromolecules* **2009**, *10*, 517–521.
54. Ross, P.D.; Subramanian, S. Thermodynamics of protein association reactions: Forces contributing to stability. *Biochemistry* **1981**, *20*, 3096–3102.
55. Takenaka, S.; Ihara, T.; Takagi, M. Bis-9-acridinyl derivative containing a viologen linker chain: Electrochemically active intercalator for reversible labelling of DNA. *J. Chem. Soc. Chem. Commun.* **1990**, *23*, 1485–1487.
56. Qiao, C.Y.; Bi, S.Y.; Sun, Y.; Song, D.Q.; Zhang, H.Q.; Zhou, W.H. Study of interactions of anthraquinones with DNA using ethidium bromide as a fluorescence probe. *Spectrochim. Acta A: Mol. Biomol. Spectrosc.* **2008**, *70*, 136–143.
57. Zhang, G.W.; Hu, X.; Zhao, N.; Li, W.B.; He, L. Studies on the interaction of aminocarb with calf thymus DNA by spectroscopic methods. *Pestic. Biochem. Phys.* **2010**, *98*, 206–212.

*Sample Availability:* Samples of the compounds are available from the authors.

© 2014 by the authors; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/3.0/>).