

## Research Article

# Harnessing *Mucor* spp. for Xylanase Production: Statistical Optimization in Submerged Fermentation Using Agro-Industrial Wastes

Amal A. Al Mousa <sup>1</sup>, Nageh F. Abo-Dahab <sup>2</sup>, Abdallah M. A. Hassane <sup>2</sup>,  
Abd El-Rahman F. Gomaa <sup>2</sup>, Jana A. Aljuriss<sup>3</sup>, and Noura D. Dahmash<sup>1</sup>

<sup>1</sup>Department of Botany and Microbiology, College of Science, King Saud University, Riyadh, P. BOX 145111, ZIP, 4545, Saudi Arabia

<sup>2</sup>Botany and Microbiology Department, Faculty of Science, Al-Azhar University, Assiut 71524, Egypt

<sup>3</sup>King Saud University, Vice Rectorate for Graduate Studies and Scientific Research, Deanship of Scientific Research, Research Assistant Internship Program, Saudi Arabia

Correspondence should be addressed to Amal A. Al Mousa; aalmosa@ksu.edu.sa

Received 25 February 2022; Revised 28 March 2022; Accepted 1 April 2022; Published 20 April 2022

Academic Editor: Walid Saibi

Copyright © 2022 Amal A. Al Mousa et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Xylan is the primary hemicellulosic polymer found in lignocellulosic agricultural wastes and can be degraded by xylanase. In the current research, *Mucor circinelloides* and *M. hiemalis* were tested for their ability to produce xylanase from tangerine peel by submerged fermentation. Experiments on five variables were designed with Box–Behnken design and response surface methodology. Analysis of variance was exercised, the xylanase output was demonstrated with a mathematical equation as a function of the five factors, and the quixotic states for xylanase biosynthesis was secured. In addition, xylanase was partially purified, characterized, and immobilized on calcium alginate beads. The optimum parameters for xylanase production by *M. circinelloides* and *M. hiemalis* were consisted of incubation temperature (30 and 20°C), pH value (9 and 7) incubation period (9 and 9 days), inoculum size (3 and 3 mL), and substrate concentration (3 and 3 g/100 mL), respectively. *M. circinelloides* and *M. hiemalis* demonstrated the highest xylanase activities after RSM optimization, with 42.23 and 35.88 U/mL, respectively. The influence of single, interchange, and quadratic factors on xylanase output was investigated using nonlinear regression equations with significant  $R^2$  and  $p$  values. The partial purification of *M. circinelloides* and *M. hiemalis* xylanase yielded 1.69- and 1.97-fold purification, and 30.74 and 31.34% recovery with 292.08 and 240.15 U/mg specific activity, respectively. Partially purified xylanase from *M. circinelloides* and *M. hiemalis* demonstrated the highest activity at neutral pH and 60 and 50°C, respectively. The immobilized *M. circinelloides* and *M. hiemalis* xylanase retained 84.02 and 79.43% activity, respectively. The production of xylanase from *M. circinelloides* and *M. hiemalis* utilizing RSM is deemed profitable for the decomposition of the agro-industrial wastes.

## 1. Introduction

Biotechnological processes on filamentous fungi have enabled the industrial utilization of their ability to produce valuable enzymes due to their ease of propagation and increased production of extracellular enzymes with specific characteristics such as stability over a wide range of pH and temperature [1–3].

The class Zygomycetes, order Mucorales, including the genus *Mucor*, are mainly saprobic fungi that can grow well on various agro-industrial wastes with the potential to produce extracellular hydrolytic enzymes that can be utilized in a variety of industrial applications [4].

The elevated dilation of agricultural junks apostasy has resulted in the cumulation of significant amounts of lignocellulosic wastes across the globe [5]. Lignocellulosic plant

TABLE 1: Box–Behnken design levels of independent factors.

No.	Factor	Variables	Units	Minimum	Range Maximum	Mean
1	A	Temperature	°C	20	40	30
2	B	pH	—	5	9	7
3	C	Incubation period	Day	5	9	7
4	D	Inoculum size	mL	1	5	3
5	E	Substrate concentration	g	1	5	3

TABLE 2: Xylanase production on different substrates under shaking and static conditions.

Substrate	Xylanase activity (U/mL)			
	<i>M. circinelloides</i>		<i>M. hiemalis</i>	
	Shaking	Static	Shaking	Static
Pomegranate peel	3.26 ± 0.41 <sup>b</sup>	2.11 ± 0.31 <sup>b</sup>	2.91 ± 0.28 <sup>b</sup>	3.73 ± 0.29 <sup>b</sup>
Tangerine peel	17.90 ± 0.68 <sup>a</sup>	21.77 ± 0.96 <sup>a</sup>	13.96 ± 0.13 <sup>a</sup>	15.28 ± 0.29 <sup>a</sup>
Wheat straw	0.69 ± 0.06 <sup>c</sup>	1.32 ± 0.23 <sup>b</sup>	2.84 ± 0.52 <sup>b</sup>	0.86 ± 0.07 <sup>c</sup>

The data were given as averages of three replicates (mean ± SD). Values followed by the different letters are significantly different at  $p < 0.05$ .

biomass mainly comprises hemicellulose, cellulose, and lignin [6]. The most crucial hemicellulosic heteroglycan is xylan, which contains a long chain of  $\beta$ -1,4-linked xylopyranose monomers and comprises a significant renovatable biocluster containing up to 20-35% of the dry mass of agricultural residues [5, 7]. Two main xylanolytic enzymes (endo- $\beta$ -1,4-xylanases and exo- $\beta$ -1,4-xylanases) are a candidate to degrade xylan into xylose and xylooligomers [8]. Xylanase is included in various industrial applications like animal food, feed, biobleaching, biofuel, pharmaceuticals, textile [9], and juice clarification [10].

The amount of fungal enzyme manufactured is determined by the conditions of the fermentation process and the need to optimize these conditions for low enzyme output cost [11]. The fungal enzyme manufacturing is prevalently implemented by submerged or solid-state fermentation [12]. Submerged fermentation, utilized in 90% enzyme industry, occurs in the presence of water excess, providing a soft handling and best control [13, 14]. Immobilization of enzymes and entrapment of enzymes in polymers such as alginate eliminate the need to isolate the enzyme from the outcome solution, facilitate enzyme recycling, enable repetitive and continuous use, and control enzyme activity [15–18].

This study aimed to evaluate the production of high-value xylanase by *Mucor* spp. using agro-industrial by-products as a cheap substrate in submerged fermentation conditions. The optimization of xylanase production, as well as characterization and immobilization of partially purified xylanase, was investigated.

## 2. Materials and Methods

**2.1. Tested Fungi.** *Mucor circinelloides* AUMC 6696.A (Accession no. MT509983) and *M. hiemalis* AUMC 6031 (Accession no. MT365791) [19] were utilized in the current search for xylanase production. Pure cultures were kept in potato dextrose agar (PDA) tubes and preserved at 4°C for further use.

### 2.2. Enzymes Preliminary Screening

**2.2.1. Screening on Agar Plates.** Czapek’s agar medium (g/L:  $\text{KH}_2\text{PO}_4$ , 1;  $\text{NaNO}_3$ , 2;  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ , 0.5 and  $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$ , 0.5) was supplemented with 10 g/L of xylan powder as a carbon source for xylanase production. The pH was set to 7, and the impregnated plates were preserved at  $28 \pm 2^\circ\text{C}$  for five days and then screened for enzymes production [20]. After immersing the cultured agar plates in iodine solution for 15 minutes, they were examined for the presence of a clear zone.

**2.2.2. Screening by Submerged Fermentation (SmF).** Pomegranate peel, tangerine peel, and wheat straw were desiccated at  $65^\circ\text{C}$  for 24 h, then squelched to grist, and used as substrates (10 g/L) in Czapek’s mineral salts broth. The broth containing flasks was adjusted to pH 7 and autoclave sterilized. The strains’ spore suspensions ( $10^7$  spores/mL) were utilized to inoculate 250-mL flasks holding 100 mL submerged broth. The impregnated flasks were preserved at  $28 \pm 2^\circ\text{C}$  for seven days on both shaking and static conditions. Subsequently, broth media were centrifuged, and supernatants were maintained for further enzymatic analysis.

**2.2.3. Quantitative Screening of Xylanase.** Xylanase efficiency was measured according to Miller [21]. In addition, 0.5 mL enzyme supernatant was added to 0.5 mL xylan (1% w/v) in acetate buffer, and the mixture was incubated at  $50^\circ\text{C}$  for 30 min. Afterward, the interaction was intercepted by appending 1 mL of 3,5-dinitrosalicylic acid reagent and incubated at  $100^\circ\text{C}$  for 10 min. After cold dishing, the absorbance was measured at 570 nm using a spectrophotometer (Jenway 7315, UK). The amount of reducing sugars was determined using xylose as a standard for plotting the calibration curve. All the tests were carried out three times, and the outputs were expressed as an average value. A unit of the enzyme was acquainted as the quantity of the enzyme per one mL required to release one  $\mu\text{mol}$  of reducing sugar from a substrate per 60 seconds under the optimum trial conditions [22].

TABLE 3: Box–Behnken design of optimization variables with experimental and predicted xylanase activity of both *M. circinelloides* and *M. hiemalis*.

Run order	Variables					Xylanase activity (U/mL)			
						<i>M. circinelloides</i>		<i>M. hiemalis</i>	
	A	B	C	D	E	Experimental response	Predicted response	Experimental response	Predicted response
1	40	7	7	5	3	17.73	18.54	17.32	17.49
2	30	7	7	5	1	27.63	27.49	17.23	18.32
3	40	7	7	3	1	31.92	31.24	28.70	28.42
4	20	7	7	3	1	28.62	28.03	30.02	28.82
5	20	9	7	3	3	38.52	38.49	29.69	31.58
6	40	9	7	3	3	23.42	23.39	24.00	24.96
7	30	9	7	3	1	31.67	33.36	28.45	29.76
8	30	9	5	3	3	23.26	22.76	25.07	25.48
9	30	7	7	3	3	27.38	28.56	19.79	23.78
10	30	5	9	3	3	28.45	27.53	32.91	32.16
11	30	5	7	5	3	27.22	26.26	28.12	27.07
12	30	5	7	3	1	32.58	33.81	26.89	27.84
13	30	7	5	3	1	28.37	28.36	27.79	26.20
14	30	9	7	3	5	34.72	34.49	19.54	19.21
15	40	7	7	3	5	15.50	14.68	16.90	16.68
16	30	7	7	3	3	29.28	28.56	23.50	23.78
17	30	5	5	3	3	31.84	30.84	25.32	26.07
18	40	7	5	3	3	25.24	24.05	26.72	26.33
19	20	5	7	3	3	30.02	31.23	34.89	36.26
20	20	7	7	1	3	30.27	29.44	30.93	29.60
21	30	7	9	3	5	33.40	33.82	24.08	24.25
22	30	7	9	3	1	34.81	33.59	26.80	24.94
23	30	7	9	1	3	25.81	25.87	21.11	22.74
24	30	5	7	3	5	28.54	27.85	29.61	28.92
25	30	7	5	3	5	21.69	23.31	16.99	17.42
26	30	9	7	5	3	30.10	29.43	24.49	22.92
27	30	9	9	3	3	42.23	41.80	26.06	24.96
28	30	7	7	3	3	26.89	28.56	23.17	23.78
29	30	7	9	5	3	32.83	32.45	28.78	28.29
30	30	5	7	1	3	26.39	26.31	32.58	31.54
31	30	7	7	1	5	24.91	25.05	17.15	18.30
32	40	5	7	3	3	23.26	24.46	27.63	28.08
33	30	9	7	1	3	29.11	29.32	29.44	27.88
34	20	7	7	5	3	28.95	29.26	30.85	32.15
35	30	7	7	3	3	28.21	28.56	25.40	23.78
36	30	7	7	1	1	28.95	28.66	31.59	33.18
37	40	7	7	1	3	18.64	18.30	31.92	29.46
38	30	7	7	5	5	25.98	26.28	23.09	23.74
39	30	7	5	1	3	23.42	24.56	28.21	30.22
40	30	7	7	3	3	29.86	28.56	25.48	23.78
41	20	7	7	3	5	40.50	39.76	32.25	31.10
42	20	7	5	3	3	22.43	21.66	26.39	24.86
43	40	7	9	3	3	17.56	18.59	18.47	20.24
44	30	7	7	3	3	29.77	28.56	25.32	23.78
45	30	7	5	5	3	17.32	18.03	15.34	15.24
46	20	7	9	3	3	41.41	42.85	35.88	36.52

TABLE 4: ANOVA for the experimental results of xylanase biosynthesis by *M. circinelloides*.

Source	Sum of squares	Degree of freedom	Mean of squares	F value	p value	Prob > F
Model	1608.62	20	80.431	58.10	0.000	Significant
Linear	787.16	5	157.431	113.73	0.000	
A	477.94	1	477.943	345.27	0.000	
B	38.28	1	38.283	27.66	0.000	
C	247.63	1	247.635	178.90	0.000	
D	0.00	1	0.004	0.00	0.958	
E	23.29	1	23.291	16.83	0.000	
Square	237.41	5	47.483	34.30	0.000	
A <sup>2</sup>	21.21	1	21.208	15.32	0.001	
B <sup>2</sup>	49.74	1	49.738	35.93	0.000	
C <sup>2</sup>	0.42	1	0.416	0.30	0.589	
D <sup>2</sup>	84.92	1	84.925	61.35	0.000	
E <sup>2</sup>	17.72	1	17.717	12.80	0.001	
2-way interaction	584.05	10	58.405	42.19	0.000	
AB	17.36	1	17.357	12.54	0.002	
AC	177.51	1	177.513	128.24	0.000	
AD	0.04	1	0.043	0.03	0.862	
AE	200.18	1	200.176	144.61	0.000	
BC	124.96	1	124.958	90.27	0.000	
BD	0.01	1	0.007	0.00	0.945	
BE	12.58	1	12.584	9.09	0.006	
CD	43.01	1	43.015	31.07	0.000	
CE	6.97	1	6.969	5.03	0.034	
DE	1.43	1	1.431	1.03	0.319	
Residual	34.61	25	1.384			
Lack-of-fit	26.62	20	1.331	0.83	0.655	Not significant
Pure error	7.99	5	1.597			
Total	1643.23	45				

$R^2$ : 0.9789; adjusted  $R^2$ : 0.9621; predicted  $R^2$ : 0.9282.

2.3. *Optimization of Enzymatic Productivity under Submerged Fermentation (SmF)*. Response surface methodology (RSM) tactic using Box–Behnken design (BBD) was exercised to determine the optimum factors for boosted xylanase production, including A, temperature; B, pH; C, incubation period; D, inoculum size; and E, substrate concentration (Table 1). Forty-six experiments with the central points were employed to satisfy the polynomial pattern established on a Box–Behnken design (BBD, 5 variables) attained by Minitab 19® software (Version19.1.1.0. LLC). At three-level and five factors, experimental BBD was examined, and the number of the tests ( $N$ ) was determined corresponding to the subsidiary equation:

$$N = 2k * (k - 1) + C0 \quad (1)$$

where  $k$  is the digit of factors and  $C0$  is the digit of central points equal to 6.

The impact of variables on the simulation ( $Y$ ) was evaluated by utilizing a second-order polynomial equation

to determine the quixotic states of the xylanase biosynthesis.

$$\begin{aligned}
 Y = & \beta_0 + \beta_1 A + \beta_2 B + \beta_3 C + \beta_4 D + \beta_5 E + \beta_{11} A^2 + \beta_{22} B^2 \\
 & + \beta_{33} C^2 + \beta_{44} D^2 + \beta_{55} E^2 + \beta_{12} AB + \beta_{13} AC + \beta_{14} AD \\
 & + \beta_{15} AE + \beta_{23} BC + \beta_{24} BD + \beta_{25} BE + \beta_{34} CD \\
 & + \beta_{35} CE + \beta_{45} DE
 \end{aligned} \quad (2)$$

where  $Y$  (response variable);  $\beta_0$  (intercept);  $\beta_1, \beta_2, \beta_3, \beta_4,$  and  $\beta_5$  (linear coefficients);  $\beta_{11}, \beta_{22}, \beta_{33}, \beta_{44},$  and  $\beta_{55}$  (square coefficients);  $\beta_{12}, \beta_{13}, \beta_{14}, \beta_{15}, \beta_{23}, \beta_{24}, \beta_{25}, \beta_{34}, \beta_{35},$  and  $\beta_{45}$  (interaction coefficients); and  $A, B, C, D, E, A^2, B^2, C^2, D^2, E^2, AB, AC, AD, AE, BC, BD, BE, CD, CE,$  and  $DE$  (levels of independent variables). The corresponding coefficients of variables, interaction variables, and contour graphs were obtained by Minitab 19® software. By analyzing the regression equation and constructing the response plots,

TABLE 5: ANOVA for the experimental results of xylanase biosynthesis by *M. hiemalis*.

Source	Sum of squares	Degree of freedom	Mean of squares	F value	p value	Prob > F
Model	1136.70	20	56.835	17.59	0.000	Significant
Linear	489.53	5	97.906	30.30	0.000	
A	219.29	1	219.287	67.86	0.000	
B	60.78	1	60.778	18.81	0.000	
C	31.01	1	31.009	9.60	0.005	
D	88.84	1	88.837	27.49	0.000	
E	89.62	1	89.617	27.73	0.000	
Square	196.46	5	39.292	12.16	0.000	
A <sup>2</sup>	85.58	1	85.581	26.48	0.000	
B <sup>2</sup>	95.63	1	95.630	29.59	0.000	
C <sup>2</sup>	0.05	1	0.055	0.02	0.898	
D <sup>2</sup>	0.61	1	0.611	0.19	0.667	
E <sup>2</sup>	3.76	1	3.762	1.16	0.291	
2-way interaction	450.72	10	45.072	13.95	0.000	
AB	0.61	1	0.614	0.19	0.667	
AC	78.65	1	78.650	24.34	0.000	
AD	52.70	1	52.705	16.31	0.000	
AE	49.17	1	49.172	15.22	0.001	
BC	10.89	1	10.889	3.37	0.078	
BD	0.06	1	0.061	0.02	0.892	
BE	33.83	1	33.827	10.47	0.003	
CD	105.49	1	105.493	32.65	0.000	
CE	16.34	1	16.341	5.06	0.034	
DE	102.97	1	102.966	31.86	0.000	
Residual	80.79	25	3.232			
Lack-of-fit	56.54	20	2.827	0.58	0.823	Not significant
Pure error	24.25	5	4.850			
Total	1217.49	45				

$R^2$ : 0.9336; adjusted  $R^2$ : 0.8806; predicted  $R^2$ : 0.7856.

the ideal values of the tested variables were secured. The coefficient of limitation  $R^2$  was used to express the fineness of profit of the polynomial equation, and the ( $F$ ) test was used to determine its statistical significance level.

#### 2.4. Partial Purification of Xylanase from *Mucor* Strains.

After the incubation period under optimum conditions, the contents of the broth culture were centrifuged, and the supernatant was utilized for enzyme assay. Xylanase activity and protein concentration were measured in the supernatant according to Miller [21] and Lowry et al. [23], respectively, utilizing the standard of bovine serum albumin to generate the calibration curve spectrophotometrically at 750 nm.

Crude enzyme solution was partially purified by precipitation using cold acetone. Precooled acetone ( $-20^{\circ}\text{C}$ ) was subjoined to the enzyme solution until the volume ratio between enzyme solution and acetone reached 1:1; 1:2; 1:3; 1:4, and 1:5 (v/v). The solution was left at  $-20^{\circ}\text{C}$  overnight to allow protein precipitation. The precipitates were gathered by centrifugation at 10000 rpm for 15 min and resuspended in a small volume of (sodium citrate buffer,


pH 4.8, 0.05 M). These samples were used for determining the activity of xylanase, purification factor, and enzyme recovery yield [24]. Protein was estimated, and suitable precipitants for characterization were selected. The following equations were used to calculate the partially purified xylanase's specific activity, yield, and purification fold.

$$\text{Specific activity (U/mg)} = \frac{\text{Total activity}}{\text{Total protein}},$$


$$\text{Yield (\%)} = \frac{\text{Total units in partially purified enzyme} \times 100\%}{\text{Total units in crude enzyme}},$$

$$\text{Purification fold} = \frac{\text{Specific activity of partially purified enzyme}}{\text{Specific activity of crude enzyme}}. \quad (3)$$


2.5. *Characterization of Partially Purified Xylanase.* The optimum temperature for partially purified xylanase activity was determined to be in the range of ( $30-90^{\circ}\text{C}$ ), and the thermal stability was determined after premaintaining the




(a)



(b)




(c)




(d)


FIGURE 1: Continued.




(e)



(f)



(g)



(h)

FIGURE 1: Continued.




FIGURE 1: Contour plot showing interactions between independent variables (a)–(j) for xylanase activity produced by *M. circinelloides*.

enzyme at each temperature degree for one hour before screening. Ideal pH estimation was carried out at optimum temperature utilizing various buffers with values (3-11), and the pH stability was assessed after preserving these pH values for 1 h before the screening. In addition, xylanase activity was estimated after maintaining the enzyme with different metal ions (10 mM of  $K^+$ ,  $Mg^{2+}$ ,  $Ba^{2+}$ , and  $Ni^{2+}$ ) for 1 h at optimum temperature and pH. Xylanase activity was assessed after processing with diverse detergents comprising tween 80 and 20 at concentrations of 1 and 5% v/v, urea (1 and 5% w/v), and  $Na_2CO_3$  (50 and 75 mM) compared to control (100% activity) [25, 26].

**2.6. Immobilization of Xylanase.** The immobilization of partially purified xylanase was performed by combining enzyme solution with an equal volume of 3.0% sodium alginate solution. Calcium alginate beads were formalized by adding the mixture drop-wise in 0.2 M  $CaCl_2$  at 4°C. Calcium alginate beads were rinsed by double-distilled  $H_2O$  to remove unstriated enzyme units. Then, the beads were desiccated and stored in phosphate buffer. The entrapped beads were activated using glutaraldehyde for covalent binding of xylanase onto the beads [27], and the assay of free and immobilized xylanase was performed. The immobilization yield was

assayed according to the equation [28].

$$\text{Immobilization yield(\%)} = \frac{A_i - A_f}{A_f} \times 100, \quad (4)$$

where  $A_i$  is the immobilized enzyme activity and  $A_f$  is the free enzyme activity.

**2.6.1. Scanning Electron Microscopy.** A scanning electron microscope (JEOL JSM 5400, Japan) was utilized to examine the outer surface forms of calcium alginate beads before and after xylanase immobilization.

**2.7. Data Analysis.** All tests and measurements were repeated three times. Using the SPSS, software program (version No. 16), and one-way ANOVA, the values were expressed as the mean  $\pm$  SD at the 0.05 significance level.

### 3. Results

**3.1. Preliminary Screening of Xylanase Production.** The preliminary screening for extracellular fungal xylanase revealed that both *M. circinelloides* and *M. hiemalis* had a high potential to produce xylanase qualitatively on a solid assay









FIGURE 2: Continued.




(e)



(f)



(g)



(h)

FIGURE 2: Continued.




FIGURE 2: Contour plot showing interactions between independent variables (a)–(j) for xylanase activity produced by *M. hiemalis*.

TABLE 6: Precipitation of xylanase produced by *M. circinelloides* and *M. hiemalis* using different acetone concentrations.

Ratio (crude : acetone)	Xylanase activity (U/mL)	
	<i>M. circinelloides</i>	<i>M. hiemalis</i>
1 : 1	6.92 ± 0.22 <sup>c</sup>	5.19 ± 0.22 <sup>c</sup>
1 : 2	5.41 ± 0.27 <sup>d</sup>	3.68 ± 0.27 <sup>d</sup>
1 : 3	7.25 ± 0.07 <sup>c</sup>	5.52 ± 0.07 <sup>c</sup>
1 : 4	11.89 ± 0.23 <sup>a</sup>	10.73 ± 0.77 <sup>a</sup>
1 : 5	9.59 ± 0.34 <sup>b</sup>	8.38 ± 0.87 <sup>b</sup>

The data were given as averages of three replicates (mean ± SD). Values followed by the different letters are significantly different at  $p < 0.05$ .

medium. They were then quantitatively assayed under SmF and produced 21.77 and 15.28 U/mL, respectively, on the tangerine peel as a substrate under static condition (Table 2).

3.2. Response Surface Methodology for Optimization of Xylanase Production. Table 1 shows the independent factors with their competent levels employed in optimization of xylanase output, while BBD of the independent factors along with predicted as well as experimental values are depicted in

Table 3. The production of xylanase by *M. circinelloides* was predicted using the following equation:

$$\begin{aligned}
 Y(U/mL) = & 16.6 + 4.495A - 15.60B - 0.51C - 1.73D \\
 & + 2.01E - 0.01559A^2 + 0.5968B^2 - 0.0546C^2 \\
 & - 0.7799D^2 + 0.3562E^2 - 0.1042AB - 0.3331AC \\
 & + 0.0052AD - 0.3537AE + 1.397BC + 0.010BD \\
 & + 0.443BE + 0.820CD + 0.330CE + 0.150DE
 \end{aligned}
 \tag{5}$$


While the production of xylanase by *M. hiemalis* was predicted by the following equation:

$$\begin{aligned}
 Y(U/mL) = & 51.0 + 0.236A - 7.99B + 4.59C - 8.71D + 2.81E \\
 & + 0.03131A^2 + 0.828B^2 + 0.020C^2 + 0.066D^2 \\
 & - 0.164E^2 + 0.0196AB - 0.2217AC - 0.1815AD \\
 & - 0.1753AE - 0.412BC - 0.031BD - 0.727BE \\
 & + 1.284CD + 0.505CE + 1.268DE
 \end{aligned}
 \tag{6}$$

The highest xylanase activity of both *M. circinelloides*

TABLE 7: Summary of specific activity, yield, and purification fold of xylanase produced by *M. circinelloides* and *M. hiemalis*.

Fungal strain	Purification steps	Total activity (U/mL)	Total protein (mg/mL)	Specific activity (U/mg)	Yield (%)	Purification fold
<i>M. circinelloides</i>	Culture supernatant	14519.60	86.05	169.09	100	1
<i>M. hiemalis</i>		12458.07	102.39	121.74	100	1
<i>M. circinelloides</i>	Acetone	4522.52	15.48	292.08	30.74	1.69
<i>M. hiemalis</i>	Acetone	3904.07	16.19	240.15	31.34	1.97

FIGURE 3: Effects of temperature and pH on activity and stability of partially purified xylanase (a) and (c) *M. circinelloides* and (b) and (d) *M. hiemalis*.

(42.23 U/mL) and *M. hiemalis* (35.88 U/mL) was obtained from runs No. 27 to 46, respectively. The run No. 27 for *M. circinelloides* consisted of incubation temperature (30°C), pH value (9), the incubation period (9 days), inoculum size (3 mL), and substrate concentration (3 g/100 mL). In comparison, the run No. 46 for *M. hiemalis* consisted of incubation temperature (20°C), pH value (7), the incubation period (9 days), inoculum size (3 mL), and substrate concentration (3 g/100 mL).

Analysis of variance (ANOVA) for the xylanase quadric model of *M. circinelloides* and *M. hiemalis* is shown in

Tables 4 and 5. Model terms with a  $p$  value  $<0.05$  were deemed significant. The model  $F$  value of 58.10 and 17.59 for *M. circinelloides* and *M. hiemalis*, respectively, indicated that the model is significant. Values of “Prob  $> F$ ”  $<0.05$  pointed model terms are significant. In this case,  $A, B, C, E, A^2, B^2, D^2, E^2, AB, AC, AE, BC, BE, CD,$  and  $CE$  are significant model terms for *M. circinelloides*, while for *M. hiemalis*,  $A, B, C, D, E, A^2, B^2, AC, AD, AE, BE, CD, CE,$  and  $DE$  are significant model terms. The “Lack of Fit  $F$ -value” of 0.83 and 0.58 for *M. circinelloides* and *M. hiemalis*, respectively, indicated that the lack of fit is insignificant concerning

TABLE 8: Effects of metal ions and detergents on activity and stability of partially purified xylanase from *M. circinelloides* and *M. hiemalis*.

Metal ions and detergents	Conc.	<i>M. circinelloides</i>		<i>M. hiemalis</i>	
		Relative activity (%)	Relative stability (%)	Relative activity (%)	Relative stability (%)
Control	0	100.00 ± 0.00	—	100.00 ± 0.00	—
K <sup>+</sup>	10 mM	48.52 ± 0.75	50.55 ± 0.47	140.34 ± 0.34	109.01 ± 1.03
Mg <sup>2+</sup>	10 mM	42.99 ± 0.47	82.08 ± 0.21	96.30 ± 0.44	90.70 ± 2.35
Ba <sup>2+</sup>	10 mM	28.37 ± 0.55	10.10 ± 0.12	77.90 ± 0.69	84.75 ± 1.11
Ni <sup>2+</sup>	10 mM	16.68 ± 0.42	35.12 ± 0.25	38.29 ± 0.96	93.44 ± 1.04
Tween 80	1% (v/v)	53.06 ± 0.83	40.94 ± 0.18	61.76 ± 0.13	35.20 ± 0.49
	5% (v/v)	55.09 ± 0.94	27.99 ± 0.54	72.15 ± 0.11	29.93 ± 0.70
Tween 20	1% (v/v)	61.90 ± 0.28	44.74 ± 0.42	62.33 ± 0.61	47.10 ± 0.68
	5% (v/v)	64.27 ± 0.40	44.57 ± 0.20	65.15 ± 1.05	40.10 ± 1.17
Urea	1% (w/v)	46.58 ± 0.38	37.74 ± 0.59	61.04 ± 0.58	42.13 ± 0.66
	5% (w/v)	59.43 ± 0.80	20.73 ± 0.09	49.89 ± 0.35	27.20 ± 0.67
Na <sub>2</sub> CO <sub>3</sub>	50 mM	30.30 ± 0.89	9.63 ± 0.02	29.37 ± 0.36	22.62 ± 0.22
	75 mM	30.49 ± 0.58	4.44 ± 0.04	40.08 ± 0.82	7.81 ± 0.51

The data were given as averages of three replicates (mean ± SD).

TABLE 9: The immobilization yield (%) of xylanase entrapped in calcium alginate beads.

Fungal strain	<i>M. circinelloides</i>	<i>M. hiemalis</i>
Yield (%)	84.02 ± 0.63	79.43 ± 0.47

The data were given as averages of three replicates (mean ± SD).

the pure error. Nonsignificant lack of fit is proper for the model to be convenient. The resulted multiple correlation coefficient ( $R^2 = 0.9789$  for *M. circinelloides* and  $R^2 = 0.9336$  for *M. hiemalis*) that nigh to 1 betokened preferable interconnection between experimental and predicted values and elucidated the model accuracy with upgrade response.

Contour plots explained the relationship between parameters and defined each factor's optimum scale for xylanase efficiency by *M. circinelloides* (Figures 1(a)–1(j)) and *M. hiemalis* (Figures 2(a)–2(j)). The response surface plot constructed any two variables, while other variables were maintained at their optimal level. Contour plots of xylanase activity by *M. circinelloides* revealed significant interactions between incubation temperature with pH, incubation period, and substrate concentration; pH with incubation period and substrate concentration; and incubation period with inoculum size and substrate concentration. In contrast, contour plots of the interactions between incubation temperature with incubation period, inoculum size, and substrate concentration; pH with substrate concentration; incubation period with inoculum size and substrate concentration; and inoculum size with substrate concentration significantly influenced xylanase production by *M. hiemalis*. The remaining interactions insignificantly influenced xylanase production.

**3.3. Partial Purification of Xylanase.** Extracellular xylanase from *M. circinelloides* and *M. hiemalis* was partially purified from broth cultures by using different acetone concentra-

tions. The highest xylanase activity (11.89 and 10.73 U/mL from *M. circinelloides* and *M. hiemalis*, respectively) was obtained via crude filtrate precipitation with acetone at ratio 1:4 (Table 6). Xylanase purification resulted in 1.69-fold purification and 30.74% xylanase recovery with a specific activity of 292.08 U/mg from *M. circinelloides*, while in 1.97-fold purification, 31.34% recovery with a specific activity of 240.15 U/mg was obtained from *M. hiemalis* xylanase (Table 7).

**3.4. Characterization of Partially Purified Xylanase.** *M. circinelloides* partially purified xylanase was highly active at 60°C (total activity 100%) and decreased gradually at 50–30°C and 70–90°C, while xylanase activity of *M. hiemalis* was highly active at 50°C (total activity 100%) and decreased gradually at 40–30°C and 60–90°C. The relative xylanase stability from both *M. circinelloides* and *M. hiemalis* was high at 30°C and decreased in the range of 40–90°C (Figures 3(a) and 3(b)). Furthermore, partially purified xylanase from both strains had the highest activity and stability (100%) at pH 7.0, and then, at higher or lower pH values, activity and stability of xylanase were reduced (Figures 3(c) and 3(d)).

After incubation with 10 mM K<sup>+</sup>, Mg<sup>2+</sup>, Ba<sup>2+</sup>, and Ni<sup>2+</sup>, xylanase activity of both strains decreased with the exception of K<sup>+</sup>, which increased *M. hiemalis* xylanase activity by 40.34% compared to the control, while stability increased except for Ba<sup>2+</sup> with *M. circinelloides* enzyme and K<sup>+</sup> and Mg<sup>2+</sup> with *M. hiemalis* enzyme. Detergents including Tween 80, Tween 20, urea, and Na<sub>2</sub>CO<sub>3</sub> at low concentrations of 1%, 1%, 1%, and 50 mM, respectively, reduced *M. circinelloides* and *M. hiemalis* partially purified xylanase activity. In contrast, there was a significant increase in activity and decreased stability at high concentrations of the tested detergents, except urea, which decreased *M. hiemalis* partially purified xylanase activity at its applied high concentration (5%) (Table 8).




FIGURE 4: Scanning electron micrographs (SEM) of calcium alginate beads with and without entrapped xylanase. (a)–(b) Micrographs of calcium alginate beads without enzyme, (c)–(e) micrographs of calcium alginate beads with immobilized xylanase from *M. circinelloides*, and (f)–(h) immobilized xylanase from *M. hiemalis* at magnification scales of 5000 $\times$  and 10,000 $\times$ , respectively.

**3.5. Xylanase Immobilization and Scanning Electron Microscopy.** *M. circinelloides* enzyme had an immobilization yield of 84.02%, while *M. hiemalis* enzyme had a yield of 79.43% (Table 9). A scanning electron microscope was utilized to investigate the surface superficial of calcium alginate beads together and without entrapped xylanase (Figure 4). The vents of calcium alginate beads were examined in the micrographs of calcium alginate beads without entrapped xylanase, whereas together with xylanase, the vents were coated with intense molecule xylanase.

#### 4. Discussion

Fungi play an excellent source for the production of various beneficial enzymes. Xylan represents the principal hemicellulosic xylopolymer found in plant cell walls. The biodegradation of xylan functions as an essential role of plant materials natural [29]. In this research, *M. circinelloides* and *M. hiemalis* were investigated as potential sources for producing xylanase.

Many fungi were proven to have multiple xylanase forms and can efficiently breakdown xylan. In addition, the

substrate type influenced the concentration and number of various expressed forms of xylanase [30, 31]. On the contrary, the substrate installation and growth factors affect the fungal growth and the biosynthesis of enzymes. Therefore, there is an urgent need to modify the biosynthesis of the enzymes by optimizing fungal production.

In the current research, the low  $p$  values, which are attained by the  $F$  test, and high  $R^2$  values indicated that the employed model attained a high significance, and its sufficiency was confirmed [32]. For xylanase production on wheat bran by solid-state fermentation by *M. indicus* and *M. hiemalis* using RSM, the optimum temperatures were 40.0 and 43.4 °C, respectively, and 51.3 and 53.2 h for incubation time, respectively, while the highest xylanase activities were 43.1 and 43.8 U/g for *M. indicus* and *M. hiemalis*, respectively [33]. Atalla et al. [34] reported a maximum xylanase activity from *A. oryzae* (0.37 U/mL) by utilizing rice straw waste, while orange peel exhibited low xylanase production (0.17 U/mL). Cui and Zhao [35] highlighted the magnitude of the substrate concentration for xylanase biosynthesis, where the enzyme activity by *Penicillium* sp. WX-Z1 demonstrated a gradual rise in wheat bran concentration. Statistical analysis of xylanase production from *Penicilliumoxalicum* ZH-30 by RSM showed that the linear, quadric terms and initial pH and temperature interaction significantly impacted the optimal conditions for raised xylanase biosynthesis at pH 7.38 and temperature 31.1 °C. Under optimal conditions, the portended and experimented xylanase activities were 14.33 and 14.50 U/mL, respectively [36].

Box–Behnken design statistical analysis by Cao et al. [37] demonstrated that the linear and quadric terms of cultivation time, pH, and substrate concentration variables significantly affected xylanase production by *Aspergillus niger* AN-13 using wheat bran under submerged fermentation. In submerged fermentation with corncob, *Aspergillus niger* KIBGE-IB36 was found to be a high producer of xylanase [38] and different spectra of xylanase synthesis using wheat bran, rice husk, orange peel, and pomegranate peel [5]. RSM method by Azzouz et al. [39] reported a 65.01% increase in xylanase output by *Aspergillus niger* strain BG on 84% humidified wheat bran with a pH of 2.5 at 37 °C and incubation for 66 h. Statistical analysis of xylanase output by *Aspergillus foetidus* in submerged fermentation with soybean scraps revealed that variables pH and the interaction of pH and temperature influenced xylanase biosynthesis, with good xylanase activity (13.98 U/mL) at pH 7.0, 28 °C, and 120 rpm for 168 hours [40]. Response surface methodology by Ramanjaneyulu and Rajasekhar [41] for maximal production of xylanase (4560 U/mL) by *Fusarium* sp. BVKT R2 demonstrated that the optimal conditions were sorbitol 1.5%, yeast extract 1.5%, pH of 5.0, temperature of 32.5 °C, and shaking of 175 rpm.

High temperature increases the solubility of reactants and products by decreasing viscosities, resulting in faster hydrolysis [42], and prolonged energetic existence would fabricate enzymes convenient for promoted and active biomass diversion. Therefore, thermostability is the most substantial property for the enzyme used under extreme bioprocessing conditions to be efficient [8]. RSM optimization of xylanase yield by *Aspergillus niger* 3-fold and

1.41-fold purification was attained with about 6.2% yield, and the highest activity of the purified xylanase was observed at pH 6 and 50 °C. The produced xylanase exhibited high thermal and pH stability, with more than 90% residual activity between 30 and 40 °C and pH 3–9 after incubation of 24 h, with half-lives of 30 min at 50 and 60 °C [43]. Liet al. [36] reported a temperature range of 50–60 °C suitable for the industrial application of xylanase from *P. oxalicum* ZH-30. Supplementation of Tween 80 as additional surfactant improved (72.4 ± 1.42 U/g) the titer of xylanase on sugarcane bagasse by *T. viride*-IR05 [44]. Regarding xylanase immobilization, the formation of an ionic bond of Ca<sup>2+</sup> with carboxylate groups of sodium alginate help to formulate mechanically firm grids of alginate gel that fulfill the demands for efficient hydrogel systems acting as carriers of bioactive molecules in several industrial applications [45]. Immobilized *Talaromyces thermophilus* xylanase covalently bound by glutaraldehyde to chitosan, chitin, amberlite, duolite, florisil, and gelatin gave immobilization yield of 89.0, 87.8, 89.3, 81.1, 96.2, and 98.8%, respectively [28].

## 5. Conclusion

The present study utilized the response surface methodology via the Box–Behnken design to improve xylanase production by *M. circinelloides* and *M. hiemalis*. The experimental results are consistent with predicted responses. The produced enzyme was partially purified, characterized, and immobilized. The optimum parameters for xylanase production by *M. circinelloides* and *M. hiemalis* consisted of incubation temperature (30 and 20 °C), pH value (9 and 7) incubation period (9 and 9 days), inoculum size (3 and 3 mL), and substrate concentration (3 and 3 g/100 mL), respectively. The partial purification of *M. circinelloides* and *M. hiemalis* xylanase yielded 1.69- and 1.97-fold purification, and the immobilized xylanase retained 84.02 and 79.43% activity, respectively. Response surface methodology was effective and satisfactory and investigated many factors simultaneously. More research is needed to scale up enzyme production for a wide range of applications.

## Abbreviations

RSM:	Response surface methodology
BBD:	Box–Behnken design
ANOVA:	Analysis of variance
SmF:	Submerged fermentation.

## Data Availability

No data were used to support this study.

## Conflicts of Interest

The authors declare that there is no conflict of interest regarding the publication of this paper.

## Authors' Contributions

All authors contributed equally to this work. All authors read and approved the final manuscript.

## Acknowledgments

The authors extend their appreciation to the Deanship of Scientific Research at King Saud University for funding this work through research group No (RG-1441-419). Also, the authors extend their appreciation to the Deanship of Scientific Research at King Saud University for the logistic support of this work through the Research Assistant Internship Program, Project no. (RAIP-4-21-2).

## References

- [1] L. H. S. Guimaraes, S. C. Peixoto-Nogueira, M. Michelin et al., "Screening of filamentous fungi for production of enzymes of biotechnological interest," *Brazilian Journal of Microbiology*, vol. 37, no. 4, pp. 474–480, 2006.
- [2] P. S. Nigam, "Microbial enzymes with special characteristics for biotechnological applications," *Biomolecules*, vol. 3, no. 4, pp. 597–611, 2013.
- [3] R. L. Torres-Barajas and G. Aguilar-Osorio, "Xylanolytic system proteins desorption from *Aspergillus flavipes* FP-500 cultures with agrowastes," *Revista Mexicana de Ingeniería Química*, vol. 12, no. 3, 2013.
- [4] M. Tako, A. Kotogan, J. Krisch, C. Vagvolgyi, K. C. Mondal, and T. Papp, "Enhanced production of industrial enzymes in *Mucoromycota* fungi during solid-state fermentation of agricultural wastes/by-products," *Acta Biologica Hungarica*, vol. 66, no. 3, pp. 348–360, 2015.
- [5] U. Javed, A. Ansari, A. Aman, and S. A. UlQader, "Fermentation and saccharification of agro-industrial wastes: a cost-effective approach for dual use of plant biomass wastes for xylose production," *Biocatalysis and Agricultural Biotechnology*, vol. 21, article 101341, 2019.
- [6] D. Mamma, E. Kourtoglou, and P. Christakopoulos, "Fungal multienzyme production on industrial by-products of the citrus-processing industry," *Bioresource Technology*, vol. 99, no. 7, pp. 2373–2383, 2008.
- [7] S. Senthilkumar, B. Ashokkumar, K. Chandraraj, and P. Gunasekaran, "Optimization of medium composition for alkali-stable xylanase production by Fxn 1 in solid-state fermentation using central composite rotary design," *Bioresource Technology*, vol. 96, no. 12, pp. 1380–1386, 2005.
- [8] A. Bhalla, K. M. Bischoff, and R. K. Sani, "Highly thermostable xylanase production from a thermophilic *Geobacillus* sp. strain WsUcF1 utilizing lignocellulosic biomass," *Frontiers in Bioengineering and Biotechnology*, vol. 3, p. 84, 2015.
- [9] S. C. Izidoro and A. Knob, "Production of xylanases by an *Aspergillus niger* strain in wastes grain," *Acta Scientiarum Biological Sciences*, vol. 36, no. 3, p. 313, 2014.
- [10] E. Rosmine, N. C. Sainjan, R. Silvester, A. Alikkunju, and S. A. Varghese, "Statistical optimisation of xylanase production by estuarine *Streptomyces* sp. and its application in clarification of fruit juice," *Journal of Genetic Engineering and Biotechnology*, vol. 15, no. 2, pp. 393–401, 2017.
- [11] G. M. Maciel, L. P. S. Vandenberghe, C. W. I. Haminiuk et al., "Xylanase production by *Aspergillus niger* LPB 326 in solid-state fermentation using statistical experimental designs," *Food Technology and Biotechnology*, vol. 46, no. 2, pp. 183–189, 2008.
- [12] M. K. Selwal and K. K. Selwal, "High-level tannase production by *Penicillium atramentosum* KM using agro residues under submerged fermentation," *Annals of Microbiology*, vol. 62, no. 1, pp. 139–148, 2012.
- [13] H. Joo and C. Chang, "Production of an oxidant and SDS-stable alkaline protease from an alkaloiphilic *Bacillus clausii* I-52 by submerged fermentation: Feasibility as a laundry detergent additive," *Enzyme and Microbial Technology*, vol. 38, no. 1-2, pp. 176–183, 2006.
- [14] R. R. Singhanian, R. K. Sukumaran, A. K. Patel, C. Larroche, and A. Pandey, "Advancement and comparative profiles in the production technologies using solid-state and submerged fermentation for microbial cellulases," *Enzyme and Microbial Technology*, vol. 46, no. 7, pp. 541–549, 2010.
- [15] S. Mazzuca, L. Giorno, A. Spadafora, R. Mazzei, and E. Drioli, "Immunolocalization of  $\beta$ -glucosidase immobilized within polysulphone capillary membrane and evaluation of its activity *in situ*," *Journal of Membrane Science*, vol. 285, no. 1-2, pp. 152–158, 2006.
- [16] A. B. EL-Tanash, A. A. Sherief, and A. Nour, "Catalytic properties of immobilized tannase produced from *Aspergillus aculeatus* compared with the free enzyme," *Brazilian Journal of Chemical Engineering*, vol. 28, no. 3, pp. 381–391, 2011.
- [17] M. F. Alkhatib, M. Z. Alam, and R. Mohammed, "Statistical modelling optimisation of cellulose enzyme immobilisation on functionalised multi-walled carbon nanotubes for empty fruit bunches degradation," *Australian Journal of Basic and Applied Sciences*, vol. 6, no. 1, pp. 30–38, 2012.
- [18] M. Gutierrez-Correa, Y. Ludena, G. Ramage, and G. K. Villena, "Recent advances on filamentous fungal biofilms for industrial uses," *Applied Biochemistry and Biotechnology*, vol. 167, no. 5, pp. 1235–1253, 2012.
- [19] H. Mohamed, A. El-Shanawany, A. M. Shah et al., "Comparative analysis of different isolated oleaginous *Mucoromycota* fungi for their  $\gamma$ -linolenic acid and carotenoid production," *Bio Med Research International*, vol. 2020, p. 3621543, 2020.
- [20] F. Meddeb-Mouelhi, J. K. Moisan, and M. Beauregard, "A comparison of plate assay methods for detecting extracellular cellulase and xylanase activity," *Enzyme and Microbial Technology*, vol. 66, pp. 16–19, 2014.
- [21] L. G. Miller, "Use of dinitrosalicylic acid reagent for determination of reducing sugar," *Analytical Chemistry*, vol. 31, no. 3, pp. 426–428, 1959.
- [22] E. N. Fatokun, U. U. Nwodo, and A. I. Okoh, "Classical optimization of cellulase and xylanase production by a marine *Streptomyces* species," *Applied Sciences*, vol. 6, no. 10, 2016.
- [23] O. H. Lowry, N. J. Rosebrough, A. L. Farr, and R. J. Randall, "Protein measurement with the Folin phenol reagent," *Journal of Biological Chemistry*, vol. 193, no. 1, pp. 265–275, 1951.
- [24] S. Roe, *Protein Purification Applications*, Oxford University Press, 2nd edition, 2001.
- [25] G. Lakshmi and N. N. Prasad, "Purification and characterization of alkaline protease from a mutant *Bacillus licheniformis* B18," *Advances in Biological Research*, vol. 9, no. 1, 2015.
- [26] B. K. M. Lakshmi, D. M. Kumar, and K. P. J. Hemalatha, "Purification and characterization of alkaline protease with novel properties from *Bacillus cereus* strain S8," *Journal of Genetic Engineering and Biotechnology*, vol. 16, no. 2, pp. 295–304, 2018.
- [27] H. U. Rehman, A. Aman, A. Silipo, S. A. U. Qader, A. Molinaro, and A. Ansari, "Degradation of complex carbohydrate: Immobilization of pectinase from *Bacillus*



- licheniformis* KIBGE-IB21 using calcium alginate as a support,” *Food Chemistry*, vol. 139, no. 1-4, pp. 1081–1086, 2013.
- [28] I. Maalej-Achouri, M. Guerfali, A. Gargouri, and H. Belghith, “Production of xylo-oligosaccharides from agro-industrial residues using immobilized *Talaromyces thermophilus* xylanase,” *Journal of Molecular Catalysis B: Enzymatic*, vol. 59, no. 1-3, pp. 145–152, 2009.
- [29] A. Kocabas, U. Bakir, and Z. B. Ogel, “Optimization of xylanase production from *Aspergillus terreus* by using renewable agricultural lignocellulosic residues,” *New Biotechnology*, vol. 25, p. S145, 2009.
- [30] H. Liao, C. Xu, S. Tan et al., “Production and characterization of acidophilic xylanolytic enzymes from *Penicillium oxalicum* GZ-2,” *Bioresource Technology*, vol. 123, pp. 117–124, 2012.
- [31] A. Pollet, T. Belien, K. Fierens, J. A. Delcour, and C. M. Courtin, “*Fusarium graminearum* xylanases show different functional stabilities, substrate specificities and inhibition sensitivities,” *Enzyme and Microbial Technology*, vol. 44, no. 4, pp. 189–195, 2009.
- [32] F. Francis, A. Sabu, K. M. Nampoothiri, S. Ramachandram, S. Ghosh, and G. Szakacs, “Use of response surface methodology for optimizing process parameters for the production of  $\alpha$ -amylase by *Aspergillus oryzae*,” *Biochemical Engineering Journal*, vol. 15, no. 2, pp. 107–115, 2003.
- [33] S. Behnam, K. Karimi, M. Khanahmadi, and Z. Salimian, “Optimization of xylanase production by *Mucor indicus*, *Mucor hiemalis*, and *Rhizopus oryzae* through solid state fermentation,” *Biological Journal of Microorganism*, vol. 4, 2016.
- [34] S. M. M. Atalla, N. E. Ahmed, H. M. Awad, N. G. El Gamal, and A. R. El Shamy, “Statistical optimization of xylanase production, using different agricultural wastes by *Aspergillus oryzae* MN894021, as a biological control of faba bean root diseases,” *Egyptian Journal of Biological Pest Control*, vol. 30, no. 1, 2020.
- [35] F. Cui and L. Zhao, “Optimization of xylanase production from *Penicillium* sp. WXZ1 by a two-step statistical strategy: Plackett-Burman and Box-Behnken experimental design,” *International Journal of Molecular Sciences*, vol. 13, no. 8, pp. 10630–10646, 2012.
- [36] Y. Li, Z. Liu, H. Zhao, Y. Xu, and F. Cui, “Statistical optimization of xylanase production from new isolated *Penicillium oxalicum* ZH-30 in submerged fermentation,” *Biochemical Engineering Journal*, vol. 34, no. 1, pp. 82–86, 2007.
- [37] Y. Cao, D. Meng, J. Lu, and J. Long, “Statistical optimization of xylanase production by *Aspergillus niger* AN-13 under submerged fermentation using response surface methodology,” *African Journal of Biotechnology*, vol. 7, no. 5, pp. 631–638, 2008.
- [38] U. Javed, A. Aman, and S. A. U. UlQader, “Utilization of corn-cob xylan as a sole carbon source for the biosynthesis of endo-1, 4- $\beta$  xylanase from *Aspergillus niger* KIBGE-IB36,” *Biore-sources and Bioprocessing*, vol. 4, no. 1, p. 19, 2017.
- [39] Z. Azzouz, A. Bettache, I. Djinni, N. Boucherba, and S. Benallaoua, “Biotechnological production and statistical optimization of fungal xylanase by bioconversion of the lignocellulosic biomass residues in solid-state fermentation,” *Bio-mass Conversion and Biorefinery*, vol. 2020, p. 96, 2020.
- [40] L. Cunha, R. Martarello, P. M. de Souza et al., “Optimization of Xylanase Production from *Aspergillus foetidus* in Soybean Residue,” *Enzyme Research*, vol. 2018, Article ID 6597017, 7 pages, 2018.
- [41] G. Ramanjaneyulu and R. B. Rajasekhar, “Optimization of xylanase production through response surface methodology by *Fusarium* sp. BVKT R2 isolated from forest soil and its application in saccharification,” *Frontiers in Microbiology*, vol. 7, 2016.
- [42] L. Viikari, M. Alapuranen, T. Puranen, J. Vehmaanpera, and M. Siika-Aho, “Thermostable enzymes in lignocellulose hydrolysis,” *Advances in Biochemical Engineering/Biotechnology*, vol. 108, pp. 121–145, 2007.
- [43] Z. Azzouz, A. Bettache, N. Boucherba et al., “Optimization of  $\beta$ -1,4-Endoxylanase production by an *Aspergillus niger* strain growing on wheat straw and application in xylooligosaccharides production,” *Molecules*, vol. 26, no. 9, p. 2527, 2021.
- [44] M. Irfan, M. Nadeem, and Q. Syed, “One-factor-at-a-time (OFAT) optimization of xylanase production from *Trichoderma viride*-IR05 in solid-state fermentation,” *Journal of Radiation Research and Applied Sciences*, vol. 7, no. 3, pp. 317–326, 2014.
- [45] K. Y. Lee, J. A. Rowley, P. Eiselt, E. M. Moy, K. H. Bouhadir, and D. J. Mooney, “Controlling mechanical and swelling properties of alginate hydrogels independently by cross-linker type and cross-linking density,” *Macromolecules*, vol. 33, no. 11, pp. 4291–4294, 2000.