

Review

The microcirculation as a functional system

Christopher G Ellis¹, Justin Jagger² and Michael Sharpe³

¹Professor, Department of Medical Biophysics, University of Western Ontario, London, Ontario, Canada

²MD/PhD, Department of Medical Biophysics, University of Western Ontario, London, Ontario, Canada

³MD, FRCPC, Professor, Department of Anesthesia and Perioperative Medicine, Program in Critical Care Medicine, The University of Western Ontario, London Health Sciences Center, London, Ontario, Canada

Corresponding author: Christopher G Ellis, chris.ellis@uwo.ca

Published online: 25 August 2005

This article is online at <http://ccforum.com/supplements/9/S4/S3>

© 2005 BioMed Central Ltd

Critical Care 2005, **9(suppl 4)**:S3-S8 (DOI 10.1186/cc3751)

Abstract

This review examines experimental evidence that the microvascular dysfunction that occurs early in sepsis is the critical first stage in tissue hypoxia and organ failure. A functional microvasculature maintains tissue oxygenation despite limitations on oxygen delivery from blood to tissue imposed by diffusion; the density of perfused (functional) capillaries is high enough to ensure appropriate diffusion distances, and arterioles regulate the distribution of oxygen within the organ precisely to where it is needed. Key components of this regulatory system are the endothelium, which communicates and integrates signals along the microvascular network, and the erythrocytes, which directly monitor and regulate oxygen delivery. During hypovolemic shock, a functional microvasculature responds to diminish the impact of a decrease in oxygen supply on tissue perfusion. However, within hours of the onset of sepsis, a dysfunctional microcirculation is, due to a loss of functional capillary density and impaired regulation of oxygen delivery, unable to maintain capillary oxygen saturation levels and prevent the rapid onset of tissue hypoxia despite adequate oxygen supply to the organ. The mechanism(s) responsible for this dysfunctional microvasculature must be understood in order to develop appropriate management strategies for sepsis.

Introduction

One of the primary functions of the microcirculation is to ensure adequate oxygen delivery to meet the oxygen demands of every cell within an organ. In order to achieve this, the healthy microvasculature will respond to changes in metabolic demand or blood flow to the organ. However, if the microvasculature is dysfunctional, as it is in sepsis, then tissue hypoxia can occur despite supranormal oxygen delivery values. In order to understand how sepsis can result in tissue hypoxia in organs remote to the initial site of injury, we first need to understand oxygen transport and the regulation of oxygen delivery under normal physiological conditions.

Normal physiology

Diffusion limitation for oxygen

More than 80 years ago, Krogh [1] published the first oxygen transport model that described diffusion of oxygen from a single capillary cross-section into the surrounding cylinder of tissue. This model highlighted the impact of diffusion limitation on tissue oxygenation and hence explained why capillary density was greater in tissues with higher oxygen consumption rates. The model also demonstrated that it is not sufficient to simply supply an adequate amount of oxygen to the organ as a whole, but that oxygen must be distributed within the organ precisely to where it is needed.

Integration of arteriolar regulation

Arterioles, which control the vascular resistance of an organ and hence its total blood flow, are also responsible for regulating the distribution of oxygen within the organ itself. To achieve this degree of control, the response of the microvasculature to changing conditions (e.g. increased oxygen demand, reduced oxygen delivery) must be highly integrated across the entire microvascular bed [2-4]. The endothelial cells play a critical role in conducting and integrating local stimulatory signals via cell-to-cell communication along the microvascular endothelium [5-7] or by responding to changes in blood flow as signal transducers of local shear stress [8]. For example, if there is a dilatory stimulus originating in one region of the capillary bed, the vascular endothelium will conduct this stimulus to the arterioles supplying these capillaries, causing them to dilate, thus increasing blood flow. Endothelium lining larger arterioles and resistance arteries further upstream will respond to the increase in shear stress by dilating to the point that local shear stress is restored back to baseline, and thus further reducing vascular resistance.

ATP = adenosine triphosphate; ENHANCE = Extended Evaluation of Recombinant Human Activated Protein C; iNOS = inducible form of nitric oxide synthase; LPS = lipopolysaccharide; NO = nitric oxide; PROWESS = Recombinant Human Activated Protein C Worldwide Evaluation in Severe Sepsis; RBC = red blood cell; SNO = S-nitrosothiol.

Without this integrated response, a local dilatory stimulus could “steal” flow from other regions of the tissue.

Precapillary fall in oxygen saturation

Thirty-five years ago, Duling and Berne [9] reported that oxygen levels diminished along the arteriolar tree and that up to two-thirds of the oxygen delivered to a tissue has already been extracted by the time blood reaches the capillary bed. Using a variety of techniques in different organs and species, numerous researchers have documented these experimental observations [10,11]. Although we do not fully understand why there is such a large precapillary decrease in oxygen, Ellsworth and Pittman [12] provided experimental evidence to show that some of the oxygen leaving the arterioles can reoxygenate red blood cells (RBCs) flowing through nearby capillaries by diffusion. If oxygen can be transported from arterioles to capillaries, it is also likely that oxygen exchange occurs between capillaries with different oxygen levels [10], and between arterioles and venules [13]. In addition, quantitative studies of microvascular blood flow have demonstrated considerable spatial heterogeneity of capillary perfusion [14,15]. The unique rheological properties of RBC flow through branching networks of small vessels (Fahreaus effect and plasma skimming at bifurcations [16]) results in wide distributions of capillary hematocrits and RBC flow rates. The heterogeneity of microvascular hematocrit, the precapillary drop in oxygen saturation, and the diffusional exchange of oxygen among microvessels mean that blood flow by itself is not a good indicator of adequate oxygen delivery to tissue. This has important implications for the regulation of the oxygen supply, particularly during disease states and the investigation of microvascular oxygen delivery *in vivo*.

The role of RBCs in local regulation of oxygen delivery

The automatic feedback system responsible for regulating local oxygen delivery must be able to monitor and regulate oxygen delivery throughout the microvascular bed. Bergfeld and Forrester [17] were the first to demonstrate that RBCs exposed to hypoxic conditions released adenosine triphosphate (ATP). Since ATP is a potent vasodilator, they proposed that RBCs flowing through a hypoxic region could stimulate local vasodilation and an increase in blood flow. Ellsworth and colleagues [18,19] demonstrated that ATP injected into arterioles results in local vasodilation that is also conducted along the arteriole, thus demonstrating the presence of purinergic receptors (P_{2y1} and P_{2y2}) on the endothelium of these vessels. ATP binding to P_{2y1} and P_{2y2} on vascular endothelium causes vasodilation of vascular smooth muscle by inducing the endothelium to produce nitric oxide (NO) [20], prostaglandin [21], or endothelium-derived hyperpolarizing factor [22,23]. Collins and colleagues [24] demonstrated that ATP injected into postcapillary venules results in vasodilation of the feeding arteriole. Dietrich and colleagues [25] showed that isolated cerebral arterioles dilate in response to a fall in oxygen in their environment only if the arterioles are perfused with RBCs, and not if they are

perfused with a physiological solution without RBCs. They also observed that this vasodilation was caused by the efflux of ATP from the RBCs [25], and demonstrated that the oxygen-dependent release of ATP occurred rapidly enough to be physiologically relevant. Jagger and colleagues [26] have shown that ATP efflux is linearly related to hemoglobin oxygen saturation and that the regulation of glycolysis by deoxyhemoglobin in RBCs is the first step in the signaling pathway for ATP release. Also, ATP injected into larger venules results in vasodilation of the paired arteriole [27-29]. Saltin and colleagues, studying exercising human volunteers, have reported that ATP released from RBCs in response to a fall in hemoglobin oxygen saturation was responsible for regulating oxygen delivery to skeletal muscle [30,31].

In 1996, Stamler and his colleagues [32] also proposed that RBCs are responsible for regulating oxygen delivery through the transport of NO, produced in the lungs, to the periphery in the form of the bioactive compound S-nitrosothiol (SNO). SNO, reported to be a potent vasodilator, is carried by hemoglobin and released as the hemoglobin oxygen saturation falls in response to local oxygen demand. Although Stamler's group have published numerous papers supporting their theory [33,34], a number of groups have questioned the physiological role of SNO *in vivo* [35,36] as well as the accuracy of measurements of SNO from biological samples [37]. In 2003, Cosby and colleagues [38] reported that deoxyhemoglobin acts as a nitrite reductase, converting nitrite to NO, and hence making it possible for RBCs to vasodilate arterioles in response to hypoxia.

The potential for hemoglobin to play a key role in regulating vascular tone and hence oxygen delivery has generated considerable excitement [39], and has elevated the RBC from a simple carrier of oxygen to a cell ideally suited to monitor and regulate oxygen delivery across the entire microvascular bed [40].

Sepsis and microvascular dysfunction

What is the cause of organ failure in sepsis? A review article from 2000 suggests that clinical and experimental evidence “clearly indicate that microcirculatory dysfunction lies at the centre of sepsis pathogenesis” [41].

Loss of capillaries in remote organs

In 1994, Lam and colleagues [42] reported that a 24-hour peritonitis model of sepsis (cecal ligation and puncture) in rats caused a decrease in the number of perfused capillaries (i.e. decrease in functional capillary density) in skeletal muscle, with increased heterogeneity of blood flow. The loss of perfused capillaries in experimental models of sepsis has been reported in the microvasculature of intestinal villi [43,44], the diaphragm [45], and the liver [46].

Maldistribution of oxygen delivery

Using intravital video microscopy, we have studied the impact of the loss of capillary density on capillary oxygen saturation

in a fluid resuscitated, normotensive, peritonitis model of sepsis similar to that used by Lam and colleagues [42]. Using a dual-wavelength system for spectrophotometric analysis of RBC oxygen saturation, video images of microvascular blood flow were analyzed for perfused capillary density, RBC hemodynamics, and the oxygen saturation levels at the entrance and exit of the capillary bed [47]. This study confirmed the presence of stopped-, normal-, and high-flow capillaries in the same field of view. We demonstrated that the loss of capillaries (from 20% to 50% stopped flow) leads to a significant fall in oxygen saturation in normally perfused capillaries (from 60% to 20% saturation) and an increase in capillary oxygen extraction [47], as shown in Fig. 1. There was no evidence that the local oxygen regulatory system was effective in redistributing oxygen supply to offset the fall in capillary oxygen saturation levels, a result that is in accordance with the reported impaired hyperemic response to exercise observed by Lam and colleagues in the same sepsis model [42].


Hypovolemic shock versus septic shock

The situation is very different if the microvasculature is still functional and able to regulate oxygen distribution within the capillary bed. Nakajima and colleagues [44] compared microvascular perfusion in intestinal villi in mouse models of septic shock and hypovolemic shock (hemorrhage). They demonstrated that, at the same level of hypotension, hemodynamic and mucosal perfusion disorders were considerably more pronounced in endotoxin-induced hypotension than in hemorrhagic hypotension. RBC velocity was maintained in hemorrhagic shock but not during septic shock. During hypovolemic shock the microvasculature was still able to regulate microvascular perfusion, but during sepsis the regulatory response was impaired.

Experiment-based mathematical model of oxygen transport in sepsis

Our simple interpretation of the increase in oxygen extraction following a loss of perfused capillaries in sepsis was that each perfused capillary would need to support a larger volume of tissue to compensate for the loss of oxygen supply from stopped-flow capillaries [47]. However, this interpretation did not take into account the possibility of an increase in oxygen consumption rate or the potential contribution of oxygen from fast-flow capillaries. To address this limitation, Goldman and colleagues [48] developed a mathematical model of capillary oxygen delivery in a three-dimensional volume of tissue that was based on our experimental data on capillary hemodynamics and oxygen saturation in sepsis. Tissue oxygen consumption rates were adjusted in the model to yield oxygen extraction values that were consistent with our experimental measurements of capillary oxygen extraction. The model predicted that oxygen consumption increases from between two- to fourfold depending upon the severity of sepsis, and that the loss of perfused capillaries leads to significant tissue hypoxia but not

Figure 1



Oxygen saturation of red blood cells at the venous end of normally perfused capillaries versus the percentage of capillaries with stopped-flow (%CDstop) in extensor digitorum longus muscle in rat. No relationships existed in the sham animals between these parameters. In animals that underwent a 24-hour peritonitis model of sepsis (cecal ligation and perforation [CLP]), there was a decrease in oxygen saturation with increasing %CDstop (linear regression: $y = 98.8 - 1.8x$; $r^2 = 0.64$; $P < 0.05$). Reproduced with permission [47].

to anoxia. Despite the loss of capillaries and increased oxygen consumption, the model predicted that the tissue is protected from zero oxygen levels by the high-flow capillaries that supply a substantial fraction of the total oxygen delivered to the tissue. However, these high-flow capillaries do have higher venular end-oxygen saturations than normal-flow capillaries, and hence "shunt" oxygen through the capillary bed, thus elevating venular oxygen saturation levels. If the excess oxygen carried by these capillaries is uniformly distributed to all perfused capillaries, then the fall in tissue oxygen levels would be less.

Implications from experimental and mathematical models of sepsis

Based on our experiments and mathematical model, we propose that loss of perfused capillaries and impaired regulation of oxygen delivery within the microcirculation leads to a maldistribution of microvascular blood flow and tissue hypoxia early in sepsis, and that this is the first step in the progression to organ failure [49]. The tissue is still capable of extracting oxygen, but oxygen is not being delivered to where it is needed. Early in sepsis, the inability of the microvasculature to compensate for a loss of functional capillary density is the critical factor that leads to tissue hypoxia and thus organ dysfunction.

Clinical relevance

Are these results from our experimental models of sepsis clinically relevant? Using orthogonal polarization spectral imaging, De Backer and colleagues demonstrated that the density of perfused capillaries in sublingual tissue was reduced in septic patients [50], similar to what we have

observed in our animal models. Recently, this group has reported that survivors of septic shock show an improvement in perfused capillary density, but those who die have a persistent loss of perfused capillaries [51]. The loss of perfused capillaries in organs remote to the initial site of inflammation occurs in septic patients and may be an important indicator of outcomes. The key questions from an oxygen transport perspective are why does capillary blood flow stop in sepsis and why has the local oxygen regulatory system not responded to the fall in capillary oxygen saturation by distributing blood flow and oxygen to where it is needed?

Mechanisms underlying the maldistribution of oxygen delivery in sepsis

Occlusion of capillaries


There are several proposed mechanisms for the occlusion of capillaries early in sepsis: stiff leukocytes, stiff RBCs, endothelial cell swelling, and platelet/fibrin clots [49].

Piper and colleagues [52] investigated the time course (from 6–48 hours) of leukocyte rolling, adhesion, and extravasation in postcapillary venules in skeletal muscle using the same peritonitis model of sepsis as that of Lam and colleagues [42]. Although Piper and colleagues observed an increase in rolling at 24 hours, they found that leukocyte adhesion in venules was reduced due to a fall in circulating white blood cell count. However, Goddard and colleagues presented evidence from endotoxemia models of sepsis that leukocytes have a prolonged capillary transit time and are retained in the coronary capillaries of pigs [53] and rabbits [54], making the leukocyte a good candidate for occluding capillaries. Although the results of Piper and colleagues might at first seem to contradict that of Goddard and colleagues, both studies support the concept that the loss of capillaries is not due to occlusion of venules by an accumulation of leukocytes but due to the direct occlusion of capillaries.

We developed a 6-hour peritonitis model of sepsis in the rat to follow the progression of remote inflammatory injury in skeletal muscle (Fig. 2). Using this model, Bateman and colleagues [55] observed that the time course for loss of RBC deformability, excess NO production, and increased numbers of stopped-flow capillaries were correlated. Treatment of the septic rats with aminoguanidine (an inhibitor of the inducible form of NO synthase [iNOS]) to maintain plasma nitrite/nitrate levels at baseline prevented the loss of RBC deformability and the loss of perfused capillaries [55]. Our report of a subpopulation of RBCs with very low deformability at 37°C [55,56] very early in sepsis was recently confirmed [57]. These results support the role of stiff RBCs in capillary plugging [49].

There is convincing evidence that disseminated intravascular coagulation plays a central role in organ failure in sepsis [58]. Treatment of severely septic patients with activated protein C, which targets both the coagulation and inflammation

Figure 2



Functional images of the same capillary bed in the extensor digitorum longus muscle of the rat at 2.5 and 3.5 hours after induction of a peritonitis model of sepsis (cecal ligation and perforation [CLP]). The functional images were generated from captured video sequences (30 seconds) and show those capillaries through which red blood cells were flowing. At 2.5 hours after CLP, most capillaries in the field of view are perfused. One hour later, individual capillary segments from within the capillary network no longer have red blood cell flow, indicating the rapid progression of the remote injury to the microvasculature of this muscle. The procedure used for generating functional (variance) images was described by Japee *et al.* [70].

pathways in sepsis, has been shown to be effective in reducing mortality [59,60]. Although the success of the Recombinant Human Activated Protein C Worldwide Evaluation in Severe Sepsis (PROWESS) and Extended Evaluation of Recombinant Human Activated Protein C (ENHANCE) trials supports the possibility of platelet/fibrin clots impairing microvascular perfusion, experimental studies are needed to further elucidate the mechanisms of action of activated protein C on the microcirculation during the early stages of sepsis.

It is likely that a combination of these mechanisms contributes to the loss of functional capillary density in sepsis. Since the loss of capillaries in remote organs begins to occur several hours after the initial injury, and hence several hours after leukocyte activation, we speculate that activation and/or injury of the microvascular endothelium in remote organs is the critical first step leading to capillary loss.

Impaired local regulation of oxygen delivery

In addition to an impaired arteriolar response to vasoactive stimuli in animal models of sepsis [61–63], Tyml and colleagues have shown that there is impaired communication of signals between endothelial cells in culture exposed to lipopolysaccharide (LPS) [64,65] and along the vascular endothelium *in vivo* in peritonitis [62] and LPS models of sepsis [64,66]. The mechanism responsible for impaired arteriolar responsiveness to stimuli appears to be excess NO production in endothelial cells via iNOS [67]. Impaired communication along the vascular endothelium is reported to be due to an LPS-induced increase in intercellular resistance [64] that may be mediated by tyrosine phosphorylation of

connexin 43, a gap-junction molecule [68,69]. The inability of the arteriolar tree to properly integrate its response to the tissue's needs may be a significant factor in the maldistribution of oxygen delivery to tissue in sepsis. We can also speculate that erythrocyte injury in sepsis, as indicated by a loss of RBC deformability, may mean that the ability of RBCs to regulate oxygen delivery through ATP release is also impaired.

Conclusion

In metabolically active tissue, diffusion limitation places strict constraints on how far cells can be from an oxygen source. This determines not only functional capillary density but also the characteristics of the microvascular control systems. Vascular endothelium and RBCs play a significant role in coordinating the response of the arteriolar tree to changes in oxygen demand or oxygen delivery to the organ. As long as the regulatory system is functional and capillary density is sufficient, the microvasculature will deliver all available oxygen to where it is needed within an organ. In hemorrhagic shock, a "functional" microvasculature reduces the impact of a decrease in oxygen supply on tissue hypoxia by efficiently distributing oxygen to where it is needed. During the early stages of sepsis, however, the loss of capillary density and the impaired ability to regulate local oxygen delivery results in the rapid onset of tissue hypoxia despite more than adequate oxygen supply to the organ. Clearly we need to understand the mechanism(s) responsible for this dysfunctional microvasculature in order to develop appropriate management strategies for sepsis.

Competing interests

CGE received reimbursement for travel expenses and an honorarium for speaking at the Global Medical Conference in Brussels 2005 and for preparing this manuscript. CGE's sepsis research is supported by an operating grant from the Canadian Institutes for Health Research (MOP-49416), and his research on the erythrocyte role in local regulation of oxygen delivery by an operating grant from Heart and Stroke Foundation of Ontario.

Acknowledgements

The authors wish to thank Graham Fraser for his comments, and Stephanie Milkovich and Karen Donais for their assistance with the video data for Fig. 2.

References

- Krogh A: **The number and the distribution of capillaries in muscle with the calculation of the oxygen pressure necessary for supplying tissue.** *J Physiol (Lond)* 1919, **52**:409-515.
- Duling BR, Hogan RD, Langille BL, Lelkes P, Segal SS, Vatner SF, Weigelt H, Young MA: **Vasomotor control: functional hyperemia and beyond.** *Fed Proc* 1987, **46**:251-263.
- Segal SS, Damon DN, Duling BR: **Propagation of vasomotor responses coordinates arteriolar resistances.** *Am J Physiol* 1989, **256**:H832-837.
- Segal SS: **Regulation of blood flow in the microcirculation.** *Microcirculation* 2005, **12**:33-45.
- Segal SS, Duling BR: **Propagation of vasodilation in resistance vessels of the hamster: development and review of a working hypothesis.** *Circ Res* 1987, **61**:1120-1125.
- Segal SS, Duling BR: **Conduction of vasomotor responses in arterioles: a role for cell-to-cell coupling?** *Am J Physiol* 1989, **256**:H838-H845.
- Dietrich HH, Tymi K: **Capillary as a communicating medium in the microvasculature.** *Microvasc Res* 1992, **43**:87-99.
- Koller A, Kaley G: **Endothelial regulation of wall shear stress and blood flow in skeletal muscle microcirculation.** *Am J Physiol* 1991, **260**:H862-H868.
- Duling BR, Berne RM: **Longitudinal gradients in periarteriolar oxygen tension. A possible mechanism for the participation of oxygen in local regulation of blood flow.** *Circ Res* 1970, **27**:669-678.
- Ellsworth ML, Ellis CG, Popel AS, Pittman RN: **Role of microvessels in oxygen-supply to tissue.** *News Physiol Sci* 1994, **9**:119-123.
- Tsai AG, Johnson PC, Intaglietta M: **Oxygen gradients in the microcirculation.** *Physiol Rev* 2003, **83**:933-963.
- Ellsworth ML, Pittman RN: **Arterioles supply oxygen to capillaries by diffusion as well as by convection.** *Am J Physiol* 1990, **258**:H1240-H1243.
- Stein JC, Ellis CG, Ellsworth ML: **Relationship between capillary and systemic venous PO₂ during nonhypoxic and hypoxic ventilation.** *Am J Physiol* 1993, **265**:H537-H542.
- Tymi K, Ellis CG, Safranyos RG, Fraser S, Groom AC: **Temporal and spatial distributions of red cell velocity in capillaries of resting skeletal muscle, including estimates of red cell transit times.** *Microvasc Res* 1981, **22**:14-31.
- Ellis CG, Wrigley SM, Groom AC: **Heterogeneity of red blood cell perfusion in capillary networks supplied by a single arteriole in resting skeletal muscle.** *Circ Res* 1994, **75**:357-368.
- Pries AR, Secomb TW, Gaehtgens P: **Biophysical aspects of blood flow in the microvasculature.** *Cardiovasc Res* 1996, **32**:654-667.
- Bergfeld GR, Forrester T: **Release of ATP from human erythrocytes in response to a brief period of hypoxia and hypercapnia.** *Cardiovasc Res* 1992, **26**:40-47.
- McCullough WT, Collins DM, Ellsworth ML: **Arteriolar responses to extracellular ATP in striated muscle.** *Am J Physiol* 1997, **272**:H1886-H1891.
- Ellsworth ML, Forrester T, Ellis CG, Dietrich HH: **The erythrocyte as a regulator of vascular tone.** *Am J Physiol* 1995, **269**:H2155-H2161.
- Rubino A, Ralevic V, Burnstock G: **Contribution of P1-(A_{2b} subtype) and P2-purinoceptors to the control of vascular tone in the rat isolated mesenteric arterial bed.** *Br J Pharmacol* 1995, **115**:648-652.
- Needham L, Cusack NJ, Pearson JD, Gordon JL: **Characteristics of the P2 purinoceptor that mediates prostacyclin production by pig aortic endothelial cells.** *Eur J Pharmacol* 1987, **134**:199-209.
- Malmstro M, Erlinge D, Hogestatt ED, Zygmunt PM: **Endothelial P2Y receptors induce hyperpolarisation of vascular smooth muscle by release of endothelium-derived hyperpolarising factor.** *Eur J Pharmacol* 1999, **364**:169-173.
- Wihlborg A-K, Malmstro M, Eijloffsson A, Gustafsson R, Jacobson K, Erlinge D: **Extracellular nucleotides induce vasodilatation in human arteries via prostaglandins, nitric oxide and endothelium-derived hyperpolarising factor.** *Br J Pharmacol* 2003, **138**:1451-1458.
- Collins DM, McCullough WT, Ellsworth ML: **Conducted vascular responses: communication across the capillary bed.** *Microvasc Res* 1998, **56**:43-53.
- Dietrich HH, Ellsworth ML, Sprague RS, Dacey RG Jr: **Red blood cell regulation of microvascular tone through adenosine triphosphate.** *Am J Physiol Heart Circ Physiol* 2000, **278**:H1294-H1298.
- Jagger JE, Bateman RM, Ellsworth ML, Ellis CG: **Role of erythrocyte in regulating local O₂ delivery mediated by hemoglobin oxygenation.** *Am J Physiol Heart Circ Physiol* 2001, **280**:H2833-H2839.
- Hammer LW, Ligon AL, Hester RL: **ATP-mediated release of arachidonic acid metabolites from venular endothelium causes arteriolar dilation.** *Am J Physiol Heart Circ Physiol* 2001, **280**:H2616-H2622.
- Hester RL, Hammer LW: **Venular-arteriolar communication in the regulation of blood flow.** *Am J Physiol Regul Integr Comp Physiol* 2002, **282**:R1280-R1285.

29. Hammer LW, Overstreet CR, Choi J, Hester RL: **ATP stimulates the release of prostacyclin from perfused veins isolated from the hamster hindlimb.** *Am J Physiol Regul Integr Comp Physiol* 2003, **285**:R193-R199.
30. Gonzalez-Alonso J, Richardson RS, Saltin B: **Exercising skeletal muscle blood flow in humans responds to reduction in arterial oxyhaemoglobin, but not to altered free oxygen.** *J Physiol (Lond)* 2001, **530**:331-341.
31. Gonzalez-Alonso J, Olsen DB, Saltin B: **Erythrocyte and the regulation of human skeletal muscle blood flow and oxygen delivery: role of circulating ATP.** *Circ Res* 2002, **91**:1046-1055.
32. Jia L, Bonaventura C, Bonaventura J, Stamler JS: **S-nitrosohaemoglobin: a dynamic activity of blood involved in vascular control.** *Nature* 1996, **380**:221-226.
33. Stamler JS, Jia L, Eu JP, McMahon TJ, Demchenko IT, Bonaventura J, Gernert K, Piantadosi CA: **Blood flow regulation by S-nitrosohemoglobin in the physiological oxygen gradient.** *Science* 1997, **276**:2034-2037.
34. Liu L, Yan Y, Zeng M, Zhang J, Hanes MA, Ahearn G, McMahon TJ, Dickfeld T, Marshall HE, Que LG, Stamler JS: **Essential roles of S-nitrosothiols in vascular homeostasis and endotoxic shock.** *Cell* 2004, **116**:617-628.
35. Patel RP, Hogg N, Spencer NY, Kalyanaraman B, Matalon S, Darley-Usmar VM: **Biochemical characterization of human S-nitrosohemoglobin. Effects on oxygen binding and transnitrosation.** *J Biol Chem* 1999, **274**:15487-15492.
36. Gladwin MT, Schechter AN: **NO contest: nitrite versus S-nitroso-hemoglobin.** *Circ Res* 2004, **94**:851-855.
37. Gladwin MT, Lancaster JR Jr, Freeman BA, Schechter AN: **Nitric oxide's reactions with hemoglobin: a view through the SNO-storm.** *Nat Med* 2003, **9**:496-500.
38. Cosby K, Partovi KS, Crawford JH, Patel RP, Reiter CD, Martyr S, Yang BK, Waclawiw MA, Zalos G, Xu X, et al.: **Nitrite reduction to nitric oxide by deoxyhemoglobin vasodilates the human circulation.** *Nat Med* 2003, **9**:1498-1505.
39. Patel RP, Gladwin MT: **Physiologic, pathologic and therapeutic implications for hemoglobin interactions with nitric oxide.** *Free Radic Biol Med* 2004, **36**:399-401.
40. Singel DJ, Stamler JS: **Chemical physiology of blood flow regulation by red blood cells: the role of nitric oxide and S-nitroso-hemoglobin.** *Annu Rev Physiol* 2005, **67**:99-145.
41. Lehr HA, Bittinger F, Kirkpatrick CJ: **Microcirculatory dysfunction in sepsis: a pathogenetic basis for therapy?** *J Pathol* 2000, **190**:373-386.
42. Lam C, Tymi K, Martin C, Sibbald W: **Microvascular perfusion is impaired in a rat model of normotensive sepsis.** *J Clin Invest* 1994, **94**:2077-2083.
43. Farquhar I, Martin CM, Lam C, Potter R, Ellis CG, Sibbald WJ: **Decreased capillary density *in vivo* in bowel mucosa of rats with normotensive sepsis.** *J Surg Res* 1996, **61**:190-196.
44. Nakajima Y, Baudry N, Duranteau J, Vicaute E: **Microcirculation in intestinal villi: a comparison between hemorrhagic and endotoxin shock.** *Am J Respir Crit Care Med* 2001, **164**:1526-1530.
45. Boczkowski J, Vicaute E, Aubier M: ***In vivo* effects of Escherichia coli endotoxemia on diaphragmatic microcirculation in rats.** *J Appl Physiol* 1992, **72**:2219-2224.
46. Gundersen Y, Corso CO, Leiderer R, Dorger M, Lilleaasen P, Aasen AO, Messmer K: **Use of selective and nonselective nitric oxide synthase inhibitors in rat endotoxemia: effects on hepatic morphology and function.** *Shock* 1997, **8**:368-372.
47. Ellis CG, Bateman RM, Sharpe MD, Sibbald WJ, Gill R: **Effect of a maldistribution of microvascular blood flow on capillary O₂ extraction in sepsis.** *Am J Physiol Heart Circ Physiol* 2002, **282**:H156-H164.
48. Goldman D, Bateman RM, Ellis CG: **Effect of sepsis on skeletal muscle oxygen consumption and tissue oxygenation: interpreting capillary oxygen transport data using a mathematical model.** *Am J Physiol Heart Circ Physiol* 2004, **287**:H2535-H2544.
49. Bateman R, Sharpe M, Ellis C: **Bench-to bedside review: Microvascular dysfunction in sepsis – hemodynamics, oxygen transport, and nitric oxide.** *Crit Care* 2003, **7**:359-373.
50. De Backer D, Creteur J, Preiser J-C, Dubois M-J, Vincent J-L: **Microvascular blood flow is altered in patients with sepsis.** *Am J Respir Crit Care Med* 2002, **166**:98-104.
51. Sakr Y, Dubois MJ, De Backer D, Creteur J, Vincent JL: **Persistent microcirculatory alterations are associated with organ failure and death in patients with septic shock.** *Crit Care Med* 2004, **32**:1825-1831.
52. Piper RD, Pitt-Hyde ML, Anderson LA, Sibbald WJ, Potter RF: **Leukocyte activation and flow behavior in rat skeletal muscle in sepsis.** *Am J Respir Crit Care Med* 1998, **157**:129-134.
53. Goddard CM, Allard MF, Hogg JC, Herbertson MJ, Walley KR: **Prolonged leukocyte transit time in coronary microcirculation of endotoxemic pigs.** *Am J Physiol Heart Circ Physiol* 1995, **269**:H1389-H1397.
54. Goddard CM, Poon BY, Klut ME, Wiggs BR, van Eeden SF, Hogg JC, Walley KR: **Leukocyte activation does not mediate myocardial leukocyte retention during endotoxemia in rabbits.** *Am J Physiol Heart Circ Physiol* 1998, **275**:H1548-H1557.
55. Bateman RM, Jagger JE, Sharpe MD, Ellsworth ML, Mehta S, Ellis CG: **Erythrocyte deformability is a nitric oxide-mediated factor in decreased capillary density during sepsis.** *Am J Physiol Heart Circ Physiol* 2001, **280**:H2848-H2856.
56. Jagger JE, Ellis CG, Sibbald WJ, Eichelbronner O: **Measurement temperature plays a pivotal role in the distribution of erythrocyte deformability after LPS.** *Biorheology* 2001, **38**:439-448.
57. Condon MR, Kim JE, Deitch EA, Machiedo GW, Spolarics Z: **Appearance of an erythrocyte population with decreased deformability and hemoglobin content following sepsis.** *Am J Physiol Heart Circ Physiol* 2003, **284**:H2177-H2184.
58. Levi M, de Jonge E, van der Poll T: **Sepsis and disseminated intravascular coagulation.** *J Thromb Thrombolysis* 2003, **16**:43-47.
59. Bernard G, Vincent J, Laterre P, LaRosa S, Dhainaut J, Lopez-Rodriguez A, Steingrub J, Garber G, Helterbrand J: **Efficacy and safety of recombinant human activated protein C for severe sepsis.** *N Engl J Med* 2001, **344**:699-709.
60. Bernard G, Macias W, Joyce D, Williams M, Bailey J, Vincent J-L: **Safety assessment of drotrecogin alfa (activated) in the treatment of adult patients with severe sepsis.** *Crit Care* 2003, **7**:155-163.
61. Hollenberg SM, Tangora JJ, Piotrowski MJ, Easington C, Parrillo JE: **Impaired microvascular vasoconstrictive responses to vasopressin in septic rats.** *Crit Care Med* 1997, **25**:869-873.
62. Tymi K, Yu J, McCormack DG: **Capillary and arteriolar responses to local vasodilators are impaired in a rat model of sepsis.** *J Appl Physiol* 1998, **84**:837-844.
63. Hollenberg SM, Broussard M, Osman J, Parrillo JE: **Increased microvascular reactivity and improved mortality in septic mice lacking inducible nitric oxide synthase.** *Circ Res* 2000, **86**:774-778.
64. Tymi K, Wang X, Lidington D, Ouellette Y: **Lipopolysaccharide reduces intercellular coupling *in vitro* and arteriolar conducted response *in vivo*.** *Am J Physiol Heart Circ Physiol* 2001, **281**:H1397-H1406.
65. Lidington D, Ouellette Y, Tymi K: **Endotoxin increases intercellular resistance in microvascular endothelial cells by a tyrosine kinase pathway.** *J Cell Physiol* 2000, **185**:117-125.
66. Lidington D, Ouellette Y, Li F, Tymi K: **Conducted vasoconstriction is reduced in a mouse model of sepsis.** *J Vasc Res* 2003, **40**:149-158.
67. Wu F, Wilson JX, Tymi K: **Ascorbate inhibits iNOS expression and preserves vasoconstrictor responsiveness in skeletal muscle of septic mice.** *Am J Physiol Regul Integr Comp Physiol* 2003, **285**:R50-R56.
68. Lidington D, Ouellette Y, Tymi K: **Communication of agonist-induced electrical responses along 'capillaries' *in vitro* can be modulated by lipopolysaccharide, but not nitric oxide.** *J Vasc Res* 2002, **39**:405-413.
69. Lidington D, Tymi K, Ouellette Y: **Lipopolysaccharide-induced reductions in cellular coupling correlate with tyrosine phosphorylation of connexin 43.** *J Cell Physiol* 2002, **193**:373-379.
70. Japee SA, Ellis CG, Pittman RN: **Flow visualization tools for image analysis of capillary networks.** *Microcirculation* 2004, **11**:39-54.