

Knorr-Rabe partial reduction of pyrroles: Application to the synthesis of indolizidine alkaloids

Brendon S Gourlay¹, John H Ryan² and Jason A Smith*¹

Address: ¹School of Chemistry, University of Tasmania, Hobart, Australia and ²CSIRO Division of Molecular and Health Technologies, Bag 10, Clayton South, Victoria, Australia

Email: Brendon S Gourlay - gourlayb@utas.edu.au; John H Ryan - Jack.Ryan@csiro.au; Jason A Smith* - Jason.Smith@utas.edu.au

* Corresponding author

Published: 15 January 2008

Received: 8 October 2007

Beilstein Journal of Organic Chemistry 2008, 4:3 doi:10.1186/1860-5397-4-3

Accepted: 15 January 2008

This article is available from: <http://bjoc.beilstein-journals.org/content/4/1/3>

© 2008 Gourlay et al; licensee Beilstein-Institut

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract


Background: The Birch reduction of electron rich pyrroles does not occur readily. However, dissolving metal reduction with zinc under acidic conditions gives 3-pyrrolines (2,5-dihydropyrroles) in reasonable yield. This dissolving metal reduction was first reported by Knorr and Rabe in 1901 but since then has only been reported for the reduction of electron rich pyrroles.

Results: The partial reduction of bicyclic α -ketopyrrole derivatives has been performed under dissolving metal conditions with zinc and hydrochloric acid to give excellent yields of hexahydroindolizidines. This reduction method has been utilised for the diastereoselective synthesis of 5-alkylindolizidines and the stereoselectivity obtained is opposite to that of catalytic hydrogenation.

Conclusion: An efficient stereoselective synthesis of indolizidine alkaloids has been developed from α -ketopyrrole intermediates using a modified version of Knorr and Rabe's pyrrole reduction.

Background


The Birch reaction for the dearomatisation of aromatic substrates is an extremely practical and important tool for synthetic chemists and is used widely as a key step for the synthesis of natural products and molecules of biological interest.[1] However, the partial reduction of pyrrole is difficult as the high electron density of these aromatic heterocycles inhibits the addition of an electron, the first step of a Birch reaction.[2] Donohoe has shown that the partial reduction of pyrroles is possible but this process generally requires the presence of at least two electron withdrawing groups that reduce the electron density of the heterocycle such that reasonable yields of the 3-pyrrolines are obtained.[3] This method was recently exploited for the elegant synthesis of the pyrrolidine alkaloid (\pm)-1-epi-australine 3 (Scheme 1).[4]



Scheme 1: Donohoe's approach to (\pm)-1-epi-australine utilising a modified Birch reduction

During our studies towards the synthesis of indolizidine alkaloids we required bicyclic 3-pyrrolines and chose to explore accessing these intermediates *via* partial reduction of the corresponding pyrrole derivatives. These substrates were far more electron rich than those of Donohoe and thus not amenable to Birch reduction methodology.


Therefore, we turned to an underutilised reaction that was reported by Knorr and Rabe [5] in 1901 and has only been reported a handful of times since. [6-9] The method employs powdered zinc in an acid media to give 3-pyrrolines, presumably by protonation of the pyrrole to give an iminium ion which is then reduced. It has been shown that reaction of 2,5-dialkylpyrroles gives predominantly the *trans* 3-pyrroline isomer (Scheme 2). [7-9]



Scheme 2: Reaction conditions i) Zn, HCl (aq)

Results and Discussion

The synthetic plan that we adopted was to construct a bicyclic pyrrole derivative by exploiting the natural reactivity of pyrrole and then to partially reduce the heterocyclic core (Scheme 3). The synthesis started with formation of the γ -pyrrolic ester 7 in high yield using an improved Clauson-Kaas synthesis, [10] followed by boron tribromide mediated cyclisation to give the known bicyclic ketone 8. [11] Upon submission of this α -ketopyrrole 8 to the modified conditions reported by Andrews and McElvain (slow addition of HCl to the substrate and Zn at 0–10°C) [5,9] we observed no reaction and starting material was returned. However, when zinc and concentrated HCl were added in small portions to a hot solution of the α -ketopyrrole in methanol over ~10 minutes the starting material was consumed to give the hexahydroindolizidine 9 as the only observable product in ~80% yield. The chemoselectivity using these modified conditions is noteworthy while the carbonyl group is fully reduced the pyrrole group is selectively and partially reduced to the 3-pyrroline. This result was confirmed by comparison of the spectral data with that reported by Huxtable who prepared 9 as an intermediate in the synthesis of lentiginosine. [12]




Scheme 3: Reaction conditions: i) ref. [10] ii) ref. [11] iii) Zn, conc. HCl(aq)

For the partial reduction of electron rich pyrroles reported previously, over reduction to give pyrrolidines is a problematic side-reaction. For example, Andrews and McElvain kept the reaction temperature below 10°C to limit pyrrolidine formation. Under our conditions, starting with the α -ketopyrrole, there was no indication of pyrro-


lidine formation. The loss of the keto group means that the product is the same as that would be obtained by reduction of the parent bicyclic pyrrole 13. The reduction of the carbonyl group resembles that of a Clemmensen reduction; however, amalgamated zinc is required for Clemmensen reaction. [13]

There are several possible mechanisms for this transformation, however, we propose the first step involves protonation of the carbonyl group to give a conjugated iminium ion 10 (Scheme 4). This species would undergo a two-electron reduction process, with associated protonation to give the α -hydroxy pyrrole 11. Acid-promoted dehydration of 11 would afford a second iminium ion 12 which could undergo further reduction and protonation to give pyrrole 13. The pyrrole could then be protonated to give a third iminium ion 14 and reduction would then give rise to the product 9.



Scheme 4: Potential mechanism for α -ketopyrrole reduction


Our reaction conditions are much harsher than those previously reported, and yet we do not see pyrrolidine products and this suggests that an alternative pathway is in operation. One possibility is that the intermediate 12 could undergo reduction to give the final product directly without the formation of the pyrrole intermediate 13 (Scheme 5).



Scheme 5: Alternative reduction pathway


To test these hypotheses we reduced the ketone 8 with NaBH₄ to give the unstable α -hydroxy pyrrole 11 which was then immediately subjected to the reduction conditions. The same result was obtained giving the 3-pyrroline 9 which lends support to the suggestion that 11 is an intermediate in the reaction. When pyrrole 13 was reacted under the same conditions 9 was formed but the ¹H NMR spectrum also showed some starting material remained. The fact that the pyrrole 13 was not observed in the reduction products from α -ketopyrrole 8 lends the support to the suggestion of an alternative pathway. At the present time the intermediacy of 13 cannot be ruled out for the reduction of ketone 8 and alcohol 11.

Due to the facile and rapid reaction of the α -ketopyrrole **8** we explored the potential tandem α -ketopyrrole reduction/catalytic hydrogenation as an alternative to catalytic hydrogenation. The catalytic hydrogenation of 5-substituted tetrahydroindolizidines proceeds with high diastereoselectivity [14,15] and has also been exploited for the synthesis of numerous indolizidine alkaloids.[16,17] The presence of a substituent at C-5 directs the hydrogenation at C-8a from the opposite, least hindered face, to give the *cis* derivative (Scheme 6).



Scheme 6: Catalytic hydrogenation


We were interested in the stereochemical outcome for C-8a using the modified Knorr-Rabe zinc reduction and synthesised the known 5-methyl derivative **18** (Scheme 7) as a model. The methyl ester of (\pm)-alanine was subjected to the modified Clauson-Kaas pyrrole synthesis to give an α -pyrrolic ester **19** which was subjected to two carbon homologation by ester reduction with DIBAL-H followed by an *in situ* Wadsworth-Emmons olefination.[18] The alkene **20** was then hydrogenated to the γ -pyrrolic ester **21** and cyclised to give α -ketopyrrole **18** in 67% overall yield from **19**. The modified Knorr-Rabe reduction of **18** gave the desired pyrroline **22** in near quantitative yield as a 9:1 mixture of diastereomers. The volatility of the compound meant that for practical purposes it was isolated as the hydrochloride salt by adding concentrated HCl to the organic extract before evaporation. Catalytic hydrogenation of the hydrochloride salt of the pyrroline gave a corresponding mixture of isomers of 5-methylindolizidine **23** but to our surprise the *trans* isomer was the major diastereomer. The stereochemical assignment of the major and minor isomers was confirmed by comparison of the ^{13}C nmr spectra with the reported spectra for both previously synthesised isomers.[19] The resonance of the carbon signals for C-8a, C-5 and C-3 are diagnostic with these carbons for the major isomer resonating at 54.9, 50.2 and 49.1 ppm respectively. This compares to 54.5, 50.0 and 49.2 ppm for the *trans* isomer and 64.8, 58.9 and 51.8 ppm for the *cis* isomer as reported in the literature.[19] This result indicates that the major product **22** from the modified Knorr-Rabe zinc reduction has the opposite C-5/C-8a stereochemistry to that typically obtained by catalytic hydrogenation.




Scheme 7: Reaction conditions: i) DIBAL-H, CH_2Cl_2 , -78°C , 1 h then triethylphosphonoacetate, NaH, THF, -78°C - rt ii) H_2 (40 psi), Pd/C, EtOH iii) BBr_3 , CH_2Cl_2 , 0°C , 10 min iv) Zn, conc. HCl(aq) v) H_2 (40 psi), Pd/C, EtOH, 2 M HCl

To explain this result we propose that the zinc complexation to the less hindered face of the indolizidine causes protonation to occur on the same side as the C-5 substituent, which results in the *trans* stereochemistry between C-5 and C-8a.

A beneficial outcome from these observations is that one can now reduce bicyclic intermediates like **18** stereoselectively to enter either diastereomeric series. Corvo has reported the synthesis of the proposed structure of indolizidine 167B by the catalytic hydrogenation of (-)-**24**, (Scheme 8)[17] and herein we report the racemic synthesis of its epimer (Scheme 9). We have reported the synthesis of the bicyclic ketone (\pm)-**24** [18] and subsection of this α -ketopyrrole to the modified Knorr-Rabe reduction conditions gave the crude 3-pyrroline **26** which was immediately subjected to catalytic hydrogenation to yield a 9:1 mixture of (\pm)-*epi*-indolizidine 167B (*trans*-(\pm)-**27**) and (\pm)-indolizidine 167B (*cis*-(\pm)-**25**) in 91% overall yield from **24**. As for the 5-methyl derivative the spectral data of the *trans* isomer **27** was dramatically different to that of the *cis* isomer **25** and is consistent with that reported previously.[20] Therefore, this method extends the flexibility of bicyclic pyrroles as intermediates for the synthesis of indolizidine alkaloids, as diastereomeric targets can be accessed simply by the choice of reagent system for reduction of the pyrrole nucleus.



Scheme 8: i) ref. [17]



Scheme 9: i) CH_3OH , Zn, conc. HCl(aq) ii) H_2 (40 psi), Pd/C, EtOH, 2 M HCl

Conclusion

In conclusion, we have discovered a modified method for the Knorr-Rabe partial reduction of electron rich pyrroles which is effective for the reduction of bicyclic α -ketopyrroles to the corresponding 3-pyrroline or hexahydroindolizidine derivatives. The reduction occurs with high diastereoselectivity with 5-alkyl derivatives and gives the opposite diastereoselectivity to that of direct catalytic hydrogenation. This complimentary method allows for the synthesis of both diastereomers of indolizidine 167B from a late-stage common intermediate.

See additional file1 for full experimental data.

Additional material

Additional file 1

Experimental details for "Knorr-Rabe partial reduction of pyrroles: Application to the synthesis of indolizidine alkaloids." Experimental details which includes experimental procedures and spectroscopic data

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1860-5397-4-3-S1.CDX>]

Acknowledgements

The authors thank the University of Tasmania and CSIRO for financial support. The authors would like to thank Peter P. Molesworth of UTAS for a sample of compound **13**. JHR thanks University of Tasmania for a Woolmers Lectureship. BSG is grateful for an Australian Postgraduate Award and a CSIRO Postgraduate Top-up Scholarship.

References

- Subba Rao GSR: *Pure and Applied Chemistry* 2003, **75**:1443-1451.
- Donohoe TJ, Guyo PM: *J Org Chem* 1996, **61**:7664-7665.
- Donohoe TJ, Guyo PM, Harji RR, Helliwell M: *Tetrahedron Lett* 1998, **39**:3075-3078.
- Donohoe TJ, Sintim HO: *Org Lett* 2004, **6**:2003-2006.
- Knorr L, Rabe P, Bufler H, Jakobi C: *Chem Ber* 1901, **34**:3491-3502.
- Hall SS, Schumacher DP: *J Am Chem Soc* 1982, **104**:6076-6080.
- Krawczyk AR, Wrobel JT: *Polish Journal of Chemistry* 1981, **55**:1363-1367.
- Evans GG: *J Am Chem Soc* 1951, **73**:5230-5234.
- Andrews LH, McElvain SM: *J Am Chem Soc* 1929, **51**:887-892.
- Gourlay BS, Molesworth PP, Ryan JH, Smith JA: *Tetrahedron Lett* 2006, **47**:799-801.

- Amos RIJ, Gourlay BS, Molesworth PP, Smith JA, Sprod OR: *Tetrahedron* 2005, **61**:8226-8230.
- Colegate SM, Dorling PR, Huxtable CR: *Aust J Chem* 1984, **37**:1503-1509.
- Clemmensen E: *Chem Ber* 1914, **47**:51-63.
- Jefford CW, Tang Q, Zaslona A: *J Am Chem Soc* 1991, **113**:3513-3518.
- Bond TJ, Jenkins R, Taylor PC: *Tetrahedron Lett* 1994, **35**:9263-9266.
- Jefford CW, Sienkiewicz K, Thornton SR: *Helv Chim Acta* 1995, **78**:1511-1524.
- Corvo MC, Pereira MMA: *Tetrahedron Lett* 2002, **43**:455-458.
- Gourlay BS, Little I, Ryan JH, Smith JA: *Natural Product Communications* 2006, **1**:831-837.
- Tehrani KA, D'hooghe M, Kimpe ND: *Tetrahedron* 2003, **59**:3099-3108.
- Polniaszek RP, Belmont SE: *J Org Chem* 1990, **55**:4688-4693.