

Taxonomic revision of *Aspergillus* section *Clavati* based on molecular, morphological and physiological data

J. Varga^{1,3}, M. Due², J.C. Frisvad² and R.A. Samson¹

¹CBS Fungal Biodiversity Centre, Uppsalalaan 8, NL-3584 CT Utrecht, The Netherlands; ²BioCentrum-DTU, Building 221, Technical University of Denmark, DK-2800 Kgs. Lyngby, Denmark; ³Department of Microbiology, Faculty of Science and Informatics, University of Szeged, H-6701 Szeged, P.O. Box 533, Hungary

*Correspondence: János Varga, j.varga@cbs.knaw.nl

Abstract: *Aspergillus* section *Clavati* has been revised using morphology, secondary metabolites, physiological characters and DNA sequences. Phylogenetic analysis of β -tubulin, ITS and calmodulin sequence data indicated that *Aspergillus* section *Clavati* includes 6 species, *A. clavatus* (synonyms: *A. apicalis*, *A. pallidus*), *A. giganteus*, *A. rhizopodus*, *A. longivesica*, *Neocarpenteles acanthosporus* and *A. clavatonanicus*. *Neocarpenteles acanthosporus* is the only known teleomorph of this section. The sister genera to *Neocarpenteles* are *Neosartorya* and *Dichotomomyces* based on sequence data. Species in *Neosartorya* and *Neocarpenteles* have anamorphs with green conidia and share the production of tryptoquivalins, while *Dichotomomyces* was found to be able to produce gliotoxin, which is also produced by some *Neosartorya* species, and tryptoquivalines and tryptoquivalones produced by members of both section *Clavati* and *Fumigati*. All species in section *Clavati* are alkalitolerant and acidotolerant and they all have clavate conidial heads. Many species are coprophilic and produce the effective antibiotic patulin. Members of section *Clavati* also produce antafumicin, tryptoquivalines, cytochalasins, sarcins, dehydrocarolic acid and kotanins (orlandin, desmethylkotanin and kotanin) in species specific combinations. Another species previously assigned to section *Clavati*, *A. ingratus* is considered a synonym of *Hemicarpenteles paradoxus*, which is phylogenetically very distantly related to *Neocarpenteles* and section *Clavati*.

Key words: Ascomycetes, *Aspergillus* section *Clavati*, β -tubulin, calmodulin, *Dichotomomyces*, Eurotiales, *Hemicarpenteles*, ITS, mycotoxin, *Neocarpenteles*, patulin, polyphasic taxonomy.

INTRODUCTION

Species in *Aspergillus* section *Clavati* are alkalitolerant, often dung-borne species that produce several mycotoxins such as patulin (Varga *et al.* 2003), cytochalasins (Demain *et al.* 1976; Steyn *et al.* 1982), tryptoquivalines and tryptoquivalones (Clardy *et al.* 1975; Büchi *et al.* 1977), and other bioactive natural products, including the sarcins (Cole & Cox 1981; Lin *et al.* 1994). Weisner (1942) and Bergel *et al.* 1943 found that *A. clavatus* produces patulin, and Florey *et al.* (1944) reported on patulin production by *Aspergillus giganteus* in 1944. Clavatol (Bergel *et al.* 1944) and ascladiol (Suzuki *et al.* 1971) were also isolates from *A. clavatus* as antibiotics. Cytochalasin E and K are also mycotoxins known from *Aspergillus clavatus* (Demain *et al.* 1976). *A. clavatus* was also reported to produce kotanin and xanthocillin X dimethylether (Büchi *et al.* 1977). Among the mycotoxins produced, patulin is receiving world-wide attention due to its frequent occurrence in apple juices (Harrison 1989; Beretta *et al.* 2000). *Aspergillus clavatus*, *A. giganteus* and *Neocarpenteles acanthosporus* isolates also produce ribotoxins, which are promising tools for immunotherapy of cancer (Martinez-Ruiz *et al.* 1999; Varga *et al.* 2003). The economically most important species of the section, *A. clavatus* is possibly a cosmopolitan fungus. It can be isolated mainly from soil and dung, but also occurs on stored products (mainly cereals) with high moisture content, e.g. inadequately stored rice, corn and millet (Flannigan & Pearce 1994). *A. clavatus* isolates appear to be particularly well adapted for growth during malting (Flannigan & Pearce 1994). *A. clavatus* was found to be responsible for an extrinsic allergic alveolitis known as malt worker's lung, and in cases of mycotoxicoses of animals fed with by-products of malting

(Flannigan & Pearce 1994; Lopez-Diaz & Flannigan 1997). The toxic syndromes observed in animals were suggested to result from the synergistic action of various mycotoxins produced by this species (Flannigan & Pearce 1994). Several species of section *Clavati* have phototrophic long conidiophores at temperatures around 20–23 °C (Fennell & Raper 1955; Trinci & Banbury 1967; Sarbhoy & Elphick 1968; Huang & Raper 1971; Yaguchi *et al.* 1993).

Aspergillus subgenus *Fumigati* section *Clavati* (Gams *et al.* 1985; Peterson 2000), formerly the *Aspergillus clavatus* group was recognised by Thom & Church (1926) with two species, *A. clavatus* and *A. giganteus*. *A. clavatonanicus* was added by Batista *et al.* (1955). After Raper & Fennell (1965) published their monograph on aspergilli, several new species or varieties assigned to section *Clavati* were described. These were summarised by Samson (1979), who recognised *A. longivesica* (Huang & Raper 1971) as the fourth species within the section. None of these have known teleomorphs. Another species, *A. rhizopodus* (Rai *et al.* 1975) was treated by Samson (1979) as a synonym of *A. giganteus*. *A. pallidus* Kamyschko has been treated as a white-spored synonym of *A. clavatus* by several authors (Peterson 2000; Varga *et al.* 2003). *A. acanthosporus* (Udagawa & Takada (1971), placed in subgenus *Ornati* (Samson 1979), was shown by Peterson (2000) to be more closely related to section *Clavati* than to section *Ornati*. Also, their major ubiquinone systems point in this direction as section *Clavati* and *A. acanthosporus* have Q10, while *H. ornatus* has Q9 ubiquinones (Tamura *et al.* 1999). Although its teleomorph was originally placed into the *Hemicarpenteles* genus, recently Udagawa & Uchiyama (2002) proposed the new ascomycete genus *Neocarpenteles* to accommodate this species, and excluded *N. acanthosporus* from section *Ornati*. Similar conclusions were drawn

Table 1. The *Aspergillus* section *Clavati* isolates examined in this study.

Species	Strain No.	Origin
<i>A. clavatus</i>	CBS 104.45	ATCC 9600; Czech Republic, Pribram
	CBS 105.45	Church, No. Ac 87
	CBS 106.45	<i>Humulus lupulus</i> (Cannabaceae), G. Smith
	CBS 114.48	Culture contaminant, Netherlands
	CBS 513.65 ^T	ATCC 1007; IMI 015949; NRRL 1; Thom 107
	CBS 514.65	ATCC 10058; IMI 321306; NRRL 4; Thom 4754.3
	CBS 470.91	Toxic feed pellets, Hungary
	CBS 116685	Milled rice, Netherlands
	CBS 118451	Medicine, Germany
	DTO 6-F8	Air, ciabatta factory, Netherlands
	DTO 27-C2	Bakery, Netherlands
	SZMC 0918	Soil, Hungary
	SZMC JV4	Stored wheat, Hungary
	SZMC JV1.1	Human mucosa, Hungary
	IMI 358435	Feed pellet, Hungary
<i>A. giganteus</i>	CBS 117.45	IMI 024256; P. Biourge
	CBS 119.48	H. Burgeff, No. 382, Germany
	CBS 118.49	Wood of ship (<i>Virola surinamensis</i>), Suriname
	CBS 122.53	Tail borad, Nigeria
	CBS 117.56	Wood in swimming pool, Netherlands
	CBS 101.64	Unknown, Poland
	CBS 515.65 ^T	ATCC 16439; IMI 235601; NRRL 7974; mouse dung, U.S.A.
	CBS 526.65	ATCC 10059; IMI 227678; NRRL 10; Thom 5581.13A
	CBS 112.27	A. Blochwitz
<i>A. rhizopodus</i>	CBS 450.75 ^T	Usar soil, India, Lucknow
	IMI 351309	Soil, Yugoslavia
<i>A. pallidus</i>	CBS 344.67 ^T	ATCC 18327; IMI 129967; soil, Moldova
	SZMC JV6	Culture contaminant, Hungary
<i>A. clavatonanicus</i>	CBS 474.65 ^T	ATCC 12413; IMI 235352; WB 4741; finger nail lesion, Brazil
<i>A. longivesica</i>	CBS 530.71 ^T	ATCC 22434; IMI 156966; soil, Nigeria
	CBS 187.77	Soil, Ivory Coast, Tai
<i>A. apicalis</i>	CBS 236.81 ^T	Wheat bran, India
<i>N. acanthosporus</i>	CBS 558.71 ^T	Solomon Islands, Bougainville Island
	CBS 445.75	Solomon Islands, Bougainville Island, Buin, Malapita
	CBS 446.75	Solomon Islands, Bougainville Island, Buin, Batubatuai
	CBS 447.75	Solomon Islands, Bougainville Island, Kieta
<i>D. cepjii</i> var. <i>cejpii</i>	CBS 761.96	spent mushroom compost, Netherlands
<i>D. cepjii</i> var. <i>cejpii</i>	CBS 779.70	Soil, Cincinatti, U.S.A.
<i>D. cepjii</i> var. <i>cejpii</i>	CBS 100192	Soil, Bratislava, Slovakia
<i>D. cepjii</i> var. <i>cejpii</i>	CBS 474.77	Soil, Egypt
<i>D. cepjii</i> var. <i>cejpii</i>	CBS 780.70	Pasturised milk, Cincinatti, U.S.A.
<i>D. cepjii</i> var. <i>cejpii</i>	CBS 397.68	Soil, South Africa
<i>D. cepjii</i> var. <i>cejpii</i>	CBS 345.68	rhizosphere of <i>Hordeum vulgare</i> , Pakistan
<i>D. cepjii</i> var. <i>cejpii</i>	CBS 159.67	Soil, Kominato, Japan
<i>D. cepjii</i> var. <i>cejpii</i>	CBS 157.66 ^T	Orchard soil, Moldova, near Tiraspol
<i>D. cepjii</i> var. <i>spinusus</i>	CBS 219.67 ^T	Soil, Kyoto, Japan

by Varga *et al.* (2003) based on sequence analysis of the internal transcribed spacer regions and the 5.8 S rRNA gene (ITS region) of isolates belonging to *Aspergillus* section *Clavati*. Another species, *A. apicalis* Mehrotra & Basu (1976) (as *A. apica*), was placed in section *Ornati* by Samson (1979) because of morphological similarities to *H. paradoxus* (small clavate blue green aspergilla). Finally, *A. ingratus* has been described by Yaguchi *et al.* (1993), who stated that this sclerotium producing species belonged to section *Clavati*.

In this study, we examined the taxonomic assignment of these alkalitolerant species characterised by clavate aspergilla using molecular, morphological and chemotaxonomical methods. We also examined the relationships among teleomorphs of *Aspergillus* subgenus *Fumigati*, including *Neocarpenteles* and *Neosartorya* species to the *Dichotomomyces* genus using molecular approaches. Although the anamorphs of *Dichotomomyces* belong to the *Polypaecilum*, ascomata and ascospores of *Dichotomomyces* species have a similar morphology as those of *Neosartorya* and *Neocarpenteles* (Samson RA, unpubl. data).

MATERIALS AND METHODS

Source of microorganisms

The fungi examined included all species allocated to *Aspergillus* section *Clavati*, and some species assigned to section *Ornati* with clavate aspergilla (the *Aspergillus ornatus* group), which could possibly be related to *A. clavatus*. The strains examined are listed in Table 1.

Morphology and physiology

The strains (Table 1) were grown for 7 d as 3-point inoculations on Czapek agar (CZA), Czapek yeast autolysate agar (CYA), creatine sucrose agar (CREA) and malt extract agar (MEA) at 25 °C in artificial daylight (medium compositions in Samson *et al.* 2004).

Analysis for secondary metabolites

The cultures were analysed according to the HPLC-diode array detection method of Frisvad & Thrane (1987, 1993) as modified by Smedsgaard (1997). The isolates were analyzed on CYA and YES agar using three agar plugs (Smedsgaard 1997). The secondary metabolite production was confirmed by identical UV spectra with those of standards and by TLC analysis using the agar plug method, the TLC plates were eluted in toluene : ethylacetate:formic acid (6:3:1) and chloroform:acetone:2-propanol (85:15:20) (Filtner *et al.* 1983; Samson *et al.* 2004). Standards of patulin, cytochalasin E, kotanin, and nortryptoquivalin known to be produced by these fungi, were also used to confirm the identity of the compounds.

Isolation and analysis of nucleic acids

The cultures used for the molecular studies were grown on malt peptone (MP) broth using 10 % (v/v) of malt extract (Brix 10) and 0.1 % (w/v) Bacto peptone (Difco), 2 mL of medium in 15 mL tubes. The cultures were incubated at 25 °C for 7 d. DNA was extracted from the cells using the Masterpure™ yeast DNA purification kit (Epicentre Biotechnol.) according to the instructions of the manufacturer. Fragments containing the ITS region were

amplified using primers ITS1 and ITS4 as described previously (White *et al.* 1990). Amplification of part of the β -tubulin gene was performed using the primers Bt2a and Bt2b (Glass & Donaldson 1995). Amplifications of the partial calmodulin gene were set up as described previously (Hong *et al.* 2005). Sequence analysis was performed with the Big Dye Terminator Cycle Sequencing Ready Reaction Kit for both strands, and the sequences were aligned with the MT Navigator software (Applied Biosystems). All the sequencing reactions were purified by gel filtration through Sephadex G-50 (Amersham Pharmacia Biotech, Piscataway, NJ) equilibrated in double-distilled water and analyzed on the ABI PRISM 310 Genetic Analyzer (Applied Biosystems). The unique ITS, β -tubulin, actin and calmodulin sequences were deposited at the GenBank nucleotide sequence database under accession numbers EU078624–EU078678 and EU076312–EU076343.

Data analysis

The sequence data was optimised using the software package Seqman from DNASTar Inc. Sequence alignments were performed by using CLUSTAL-X (Thompson *et al.* 1997) and improved manually. The neighbour-joining (NJ) method was used for the phylogenetic analysis. For NJ analysis, the data were first analysed using the Tamura–Nei parameter distance calculation model with gamma-distributed substitution rates (Tamura & Nei 1993), which were then used to construct the NJ tree with MEGA v. 3.1 (Kumar *et al.* 2004). To determine the support for each clade, a bootstrap analysis was performed with 1000 replications.

For parsimony analysis, the PAUP v. 4.0 software was used (Swofford 2002). Alignment gaps were treated as a fifth character state and all characters were unordered and of equal weight. Maximum parsimony analysis was performed for all data sets using the heuristic search option with 100 random taxa additions and tree bisection and reconstruction (TBR) as the branch-swapping algorithm. Branches of zero length were collapsed and all multiple, equally parsimonious trees were saved. The robustness of the trees obtained was evaluated by 1000 bootstrap replications (Hillis & Bull 1993). A *Neosartorya fischeri* isolate was used as outgroup in these experiments.

RESULTS AND DISCUSSION

Phylogeny

We examined the genetic relatedness of section *Clavati* isolates and their presumed relatives using sequence analysis of the ITS region of the ribosomal RNA gene cluster, and parts of the calmodulin and β -tubulin genes. During analysis of part of the β -tubulin gene, 468 characters were analyzed. Among the 174 polymorphic sites, 102 were found to be phylogenetically informative. The Neighbour-joining tree based on partial β -tubulin genes sequences is shown in Fig. 1. The topology of the tree is the same as one of the more than 10⁵ maximum parsimony trees constructed by the PAUP program (length: 233 steps, consistency index: 0.8798, retention index: 0.9728). The ITS data set included 448 characters with 8 parsimony informative characters. The Neighbour-joining tree shown in Fig. 2 has the same topology as one of the 4 maximum parsimony trees (tree length: 25, consistency index: 0.9600, retention index: 0.9896).

Phylogenetic analysis of β -tubulin sequence data indicated that *Aspergillus* section *Clavati* includes six species, namely: *A. clavatus*




Fig. 1. Neighbour-joining tree based on β -tubulin sequence data of *Aspergillus* section *Clavati*. Numbers above branches are bootstrap values. Only values above 70 % are indicated.

(synonyms: *A. pallidus*, *A. apicalis*), *A. giganteus*, *A. longivesica*, *A. rhizopodus*, *A. clavatonanicus* and *N. acanthosporus*. Some misidentifications have also been clarified: isolates previously identified as *A. clavatus* (CBS 105.45) and *A. clavatonanicus* (CBS 112.27) were found to belong to the *A. giganteus* species, while one isolate originally identified as *A. clavatus* (IMI 351309) was found to belong to the *A. rhizopodus* species. The ITS sequences of *A. clavatonanicus* and *A. rhizopodus* isolates, and *A. giganteus* and *A. longivesica* isolates, respectively, were identical, indicating their close relationship.

A. ingratus (Yaguchi *et al.* 1993) was found to be the synonym of *H. paradoxus* based on sequence data, so it was excluded from section *Clavati* (data not shown). *H. paradoxus* isolates are only distantly related to section *Clavati*, with affinities to some *Penicillium* species (to be published elsewhere).

Chemotaxonomy

The extralites produced by species of *Aspergillus* section *Clavati* are listed in Table 2. Based on the common production of patulin, tryptoquivalins, tryptoquivalons and kotanins, most of the species appear to be closely related. *A. clavatus* produces patulin (=

clavatin = clavacin) (Weisner 1942; Waksman *et al.* 1942, 1943; Hooper *et al.* 1944) and has been reported to cause mycotoxicosis in calves as early as 1954 (Forgacs *et al.* 1954). This mycotoxin was detected on YES agar in all isolates of *A. clavatus*, *A. giganteus* and *A. longivesica*. Previously the presence of the isoeopoxydon dehydrogenase gene taking part in the biosynthesis of patulin has also been proved for *A. clavatonanicus* and *A. pallidus* isolates using primer pairs developed by Paterson *et al.* (2000) to identify potential patulin producing *Penicillia* (Varga *et al.* 2003). Other interesting metabolites produced by species of section *Clavati* are ribotoxins. Ribotoxins are a family of ribosome-inactivating proteins that have specific ribonucleolytic activity against a single phosphodiester bond in the conserved sarcin/ricin domain of 26 S rRNA (Martinez Ruiz *et al.* 1999). Ribotoxins have recently been found in a number of *Aspergillus* species including *A. clavatus*, *A. giganteus*, *A. viridinutans*, *A. fumigatus*, *A. restrictus*, *A. oryzae* var. *effusus*, *A. tamarii* and *A. ostianus*. Anamorphs of *Neosartorya fischeri*, *N. glabra* and *N. spinosa* also produced ribotoxins (Lin *et al.* 1994; Martinez-Ruiz *et al.* 1999). Using the PCR probe developed by Lin *et al.* (1994), Varga *et al.* (2003) examined the presence of ribotoxin genes in isolates of *Aspergillus* section *Clavati*; a DNA fragment of about 600 bp was amplified in some *A. clavatus*, *A.*




Fig. 2. Neighbour-joining tree based on ITS sequence data of *Aspergillus* section *Clavati*. Numbers above branches are bootstrap values. Only values above 70 % are indicated.

giganteus, *A. pallidus* and *N. acanthosporus* isolates, indicating that these isolates are able to synthesize ribotoxins (Varga *et al.* 2003). *Hemicarpenales paradoxus*, however, including its synonym *A. ingratus* produces no secondary metabolites in common with these core species and appear to more distantly related to section *Clavati*. Thus this species appears to occupy a unique position in the *Aspergillus* genus with no obvious closely related species.

Morphology

All the isolates except the ex type culture of *A. clavatonanicus*, produced numerous conidiophores with blue green conidia, hyaline conidiophore stipes and clavate aspergilla. The isolates in three species were phototropic producing very long conidiophores: *A. giganteus*, *A. rhizopodus* and *A. longivesica*. Another common phenotypic similarity was the alkalophilic tendency already described for *A. rhizopodus* which was isolated from soil with pH 8.5–9 and other species in the group (Raper & Fennell 1965; Rai *et al.* 1975). Several species have been isolated from dung which is also an alkaline substrate. This is further confirmed by the strong growth of all isolates on creatine-sucrose agar. This medium has an initial pH of 8 and creatine is an alkaline amino acid. Morphological and

physiological data confirmed that *Neocarpentales acanthosporus* and *Aspergillus* section *Clavati* are closely related.

Teleomorph relationships in *Aspergillus* subgenus *Fumigati*

Aspergillus subgenus *Fumigati* includes section *Clavati* with the *N. acanthosporus* teleomorph, and section *Fumigati* with *Neosartorya* teleomorphs. We examined the relationships of these teleomorphs taxa to another ascomycete genus, *Dichotomomyces*. *Dichotomomyces cejpaii* was originally described by Saito (1949) as *D. albus*, later validated as *D. cejpaii* by Scott (1970). This species belongs to the Trichocomaceae family (although Malloch & Cain (1971) placed it to Onygenaceae). This species is characterised by the production of aleurioconidia on short branched conidiophores, and ascospores embedded in cleiethothecia (Scott 1970; Udagawa 1970). Isolates of *D. cejpaii* are highly heat resistant and can be found world-wide in soil, heat treated products and marine environments (Pieckova *et al.* 1994; Jesenska *et al.* 1993; Mayer *et al.* 2007). *D. cejpaii* isolates has been claimed to produce a range of secondary metabolites including gliotoxin (Seigle-Murandi *et al.* 1990), xanthocillin X (Kitahara & Endo 1981), and several metabolites with




Fig. 3. Neighbour-joining tree based on β -tubulin sequence data of *Neosartorya*, *Neocarpentales*, *Dichotomomyces* species and their asexual relatives. Numbers above branches are bootstrap values. Only values above 70 % are indicated.

Table 2. Extrolite production of species assigned to *Aspergillus* section *Clavati* and *D. cejpai*. These toxins were all verified or found for the first time in the species listed, the ribotoxins (including α -sarcin) and xanthocillin X in *D. cejpai* were not verified, however.

Species	Extrolites
<i>A. clavatonanicus</i>	antafumicins, glyanthrypine, kotanins, tryptoquivalines, tryptoquivalones
<i>A. clavatus</i>	patulin, cytochalasin E & K, kotanins, antafumicin, (dehydrocarolic acid), tryptoquivalones, tryptoquivalines, ascladiol, ribotoxins
<i>A. giganteus</i>	patulin, antafumicin, ascladiol, tryptoquivalones; tryptoquivalines, glyanthrypine, pyripyropen, α -sarcin and other ribotoxins
<i>A. longivesica</i>	patulin, tryptoquivalones, tryptoquivalines, antafumicins, pyripyropen
<i>A. rhizopodus</i>	pseurotins, dehydrocarolic acid, tryptoquivalines, tryptoquivalones, kotanins, cytochalasins
<i>N. acanthosporus</i>	kotanins, tryptoquivalines, tryptoquivalones, ribotoxins
<i>D. cejpai</i>	gliotoxin, tryptoquivalones, rubratoxins, (xanthocillin X)




Fig. 4. Neighbour-joining tree based on ITS sequence data of *Neosartorya*, *Neocarpenteles*, *Dichotomomyces* species and their asexual relatives. Numbers above branches are bootstrap values. Only values above 70 % are indicated.

antibiotic and ciliostatic properties (Pieckova & Jesenska 1997a, 1997b; Pieckova & Roeijmans 1999).

We examined the genetic variability and relationships of *Aspergillus* section *Clavati* and *Fumigati* isolates, *D. cejpaii* var. *cejpaii* and *D. cejpaii* var. *spinosus* (Malloch & Cain 1971; originally described as *D. albus* var. *spinosus*; Udagawa 1970). Both the ITS region and part of the β -tubulin gene were amplified and sequenced, and phylogenetic analyses were carried out as described above. The trees based on both ITS and β -tubulin data indicate that *D. cejpaii* forms a sister group with *Neosartorya* and *Neocarpenteles* species (Figs 3–4). During analysis of part of the β -tubulin gene, 469 characters were analyzed. Among the 270 polymorphic sites, 214 were found to be phylogenetically informative. The Neighbour-joining tree based on partial β -tubulin genes sequences is shown in Fig. 3. The topology of the tree is the same as one of the 22 maximum parsimony trees constructed by the PAUP program (length: 738 steps, consistency index: 0.6233, retention index: 0.8614). The ITS data set consisted of 446 nucleotides, with 45 parsimony informative sites. The topology of the Neighbour joining tree depicted in Fig. 4 was the same as one of the more than 105 maximum parsimony trees (length: 124 steps, consistency index: 0.7419, retention index: 0.9229). Both trees indicate that the *Dichotomomyces* genus should be transferred to *Aspergillus* subgenus *Fumigati*. Similar results were obtained during phylogenetic analysis of partial calmodulin gene sequences (data not shown). *D. cejpaii* isolates have been found to produce gliotoxin in common with several species assigned to section *Fumigati* including some *Neosartorya* species (Larsen *et al.* 2007), tryptoquivalones also produced by

several species assigned to sections *Clavati* and *Fumigati* (Hong *et al.* 2005), and rubratoxins, which are hepatotoxic mycotoxins produced by *P. crateriforme* (Frisvad 1989; Sigler *et al.* 1996; Richer *et al.* 1997) [misidentified as *Penicillium purpurogenum* (Natori *et al.* 1970) or *P. rubrum* (Moss *et al.* 1968)]. *D. cejpaii* has also been claimed to produce xanthocillin X (Kitahara & Endo 1981), even though it could not be confirmed in our analyses. Xanthocillin and related compounds have also been found in *H. paradoxus* (Frisvad JC, unpubl. data) *A. candidus* (Rahbaek *et al.* 2000), *Eupenicillium crustaceum* (Turner & Aldridge 1983), *E. egyptiacum* (Vesonder 1979), *P. italicum* (Arai *et al.* 1989), *P. flavigenum* (Frisvad *et al.* 2004) and *P. chrysogenum* (Hagedorn *et al.* 1960; Achenbach *et al.* 1972; Pfeiffer *et al.* 1972; Frisvad *et al.* 2004; de la Campa *et al.* 2007). Since the anamorph of *Dichotomomyces* was earlier found to belong to *Polypaecilum*, further morphological and molecular studies are needed to clarify the significance of the morphology of the anamorph in the taxonomic placement of these species, and to clarify the taxonomy of *Polypaecilum* species.

In conclusion, the polyphasic approach applied to clarify the taxonomy of *Aspergillus* section *Clavati* led to the assignment of six species, namely: *A. clavatus* (synonyms: *A. pallidus*, *A. apicalis*), *A. giganteus*, *A. longivesica*, *A. rhizopodus*, *A. clavatonanicus* and *N. acanthosporum* to this section. *Hemicarpenteles paradoxus* (synonym: *A. ingratus*) was found to be unrelated to section *Clavati*, but more closely related to *Penicillium*. *Dichotomomyces* and *Neosartorya* were found to be sister clades to the genus *Neocarpenteles*. Further studies are needed to clarify the taxonomic status of *Dichotomomyces* species with *Polypaecilum* anamorphs.

Aspergillus clavatonanicus Batista, Maia & Alecrim, Anais Fac. Med. Univ. Recife 15: 197. 1955. Fig. 5.

Type: CBS 474.65, from finger nail lesion, Recife, Brazil

Other no. of the type: ATCC 12413; DMUR 532; IMI 235352; WB 4741

Description

Colony diam (7 d): CYA25: 50–82 mm, MEA25: 45–78 mm, YES25: 57–82 mm, OA25: 49–60 mm, CYA37: 8–17 mm, CREA: very good growth and acid production in the margin of the colony

Colony colour: greyish blue green

Conidiation: abundant

Reverse colour (CZA): uncoloured to light brownish

Colony texture: floccose

Conidial head: clavate, up to 145–360 × 120–180 µm

Stipe: 40–470 × 6–16 µm, rough walled

Vesicle diam/shape: 22–125 × 5–22 µm, clavate

Conidium size/shape/surface texture: 5–8.5 × 5–6.5 µm, ellipsoid or cylindrical, smooth

Cultures examined: CBS 474.65 = IBT 12370 = IBT 24678, CBS 112.27 = IBT 12369 = IBT 24677

Diagnostic features: conidial heads smaller than 1 mm

Similar species: *A. clavatus*

Distribution: Brazil

Ecology and habitats: human

Extrolites: antafumicins, glyanthrypine, kotanin, tryptoquivalins, tryptoquivalons

Pathogenicity: isolated from nail lesion (Batista *et al.* 1955)

Aspergillus clavatus Desmazières, Ann. Sci. Nat., Bot. 2: 71, 1834. Fig. 6.

= *Aspergillus pallidus* Kamyschko (1963)

= *Aspergillus apicalis* Mehrotra & Basu (1976)

Type: CBS 513.65, J. Westerdijk > 1909, C. Thom > NRRL

Other no. of the type: ATCC 1007; ATCC 9602; ATCC 9598; CECT 2674; DSM 816; IMI 015949; IMI 015949v; IMI 015949iv; IMI 015949iii; LSHB Ac86; LSHB Ac95; NCTC 978; NCTC 3887; NRRL 1; NRRL 1656; QM 1276; QM 7404; WB 1

Description

Colony diam (7 d): CYA25: 28–45 mm; MEA25: 25–44 mm, YES25: 29–45 mm, OA25: 31–47 mm, CYA37: 9–26 mm, CREA25: very good growth and moderate to very strong acid production (exceptions: CBS 514.65, NRRL 2, NRRL 8 and NRRL 2254 grow poorly on CREA and produce no or very little acid)

Colony colour: blue-green

Conidiation: abundant

Reverse colour (CZA): uncoloured to somewhat brownish with age in some isolates

Colony texture: velvety

Conidial head: clavate, commonly ranging from 300 to 400 µm by 150 to 200 µm when young, in age commonly splitting into two, three, or more divergent columns

Stipe: 1500–3000 × 20–30 µm

Vesicle diam/shape: 200–250 × 40–60 µm, clavate

Conidium size/shape/surface texture: 3–4.5 × 2.5–3 µm, elliptical, smooth

Cultures examined: CBS 104.45, CBS 105.45, CBS 106.45, CBS 114.48, CBS 513.65, CBS 514.65, CBS 470.91, CBS 116685, CBS 118451, DTO 6-F8, DTO 27-C2, SZMC 0918, SZMC JV4, SZMC JV1.1, IMI 351309, IMI 358435, CBS 117.45, CBS 119.48, CBS 118.49, CBS 122.53, CBS 117.56, CBS 101.64, CBS 515.65, CBS 526.65

Diagnostic features: conidial heads up to 4 mm in size

Similar species: *A. clavatonanicus*

Distribution: worldwide, mainly in tropical, subtropical and Mediterranean regions

Ecology and habitats: soil, cereals, malt, dung

Extrolites: Patulin, cytochalasin E, kotanins, antafumicin, (dehydrocarolic acid), tryptoquivalone, tryptoquivalines, ascladiol (all found in this study), ribotoxins (Lin *et al.* 1995, Huang *et al.* 1997)

Pathogenicity: caused endocarditis (Opal *et al.* 1986), responsible for an extrinsic allergic alveolitis known as malt worker's lung (Grant *et al.* 1976; Lopez-Diaz & Flannigan 1997; Flannigan & Pearce 1994), and various toxic syndromes including neurological disorders (Shlosberg *et al.* 1991; McKenzie *et al.* 2004; Loretto *et al.* 2003; Gilmour *et al.* 1989; Kellerman *et al.* 1976) and other mycotoxicosis-related diseases (Byth & Lloyd 1971) observed in animals

Notes: some isolates carry dsRNA mycoviruses 35–40 nm in size (Varga *et al.* 2003)

Aspergillus giganteus Wehmer, Mem. Soc. Phys. Genève 33 (2): 85. 1901. Fig. 7.

Type: CBS 526.65, dung of bat in cave, Yucatan, Mexico

Other no. of the type: ATCC 10059; DSM 1146; IFO 5818; IMI 227678; NRRL 10; QM 1970; WB 10; IBT 12368

Description

Colony diam: CYA25: (26–) 40–65 mm, MEA25: (29–) 43–65 mm, YES25: 40–80 mm, OA25: 31–75 mm, CYA37: 10–29 mm, CREA: very good growth and poor or no acid production

Colony colour: first white, becoming pale blue-green near light celandine green to slate-olive

Conidiation: usually abundant

Reverse colour (CZA): dull tan

Colony texture: velvety

Conidial head: splitting into 2 or more columns with age, blue green

Stipe: two types: 2–3(–4) mm; or several cm in length

Vesicle diam/shape: two types: 100–250 × 30–50 µm on short conidiophores, 400–600 × 120–180 µm on long ones, clavate

Conidium size/shape/surface texture: 3.5–4.5 × 2.4–3 µm, elliptical, thick-walled, smooth

Cultures examined: CBS 117.45, CBS 119.48, CBS 118.49, CBS 122.53, CBS 117.56, CBS 101.64, CBS 515.65

Diagnostic features: produces clavate vesicles in contrast with the elongate ones of *A. longivesica*; do not produce rhizoidal foot




Fig. 5. *Aspergillus clavatonanicus*. A–B. Colonies after 7 d at 25 °C. A. CYA. B. MEA. C–J. Conidiophores. K. Conidia. Scale bars = 10 µm.




Fig. 6. *Aspergillus clavatus*. A. Colonies after 7 d at 25 °C on CYA. B–C. Macrophotograph of conidiophores. D–I. Conidiophores. J. Conidia. Scale bars = 10 µm, except D and E = 30 µm.




Fig. 7. *Aspergillus giganteus*. A. Colonies after 7 d at 25 °C on CYA. B–C. Macrophotograph of conidiophores. D–I. Conidiophores. J. Conidia. Scale bars = 10 µm, except D and E = 30 µm.

cells characteristic to *A. rhizopodus*; conidial heads can be up to 1–5 cm long

Similar species: *A. rhizopodus*, *A. longivesica*

Distribution: Nigeria, U.S.A., Egypt, Mexico, Panama, Germany, Suriname, Netherlands, Poland

Ecology and habitats: dung, soil, wood

Extrolites: patulin, antafumicin, ascladiol, tryptoquivalone; tryptoquivalines, glyanthrypine, pyripropen (found in this study), α -sarcin and other ribotoxins (Olson & Goerner 1965; Olson *et al.* 1965; Lin *et al.* 1995; Wirth *et al.* 1997; Martinez-Ruiz *et al.* 1999). Carotens are also produced (van Eijk *et al.* 1979)

Pathogenicity: not reported

Note: two types of conidial structures: (1) conidiophores commonly 2 to 3 mm, rarely exceeding 4 mm in height, bearing clavate heads 200 to 350 μ m in length; (2) conidiophores one to several centimeters in length, bearing heads up to 1 mm in length; longer conidiophores are phototropic, and only elongate in the presence of light

Aspergillus longivesica Huang & Raper, *Mycologia* 63(1): 53. 1971. Fig. 8.

Type: CBS 530.71, from soil, rain forest, Nigeria

Other no. of the type: ATCC 22434; IMI 156966; QM 9698

Description

Colony diam: CYA25: 31–51 mm; MEA25: 48–56 mm; YES25: 60–74 mm; OA25: 52–60 mm, CYA37: 0 mm, CREA25: weak growth and no acid production (CBS 187.77 grow very well on CREA, however)

Colour: white to cream

Conidiation: abundant, rarely less abundant

Reverse colour (CYA): pale cinnamon buff

Colony texture: thin floccose

Conidial head: elongate, splitting into divergent columns with age, greyish blue green

Stipe: two types: 80–420 \times 7–11.2 μ m, or 1.5–4.5 cm long, thick walled (5.6–7 μ m)

Vesicle diam/shape: two types: 2.2–3.2 mm \times 130–200 μ m, elongate, clavate, thick-walled, or 18–36 μ m, globose to flask-shaped, thin-walled

Conidia length/shape/surface texture: two types: 4.2–16.8 \times 2.8–7 μ m, globose to elliptical, or 3.5–5.2 \times 2.5–3.5 μ m, elliptical or pyriform

Cultures examined: CBS 530.71, CBS 187.77

Diagnostic features: produces longer and wider conidiophores, longer vesicles and larger conidia than *A. giganteus*; vesicles are elongate to fusoid-clavate for the long conidiophore and globose for the smaller ones, while those of *A. giganteus* are clavate

Similar species: *A. giganteus*

Distribution: Nigeria, Ivory Coast

Ecology and habitats: soil

Extrolites: patulin, tryptoquivalone, tryptoquivalines, antafumicins, pyripropens (found in this study)

Pathogenicity: not reported

Note: longer conidiophores are phototropic, and only elongate in the presence of light

Aspergillus rhizopodus Rai, Wadhvani & Agarwal, *Trans. Br. Mycol. Soc.* 64: 515. 1975. Fig. 9

Type: CBS 450.75, from usar soil, Lucknow, India

Other no. of the type: IMI 385057; WB5442

Description

Colony diam (7 d): CZA30: 40 mm; CYA25: 38–42 mm; MEA25: 50–55 mm; YES25: 68–72 mm; OA25: 43–47 mm; CYA37: 17–19 mm; CREA25: rather good growth and no acid production

Colony colour: blue green

Conidiation: abundant

Reverse colour (CZA): colourless

Colony texture: slightly furrowed

Conidial head: short columnar

Stipe: two types: 208–800 \times 11–32 μ m, or 5–22 mm \times 36 μ m, thick walled, smooth

Vesicle diam/shape: two types: 40–176 \times 11–32 μ m, or 288 \times 79 μ m, clavate

Conidium size/shape/surface texture: 4–5.5 \times 2.5–3 μ m, ellipsoidal, smooth

Cultures examined: CBS 450.75, IMI 351309

Diagnostic features: produces variously shaped foot cells with finger-like projections

Similar species: *A. giganteus*, *A. longivesica*

Distribution: India, Yugoslavia

Ecology and habitats: soil

Extrolites: pseurotins, dehydrocarolic acid, tryptoquivalines, tryptoquivalones, kotanins and cytochalasin (found in this study)

Pathogenicity: not reported

Note: large conidial heads formed only in the presence of light

Dichotomomyces cejpilii (Milko) D.B. Scott, *Trans. Brit. Mycol. Soc.* 47: 428, 1970. Fig. 10.

= *Talaromyces cejpilii* Milko (1964)

= *Dichotomomyces albus* Saito (1949)

= *Royella albida* Dwiveli (1960)

Type: CBS 157.66, from orchard soil, near Tiraspol, Moldova

Description

Colony diam (7 d): CYA25: 25–47 mm; MEA25: 35–58 mm; YES25: 47–50 mm; OA25: 38–48; CYA37: 24–32 mm; CREA: poor growth and no acid production

Colony colour: white to cream coloured

Conidiation: sparse

Reverse colour (CZA):

Colony texture: floccose, granular

Conidium size/shape/surface texture: 5–10 μ m, subglobose to pyriform, smooth

Homothallic

Cleistothecia: variable in size, spherical, white to cream coloured




Fig. 8. *Aspergillus longivesica*. A. Colonies after 10 d at 25 °C on CYA. B–C. Macrophotograph of conidiophores. D–I. Conidiophores. J. Conidia. Scale bars = 10 µm, except D and E = 30 µm.




Fig. 9. *Aspergillus rhizopodus*. A. Colonies after 10 d at 25 °C on CYA. B. Macrophotograph of conidiophores. C–I. Conidiophores. J. Conidia. Scale bars = 10 µm, except D and E = 30 µm.




Fig. 10. *Dichotomomyces cejpii*. A–B. Ascomata on MEA after 10 d at 25 °C. C. Ascomata wall. D–E. Asci and ascospores. F–I conidiophores and conidia. Scale bars = 10 μ m, except B and C = 30 μ m.

Ascospores: 3–3.5 × 4–4.5 µm, lenticular, with two closely appressed very thin equatorial crests and convex walls smooth

Cultures examined: CBS 761.96, CBS 779.7, CBS 219.67, CBS 100192, CBS 474.77, CBS 780.70, CBS 397.68, CBS 345.68, CBS 159.67, CBS 157.66, CBS 212.50

Diagnostic features: conidiophore apices are dichotomously branched, and conidia are produced from these branches (*Polypaecilum* anamorph); racquet hyphae are frequently produced; vegetative hyphae often bear rhizomorphs

Similar species: -

Distribution: Slovakia, Netherlands, Egypt, U.S.A., South Africa, Pakistan, Japan, Moldova, India

Ecology and habitats: soil, compost, pasteurised products

Extrolites: gliotoxin (Seigle-Murandi *et al.* 1990, confirmed in this study), tryptoquivalons (found in this study), rubratoxins (found in this study), xanthocillin X (Kitahara & Endo 1981; could not be confirmed in this study), and several metabolites with antibiotic and ciliostatic properties (Pieckova & Roelijmans 1999; Pieckova & Jesenska 1997a, 1997b)

Pathogenicity: not reported

Note: this species is reported as a heat resistant fungus causing food spoilage (Pieckova *et al.* 1994; Jesenska *et al.* 1993; Mayer *et al.* 2007)

Neocarpenteles acanthosporus (Udagawa & Takada) Udagawa & Uchiyama [anamorph: *A. acanthosporus* Udagawa & Takada], *Mycoscience* 43(1): 4. 2002.
= *Hemicarpenteles acanthosporus* Udagawa & Takada (1971)

Type: CBS 558.71, from soil, Bougainville Island (Solomon Islands), Papua New Guinea

Other no. of the type: ATCC 22931; IMI 164621; NHL 2462

Description

Colony diam (7 d): CYA25: 37–47 mm; MEA25: 72–85 mm; YES25: 62–82; OA25: 40–49 mm; CYA37: 0 mm; CREA: poor growth and no acid production

Colour: white to brownish orange

Conidiation: sparse

Reverse colour (CYA): greyish-orange

Colony texture: floccose

Conidial head: radiate to loosely columnar

Stipe: (50–)100–400 × 5–12 µm, smooth, septate

Vesicle diam /shape: 10–26 µm, flask shaped

Conidia length/ shape/ surface texture: 4.5–7 µm, globose to subglobose, spinulose

Homothallic

Cleistothecia: 350–1000 × 250–850 µm, sclerotoid, subglobose to ovoid, fawn, covered with dense aerial hyphae

Ascospores: 4–4.5 × 3.5–4 µm, lenticular, with two thin equatorial crests and convex walls ornamented with raised flaps

Cultures examined: CBS 558.71, CBS 445.75, CBS 446.75, CBS 447.75

Diagnostic features: small dull green radiate conidial heads, short conidiophores with small flask-shaped vesicle, production

of ascospores, and large globose conidia distinguish this species from other members of section *Clavati*

Distribution: Papua New Guinea (Bougainville Island), Japan

Ecology and habitats: soil

Extrolites: kotanins, tryptoquivalines, tryptoquivalones (found in this study), ribotoxins (Varga *et al.* 2003). (+)-isoeopoxydon has also been reported (Kontani *et al.* 1990)

Pathogenicity: not reported

Note: not illustrated here, for detailed description and illustration see Udagawa & Takada (1971); no growth at 37 °C

ACKNOWLEDGEMENTS

We thank Ellen Kirstine Lyhne for technical assistance and the Danish Technical Research Council for financial support to Programme for Predictive Biotechnology.

REFERENCES

- Achenbach H, Strittmatter H, Kohl W (1972). Die Strukturen der Xanthocilline Y1 und Y2. *Chemisches Berichte* **105**: 3061–3066.
- Arai K, Miyajima H, Mushiroda T, Yamamoto Y (1989). Metabolites of *Penicillium italicum* Wehmer: isolation and structures of new metabolites including naturally occurring 4-ylideneacyltetronic acids, italicinic acid and italicic acid. *Chemical and Pharmaceutical Bulletin* **37**: 3229–3235.
- Batista AC, Maia HS, Alecrim IC (1955). Onicomocose produzida por "*Aspergillus clavato-nanica*" n. sp. *Anais da Faculdade de Medicina da Universidade do Recife* **15**: 197–203.
- Bergel F, Klein R, Morrison AL, Mioss AR, Rindetrknecht H, Ward JL (1943) An antibacterial substance from *Aspergillus clavatus* and *Penicillium claviforme* and its probable identity with patulin. *Nature* **152**: 750.
- Bergel F, Morrison AL, Moss AR, Rinderknecht H (1944) An antibacterial substance from *Aspergillus clavatus*. *Journal of the Chemical Society* **1944**: 415–421.
- Beretta B, Gaiaschi A, Galli CL, Restani P (2000). Patulin in apple-based foods: occurrence and safety evaluation. *Food Additives and Contaminants* **17**: 399–406.
- Blyth W, Lloyd MM (1971). Granulomatous and mycotoxic syndromes in mice due to *Aspergillus clavatus* Desm. *Sabouraudia* **9**: 263–272.
- Büchi G, Luk KC, Kobbe B, Townsend JM (1977). Four new mycotoxins from *Aspergillus clavatus* related to tryptoquivaline. *Journal of Organic Chemistry* **42**: 244–246.
- Campa R de la, Seifert K, Miller JD (2007). Toxins from strains of *Penicillium chrysogenum* isolated from buildings and other sources. *Mycopathologia* **163**: 161–168.
- Clardy J, Springer JP, Büchi G, Matsuo K, Wrightman R (1975). Tryptoquivaline and tryptoquivalone, two new tremorgenic metabolites of *Aspergillus clavatus*. *Journal of the American Chemical Society* **97**: 663–665.
- Cole RJ, Cox RH (1981). *Handbook of toxic fungal metabolites*. New York: Academic Press.
- Demain AL, Hunt NA, Malik V, Kobbe B, Hawkins H, Matsuo K, Wogan GN (1976). Improved procedure for production of cytochalasin E by *Aspergillus clavatus*. *Applied and Environmental Microbiology* **31**: 138–140.
- Eijk GW van, Mummery RS, Roeymans HJ, Valadon LRG (1979). A comparative study of carotenoids of *Aschersonia aleyroides* and *Aspergillus giganteus*. *Antonie van Leeuwenhoek* **83**: 191–200.
- Fennell DI, Raper KB (1955). New species and varieties of *Aspergillus*. *Mycologia* **47**: 68–89.
- Filtenborg O, Frisvad JC, Svendsen JA (1983). Simple screening method for molds producing intracellular mycotoxins in pure culture. *Applied and Environmental Microbiology* **45**: 581–585.
- Flannigan B, Pearce AR (1994). *Aspergillus* spoilage: spoilage of cereals and cereal products by the hazardous species *Aspergillus clavatus*. In: *The genus Aspergillus. From taxonomy and genetics to industrial application*. (Powell, KA, Renwick A, Peberdy JF, eds). New York. Plenum Press: 55–62.
- Florey HW, Jennings MA, Philpot FJ (1944). Claviformin from *Aspergillus giganteus* Wehm. *Nature* **153**: 139.

- Forgacs J, Carll WT, Herring AS, Mahlandt BG (1954). A toxic *Aspergillus clavatus* isolated from feed pellets. *American Journal of Hygiene* **60**: 15–26.
- Frisvad JC (1989) The connection between the Penicillia and aspergilli and mycotoxins with special emphasis on misidentified isolates. *Archives of Environmental Contamination and Toxicology* **18**: 452–467.
- Frisvad JC, Smedsgaard J, Larsen TO, Samson RA (2004). Mycotoxins, drugs and other extrolites produced by species in *Penicillium* subgenus *Penicillium*. *Studies in Mycology* **49**: 201–241.
- Frisvad JC, Thrane U (1987). Standardized high performance liquid chromatography of 182 mycotoxins and other fungal metabolites based on alkylphenone retention indices and UV-VIS spectra (diode array detection). *Journal of Chromatography A* **404**: 195–214.
- Frisvad JC, Thrane U (1993). Liquid chromatography of mycotoxins. In: Betina V (ed.). *Chromatography of mycotoxins: techniques and applications*. Journal of Chromatography Library **54**. Amsterdam: Elsevier: 253–372.
- Gams W, Christensen M, Onions AH, Pitt JI, Samson RA (1985). Infrageneric taxa of *Aspergillus*. In: *Advances in Penicillium and Aspergillus Systematics*. Samson RA, Pitt JI, eds. New York: Plenum Press: 55–62.
- Gilmour JS, Inglis DM, Robb J, Maclean M (1989). A fodder mycotoxicosis of ruminants caused by contamination of a distillery by-product with *Aspergillus clavatus*. *The Veterinary Record* **124**: 133–135.
- Glass NL, Donaldson GC (1995). Development of primer sets designed for use with the PCR to amplify conserved genes from filamentous ascomycetes. *Applied and Environmental Microbiology* **61**: 1323–1330.
- Grant IW, Blackadder ES, Greenberg M, Blyth W (1976). Extrinsic allergic alveolitis in Scottish maltworkers. *British Medical Journal* **1**: 490–493.
- Hagedorn I, Eholzer U, Luttringhaus A (1960). Beiträge zur Konstitutionsermittlung des Antibiotikums Xanthocillin. *Chemisches Berichte* **93**: 1584–1590.
- Harrison MA (1989). Presence and stability of patulin in apple products: a review. *Journal of Food Protection* **9**: 147–153.
- Hillis DM, Bull JJ (1993). An empirical test of bootstrapping as a method for assessing confidence in phylogenetic analysis. *Systematic Biology* **42**: 182–192.
- Hong SB, Cho HS, Shin HD, Frisvad JC, Samson RA (2006). Novel *Neosartorya* species isolated from soil in Korea. *International Journal of Systematic and Evolutionary Microbiology* **56**: 477–486.
- Hong SB, Go SJ, Shin HD, Frisvad JC, Samson RA (2005). Polyphasic taxonomy of *Aspergillus fumigatus* and related species. *Mycologia* **97**: 1316–1329.
- Hooper IR, Anderson HW, Skell P, Carter HE (1944). The identity of clavacin with patulin. *Science* **99**: 16.
- Huang KC, Hwang YY, Hwu L, Lin A (1997). Characterization of a new ribotoxin gene (c-sar) from *Aspergillus clavatus*. *Toxicon* **35**: 383–392.
- Huang LH, Raper KB (1971). *Aspergillus longivesica*, a new species from Nigerian soil. *Mycologia* **63**: 50–57.
- Jesenska Z, Pieckova E, Bernat D (1993). Heat resistance of fungi from soil. *International Journal of Food Microbiology* **19**: 187–192.
- Kellerman TS, Pienaar JG, van der Westhuizen GC, Anderson GC, Naude TW (1976). A highly fatal tremorgenic mycotoxicosis of cattle caused by *Aspergillus clavatus*. *Onderstepoort Journal of Veterinary Research* **43**: 147–154.
- Kitahara N, Endo A (1981). Xanthocillin X monomethyl ether, a potent inhibitor of prostaglandin biosynthesis. *Journal of Antibiotics (Tokyo)* **34**: 1556–1561.
- Kontani M, Fukushima Y, Sakagami Y, Marumo S (1990). Inhibitors of β -glucan biosynthesis in fungal metabolites. *Tennen Yuki Kagobutsu Toronkai Koan Yoshishu* **32**: 103–110.
- Kumar S, Tamura K, Nei M (2004). MEGA3: Integrated Software for Molecular Evolutionary Genetics Analysis and Sequence Alignment. *Briefings in Bioinformatics* **5**: 150–163.
- Larsen TO, Smedsgaard J, Nielsen KF, Hansen MA, Samson RA, Frisvad JC (2007). Production of mycotoxins by *Aspergillus lentulus* and other medically important and closely related species in section *Fumigati*. *Medical Mycology* **45**: 225–232.
- Lin A, Huang K, Hwu L, Tzean SS (1994). Production of type II ribotoxins by *Aspergillus* species and related fungi in Thailand. *Toxicon* **33**: 105–110.
- Lopez-Diaz TM, Flannigan B (1997). Production of patulin and cytochalasin E by *Aspergillus clavatus* during malting of barley and wheat. *International Journal of Food Microbiology* **35**: 129–136.
- Loretti AP, Colodel EM, Driemeier D, Correa AM, Bangel JJ Jr, Ferreira L (2003). Neurological disorder in dairy cattle associated with consumption of beer residues contaminated with *Aspergillus clavatus*. *Journal of Veterinary Diagnostic Investigation* **15**: 123–132.
- Malloch D, Cain RF (1971). New genera of Onygenaceae. *Canadian Journal of Botany* **49**: 839–846.
- Martínez-Ruiz A, Kao R, Davies J, Martínez del Pozo A (1999). Ribotoxins are a more widespread group of proteins within the filamentous fungi than previously believed. *Toxicon* **37**: 1549–1563.
- Mayer KM, Ford J, Macpherson GR, Padgett D, Volkmann-Kohlmeyer B, Kohlmeyer J, Murphy C, Douglas SE, Wright JM, Wright JLC (2007). Exploring the diversity of marine-derived fungal polyketide synthases. *Canadian Journal of Microbiology* **53**: 291–302.
- McKenzie RA, Kelly MA, Shivas RG, Gibson JA, Cook PJ, Widderick K, Guilfoyle AF (2004). *Aspergillus clavatus* tremorgenic neurotoxicosis in cattle fed sprouted grains. *Australian Veterinary Journal* **82**: 635–638.
- Mehrotra BS, Basu M (1976). Some interesting new isolates of *Aspergillus* from stored wheat. *Nova Hedwigia* **27**: 597–607.
- Moss MO, Robinson FV, Wood AB (1968). Rubratoxin B, a toxic metabolite of *Penicillium rubrum*. *Chemistry & Industry* **18**: 587–588.
- Natori S, Sakaki S, Kurata H, Udagawa SI, Ichinoe M (1970). Production of rubratoxin B by *Penicillium purpurogenum* Stoll. *Applied Microbiology* **19**: 613–617.
- Olson BH, Goerner GL (1965). Alpha sarcin, a new antitumor agent. I. Isolation, purification, chemical composition, and the identity of a new amino acid. *Applied Microbiology* **13**: 314–321.
- Olson BH, Jennings JC, BH, Roga V, Junek AJ, Schuurmans DM (1965). Alpha sarcin, a new antitumor agent. II. Fermentation and antitumor spectrum. *Applied Microbiology* **13**: 322–326.
- Opal SM, Reller LB, Harrington G, Cannady P Jr (1986). *Aspergillus clavatus* endocarditis involving a normal aortic valve following coronary artery surgery. *Reviews of Infectious Diseases* **8**: 781–785.
- Paterson RRM, Archer S, Kozakiewicz Z, Lea A, Locke T, O'Grady E (2000). A gene probe for the patulin metabolic pathway with potential for use in patulin and novel disease control. *Biocontrol Science and Technology* **10**: 509–512.
- Peterson SW (2000). Phylogenetic relationships in *Aspergillus* based on rDNA sequence analysis. In: *Integration of modern taxonomic methods for Penicillium and Aspergillus classification*. (Samson RA, Pitt JI, eds). Amsterdam. Harwood Academic Publishers: 323–355.
- Pfeiffer S, Bar H, Zarnack J (1972). Über Stoffwechselprodukte der Xanthocillin-bildende Mutante der *Penicillium notatum* Westl. *Pharmazie* **27**: 536–542.
- Pieckova E, Bernat D, Jesenska Z (1994). Heat resistant fungi isolated from soil. *International Journal of Food Microbiology* **22**: 297–299.
- Pieckova E, Jesenska Z (1997a). *Dichotomomyces cejpaii* – some characteristics of strains isolated from soil in the Slovak Republic. *Czech Mycology* **49**: 229–237.
- Pieckova E, Jesenska Z (1997b). Toxinogenicity of heat-resistant fungi detected by a bio-assay. *International Journal of Food Microbiology* **36**: 227–229.
- Pieckova E, Roeijmans H (1999). Antibiotic secondary metabolites of *Dichotomomyces cejpaii*. *Mycopathologia* **146**: 121–126.
- Rahbaek L, Frisvad JC, Christophersen C (2000). An amendment of *Aspergillus* section *Candidi* based on chemotaxonomical evidence. *Phytochemistry* **53**: 581–586.
- Rai JN, Wadhvani K, Agarwal SC (1975). *Aspergillus rhizopodus* sp.nov. from Indian alkaline soils. *Transactions of the British Mycological Society* **64**: 515–517.
- Raper KB, Fennell DI (1965). *The genus Aspergillus*. Baltimore: Williams & Wilkins.
- Richer L, Sigalet D, Kneteman N, Joens A, Scott RB, Asgbourne R, Sigler L, Frisvad J, Smith L (1997). Fulminant hepatic failure following ingestion of mouldy homemade rhubarb wine. *Gastenterology* **112**: A1366.
- Saito K (1949). Studies on the fungi in the Orient. *Journal of Fermentation Technology* **27**: 120–122.
- Samson RA (1979). A compilation of the aspergilli described since 1965. *Studies in Mycology* **18**: 1–38.
- Samson RA, Hoekstra ES, Frisvad JC (eds) (2004). *Introduction to food and airborne fungi*. 7th ed. Utrecht: Centraal Bureau voor Schimmelcultures.
- Sarbhoj AK, Elphick JJ (1968). *Hemicarpenales paradoxus* gen. & sp. nov.: The perfect state of *Aspergillus paradoxus*. *Transactions of the British Mycological Society* **51**: 155–157.
- Scott DB (1970). *Dichotomomyces cejpaii* (Milko) comb. nov. *Transactions of the British Mycological Society* **55**: 313–316.
- Seigle-Murandi F, Krivobok S, Steiman R, Marzin D (1990). Production, mutagenicity, and immunotoxicity of gliotoxin. *Journal of Agricultural and Food Chemistry* **38**: 1854–1856.
- Shlosberg A, Zadikov I, Perl S, Yakobson B, Varod Y, Elad D, Rapoport E, Handji V (1991). *Aspergillus clavatus* as the probable cause of a lethal mass neurotoxicosis in sheep. *Mycopathologia* **114**: 35–39.
- Sigler L, Abbott SP, Frisvad JC (1996). Rubratoxin mycotoxicosis by *Penicillium crateriforme* following ingestion of homemade rhubarb wine. Abstracts, 96th ASM, New Orleans. F-22, p. 77.
- Smedsgaard J (1997). Micro-scale extraction procedure for standardized screening of fungal metabolite production in cultures. *Journal of Chromatography A* **760**: 264–270.
- Steyn PS, van Heerden FR, Rabie CJ (1982). Cytochalasins E and K, toxic metabolites from *Aspergillus clavatus*. *Journal of the Chemical Society Perkin Transactions I* **1982**: 541–544.
- Suzuki T, Takeda M, Tanabe H (1971). A new mycotoxin produced by *Aspergillus clavatus*. *Chemical and Pharmaceutical Bulletin* **19**: 1786–1788.
- Swofford T (2000). *PAUP: Phylogenetic analysis using parsimony*. v. 4.0. Sunderland: Sinauer Associates.

- Tamura K, Nei M (1993). Estimation of the number of nucleotide substitutions in the control region of mitochondrial DNA in humans and chimpanzees. *Molecular Biology and Evolution* **10**: 512–526.
- Tamura M, Hamamoto M, Cañete-Gibat CF, Sugiyama J, Nakase T (1999). Genetic relatedness among species in *Aspergillus* section *Clavati* as measured by electrophoretic comparison of enzymes, DNA base composition, and DNA-DNA hybridization. *Journal of General and Applied Microbiology* **45**: 77–83.
- Thom C, Church MB (1926). *The aspergilli*. Baltimore: Williams & Wilkins.
- Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG (1997). The ClustalX windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Research* **25**: 4876–4882.
- Trinci APJ, Banbury GH (1967). A study of the growth of the tall conidiophores of *Aspergillus giganteus*. *Transactions of the British Mycological Society* **50**: 525–538.
- Turner WB, Aldridge DC (1983). *Fungal Metabolites II*. London: Academic Press.
- Udagawa S (1970). Notes on some Japanese Ascomycetes IX. *Transactions of the Mycological Society of Japan* **10**: 103–109.
- Udagawa S, Takada M (1971). Mycological reports from New Guinea and the Solomon Islands. 10. Soil and coprophilous microfungi. *Bulletin of the National Science Museum Tokyo* **14**: 501–546.
- Udagawa S, Uchiyama S (2002). *Neocarpenteles*: a new ascomycete genus to accommodate *Hemicarpenteles acanthosporus*. *Mycoscience* **43**: 3–6.
- Varga J, Rigó K, Molnár J, Tóth B, Szencz S, Téren J, Kozakiewicz Z (2003). Mycotoxin production and evolutionary relationships among species of *Aspergillus* section *Clavati*. *Antonie van Leeuwenhoek* **83**: 191–200.
- Vesonder RF (1979). Xanthocillin, a metabolite of *Eupenicillium egyptiacum* NRRL 1022. *Journal of Natural Products* **42**: 232–233.
- Waksman SA, Horning ES, Spencer EL (1942). The production of two antibacterial substances, fumigacin and clavacin. *Science* **96**: 202.
- Waksman SA, Horning ES, Spencer EL (1943). Two antagonistic fungi, *Aspergillus fumigatus* and *Aspergillus clavatus*, and their antibiotic substances. *Journal of Bacteriology* **45**: 233–248.
- Weisner BP (1942). Bactericidal effects of *Aspergillus clavatus*. *Nature* **149**: 356.
- White TJ, Bruns T, Lee S, Taylor J (1990). Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: *PCR Protocols: A guide to methods and applications*. (Innis MA, Gelfand DH, Sninsky JJ, White TJ, eds). New York: Academic Press: 315–322.
- Wirth J, Martinez del Pozo A, Mancheno JM, Martinez-Ruiz A, Lacadena J, Onaderra M, Gavilanes JG (1997). Sequence determination and molecular characterization of gigantins, a cytotoxic protein produced by the mould *Aspergillus giganteus* IFO 5818. *Archives of Biochemistry and Biophysics* **343**: 188–193.
- Yaguchi T, Someya A, Miyadoh S, Udagawa S (1993). *Aspergillus ingratus*, a new species in *Aspergillus* section *Clavati*. *Transactions of the Mycological Society of Japan* **34**: 305–310.