

RESEARCH

Open Access



# Psychosocial health in pregnancy and postpartum among women living with - and without HIV and non-pregnant women living with HIV living in the Nordic countries – Results from a longitudinal survey study

Ellen Moseholm<sup>1,2\*</sup>, Inka Aho<sup>3</sup>, Åsa Mellgren<sup>4,5</sup>, Gitte Pedersen<sup>6</sup>, Terese L. Katzenstein<sup>7</sup>, Isik S. Johansen<sup>8</sup>, Diana Bach<sup>9</sup>, Merete Storgaard<sup>10</sup> and Nina Weis<sup>1,11</sup>

## Abstract

**Background:** The success of antiretroviral therapy has normalized pregnancy among women living with HIV (WWH) with a very low risk of perinatal transmission of HIV. Despite these advances, WWH still face complex medical and psychosocial issues during pregnancy and postpartum. The aim of this study was to assess differences in psychosocial health outcomes between pregnant WWH, non-pregnant WWH, and pregnant women without HIV, and further identify factors associated with probable depression in the third trimester and postpartum.

**Methods:** In a longitudinal survey study, participants were included from sites in Denmark, Finland, and Sweden during 2019–2020. Data was collected in the 3rd trimester, 3 and 6 months postpartum using standardized questionnaires assessing depression, perceived stress, loneliness, and social support. Mixed regression models were used to assess changes over time within and between groups. Logistic regression models were used to identify factors associated with depression in pregnancy and postpartum.

**Results:** A total of 47 pregnant WWH, 75 non-pregnant WWH, and 147 pregnant women without HIV were included. The prevalence of depression was high among both pregnant and non-pregnant WWH. There was no significant difference between pregnant and non-pregnant WWH in depression scores, perceived stress scores, or social support scores at any time point. Compared to pregnant women without HIV, pregnant WWH reported worse outcomes on all psychosocial scales. Social support and loneliness were associated with an increased odds of depressive symptoms in the adjusted analysis.

**Conclusions:** A high burden of adverse psychosocial outcomes was observed in both pregnant and non-pregnant women living with HIV compared to pregnant women without HIV. Loneliness and inadequate social support were associated with increased odds of depression in pregnancy and should be a focus in future support interventions.

**Keywords:** Psychosocial health, depression, pregnancy, postpartum, women with HIV, 2BMOM

\*Correspondence: [ellen.froesig.moseholm.larsen@regionh.dk](mailto:ellen.froesig.moseholm.larsen@regionh.dk)

<sup>1</sup> Department of Infectious Diseases, Copenhagen University Hospital, Kettegård Alle 30, 2650 Hvidovre, Hvidovre, Denmark

Full list of author information is available at the end of the article



## Background

The success of combination antiretroviral therapy (cART) has resulted in a dramatic decrease in perinatal transmission of human immunodeficiency virus (HIV) to less than 1% in most Western countries, in addition to a normalization of pregnancy [1, 2]. This has resulted in an increase in the annual number of HIV pregnancies, both internationally and in Nordic countries [1, 3, 4]. However, despite these advancements, women living with HIV (WWH) still face complex psychosocial and medical issues during pregnancy, childbirth, and postpartum [5, 6].

Pregnancy and the postpartum period are times of significant biological, social, and psychological changes for a woman and HIV adds another layer of complexity [7–9]. For WWH, pregnancy may be a time of increased psychological vulnerability due to different contextual factors, disclosure issues, and HIV-related stigma [6, 9, 10]. Moreover, in addition to the usual stresses of new motherhood, WWH must also cope with stressors that include their own health, the unknown infectious states of their infants, and attending to their infants' unique needs such as the administration of prophylactic antiretroviral medication [6, 11].

Psychosocial and emotional well-being during pregnancy is essential to develop the capacity to parent effectively and facilitate attachment to the new-born [12]. It is also well documented that maternal mental health during pregnancy and postpartum has a big impact child development and behaviour [13]. Research conducted in North America and in low- and middle-income settings [14] have reported a high prevalence of depression in pregnant and postpartum WWH [14–16]. A recent meta-analysis reported a significantly increased odds ratio of both antenatal and postnatal depressive symptoms in WWH compared to women without HIV (WWOH) [17]. Depression is one of the most common distress conditions in pregnancy and may be associated with the physical and emotional wellbeing of the mother, in addition to a range of adverse behavioural and emotional outcomes for the child [18]. For WWH adverse mental health outcomes may also affect HIV disease management during pregnancy or postpartum [14]. In a recent meta-analysis by Zhu et al. [17], optimal ART adherence was reported in 76% of pregnant and 53% of postpartum WWH worldwide, and depression and emotional stress has been identified as barriers to adherence [19].

The broader social and environmental context may be important determinants for mental health problems during pregnancy and postpartum. Previous research among WWOH have reported a significant association between poor maternal social support and depressive

symptoms in pregnancy [20, 21] and the postpartum period [22]. WWH have been found to have lower levels of social support when compared to WWOH [23], and research among pregnant WWH have shown that perceived stress and social isolation may be important factors associated with higher levels of depression [24–26].

There is a lack of research exploring WWH's varied experiences of pregnancy and motherhood in a Western context, where medical systems allow for appropriate medical treatment to all WWH.

Using quantitative data from the 2BMOM Study, a multicentre longitudinal mixed methods study among pregnant WWH, non-pregnant WWH, and pregnant WWOH in the Nordic countries Denmark, Finland, and Sweden [27], we aim to explore psychosocial health outcomes of WWH across the pregnancy–postpartum trajectory, assess differences in psychosocial health outcomes between pregnant WWH, non-pregnant WWH and pregnant WWOH, and identify factors associated with depression in the third trimester and postpartum for WWH and WWOH.

## Methods

### Setting

There are approximately 1600, 1000, and 2800 WWH in Denmark, Finland, and Sweden, respectively [4, 28, 29]. The majority of WWH in Scandinavia are immigrants, mainly from sub-Saharan Africa, and mainly infected with HIV by sexual contact [4, 30]. The healthcare system in the Nordic countries is tax-based and ensures universal access to medical health care, including antenatal, perinatal, and postpartum care, and many social support services [31]. cART is provided free of charge and people with HIV in Nordic countries are generally well treated with life expectancies approaching those of the general population [4, 29, 32].

### The 2BMOM study

The 2BMOM study is a multicenter longitudinal mixed-methods study investigating psychosocial outcomes and experiences of WWH in Nordic countries during pregnancy and early motherhood. The study recruited pregnant WWH, non-pregnant WWH, and pregnant WWOH from seven sites in Denmark, Finland, and Sweden between January 2019 and December 2020. Quantitative data was collected by self-administered electronic questionnaires, with a sub-sample of pregnant WWH taking part in semi-structured qualitative interviews ( $n=31$ ). Methods are described in detail elsewhere [27]. All women gave informed consent to participate and

the study was approved by the Danish Data Protection Agency (VD-2018-253), and the Finnish and Swedish Ethics Committees (HUS/1330/2019 and Dnr: 2019–04451, respectively).

### Study population

Pregnant and non-pregnant WWH were consecutively recruited from the participating sites (Departments of Infectious Diseases at Copenhagen University Hospitals, Hvidovre and Rigshospitalet; Odense -, Aalborg – and Aarhus University Hospitals in Denmark, Department of Infectious Diseases, Helsinki University Hospital, Finland, and Department of Infectious Diseases, Sahlgrenska University Hospital, Sweden) by the medical staff during routine clinical appointments. Pregnant WWH were asked to participate if they were 18 years of age or older, pregnant, anticipated birth of a viable infant without life-threatening conditions or congenital anomalies, and could speak and read Danish, Finnish, Swedish or English. Non-pregnant WWH were asked to participate if they were between 18 and 45 years of age, not pregnant or planning to become pregnant, and could speak and read Danish, Finnish, Swedish, or English. Pregnant WWOH were consecutively recruited from the Department of Obstetrics at Copenhagen University Hospital, Hvidovre, Denmark, which has the largest maternity ward in Denmark with approximately 7000 deliveries per year, and the Department of Obstetrics at Helsinki University Hospital, Finland. Women were asked to participate if they were 18 years of age or older, pregnant, not living with HIV, anticipated birth of a viable infant without life-threatening conditions or congenital anomalies, had no known chronic or psychiatric illness associated with increased surveillance or adverse pregnancy and birth outcome, and could speak and read Danish, Finnish or English.

### Data collection

Participants answered electronic standardized questionnaires via REDCap® at three time points; in the 3rd trimester, 3 and 6 months postpartum (T1, T2, and T3, respectively). Non-pregnant WWH followed the same timeframe, i.e. baseline following enrolment, 3–4 months, and 6–7 months. At each timepoint, a survey link was sent to the participants, who then completed the survey at home.

### Demographics and medical information

The demographic variables were collected at baseline (T1) and included self-reported marital status, education,

and employment status. Information on clinical variables (maternal country of birth, maternal age at delivery, parity, CD4 cell count, HIV RNA viral load, co-morbidity, cART, non-HIV medication, smoking, alcohol, and drug use) was obtained from the patients' medical records.

### Survey instrument

Information on different psychosocial health outcomes were collected using the following validated scales:

#### Depression

The Edinburgh Postnatal Depression Scale (EPDS) was used to assess symptoms of perinatal depression [33]. The scale has 10-items with responses on a 4-point Likert scale ranging from 0 (absence of depressive moods) to 3 (worst mood). A total score ranging from 0 to 30 is calculated, and a cut-off point of  $\geq 12$  indicates an increased likelihood of clinical depression [33]. The EPDS has been translated into Danish, Finnish, and Swedish, has a well-documented validity and reliability, and has been shown to be sensitive to changes over time [33, 34]. The scale does not mention the words pregnancy, child, birth or infant, and has also been validated in a non-pregnant population [35, 36].

#### Perceived stress

Perceived stress was measured using the Perceived Stress Scale – 10 item (PSS-10), a self-administered scale that was developed to measure “the degree to which situations in one’s life are appraised as stressful” during the past month [37]. Responses are on a Likert-type five-point format that ranges from 0 (never) to 4 (very often). The range for scores is 0 to 40 with higher scores reflecting greater perceptions of stress. The PSS-10 has been translated and validated in Danish, Finnish, and Swedish [38–40].

#### Social isolation

Perceived social isolation was assessed using the short version of the 3rd version of the University of California, Los Angeles (UCLA) loneliness scale [41, 42]. The scale consists of three items with Likert scale responses ranging from 1 (hardly ever) to 3 (often). Responses are summed into a total score (ranging from 3 to 9), where higher scores indicate a greater degree of loneliness [41, 42]. A score  $> 7$  has been suggested as a cut-off for severe loneliness [43]. The UCLA loneliness scale is a widely used self-report scale measuring loneliness, with acceptable validity and reliability [41, 44]. The scale has been translated and validated in a Danish and Finnish context, respectively [45, 46]. The scale was translated into Swedish using a translation–back-translation procedure before data collection [47].

### Social support

The Multidimensional Scale of Perceived Social Support (MSPSS), a 12-item scale, was used to assess social support [48, 49]. The scale consists of three subscales; Family, Friends, and Significant others. Each item is answered from 1 (strongly disagree) to 5 (strongly agree). A total score ranging from 1 to 7 is calculated, with higher scores suggesting a greater level of perceived social support. The MSPSS is a frequently used measure of social support in somatic illness, and good reliability and validity have been reported, also in a Scandinavian context [50–53].

### Statistical analysis

Differences in baseline demographic characteristics were summarized and compared between pregnant WWH, and non-pregnant WWH and pregnant WWOH, respectively, using the Pearson's  $\chi^2$ -test or Student's unpaired T-test, as appropriate. Categorical variables were described as counts (%), and continuous variables were described as means (95% confidence intervals (CI)). Linear mixed effects models for repeated measures were used to assess changes in mean scores of the different psychosocial health outcomes (depression, stress, loneliness, and social support) over time and between groups. These models account for correlations between repeated measurements of the same women over time and also include all available data, i.e. if a woman had missing data at one time point they were not deleted from the analysis, and the available data from previous time points was included resulting in more precise estimates. Within-subject residuals were modeled with an unstructured variance-covariance structure. All mixed models were bootstrapped with 2000 repetitions to account for the nonparametric distribution of data. Univariate and multivariate logistic regression models were completed to identify factors associated with probable depression in pregnancy and postpartum, respectively. This analysis was restricted to pregnant WWH and pregnant WWOH with a pregnancy depression score ( $n=45$  and  $n=146$ ) and a postpartum depression score ( $n=36$  and  $n=122$ ). Probable depression was defined using the validated cut-off score on the EPDS scale (score  $\geq 12$ ). The variables included in the multivariate model were chosen based on the results from the univariate analysis using a step-wise selection to choose the most parsimonious model. The final model selection was based on Akaike's Information Criterion. All models included living with HIV, perceived stress, loneliness, and social support in pregnancy, relationship status, country of birth, and education. Unknown/other/missing categories were included in the analysis. The model with probable depression

postpartum as the dependent variable also included depression score in pregnancy. Analyses were performed using STATA 17 software and all reported  $p$ -values are two-sided using a significance level of 0.05.

### Results

In total, 71 pregnant WWH fulfilled the inclusion criteria during the study period; 57 agreed to participate. However, six women did not complete the baseline survey before delivery, two had a preterm delivery, and two women withdrew their consent. Thus, a total of 47 pregnant WWH were included in the study giving a response rate of 66%. The main reasons for non-participation were language barriers and major psychiatric or social complications. The number of pregnant WWH who also completed the follow-up surveys T2 and T3 were 38 (80%) and 37 (79%), respectively. A total number of 112 non-pregnant WWH agreed to participate, of which 75 (67%) completed T1 and were thus included, of these 50 (68%) women completed T2 and 49 (65%) completed T3. A total number of 168 pregnant WWOH agreed to participate, of which 147 (88%) completed T1, 132 (89%) completed T2, and 125 (85%) completed T3. Enrollment and data collection were for some women conducted during the COVID-19 pandemic (see Supplementary 1).

### Characteristics

The demographics of the study population are presented in Table 1. Compared to non-pregnant WWH, pregnant WWH were significantly younger, were more likely to be married or living with a partner, less likely to be employed part or full time, had fewer comorbidities, and higher viral loads at inclusion. None of the participants used illicit drugs, and none of the pregnant populations had any alcohol use. Two non-pregnant WWH had an increased use of alcohol. All pregnant WWH were virally suppressed at the time of delivery. Compared to pregnant WWOH, pregnant WWH were significantly more likely to be of non-Nordic origin, less likely to be married or living with a partner, to have a high level of education, to be employed part or full time, to be nulliparous, and more likely to deliver by a planned cesarean section. A larger proportion of WWH also had a preterm delivery.

### Difference in psychosocial outcomes over time and between groups

The mean scores of the psychosocial health outcomes for each group and at each time point are presented in Table 2, Figs. 1 and 2.

The mean EPDS score among pregnant WWH was in the third trimester 7.68 (95% CI: 6.49: 8.93). There was a significant decrease in depression scores over

**Table 1** Baseline characteristics

	WWH (n = 47)	Non-pregnant WWH (n = 75)	Pregnant WWOH (n = 147)	p-value
<b>Age, mean (95% CI)</b>	33.65 (32.14: 35.15)	36.01 (34.45: 37.57)	32.25 (31.57: 33.93)	<b>0.04*</b>
<b>Relationship status, n (%)</b>				<b>&lt; 0.01***</b>
Married/living with a partner	36 (76)	39 (52)	142 (97)	
Have a partner, but not living together	5 (11)	11 (15)	0	
Do not have a current partner	4 (9)	23 (31)	5 (3)	
Unknown/prefer not to say	<3 (4)	<3 (2)	0	
<b>Country of birth, n (%)</b>				<b>&lt; 0.001**</b>
Nordic country (Denmark; Finland or Sweden)	12 (26)	31 (41)	135 (91)	
Asia	0	4 (5)	0	
Africa	25 (53)	27 (36)	< 3 (1)	
Eastern Europe/Russia	6 (13)	8 (11)	0	
Other	4 (8)	5 (7)	11 (7)	
<b>Education, n (%)</b>				<b>&lt; 0.001**</b>
Primary school	3 (6)	6 (8)	3 (2)	
Secondary school	13 (28)	18 (24)	14 (10)	
Higher education (college/university)	27 (57)	50 (67)	129 (87)	
Unknown/Missing	4 (9)	< 3 (1)	< 3 (1)	
<b>Employment, n (%)</b>				<b>&lt; 0.01***</b>
Yes, part or full time	29 (62)	61 (81)	126 (86)	
<b>Smoking, n (%)</b>				0.09
During pregnancy	< 3 (2)	7 (9)	0	
Missing	0	< 3 (3)	8 (5)	
<b>Comorbidities, n (%)</b>	6 (13)	24 (32)	21 (14)	<b>0.02*</b>
<b>Number of children, n (%)</b>				<b>&lt; 0.001**</b>
0	18 (38)	26 (35)	85 (58)	
1	18 (38)	21 (28)	57 (39)	
≥ 2	11 (24)	28 (37)	5 (3)	
<b>Years since HIV diagnosis, mean (95% CI)</b>	9.45 (7.28: 11.61)	11.58 (10.04: 13.13)		0.10
<b>HIV diagnosis in pregnancy, n (%)</b>				
Yes	3 (6)			
<b>Mode of HIV transmission, n (%)</b>				0.71
Sexual	42 (89)	65 (87)		
Perinatal transmission	5 (11)	6 (8)		
Unknown/other	0	4 (5)		
<b>ART treatment<sup>+</sup>, n (%)</b>				0.36
NRTIs + NNRTI	11 (24)	10 (13)		
NRTIs + PI	16 (34)	22 (29)		
NRTIs + InSTI	18 (38)	39 (52)		
Other	< 3 (4)	4 (5)		
<b>CD4 cell count<sup>+</sup>, n (%)</b>				0.12
> 500 cells/mL	36 (76)	66 (88)		
200–500 cells/mL	6 (13)	7 (9)		
< 200 cells/mL	5 (11)	< 3 (3)		
<b>HIV viral load<sup>+</sup>, n (%)</b>				<b>0.02*</b>
< 50 copies/mL	39 (83)	73 (97)		
≥ 50 copies/mL	8 (17)	< 3 (3)		
<b>Mode of delivery, n (%)</b>				<b>&lt; 0.01**</b>
Vaginal delivery	32 (68)		104 (71)	

**Table 1** (continued)

	WWH ( <i>n</i> = 47)	Non-pregnant WWH ( <i>n</i> = 75)	Pregnant WWOH ( <i>n</i> = 147)	<i>p</i> -value
Elective caesarean delivery	9 (19)		7 (5)	
Acute caesarean delivery	4 (9)		19 (13)	
Unknown/missing	<3 (4)		17 (11)	
<b>Complications during delivery, <i>n</i> (%)</b>	12 (26)		36 (24)	0.58
Unknown/missing	7 (15)		21 (14)	
<b>Gestational age &lt; 37 weeks, <i>n</i> (%)</b>	4 (9)		<3 (1)	<b>0.03**</b>

*P* values are based on the Pearson's  $\chi^2$ -test, Fischers exact or unpaired T-test, as appropriate

\*Significant difference between pregnant WWH and non-pregnant WWH

\*\*Significant difference between pregnant WWH and pregnant WWOH

\*\*\*Significant difference between pregnant WWH and non-pregnant WWH, and significant difference between pregnant WWH and pregnant WWOH

<sup>†</sup> At time of inclusion

WWH Women living with HIV, WWOH: Women without HIV

time among pregnant WWH (mean difference between T1 and T3 EPDS score: -1.65 (95% CI: -2.99: -0.30), *p*-value=0.02). Using the recommended cutoff, the prevalence of probable depression among pregnant WWH was 24% in the third trimester, 36% at 3 months postpartum, and 22% at 6 months postpartum. There was no difference in depression scores or perceived stress scores between pregnant and non-pregnant WWH at any time point. Pregnant WWH reported significantly higher depression scores and more perceived stress at all time points compared to pregnant WWOH. Pregnant WWOH reported significantly less stress over time (mean difference between T1 and T3 PSS score: -1.09 (95% CI: -1.73: -0.44), *p*-value=<0.01) while there was no change in perceived stress in any of the WWH groups.

Pregnant WWH reported significantly lower scores on the UCLA loneliness scale at T1 compared to non-pregnant WWH and significantly higher loneliness scores at T1 compared to pregnant WWOH. However, over time there was a significant increase in loneliness scores among pregnant WWH (mean difference between T1 and T3 UCLA score: 0.46 (95% CI: 0.06: 0.85), *p*-value=0.03) and among pregnant WWOH over time (mean difference between T1 and T3 UCLA score: 0.70 (95% CI: 0.41: 0.98), *p*-value=<0.001). Thus, the proportion of women with a loneliness score above the cut-off increased over time, although the absolute numbers are low. There was no significant difference in loneliness scores between pregnant and non-pregnant WWH, and between pregnant WWH and pregnant WWOH at T2 and T3 (Fig. 1).

Pregnant and non-pregnant WWH reported similar MSPSS social support scores over time and there was no significant difference within or between groups (Fig. 2).

Compared to the pregnant WWOH, pregnant WWH at all times and in all sub-scales reported significantly less social support (Fig. 2). Pregnant WWOH reported significantly lower support from friends over time (mean difference between T1 and T3 MSPSS friends sub-scale: -0.16 (95% CI: -0.28: -0.05), *p*-value=0.03).

#### Factors associated with probable depression

The results of the univariate and multivariate analysis are presented in Table 3. In the univariate analysis, living with HIV, perceived stress, loneliness, social support, being born in a non-Nordic country, having a primary/secondary education and not being employed were associated with increased odds of probable depression in pregnancy. When adjusting for HIV status, different maternal characteristics, and pregnancy psychosocial outcomes, only loneliness score in pregnancy remained significantly associated with increased odds of probable depression in pregnancy (OR 1.56 (95% CI: 1.01: 2.41), *p*-value 0.04) while social support in pregnancy was associated with a decreased odds of probable depression in pregnancy (OR 0.50 (95% CI: 0.27: 0.95), *p*-value 0.03).

Factors associated with increased odds of probable postpartum depression were in the univariate analysis living with HIV, depression score in pregnancy, postpartum perceived stress, loneliness, and social support. Only postpartum loneliness remained significantly associated with probable postpartum depression in the adjusted analysis (OR 1.58 (95% CI 1.08: 2.31), *p*-value 0.02).

#### Discussion

Using quantitative data from the 2BMOM study [27] this study found a high prevalence of probable depression and other psychosocial challenges among WWH compared to WWOH throughout the pregnancy – postpartum


**Table 2** Psychosomatic health outcome at T1-T3 between pregnant women living with HIV (WWH), non-pregnant women living with HIV (WWH) and pregnant women without HIV (WWOH)

	Pregnant women with HIV			Non-pregnant women with HIV			Pregnant women without HIV			Difference between pregnant WWH and non-pregnant WWH			Difference between pregnant WWH and pregnant WWOH		
	T1	T2	T3	T1	T2	T3	T1	T2	T3	T1	T2	T3	T1	T2	T3
<b>Edinburg depression scale (EPDS)</b>															
Total score, mean (95% CI)	n = 45 7.68 (6.49: 8.93)	n = 36 7.00 (5.53: 8.47)	n = 36 6.04 (4.82: 7.25)	n = 74 8.19 (7.22: 9.15)	n = 49 8.83 (7.65: 10.01)	n = 46 7.58 (6.18: 8.97)	n = 146 4.34 (3.89: 4.79)	n = 130 4.53 (4.12: 4.94)	n = 122 4.07 (3.56: 4.57)	0.53	0.06	0.10	<0.01	<0.01	<0.01
Likely clinical depression (score ≥ 12), n (%)	11 (24)	13 (36)	8 (22)	14 (27)	20 (41)	17 (39)	13 (9)	11 (8)	10 (8)						
<b>Perceived Stress (PSS)</b>															
Total score, mean (95% CI)	n = 44 22.55 (21.24: 23.45)	n = 36 22.26 (20.63: 23.89)	n = 36 22.45 (21.28: 23.61)	n = 73 22.73 (21.97: 23.50)	n = 49 22.08 (20.90: 23.26)	n = 46 22.16 (21.08: 23.24)	n = 146 19.59 (19.04: 20.15)	n = 131 19.32 (18.61: 20.03)	n = 124 18.50 (17.94: 19.02)	0.80	0.86	0.74	<0.001	<0.01	<0.001
Low stress (score 0-13), n (%)	2 (5)	1 (3)	1 (3)	2 (3)	1 (2)	1 (2)	7 (5)	3 (2)	9 (7)						
Moderate stress (score 14-26), n (%)	33 (75)	29 (81)	29 (81)	52 (71)	39 (80)	38 (83)	128 (88)	125 (95)	112 (90)						
High stress (score 27-40), n (%)	9 (20)	6 (17)	6 (17)	19 (26)	9 (18)	7 (15)	11 (8)	3 (2)	3 (2)						
<b>Loneliness (UCLA)</b>															
Total score, mean (95% CI)	n = 44 4.30 (3.98: 4.61)	n = 36 4.63 (4.28: 4.98)	n = 36 4.70 (4.29: 5.11)	n = 72 5.01 (4.72: 5.31)	n = 47 5.07 (4.71: 5.43)	n = 46 4.89 (4.42: 5.36)	n = 146 3.71 (3.54: 3.87)	n = 130 4.19 (3.94: 4.44)	n = 123 4.40 (4.13: 4.68)	<0.01	0.11	0.59	<0.01	0.05	0.19
Lonely (score > 7), n (%)	0	2 (6)	3 (8)	8 (11)	3 (6)	6 (13)	1 (1)	6 (5)	8 (6)						


**Table 2** (continued)

	Pregnant women with HIV			Non-pregnant women with HIV			Pregnant women without HIV			Difference between pregnant WHH and non-pregnant WHH			Difference between pregnant WHH and pregnant WWOH		
	T1	T2	T3	T1	T2	T3	T1	T2	T3	T1	T2	T3	T1	T2	T3
<b>Social support (MSPSS)</b>															
Total score, mean (SD)	n=44 5.67 (5.33: 6.02)	n=36 5.65 (5.30: 5.99)	n=36 5.70 (5.43: 5.98)	n=72 5.62 (5.33: 5.91)	n=47 5.53 (5.23: 5.82)	n=46 5.67 (5.43: 5.89)	n=146 6.57 (6.37: 6.63)	n=130 6.54 (6.47: 6.61)	n=123 6.46 (6.40: 6.53)	0.83	0.61	0.87	<0.001	<0.001	<0.001
Significant other, mean (SD)	1-7 6.32 (6.13: 6.52)	6.24 (5.80: 6.68)	6.34 (6.11: 6.57)	6.09 (5.91: 6.28)	5.99 (5.69: 6.30)	5.91 (5.68: 6.14)	6.85 (6.77: 6.94)	6.79 (6.73: 6.85)	6.77 (6.69: 6.85)	0.10	0.31	0.06	<0.01	0.02	<0.001
Family, mean (SD)	1-7 5.65 (5.23: 6.06)	5.58 (5.16: 6.00)	5.54 (5.10: 5.97)	5.50 (5.14: 5.86)	5.40 (5.07: 5.73)	5.64 (5.35: 5.92)	6.52 (6.42: 6.61)	6.51 (6.40: 6.62)	6.45 (6.33: 6.56)	0.65	0.53	0.73	<0.001	0.001	<0.01
Friends, mean (SD)	1-7 5.05 (4.68: 5.43)	5.13 (4.76: 5.49)	5.22 (4.82: 5.62)	5.27 (4.91: 5.62)	5.20 (4.73: 5.68)	5.44 (5.09: 5.79)	6.33 (6.20: 6.46)	6.31 (6.16: 6.47)	6.17 (6.01: 6.33)	0.47	0.77	0.50	<0.001	<0.001	<0.001





**Fig. 1** Depression, Perceived stress and Loneliness among pregnant women living with HIV (WWH), non-pregnant WWH, and pregnant women without HIV (WWOH) over time. Figure legend: EPDS Depression scale: A higher score = more depressive symptoms. PSS Perceived stress scale: A higher score = more perceived stress. UCLA Loneliness scale: A higher score = more loneliness



trajectory. There were no significant differences in depression scores, perceived stress scores, or social support scores between pregnant WWH and non-pregnant WWH at any time point. Loneliness was significantly associated with increased odds of probable depression in pregnancy and postpartum, while social support was associated with a decreased odds of probable depression in pregnancy.

Both pregnant and non-pregnant WWH had a high prevalence of probable depression (24% and 27% at T1, respectively) compared to pregnant WWOH (9% at T1). A high prevalence of depression among both pregnant and non-pregnant WWH is well described in the literature [15, 17, 23, 54, 55]. Pregnant WWH reported significantly higher depression scores at all time points compared to pregnant WWOH. This is contrary to findings from several US-based studies reporting no difference in depression scores between pregnant WWH and WWOH [15, 23, 56]. However, in these studies depressive symptoms were overall high, making it difficult to detect significant differences between groups.

Pregnant WWH had significantly lower depression scores over time. Pregnancy and becoming a mother may have a beneficial effect on women's mental health and may act as a buffer to some of the negative effects of living with HIV [6, 57, 58]. A small study among WWH in Spain found that although pregnancy was associated with more negative emotions such as anxiety, fear, guilt, and sadness compared to WWOH, pregnancy was also associated with positive emotions, such as happiness [57]. This is supported by other studies, where motherhood is found to be one factor that lightens the burden of living with HIV [6, 58, 59]. We also found that pregnant WWH reported significantly lower loneliness scores in pregnancy compared to non-pregnant WWH.

Loneliness and perceived social support have been described as major risk factors for depression in pregnancy and postpartum in both WWH and WWOH [25, 60]. In our analysis, loneliness and low social support were significantly associated with increased odds of probable depression in pregnancy, while postpartum loneliness was significantly associated with increased

**Table 3** Logistic regression analysis of factors associated with probable depression in pregnancy and depression postpartum among pregnant women living with HIV and pregnant women without HIV

	Depression in pregnancy				Depression postpartum			
	Univariate		Adjusted <sup>a</sup>		Univariate		Adjusted <sup>a</sup>	
	OR (95% CI)	p-value	OR (95% CI)	p-value	OR (95% CI)	p-value	OR (95% CI)	p-value
<b>Living with HIV</b>	2.83 (1.19: 6.71)	<b>0.02</b>	0.75 (0.14; 4.17)	0.75	3.2 (1.16: 8.85)	<b>0.03</b>	2.39 (0.34: 16.76)	0.38
<b>Pregnancy depression score (EPDS score)</b>					4.55 (1.47: 14.07)	<b>&lt;0.01</b>	1.99 (0.46: 8.73)	0.92
<b>Stress (PSS score)</b>	1.15 (1.05: 1.27)	<b>&lt;0.01</b>	1.23 (1.00: 1.27)	0.05	1.20 (1.06: 1.35)	<b>&lt;0.01</b>	1.14 (0.96: 1.34)	0.14
<b>Loneliness (UCLA score)</b>	2.37 (1.67: 3.35)	<b>&lt;0.001</b>	1.56 (1.01: 2.41)	<b>0.04</b>	1.62 (1.23: 2.15)	<b>&lt;0.01</b>	1.58 (1.08: 2.31)	<b>0.02</b>
<b>Social support (MSPSS score)</b>	0.34 (0.22: 0.54)	<b>&lt;0.001</b>	0.50 (0.27: 0.95)	<b>0.03</b>	0.40 (0.24: 0.67)	<b>&lt;0.001</b>	0.58 (0.28: 1.19)	0.14
<b>Age</b>	1.03 (0.22: 0.54)	0.57			0.93 (0.82: 1.05)	0.22		
<b>Relationship status</b>								
Married/living with a partner	ref		ref		ref		ref	
Have a partner, but not living together	2.33 (0.23: 23.43)	0.47	0.57 (0.04: 7.92)	0.68	4.13 (0.35: 48.08)	0.26	3.20 (0.02: 449.36)	0.65
Do not have a current partner/prefer not to say	2.63 (0.65: 10.65)	0.18	0.65 (0.06: 6.74)	0.72	1.38 (0.16: 12.16)	0.78	1.00 (0.09: 11.33)	0.99
<b>Country of birth</b>								
Nordic country (Denmark; Finland or Sweden)	ref		ref		ref		ref	
Not Nordic country	3.19 (1.31: 7.79)	<b>0.01</b>	1.16 (0.20: 0.86)	0.87	2.54 (0.87: 7.46)	0.09	0.28 (0.03: 2.56)	0.26
<b>Education</b>								
Higher education (college/university)	ref		ref		ref			
Primary/secondary school	3.5 (1.43: 8.59)	<b>&lt;0.01</b>	1.74 (0.53: 5.63)	0.36	2.06 (0.67: 6.34)	0.21	0.31 (0.05: 1.74)	0.18
<b>Employment</b>								
Yes, part or full time	0.30 (0.12: 0.73)	<b>&lt;0.01</b>			0.81 (0.25: 2.68)	0.73		
<b>Number of children</b>								
0	ref				ref			
≥ 1	1.97 (0.84: 4.59)	0.12			0.95 (0.35: 2.55)	0.92		
<b>Mode of delivery</b>								
Vaginal delivery	ref				ref			
Caesarean delivery	1.02 (0.35: 2.99)	0.97			0.87 (0.23: 3.27)	0.83		
<b>Preterm (gestational age &lt; 37 weeks)</b>	3.35 (0.58: 19.32)	0.18			–			

<sup>a</sup> Adjusted for living with HIV, perceived stress, loneliness, and social support in pregnancy, relationship status, country of birth, and education. The postpartum depression model also included depression score in pregnancy

odds of probable postpartum depression. Living with HIV was not significantly associated with depression in the multivariate analysis, suggesting that social and/or psychological factors and not HIV per se may be the main contributor to depression. Although not significant in the multivariate analysis, stress was associated with increased odds of probable depression in pregnancy. Living with HIV can contribute to women's experience of shame, exclusion, and rejection from family, partners, and even healthcare providers [6, 58, 61]. This can lead to an increased feeling of isolation if support is limited, in addition to stress related to lack of control or limited feeling of resilience [25].

Both WWH and pregnant WWOH reported an increase in loneliness scores postpartum. This is in line with other studies [25, 60], suggesting that the postpartum period may be a period of increased parenting responsibility and limited social activities which may increase the feeling of loneliness. Moreover, WWH experienced lower social support on all domains compared to pregnant WWOH. Studies among pregnant and postpartum WWH have reported an increased risk of depression with poorer social support [23, 54]. Pregnant WWH were more likely to not be married or living with a partner than pregnant WWOH. Research among WWOH suggests that increases in both partner and family support

may be powerful protective factors for decreasing mental health difficulties in pregnancy and postpartum, highlighting the importance of targeting and increasing this type of support from pregnancy to the postpartum period [62]. Other supportive conditions for perinatal wellbeing include having access to high-quality information on HIV and being ensured the same rights as other women [63].

To our knowledge, this is the first study examining psychosocial health across the pregnancy – postpartum trajectory among pregnant WWH compared to pregnant WWOH and non-pregnant WWH. Hence, the results of this study provide a comprehensive knowledge of the psychosocial experiences and emotional care needs of WWH in a Western setting. Another strength is the use of validated scales to assess the different psychosocial outcomes. Although we included a nationwide sample of pregnant WWH in Denmark, in addition to a sample of WWH in both Finland and Sweden, the small number of pregnant WWH and the fact that they all live in Nordic countries may limit the generalizability of the results. The main reason for non-participation were language barriers and major psychiatric or social complications. Hence, a vulnerable group of women were not included, and the results may therefore reflect women with more resilience. The participants were followed longitudinally allowing us to capture changes in experiences over time. However, there was a gradual loss to follow-up over time, which further limited our sample. Enrollment and data collection was for some women conducted during the COVID-19 pandemic, which could potentially have influenced our findings in a more adverse direction [64]. However, the proportion of women who completed the survey during the COVID-19 pandemic were similar across groups, and the within-group change over time was limited and consistent with the literature.

## Conclusion

Understanding and responding to the experiences of pregnant and postpartum women living with HIV is important so that goal-oriented interventions supporting these women can be developed. Women living with HIV reported more depressive symptoms, perceived stress, and loneliness, and less social support across the pregnancy – postpartum trajectory compared to pregnant women without HIV. Compared to non-pregnant women living with HIV, pregnant women living with HIV were less lonely while pregnant. The prevalence of probable depression was high among both pregnant and non-pregnant women living with HIV. Loneliness and inadequate social support are important factors

associated with an increased odds of depressive symptoms during pregnancy and should be a focus in future support interventions.

## Abbreviations

cART: Combination Antiretroviral Therapy; EPDS: The Edinburgh Postnatal Depression Scale; HIV: Human Immunodeficiency Virus; MSPSS: The Multidimensional Scale of Perceived Social Support; PSS-10: Perceived Stress Scale – 10 item; UCLA Loneliness Scale: University of California, Los Angeles (UCLA) loneliness scale; WWH: Women living with HIV; WWOH: Women without HIV.

## Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12884-021-04357-5>.

**Additional file 1: Supplementary 1.** Number of women who completed the survey during the COVID-19 pandemic (after 1 March 2020) by group and timepoint

## Acknowledgements

We thank all the participants for contributing to this study and for taking the time to complete the survey over time. We also thank the doctors, nurses, and midwives at the participating sites in Denmark, Finland, and Sweden for their invaluable help with the enrollment of participants.

## Authors' contributions

EM and NW had the original concept for the study. All authors contributed to study design, data collection, data interpretation, writing the report, and approved the final version. EM had full access to the data and did the statistical analysis.

## Funding

The 2BMOM study was funded by The Novo Nordisk Foundation (Grant Number: NNF17OC0029508 and NNF18OC0052512) and Gilead Sciences (Grant Number: 220002078). The funders had no role in the study design, data collection, data analysis, data interpretation, or writing of the report. The corresponding author had full access to all the data in the study and had final responsibility for the decision to submit for publication.

## Availability of data and materials

The data that support the findings of this study are available upon approval from the relevant Regulatory agencies (the Danish Data Protection Agency and the Finnish and Swedish Ethics Committees), but restrictions apply to the availability of these data, which were used under license for the current study, and so are not publicly available. Data are however available from the authors upon reasonable request and with permission of the Danish Data Protection Agency, and the Finnish and Swedish Ethics Committees.

## Declarations

### Ethics approval and consent to participate

All women gave informed consent to participate and the study was approved by the Danish Data Protection Agency (VD-2018-253), and the Finnish and Swedish Ethics Committees (HUS/1330/2019 and Dnr: 2019-04451, respectively). All methods were performed in accordance with the Helsinki Declaration and the relevant guidelines and regulations.

### Consent for publication

Not applicable.

### Competing interests

EM reports personal fees from Gilead, outside the submitted work: honorarium paid to her institution. IA reports personal fees from Abbvie, Merck Sharp Dohme, Gilead, and Glaxo Smith Kline, and a grant from Gilead, outside the submitted work. ÅM received honoraria paid to her institution from Gilead

and Viiv/ GSK. TLK reports personal fees and grants from Viiv/Glaxo Smith Kline, Gilead, CLS Behring, and Baxalta, outside of the submitted work. NW reports personal fees from Abbvie, Bristol-Myers-Squibb, Merck Sharp Dohme, Gilead, Glaxo Smith Kline, outside the submitted work; honorarium paid to her institution, and unrestricted research grants from Abbvie and Gilead with no relation to the submitted work. The remaining authors (GP, ISJ, DB, and MS) declare no conflicts of interest.

#### Author details

<sup>1</sup>Department of Infectious Diseases, Copenhagen University Hospital, Kettegård Alle 30, 2650 Hvidovre, Hvidovre, Denmark. <sup>2</sup>Department of Public Health, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark. <sup>3</sup>Department of Infectious Diseases, Helsinki University Hospital, Helsinki, Finland. <sup>4</sup>Department of Infectious Diseases, Region Västera Gotland, Sahlgrenska University Hospital, Gothenburg, Sweden. <sup>5</sup>Department of Infectious Diseases, Institute of Biomedicine, Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden. <sup>6</sup>Department of Infectious Diseases, Aalborg University Hospital, Aalborg, Denmark. <sup>7</sup>Department of Infectious Diseases, Copenhagen University Hospital, Rigshospitalet, Copenhagen, Denmark. <sup>8</sup>Department of Infectious Diseases, Odense University Hospital, Odense, Denmark. <sup>9</sup>Department of Gynecology and Obstetrics, Copenhagen University Hospital, Hvidovre, Hvidovre, Denmark. <sup>10</sup>Department of Infectious Diseases, Aarhus University Hospital, Aarhus, Denmark. <sup>11</sup>Department of Clinical Medicine, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark.

Received: 8 September 2021 Accepted: 22 December 2021

Published online: 07 January 2022

#### References

- von Linstow ML, Rosenfeldt V, Lebeck AM, Storgaard M, Hornstrup T, Katzenstein TL, et al. Prevention of mother-to-child transmission of HIV in Denmark, 1994–2008. *HIV Med.* 2010;11:448–56.
- Townsend CL, Byrne L, Cortina-Borja M, Thorne C, de Ruiter A, Lyall H, et al. Earlier initiation of ART and further decline in mother-to-child HIV transmission rates, 2000–2011. *AIDS.* 2014;28:1049–57.
- French CE, Cortina-Borja M, Thorne C, Tooke PA. Incidence, patterns, and predictors of repeat pregnancies among HIV-infected women in the United Kingdom and Ireland, 1990–2009. *J Acquir Immune Defic Syndr.* 2012;59:287–93.
- Aho I, Kivela P, Kajjama M, Surcel HM, Ristola M, Heikinheimo O, et al. Comprehensive nationwide analysis of mother-to-child HIV transmission in Finland from 1983 to 2013. *Epidemiol Infect.* 2018;146:1301–7.
- Greene S, Ion A, Kwaramba G, Smith S, Loutfy MR. “Why are you pregnant? What were you thinking?”: How women navigate experiences of HIV-related stigma in medical settings during pregnancy and birth. *Soc Work Health Care.* 2016;55:161–79.
- Ion A, Greene S, Mellor K, Kwaramba G, Smith S, Barry F, et al. Perinatal care experiences of mothers living with HIV in Ontario, Canada. *J HIV/AIDS Soc Serv.* 2016;15:180–201.
- Kendall CE, Wong J, Taljaard M, Glazier RH, Hogg W, Younger J, et al. A cross-sectional, population-based study measuring comorbidity among people living with HIV in Ontario. *BMC Public Health.* 2014;14:161.
- Slot M, Sodemann M, Gabel C, Holmskov J, Laursen T, Rodkjaer L. Factors associated with risk of depression and relevant predictors of screening for depression in clinical practice: a cross-sectional study among HIV-infected individuals in Denmark. *HIV Med.* 2015;16:393–402.
- Waldron EM, Burnett-Zeigler I, Wee V, Ng YW, Koenig LJ, Pederson AB, et al. Mental Health in Women Living With HIV: The Unique and Unmet Needs. *J Int Assoc Provid AIDS Care.* 2021;20:2325958220985665.
- Ion A, Wagner AC, Greene S, Loutfy MR. HIV Mothering Study Team. HIV-related stigma in pregnancy and early postpartum of mothers living with HIV in Ontario, Canada. *AIDS Care.* 2017;29:137–44.
- Moseholm E, Helleberg M, Nordly SB, Rosenfeldt V, Storgaard M, Pedersen G, et al. Hospital admission among HIV-exposed uninfected children compared with HIV-unexposed children. *AIDS.* 2016;30:2697–706.
- Sanders LB. Women's voices: the lived experience of pregnancy and motherhood after diagnosis with HIV. *J Assoc Nurses AIDS Care.* 2008;19:47–57.
- Stein A, Pearson RM, Goodman SH, Rapa E, Rahman A, McCallum M, et al. Effects of perinatal mental disorders on the fetus and child. *Lancet.* 2014;384:1800–19.
- Kapetanovic S, Dass-Brailsford P, Nora D, Talisman N. Mental health of HIV-seropositive women during pregnancy and postpartum period: a comprehensive literature review. *AIDS Behav.* 2014;18:1152–73.
- Rubin LH, Cook JA, Grey DD, Weber K, Wells C, Golub ET, et al. Perinatal depressive symptoms in HIV-infected versus HIV-uninfected women: a prospective study from preconception to postpartum. *J Women's Health (Larchmt).* 2011;20:1287–95.
- Malee KM, Mellins CA, Huo Y, Tassiopoulos K, Smith R, Sirois PA, et al. Prevalence, incidence, and persistence of psychiatric and substance use disorders among mothers living with HIV. *J Acquir Immune Defic Syndr.* 2014;65:526–34.
- Zhu Q-Y, Huang D-S, Lv J-D, Guan P, Bai X-H. Prevalence of perinatal depression among HIV-positive women: a systematic review and meta-analysis. *BMC Psychiatry.* 2019;19:330.
- Goodman SH, Rouse MH, Connell AM, Broth MR, Hall CM, Heyward D. Maternal depression and child psychopathology: a meta-analytic review. *Clin Child Fam Psychol Rev.* 2011;14:1–27.
- Nachega JB, Uthman OA, Anderson J, Peltzer K, Wampold S, Cotton MF, et al. Adherence to antiretroviral therapy during and after pregnancy in low-income, middle-income, and high-income countries: a systematic review and meta-analysis. *AIDS.* 2012;26:2039–52.
- Wajid A, van Zanten SV, Mughal MK, Biringner A, Austin M-P, Vermeyden L, et al. Adversity in childhood and depression in pregnancy. *Arch Womens Ment Health.* 2020;23:169–80.
- Westdahl C, Milan S, Magriples U, Kershaw TS, Rising SS, Ickovics JR. Social support and social conflict as predictors of prenatal depression. *Obstet Gynecol.* 2007;110:134–40.
- Schwab-Reese LM, Schafer EJ, Ashida S. Associations of social support and stress with postpartum maternal mental health symptoms: Main effects, moderation, and mediation. *Women Health.* 2017;57:723–40.
- Bonacquisti A, Geller PA, Aaron E. Rates and predictors of prenatal depression in women living with and without HIV. *AIDS Care.* 2014;26:100–6.
- Blaney NT, Fernandez MI, Ethier KA, Wilson TE, Walter E, Koenig LJ, et al. Psychosocial and behavioral correlates of depression among HIV-infected pregnant women. *AIDS Patient Care STDs.* 2004;18:405–15.
- Khan S, Ion A, Alyass A, Greene S, Kwaramba G, Smith S, et al. Loneliness and perceived social support in pregnancy and early postpartum of mothers living with HIV in Ontario. *Canada AIDS Care.* 2019;31:318–25.
- Bailey JL, Molino ST, Vega AD, Badowski M. A Review of HIV Pre-Exposure Prophylaxis: The Female Perspective. *Infect Dis Ther.* 2017;6:363–82.
- Moseholm E, Fetters MD, Aho I, Mellgren Å, Johansen IS, Storgaard M, et al. Study protocol: becoming and being a mother living with HIV - a multicentre longitudinal mixed methods study among pregnant women living with HIV, non-pregnant women living with HIV and pregnant women not living with HIV in a high-income setting (the 2B MOM study). *BMJ Open.* 2019;9:e027761.
- The Danish HIV Cohort. National Report 2020. Available at: [https://www.sundhed.dk/content/cms/63/97963\\_danhiv\\_2020.pdf](https://www.sundhed.dk/content/cms/63/97963_danhiv_2020.pdf). Accessed 5 Jan 2022.
- Marrone G, Mellgren Å, Eriksson LE, Svedhem V. High Concordance between Self-Reported Adherence, Treatment Outcome and Satisfaction with Care Using a Nine-Item Health Questionnaire in InfCareHIV. *PLoS One.* 2016;11:e0156916.
- Orbaek M, Thorsteinsson K, Helleberg M, Moseholm E, Katzenstein TL, Storgaard M, et al. Assessment of mode of delivery and predictors of emergency caesarean section among women living with HIV in a matched-pair setting with women from the general population in Denmark, 2002–2014. *HIV Med.* 2017; Journal Article. <https://doi.org/10.1111/hiv.12519>.
- Tynkkynen L-K, Alexandersen N, Kaarbøe O, Anell A, Lehto J, Vrangbæk K. Development of voluntary private health insurance in Nordic countries - An exploratory study on country-specific contextual factors. *Health Policy.* 2018;122:485–92.
- Obel N, Omland LH, Kronborg G, Larsen CS, Pedersen C, Pedersen G, et al. Impact of non-HIV and HIV risk factors on survival in HIV-infected patients on HAART: a population-based nationwide cohort study. *PLoS One.* 2011;6:e22698.

33. Cox JL, Holden JM, Sagovsky R. Detection of postnatal depression. Development of the 10-item Edinburgh Postnatal Depression Scale. *Br J Psychiatry*. 1987;150:782–6.
34. Tendais I, Costa R, Conde A, Figueiredo B. Screening for depression and anxiety disorders from pregnancy to postpartum with the EPDS and STAI. *Span J Psychol*. 2014;17:E7.
35. Santos IS, Tavares BF, Munhoz TN, Manzolli P, de Ávila GB, Jannke E, et al. Patient Health Questionnaire-9 versus Edinburgh Postnatal Depression Scale in screening for major depressive episodes: a cross-sectional population-based study. *BMC Res Notes*. 2017;10:57.
36. Matijasevich A, Munhoz TN, Tavares BF, Barbosa APPN, da Silva DM, Abitante MS, et al. Validation of the Edinburgh Postnatal Depression Scale (EPDS) for screening of major depressive episode among adults from the general population. *BMC Psychiatry*. 2014;14:284.
37. Cohen S, Kamarck T, Mermelstein R. A global measure of perceived stress. *J Health Soc Behav*. 1983;24:385–96.
38. Eskildsen A, Dalgaard VL, Nielsen KJ, Andersen JH, Zachariae R, Olsen LR, et al. Cross-cultural adaptation and validation of the Danish consensus version of the 10-item Perceived Stress Scale. *Scand J Work Environ Health*. 2015;41:486–90.
39. Järvelä-Reijonen E, Karhunen L, Sairanen E, Rantala S, Laitinen J, Puttonen S, et al. High perceived stress is associated with unfavorable eating behavior in overweight and obese Finns of working age. *Appetite*. 2016;103:249–58.
40. Eklund M, Bäckström M, Tuveson H. Psychometric properties and factor structure of the Swedish version of the Perceived Stress Scale. *Nord J Psychiatry*. 2014;68:494–9.
41. Russell DW. UCLA Loneliness Scale (Version 3): reliability, validity, and factor structure. *J Pers Assess*. 1996;66:20–40.
42. Hughes ME, Waite LJ, Hawley LC, Cacioppo JT. A Short Scale for Measuring Loneliness in Large Surveys: Results From Two Population-Based Studies. *Res Aging*. 2004;26:655–72.
43. Lasgaard M, Friis K, Shevlin M. “Where are all the lonely people?” A population-based study of high-risk groups across the life span. *Soc Psychiatry Psychiatr Epidemiol*. 2016;51:1373–84.
44. Hartshorne TS. Psychometric properties and confirmatory factor analysis of the UCLA loneliness scale. *JPersAssess*. 1993;61:182–95.
45. Lasgaard M. Reliability and validity of the Danish version of the UCLA Loneliness Scale. *Personal Individ Differ*. 2007;42:1359–66.
46. Schuez-Havupalo L, Lahti E, Junttila N, Toivonen L, Aromaa M, Rautava P, et al. Parents’ depression and loneliness during pregnancy and respiratory infections in the offspring: A prospective birth cohort study. *PLoS One*. 2018;13:e0203650.
47. Garcia-Castillo D, Fetters MD. Quality in medical translations: a review. *J Health Care Poor Underserved*. 2007;18:74–84.
48. Zimet GD, Dahlem NW, Zimet SG, Farley GK. The Multidimensional Scale of Perceived Social Support. *JPersAssess*. 1988;52:30–41.
49. Zimet GD, Powell SS, Farley GK, Werkman S, Berkoff KA. Psychometric characteristics of the Multidimensional Scale of Perceived Social Support. *J Pers Assess*. 1990;55:610–7.
50. Shumaker SC, Frazier SK, Moser DK, Chung ML. Psychometric Properties of the Multidimensional Scale of Perceived Social Support in Patients With Heart Failure. *JNursMeas*. 2017;25:90–102.
51. Pedersen SS, Spinder H, Erdman RAM, Denollet J. Poor perceived social support in implantable cardioverter defibrillator (ICD) patients and their partners: cross-validation of the multidimensional scale of perceived social support. *Psychosomatics*. 2009;50:461–7.
52. Ekback M, Benzein E, Lindberg M, Arestedt K. The Swedish version of the multidimensional scale of perceived social support (MSPSS)—a psychometric evaluation study in women with hirsutism and nursing students. *HealthQualLifeOutcomes*. 2013;11:Journal Article:168–7525–11–168.
53. Karukivi M, Joukamaa M, Hautala L, Kaleva O, Haapasalo-Pesu K-M, Liuksila P-R, et al. Does perceived social support and parental attitude relate to alexithymia? A study in Finnish late adolescents. *Psychiatry Res*. 2011;187:254–60.
54. Bailey H, Malyuta R, Semenenko I, Townsend CL, Cortina-Borja M, Thorne C, et al. Prevalence of depressive symptoms in pregnant and postnatal HIV-positive women in Ukraine: a cross-sectional survey. *Reprod Health*. 2016;13:27.
55. Arseniou S, Arvaniti A, Samakouri M. HIV infection and depression. *Psychiatry Clin Neurosci*. 2014;68:96–109.
56. Aaron E, Bonacquisti A, Geller PA, Polansky M. Perinatal Depression and Anxiety in Women with and without Human Immunodeficiency Virus Infection. *Womens Health Issues*. 2015;25:579–85.
57. Pereira M, Canavarro MC. Quality of life and emotional distress among HIV-positive women during transition to motherhood. *Span J Psychol*. 2012;15:1303–14.
58. Sandelowski M, Barroso J. Motherhood in the context of maternal HIV infection. *Res Nurs Health*. 2003;26:470–82.
59. Carlsson-Laloo E, Rusner M, Mellgren Å, Berg M. Sexuality and Reproduction in HIV-Positive Women: A Meta-Synthesis. *AIDS Patient Care STDs*. 2016;30:56–69.
60. Luoma I, Korhonen M, Puura K, Salmelin RK. Maternal loneliness: concurrent and longitudinal associations with depressive symptoms and child adjustment. *Psychol Health Med*. 2019;24:667–79.
61. Paudel V, Baral KP. Women living with HIV/AIDS (WLHA), battling stigma, discrimination and denial and the role of support groups as a coping strategy: a review of literature. *Reprod Health*. 2015;12:53.
62. Racine N, Plamondon A, Hentges R, Tough S, Madigan S. Dynamic and bidirectional associations between maternal stress, anxiety, and social support: The critical role of partner and family support. *J Affect Disord*. 2019;252:19–24.
63. Carlsson-Laloo E, Mellgren Å, Berg M, Rusner M. Supportive conditions for sexual and perinatal wellbeing in women living with HIV in Sweden: A phenomenological study. *Sex Reprod Healthc*. 2021;29:100640.
64. Wastnedge EAN, Reynolds RM, van Boeckel SR, Stock SJ, Denison FC, Maybin JA, et al. Pregnancy and COVID-19. *Physiol Rev*. 2021;101:303–18.

## Publisher’s Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

