

RESEARCH

Open Access

The interaction of celecoxib with MDR transporters enhances the activity of mitomycin C in a bladder cancer cell line

Vincenzo Pagliarulo^{1*}, Patrizia Ancona¹, Mauro Niso², Nicola Antonio Colabufo², Marialessandra Contino², Luigi Cormio³, Amalia Azzariti⁴ and Arcangelo Pagliarulo¹

Abstract

Background: An *in vitro* model was developed to understand if celecoxib could synergize with Mitomycin C (MMC), commonly used for the prevention of non-muscle invasive bladder cancer recurrence, and eventually elucidate if the mechanism of interaction involves multi drug resistance (MDR) transporters.

Methods: UMUC-3, a non COX-2 expressing bladder cancer cell line, and UMUC-3-CX, a COX-2 overexpressing transfectant, as well as 5637, a COX-2 overexpressing cell line, and 5637si-CX, a non COX-2 expressing silenced 5637 cell line, were used in the present study. The expression of COX-2 and MDR pumps (P-gp, MDR-1 and BCRP) was explored through western blot. The anti-proliferative effect of celecoxib and MMC was studied with MTT test. Three biological permeability assays (Drug Transport Experiment, Substrate Transporter Inhibition, and ATP cell depletion) were combined to study the interaction between MDR transporters and celecoxib. Finally, the ability of celecoxib to restore MMC cell accumulation was investigated.

Results: The anti-proliferative effect of celecoxib and MMC were investigated alone and in co-administration, in UMUC-3, UMUC-3-CX, 5637 and 5637si-CX cells. When administered alone, the effect of MMC was 8-fold greater in UMUC-3. However, co-administration of 1 μ M, 5 μ M, and 10 μ M celecoxib and MMC caused a 2,3-fold cytotoxicity increase in UMUC-3-CX cell only. MMC cytotoxicity was not affected by celecoxib co-administration either in 5637, or in 5637si-CX cells. As a result of all findings from the permeability experiments, celecoxib was classified as P-gp unambiguous substrate: celecoxib is transported by MDR pumps and interferes with the efflux of MMC. Importantly, among all transporters, BCRP was only overexpressed in UMUC-3-CX cells, but not in 5637 and 5637si-CX.

Conclusions: The UMUC-3-CX cell line resembles a more aggressive phenotype with a lower response to MMC compared to the *wt* counterpart. However, the administration of celecoxib in combination to MMC causes a significant and dose dependent gain of the anti-proliferative activity. This finding may be the result of a direct interaction between celecoxib and MDR transporters. Indeed, BCRP is overexpressed in UMUC-3-CX, but not in UMUC-3, 5637, and 5637si-CX, in which celecoxib is ineffective.

Keywords: Cyclooxygenase-2, Multidrug resistance, Bladder cancer

* Correspondence: vincenzo.pagliarulo@uniba.it

¹Sezione di Urologia e Andrologia, Dipartimento dell'Emergenza e dei Trapianti di Organi (DETO), Università "Aldo Moro" di Bari, Piazza G. Cesare 11, Bari 70124, Italy

Full list of author information is available at the end of the article

Background

The natural history of non-muscle invasive bladder carcinoma (NMIBC) may be characterized by multiple recurrences according to patient risk stratification [1]. Following transurethral resection (TUR), endovesical chemotherapy is recommended by clinical practice guidelines in intermediate and high-risk patients [2]. Mitomycin C (MMC) is a potent DNA cross-linker commonly used as a bladder instillation to reduce the likelihood of bladder cancer recurrence and/or progression [3]. Nonetheless, early recurrence is common after MMC treatment and may recognize several causes, such as disease phenotype [1], the presence of residual tumor following a TUR [4], as well as downstaging [5]. Another possible cause may be the failure of cancer treatment due to resistance to structurally unrelated chemotherapeutic agents, also defined Multi Drug Resistance (MDR) [6,7]. MDR is usually associated with a decreased intracellular concentration of cytostatic drugs and with the overexpression of ATP-Binding Cassette (ABC) transporters, such as P-glycoprotein (P-gp), Breast Cancer Resistance Proteins (BCRP) and Multidrug Resistance associated Proteins (MRPs) [8-10]. These transporters cause the efflux of drugs and xenobiotics out from the cells [11]. Interestingly, their overexpression has been observed in urothelial cancer cells [12] and may be useful for selecting patients with bladder cancer to be candidates for neoadjuvant chemotherapy [13].

Cyclooxygenases (COX) are enzymes required for the conversion of arachidonic acid to prostaglandins. As COX-1 is constitutively expressed, COX-2 is highly induced in response to inflammatory signals [14]. Although specific COX-2 inhibitors were generated for pain relief and for their anti-inflammatory properties, experimental and translational studies have shown that COX-2 is involved in carcinogenesis, finally encouraging clinical testing [15]. Recently, COX-2 inhibitors have gained attention as chemosensitizers when combined with other agents [16,17]. This property may be exerted by interference with the activity of membrane proteins involved in MDR [18,19].

In the urinary bladder, COX-2 plays an important role in the development of transitional cell hyperplasia and carcinoma [20]; COX-2 expression in the urothelium is associated with high tumor grade and stage, and is an independent predictor of disease progression and survival [21,22]. Although treating these patients with endovesical MMC causes a reduction in the recurrence and progression rates of NMIBC, disease relapse is still high [2]. Since COX-2 inhibitors, such as celecoxib, may sensitize cells to antineoplastic agents we sought to investigate if there is an interaction between celecoxib, MMC and MDR transporters in a human bladder cancer cell line.

Methods

Cell lines and cell culture

UMUC-3, 5637, TCC sup cells and J82 (ATCC, Manassas, VA, USA), derived from human transitional cell carcinoma of the bladder, were routinely cultured in MEM supplemented with 10% fetal bovine serum, 2 mM glutamine, 100 U/mL penicillin, 100 µg/mL streptomycin, 1X NEAA and 1 mM sodium pyruvate. All components were purchased from Invitrogen Corporation (Cergy Pontoise, France), in a humidified incubator at 37°C with a 5% CO₂ atmosphere. Caco-2 (IRCCS "S. De Bellis", Castellana Grotte, Italy), MDCK-MDR1, MDCK-MRP1 (gift of Prof. P. Borst, NKI-AVL Institute, Amsterdam, Netherlands) and MDCK-BCRP cells (gift of Dr. A. Schinkel, NKI-AVL Institute, Amsterdam, Netherlands) were grown in DMEM medium with 10% heat-inactivated fetal calf serum, 100 U/mL penicillin, 100 µg/mL streptomycin and 2 mM L-glutamine (Invitrogen) in a humidified incubator at 37°C with a 5% CO₂ atmosphere. The cells were trypsinized twice a week with trypsin/ethylenediaminetetraacetic acid (EDTA) (0.05%/0.02%) and the medium was changed twice a week.

Cell lines transfection

The pSG5-COX-2 plasmid, which contains a full-length human COX-2 cDNA in the pSG expression vector [23], was used for COX-2 transfection. The plasmid DNA was introduced into UMUC-3 cells using Lipofectamine LTX (Invitrogen). Briefly, 2.5 µg of plasmid DNA and 6.25 µl of Lipofectamine LTX were combined in 500 µl of OpTI-MEM*1 reduced serum medium (Invitrogen) and allowed to stand at room temperature for 30 minutes. UMUC-3 cells were plated at 1X10⁵ cells/well in six-well plates. After overnight attachment, the Lipofectamine mixture was applied to cells. After 5 hours of transfection, the medium was removed and complete medium was added. After 48 hours, geneticin G418 (Invitrogen) was added to cells at a concentration of 800 mg/ml. Cell lines were obtained from individual colonies using cloning cylinders. UMUC-3-CX overexpressing cells were obtained and were continuously cultured in complete medium with the addition of 800 µg/ml G418.

Cell lines siRNA transfection

The COX-2 si RNA, target-specific 19–25 nt siRNA designed to knock down gene expression, was used for the inhibition of COX-2 expression in 5637 bladder cells. The COX-2 siRNA (Santa Cruz Biotech., CA, USA) was introduced into 5637 cells using siRNA Transfection Reagent (Santa Cruz Biotech.) Briefly, 5 µl of siRNA and 5 µl of siRNA Transfection Reagent were combined in 1 ml of siRNA Transfection Medium (Santa Cruz) and allowed to stand at room temperature for 30 minutes. 5637 cells were plated at 2X10⁵ cells/well in six-well

plates. After overnight attachment, the siRNA Transfection Reagent mixture was applied to cells. After 5 hours of transfection, the medium was removed and complete medium was added.

Prostaglandin E₂ (PGE₂) quantification

PGE₂ concentration was determined in the culture medium of UMUC-3, UMUC-3-CX, 5637, 5637si-CX, TCC sup and J82, cells. The cells were seeded into 24-well plates in the absence and presence of celecoxib (1- 5–50 μM) into a final volume of 500 μl/well of standard growth medium and incubated at 37°C for 24–48 h. After incubation time, the culture media were removed to determine PGE₂ levels with the PGE₂ monoclonal enzyme immunoassay kit (Cayman Chemicals) according to the manufacturer's protocol and results are expressed in pg of PGE₂/ml of medium.

Western immunoblotting

UMUC-3, UMUC-3-CX, 5637, 5637si-CX, TCC sup, and J82 cells were washed with 10 mL phosphate-buffered saline (PBS), scraped in 1 mL PBS and centrifuged for 1 min at 11,000g. Proteins were extracted from cells by homogenization in radioimmunoprecipitation (RIPA) buffer [0.5 M NaCl, 1% Triton X100, 0.5% NP40, 1% deoxycolic acid, 3.5 mM sodium dodecyl sulfate (SDS), 8.3 mM Tris HCl pH 7.4, 1.6 mM Tris base] and treated with 20% protease inhibitor cocktail (Sigma Aldrich, St. Louis, MO, USA). They were sonicated and centrifuged at 14,000 g for 15 min at 4°C and the protein content in the supernatant was measured using the Bradford method. 25–50 μg of protein extract was separated electrophoretically on 8% SDS–polyacrylamide gel electrophoresis (SDS–PAGE) and proteins electroblotted onto a nitrocellulose membrane (Bio-Rad, Hercules, CA, USA). Membranes were stained with 0.5% ponceau in 1% acetic acid to confirm equal loading. After overnight incubation of the membranes in blocking buffer (5% non-fat dried milk, 0.1% Tween 20 in Tris-buffered salt solution, TBS) they were incubated overnight with the respective primary antibody directed against COX-2 (1:500 mouse monoclonal), P-gp (1:1000 mouse monoclonal), MRP1 (1:1000 mouse monoclonal), BCRP (1:1000 mouse monoclonal) diluted in blocking buffer. The COX-2 antibody was purchased from Cayman Chemical (Ann Arbor, MI, USA), all other antibodies from Sigma-Aldrich. Membranes were washed four times with 0.1% Tween 20 in TBS and then incubated with a peroxidase-conjugated secondary antibody (Amersham, Buckinghamshire, UK) for 1 hr. After extensive rinsing in 0.1% Tween 20 in TBS, protein–antibody complexes conjugated with peroxidase were treated with enhanced chemoluminescence (ECL-Plus, Amersham) according to the manufacturer's protocol and exposed to a chemoluminescence film. The expression level

was evaluated by densitometric analysis using Quantity One software (Bio-Rad) and β-actin expression level was used to normalize the sample values.

Flow cytometry

UMUC-3 and UMUC-3-CX cells were harvested, washed twice in ice-cold PBS (pH 7.4), fixed in 4.5 mL of 70% ethanol, and stored at –20°C. Fixed cells were washed in ice-cold PBS once and incubated in 0.5 mL of 0.1% Tween 20 in PBS for 15 minutes at 25°C. To analyze P-gp/MRP1/BCRP expression, UMUC-3 and UMUC-3-CX cells were incubated O/N at 4°C with monoclonal anti-BCRP in 0.5% Tween 20 and 1% FBS in PBS. To determine the non-specific fluorescence due to the fluorescein conjugated secondary antibody, UMUC-3 and UMUC-3-CX cells were incubated with an appropriate isotype control (50 Ag/106 cells) in the same experimental conditions (isotype control). After 15 minute incubation with 0.5 mL of 0.5% FBS in PBS, cells were centrifuged and washed once in 0.5 mL of 0.5% FBS in PBS. The pellet was resuspended in 0.5% FBS in PBS in the presence of the goat anti-mouse IgG fluorescein-conjugated affinity-purified secondary antibody (Millipore, Billerica, MA, USA; 1:50) and incubated for 1 hour at 4°C. After a wash step with 0.5 mL of 0.5% FBS in PBS, cells were centrifuged and incubated in 5 Ag/mL propidium iodide overnight at 4°C. BCRP protein determination was done using a FACScan flow cytometer (Becton Dickinson, Franklin Lakes, NJ, USA). Fluorescence analysis was gated to include single cells based on forward and side light scatter and was based on the acquisition of data from 10,000 cells. Log fluorescence was collected and displayed as single variable histograms. The data analysis was carried out with the CellQuest software (Becton Dickinson).

Cell anti-proliferative effect

The cells were seeded into 96-well plates in the absence and presence of known concentrations (0,1-0,5-1-5-10-30-50 μM) of celecoxib (Pfizer, NY, USA) or MMC (Kyowa, Dusseldorf, DE), alone and in co-administration, added to a final volume of 200 μl/well of standard growth medium and incubated at 37°C for 48 h. Afterwards, 10 μl of 3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyltetrazoliumbromide (MTT) freshly prepared solution (5 mg/ml in PBS) (Sigma) was added in each well, and the plate was incubated in a humidified atmosphere 5% CO₂ at 37° for 3–4 h. MTT solution was removed and 200 μL of EtOH/DMSO (1:1) was added to each well to dissolve the blue formazan solid crystals. The optical density was measured at 570 nm and 650 nm wavelengths using Victor3 (PerkinElmer, Waltham, MA, USA). The results are expressed as EC₅₀ values, obtained from non-linear iterative curve fitting by Prism v.3.0, GraphPad software (GraphPad Software, Inc. San Diego, USA).

Permeability experiments

In the following three experiments a Caco-2 cell monolayer was used to evaluate P-gp transporter's interaction with a given compound. In the "drug transport experiment" the ability of a compound to permeate a cellular monolayer is tested. Both apical (AP)→basolateral (BL) flux and BL→A flux are evaluated since their ratio determines the permeability value: compounds with a permeability ratio > 2 do not permeate, compared to compounds with a ratio < 2. In the "[3H]Substrate transport inhibition" assay, a given compound competes for P-gp with its known substrate, [3H]vinblastine. If our compound (i.e. celecoxib) binds P-gp, [3H]vinblastine will be displaced reducing residual radioactivity. Finally, since P-gp is an ATPase pump, the "Cell ATP availability assay" will test if cells consume ATP in the presence of a given compound. P-gp binding compounds are transported out of the cell and produce ATP consumption. Taken together, these assays allow us to define if a compound is either a transporter's substrate, inhibitor, or modulator, according to Polli's classification [24].

Preparation of caco-2 monolayer

Caco-2 cells were seeded onto a Millicell® assay system (Millipore), where a cell monolayer is set in between a filter cell and a receiver plate, at a density of 10,000 cells/well. The culture medium was replaced every 48 h and the cells kept for 21 days in culture. The Trans Epithelial Electrical Resistance (TEER) of the monolayers was measured daily, before and after the experiment, using an epithelial volttohometer (Millicell® -ERS). Generally, TEER values greater than 1000 Ω for a 21day culture, are considered optimal.

Drug transport experiment

After 21 days of Caco-2 cell growth, the medium was removed from filter wells and from the receiver plate, which were filled with fresh HBSS buffer (Invitrogen). This procedure was repeated twice, and the plates were incubated at 37°C for 30 min. After incubation time, the HBSS buffer was removed and drug solutions of celecoxib, MMC and reference compounds, were added to the filter well at various concentrations (1–100 μM), while fresh HBSS was added to the receiver plate. The plates were incubated at 37°C for 120 min. Afterwards, samples were removed from the apical (filter well) and basolateral (receiver plate) side of the monolayer to measure the permeability [24]. The apparent permeability (Papp), in units of nm/second, was calculated using the following equation:

$$P = \left[\frac{V_A}{\text{Area} \times \text{time}} \right] \times \left[\frac{[\text{drug}]_{\text{acceptor}}}{[\text{drug}]_{\text{initial}}} \right]$$

VA = the volume (in mL) in the acceptor well;

Area = the surface area of the membrane (0.11 cm² of the well);

time = the total transport time in seconds (7200 sec);

[drug]_{acceptor} = the concentration of the drug measured by ESI-MS analyses or U.V. spectroscopy;

[drug]_{initial} = the initial drug concentration (1 × 10⁻⁴ M) in the apical or basolateral wells.

[3H]Substrate transport inhibition

20 nM of [3H]vinblastine were added in each well to the BL compartment, in the absence and in the presence of P-gp inhibitors (from 200 nM to 400 μM). After 120 min at 37°C, [3H]vinblastine's appearance was monitored at the AP compartment. At 120 min a 20 μL sample was taken from the donor compartment to determine the concentration of the residual radioligand at the end of the experiment. Samples were analyzed using LS6500 Beckman-Counter. For each compound, [3H]vinblastine transport inhibition was calculated as radioactivity difference between radioligand in the presence and absence of the given compound. These differences were expressed as inhibition rate at each drug concentration. Finally, the half maximal effective concentration (EC₅₀) values were determined [25].

Cell ATP availability assay

Caco-2 cells were seeded into 96-well microplate in 100 μL of complete medium at a density 2×10⁴ cells/well. The plate was incubated O/N in a humidified atmosphere (5% CO₂ at 37°C). The medium was removed and 100 μL of complete medium were added, in the presence or absence of different concentrations of test compounds. The plate was again incubated for 2 h in a humidified atmosphere. Then, 50 μL of mammalian cell lysis solution were added to all wells and the plate stirred for 5 min in an orbital shaker. In all wells, 50 μL of substrate solution were added, and the plate stirred for 5 min as above reported. The plate was dark adapted for 10 min and the luminescence measured in Victor3 (PerkinElmer) [26].

Calcein-AM experiment

The ability of celecoxib to interact with each transporter and restore calcein-acetoxymethylester (calcein-AM) was studied in a cell system represented by MDCK cells monolayers, specifically prepared to overexpress P-gp, MRP1 or BCRP (MDCK-P-gp, MDCK-MRP1, and MDCK-BCRP). The method was adapted from Eneroth et al. [27] and Korjamo et al. [28] with minor modifications. Briefly, each cell line was seeded into a black Cultureplate (PerkinElmer) 96/wells plate with 100 μL medium and allowed to become confluent overnight. Test compounds were solubilized in 100 μL of culture medium and added. After incubation at 37°C for 30 min. Calcein-AM, a fluorescent dye, was added in 100 μL of PBS to yield a final concentration of 2.5 μM.

After 30 min, each well was washed with PBS and the plate was read using Victor3 at excitation and emission wavelengths of 485 nm and 535 nm, respectively. After definition of the fluorescence basal level in untreated cells, calcein-AM cell accumulation, was measured in the presence of the tested compound (celecoxib). In treated wells the increase of fluorescence as compared to basal level was measured. The half maximal effective concentration (EC50) values were determined by fitting the fluorescence increase percentage versus log[dose] [29].

Intracellular mitomycin C accumulation

The time course of MMC intracellular accumulation and its modulation by celecoxib were evaluated by flow cytometry. For the purpose of the experiment, MMC and celecoxib were added to UMUC-3 and UMUC-3-CX cells at EC50 concentrations of the corresponding cell lines (Table 1). After incubation, the cell medium was removed and trypsin/EDTA was used to detach the cells from the plates. Cells were harvested, washed twice in ice-cold PBS (pH 7.4), and placed on ice (<1 hour) until analysis. Analysis was performed using a 530/30 filter (FL1-H [height of fluorescence intensity]). Fluorescence measurements of individual cells were done with a Becton Dickinson FACScan equipped with an UV argon laser. Analysis was gated to include single cells, based on forward and side light scatter, and was based on the acquisition of data from 10,000 cells [30].

Statistical methodology

All the reported values are expressed as means \pm standard deviation (SD) from triplicate experiments. EC50 values were obtained from non-linear interactive curve fitting by GraphPad, Prism. For variance estimation, *t*-test and Wilcoxon test were used. Statistical differences in Table 1 and in Table 2 were determined by Mann–Whitney unpaired test. Differences were considered statistically significant when P values were <0.05. All the research carried out and reported in the present manuscript was reviewed and approved by the ethical committee of the Ospedale Policlinico Consorziale of Bari, Italy.

Results

Expression of COX-2 and ABC transporters in bladder cancer cells

COX-2 and ABC transporters expression was studied via western blot in several bladder cancer cell lines (Figure 1). As previously reported [15], UMUC-3 lacks COX-2 expression and was used to create UMUC-3-CX, a COX-2 overexpressing transfectant. Moreover, 5637 showed the highest COX-2 levels among the tested bladder cancer cells and was used to obtain a silenced non-expressing 5637si-CX cell line. Successful UMUC-3 transfection and 5637 silencing are shown in Figure 1. Among the ABC transporters, P-gp expression was similar in wild type and in transfected UMUC-3, but undetectable in the other cell lines. MRP1 protein was neglectable in all cell lines. Finally, BCRP levels were increased in UMUC-3-CX as compared to UMUC-3, while remained undetectable in all other cell lines. Additionally, COX-2 functional activity was tested in all cells by measuring PGE2 production. As expected, celecoxib administration caused a dose dependent reduction of endogenous PGE2 secretion in all, except COX-2 non-expressing cells (UMUC-3 and 5637si-CX) (Figure 2). Short-term (24 and 48 h) exposure of our cell lines to celecoxib (0.1 μ M to 50 μ M) caused no significant change in transporter's expression as studied by western blot (data not shown).

Mitomycin C anti-proliferative effect and intracellular accumulation in UMUC-3 and UMUC-3-CX cells

The anti-proliferative effect of celecoxib and MMC were investigated alone and in co-administration at 48 h, in 5637 and 5637si-CX cells (Figure 3), and in UMUC-3 and UMUC-3-CX cells (Figure 4). Compounds were tested at escalating doses, starting from 0.1 μ M to 50 μ M. Results are expressed as EC50 and are shown in Table 1. Dose response curves are shown in the corresponding figures. When administered alone, MMC had a comparable anti-proliferative effect both in 5637 and 5637si-CX cells; in these cells celecoxib was unable to affect MMC killing (Figure 3). However, in the same experimental conditions, the effect of MMC was eight fold greater in

Table 1 Antiproliferative effect of CLX and MMC, alone and in co-administration at 48 h, in bladder cancer cell lines

	UMUC3	UMUC-3- CX	5637	5637 si-CX	TCC sup	J82
	EC ₅₀ (μ M)					
CLX	14.2 \pm 0.7	16.3 \pm 0.8	> 50	> 50	15.7 \pm 0.50	11.6 \pm 0.31
MMC	1.69 \pm 0.09	13.0 \pm 1.0	0.93 \pm 0.04	0.62 \pm 0.05	0.83 \pm 0.07	0.64 \pm 0.09
CLX 1 μ M + MMC	1.54 \pm 0.1	6 \pm 0.3	1.46 \pm 0.10	0.59 \pm 0.03	0.74 \pm 0.08	0.48 \pm 0.03
CLX 5 μ M + MMC	1.75 \pm 0.07	5.7 \pm 0.2	1.31 \pm 0.06	0.47 \pm 0.01	0.68 \pm 0.04	1.04 \pm 0.10
CLX 10 μ M + MMC	1.9 \pm 0.1	4 \pm 0.18	1.68 \pm 0.11	1.01 \pm 0.20	1.08 \pm 0.20	1.09 \pm 0.10

EC₅₀ = the concentration of a compound where 50% of its maximal effect is observed.
 The EC₅₀ values were obtained from non-linear iterative curve fitting by Prism v.3.0, GraphPad software.
 Compounds were tested at escalating doses, starting from 0.1 μ M to 50 μ M. Results are expressed as EC₅₀.

Table 2 Biological evaluation of MDR inhibitors and reference compounds

Compounds	EC ₅₀ ± SEM, ^a (μM)			
	Caco-2 [³ H] vinblastine transport inhibition	MDCK-MDR1 Calcein-AM	MDCK-MRP1 Calcein-AM	MDCK-BCRP Calcein-AM
Celecoxib	30 ± 2.0	46.9 ± 2.5	10.1 ± 0.5	24.1 ± 1.2
Mitomycin C	10 ± 0.5	N.D. ^b	N.D. ^b	N.D. ^b
MK-571 ^c			2.85 ± 0.25	
Verapamil ^c	20 ^d			3.65 ± 0.2

^aData ± SEM are the mean of three independent determinations (samples in triplicate).

^bMitomycin C activity was not determined because in the experimental conditions quenched the spectroscopic properties of the probe.

^cReference compounds for MDR pumps.

^dSee Colabufo et al. [40].

UMUC-3 as compared to UMUC-3-CX (Figure 4). Importantly, the co-administration of 1 μM, 5 μM, and 10 μM celecoxib to MMC did not cause a significant increase of MMC cytotoxicity in UMUC-3 cells (Figure 4C). By contrast, in UMUC-3-CX cells, the anti-proliferative activity of MMC was 2–3 fold improved by the co-administration of celecoxib compared to MMC alone (Table 1, and Figure 4D).

Since UMUC-3 transfection caused an overexpression of BCRP in UMUC-3-CX cells, we sought to explore the ability of celecoxib to interfere with the efflux of MMC. For this purpose, flow cytometry was performed to study MMC intracellular accumulation, both in UMUC-3 and UMUC-3-CX cells. The cells were either treated one day with MMC or celecoxib alone, or with both in co-administration using the following sequential schedule: 1 day celecoxib, and afterwards, 1 day MMC and celecoxib added to the same plate (Figure 5). Importantly, celecoxib was able to produce an increase of intracellular MMC concentration only in UMUC-3-CX cells, as shown by the right shift of the d curve as compared to the c curve (Figure 5B). This effect was not seen in UMUC-3 cells (Figure 5A).

Celecoxib-transporter interacting mechanism

Three biological assays (Drug Transport Experiment, Substrate Transporter Inhibition, and ATP cell depletion)

were combined to establish if there was an interaction between P-gp and celecoxib. In the drug transport experiment, a Caco-2 cell monolayer was used to determine the apparent permeability (Papp) of celecoxib, both as basolateral-apical flux (B→A) and apical-basolateral flux (A→B). Celecoxib displayed Papp A→B = 137 nm/sec, Papp B→A = 1027 nm/sec, and a 7.5 BA/AB ratio. This result suggests that celecoxib is effluxed by P-gp. Indeed, compounds displaying BA/AB ratio > 2 are linked and flipped to the extra cellular compartment by pumps, while drugs with BA/AB ratio < 2 are not transported. Moreover, the Substrate Transporter Inhibition experiment was performed to study the modulating effect of celecoxib on [3H]vinblastine transport. Compared to Verapamil, which is the reference compound for this experiment, celecoxib had a similar EC50 value (EC50 = 30 ± 2.0 μM) (Table 2). Finally, when tested in Caco-2 cells to monitor ATP cell depletion, celecoxib decreased ATP levels in a time- and dose-dependent fashion. As a result of all finding combined, celecoxib should be classified as P-gp unambiguous substrate [24].

Celecoxib in MDCK cells overexpressing P-gp or BCRP or MRP1 pumps

In MDCK-P-gp, in MDCK-MRP1 and in MDCK-BCRP cells the ability of celecoxib to restore calcein-AM cell accumulation was tested. This dye, calcein-acetoxymethylester,




Figure 1 Protein expression levels of COX-2, P-gp, MRP1 and BCRP in UMUC3, UMUC-3-CX, 5637, 5637si-CX, TCCsup, and J82, as determined by western blot. Cell lysates were obtained from exponentially growing cells and subjected to immunoblotting with appropriate antibodies. Immunoblotting with an antibody to β-actin was used to ensure equal loading of proteins in each lane (bottom).




Figure 2 Secretion of prostaglandin E2 in the culture medium of a selection of bladder cancer cell lines. PGE2 in cell supernatant was determined by enzyme immunoassay after treatment with or without known concentrations (0 - 1- 5-50 μM) of Celecoxib for 24 and 48 h.

is a non fluorescent lipophilic P-gp/MRP1/BCRP substrate that diffuses across the plasma membrane into the cell where is hydrolyzed into highly fluorescent calcein by endogenous cytoplasmic esterases. This fluorescent compound is not effluxed by P-gp, MRP1, BCRP and it cannot cross the cell membrane via passive diffusion because of its hydrophilie. Therefore, in the presence of an MDR transporter modulator, calcein retention and a rapid fluorescence increase is monitored (Figure 6). In Table 2, calcein cell accumulation is plotted vs Log[celecoxib] and the following results are displayed: in MDCK-P-gp cells $EC_{50} = 46.9 \pm 2.5 \mu M$, in MDCK-MRP1 $EC_{50} = 10.1 \pm 0.5 \mu M$, and in MDCK-BCRP $EC_{50} = 24.1 \pm 1.2 \mu M$. Compared to the reference compounds, these findings confirm that celecoxib is a substrate for P-gp, MRP1 and BCRP pumps.

Discussion

In recent years, there has been an interest to investigate the potential link between COX-2 expression and the development of MDR in multiple tumors, including urothelial cancer [21,22]. A direct causal link between the activation of the COX-2/PGE2 signal pathway and the up-regulation of all three ABC transporters has been documented [31-35] and supports the strong correlation between COX-2 and MDR1/P-gp expression seen in several tumor specimens [36-38]. Overexpression of P-gp has been found in human bladder cancer cells selected by drug resistance against P-gp-targeting drugs [39-41]. In patients with bladder cancers, expression of P-gp is often increased after chemotherapeutic treatment [42]. Prophylactic intravesical instillation of MMC, doxorubicin and epirubicin has been useful for reducing recurrence of NMIBC; however, recurrence rates are still high and





Figure 3 Antiproliferative effect of celecoxib and MMC administered alone at increasing doses (0,1 - 0,5 - 1-5 - 10-30 - 50 μ M) (A, B). Antiproliferative effect of MMC alone and in co-administration with known concentrations of celecoxib (C, D). All experiments conducted after 48 h of incubation in 5637 (A, C) and in 5637si-CX (B, D) cells.



progression to more invasive disease is not affected even after intensive intravesical chemotherapy [43].

In this study we developed an *in vitro* bladder cancer model to study if COX-2 inhibitors can modulate tumor resistance to MMC by interfering with the activity of membrane transporter proteins of the ABC family. For this purpose we used UMUC-3 cells, constitutively lacking COX-2 expression, and UMUC-3-CX cells, in which COX-2 is overexpressed. When MMC was administered alone, UMUC-3-CX cells resulted resistant to MMC killing. However, for the first time we showed that pre-treatment with a selective COX-2 inhibitor, celecoxib, caused a significant and dose dependent increase in the cytotoxic activity of MMC. Interestingly, in UMUC-3 cells MMC activity was not affected by celecoxib. Moreover, compared to UMUC-3, we found that forced COX-2 overexpression in UMUC-3-CX cells increased PGE2 production and up-regulated BCRP, one of the transporters involved in MDR. These data were confirmed by the observation of an increase in intracellular concentration of MMC when UMUC-3-CX cells were co-treated with celecoxib. Again, intracellular MMC concentration was not affected by celecoxib in UMUC-3 cells. Although several causes may be taken into account it has been

shown that ABC transporters, such as BCRP, induce drug resistance by promoting drug efflux out of the cells [44]. Indeed, when the cytotoxicity properties of MMC were studied in a cell line completely lacking any ABC transporter expression, such as 5637 and 5637si-CX cells, celecoxib administration was unable to affect MMC killing. Assuming a causal link between COX-2 expression and MDR, COX-2 inhibitors would be expected to prevent ABC transporters induction and sensitize cells to antineoplastic agents. This has been previously shown in Caco-2 cells where indomethacin, nimesulide and naproxen directly reduced MRP1 expression and P-gp relative amount and function [45]. Similarly, in human lung cancer cells celecoxib was shown to downregulate the expression of MRP1 [19].

Although COX-2 enzyme inhibition could not be excluded in our experimental model, we sought to investigate if the effect seen in UMUC-3-CX cells after celecoxib administration could be the result of a direct interaction between celecoxib and any of the three transporters involved in MDR. To explore this hypothesis specific biological assays were performed to demonstrate that celecoxib is a substrate for the MDR transporters explored in this study. Our data




Figure 5 Flow cytometry analysis to study the time course of MMC intracellular accumulation and its modulation by celecoxib in **UMUC-3 (A)** and **UMUC-3-CX (B)**. In both cases, cells were treated one day with MMC (1.69 μ M) or celecoxib (14.2 μ M) alone, or with both in co-administration (MMC 1.69 μ M + celecoxib 14.2 μ M). Celecoxib causes an increase in the intracellular concentration of MMC in UMUC-3-CX, but not in UMUC-3 cells.

demonstrate that celecoxib is effluxed by P-gp, BCRP and MRP1 pumps and causes a time- and dose-dependent ATP cell depletion in Caco-2 cells. Further, celecoxib competes with and may inhibit the transport of other reference drugs (vinblastine in our experimental model) (Table 2). Finally, the ability of celecoxib to restore Calcein-AM cell accumulation in MDCK-P-gp, MDCK-MRP1 and MDCK-BCRP cells suggests that celecoxib is a substrate for all the transporters tested in our study. Thus, the increase in MMC

concentration seen in UMUC-3-CX after co-administration with celecoxib may be the result of a transporter-celecoxib interaction. As a hypothesis, since BCRP was the only transporter overexpressed by UMUC-3-CX cells, a BCRP-celecoxib interaction may justify the greater anti-proliferative activity obtained when MMC and celecoxib were co-administered (Figures 3 and 4).

Our findings are in agreement with previous studies in which COX-2 inhibitors have shown to produce




Figure 6 Schematic view of the mechanism of calcein cellular retention in the absence (A) and in the presence (B) of an MDR pump substrate, such as celecoxib.

MDR-regulating effects that are COX-2 independent. Van Wijngaarden and co-workers proposed that the effects of celecoxib were most likely mediated by inhibition of NF- κ b and not related to COX-2 or the activity of the ABC transporters P-gp, MRP1 and ABCG2 [46]. Likewise, Ye et al. found that indomethacin and SC236, a COX-2-selective inhibitor, sensitized human hepatocellular carcinoma HepG2 cells to the cytotoxicity of doxorubicin by significantly increasing doxorubicin intracellular accumulation; however, the effects were not reversed by prostaglandin E₂, implicating a COX-independent mechanism [47]. Taken together, these findings suggest that COX-2-selective inhibitors may enhance the effect of certain anti-cancer agents and overcome drug resistance through mechanisms that by-pass the COX-2 cascade.

Many reports examining the clinical benefits of COX-2 inhibitors and nonsteroidal anti-inflammatory drugs (NSAIDs) have addressed the role of these compounds in bladder chemoprevention. *In vitro* and *in vivo* research suggests that NSAIDs hinder growth and survival of bladder cancer cells [48,49], however, epidemiologic studies investigating the association between non selective NSAIDs and bladder cancer have been conflicting, with two large cohort studies suggesting that non-aspirin NSAIDs, but not aspirin, may protect from bladder cancer [50,51]. Such conflicting data have discouraged the scientific community to promote prospective clinical testing of aspirin or acetic acids (indomethacin, sulindac) in patients with NMIBC. On the other hand, the body of evidence pointing to the role of COX-2 pathway in the development of bladder cancer is compelling and supported by several epidemiologic studies [51]. More recently, COX-2 selective inhibitors have been the focus of much scrutiny in terms of cardiovascular risk. Of such inhibitors, celecoxib is thought to have a better safety profile, and has been tested in patients with NMIBC, and

clinical trials are under way. Sabichi et al. recently reported the results of a phase IIb randomized controlled trial of celecoxib to prevent recurrence following TUR in 146 patients with NMIBC [52]. Although the primary endpoint (time to recurrence) did not reach statistical significance, the results support a beneficial effect of celecoxib in such patients. For the ultimate interpretation of these results, the ongoing phase III randomized controlled Bladder COX Inhibition Trial (BOXIT) of the same daily dose of celecoxib versus placebo is being conducted [www.cancerresearchuk.org].

We recognize this is a pilot study, in which we show that celecoxib is a substrate for ABC transporters and may enhance the activity of cytotoxic drugs such as MMC. However, translational studies and, more importantly, ongoing and future clinical trials will have to confirm the preclinical potential of celecoxib and other COX-2 inhibitors in NMIBC.

Conclusions

Compared to the *wt* counterpart, the UMUC-3-CX cell line resembles a more aggressive phenotype with a higher BCRP protein expression and a low response to MMC. Interestingly, the cytotoxic activity is regained when MMC is administered in combination with celecoxib. Although COX-2 enzyme inhibition cannot be excluded, this result may partly depend on a direct interaction between celecoxib and any of the transporters involved in MDR. Indeed, we have shown that celecoxib is a substrate for P-gp/BCRP/MRP1 transporters and may modulate their activity. These findings and others from recent reports imply that the therapeutic approach of combining conventional chemotherapy with selective COX-2 inhibitors seems promising and warrants prospective clinical evaluation in patients with NMIBC in which COX-2 is overexpressed.

Competing interests

The authors declare that they have no competing interests.

Authors' contribution

VP substantially contributed the study conception and design, to the analysis and interpretation of the data, to the manuscript drafting. PA carried out all cell line studies, including western blot, PGE2 assays and MTT tests. MN carried out all permeability experiments and contributed to data analysis and interpretation. NAC participated in the design of the study and performed the statistical analysis. MC and AA performed all flow cytometry studies. LC and AP participated in coordinating the study and helped to draft the manuscript. All authors read and approved the final manuscript.

Author details

¹Sezione di Urologia e Andrologia, Dipartimento dell'Emergenza e dei Trapianti di Organi (DETO), Università "Aldo Moro" di Bari, Piazza G. Cesare 11, Bari 70124, Italy. ²Dipartimento Farmacochimico, Università "Aldo Moro" di Bari, Via Orabona, 4, Bari 70125, Italy. ³Sezione di Urologia e Trapianti di Reni, Università di Foggia, Viale Pinto 1, Foggia 71100, Italy. ⁴Istituto Tumori "Giovanni Paolo II" IRCCS Ospedale Oncologico, Viale Orazio Flacco 65, Bari 70124, Italy.

Received: 26 October 2012 Accepted: 2 May 2013

Published: 24 May 2013

References

- Sylvester RJ, van der Meijden APM, Oosterlinck W, Witjes JA, Bouffoux C, Donald LD, Newling WW: **Predicting recurrence and progression in individual patients with stage Ta T1 bladder cancer using EORTC risk tables: a combined analysis of 2596 patients from seven EORTC trials.** *Eur Urol* 2006, **49**:466–477.
- Babjuk M, Oosterlinck W, Sylvester R, Kaasinen E, Böhle A, Palou-Redorta J, Roupřet M: **European Association of Urology (EAU), EAU guidelines on non-muscle-invasive urothelial carcinoma of the bladder, the 2011 update.** *Eur Urol* 2011, **59**:997–1008.
- Pawinski A, Sylvester R, Kurth KH, Bouffoux C, van der der Meijden A, Parmar MK, Bijmans L: **A combined analysis of european organization for research and treatment of cancer, and medical research council randomized clinical trials for the prophylactic treatment of TaT1 bladder cancer. European organization for research and treatment of cancer genitourinary tract cancer cooperative group and the medical research council working part on superficial bladder cancer.** *J Urol* 1996, **156**:1934–1941.
- Klan R, Loy V, Huland H: **Residual tumor discovered in routine second transurethral resection in patients with stage T1 transitional cell carcinoma of the bladder.** *J Urol* 1991, **146**:316–318.
- Divrik RT, Yildirim U, Zorlu F, Ozen H: **The effect of repeat transurethral resection on recurrence and progression rates in patients with T1 tumors of the bladder who received intravesical mitomycin: a prospective, randomized clinical trial.** *J Urol* 2006, **175**:1641–1644.
- Hayes MC, Birch BR, Cooper AJ, Primrose JN: **Cellular resistance to mitomycin C is associated with overexpression of MDR-1 in a urothelial cancer cell line (MGH-U1).** *Br J Urol* 2001, **87**:245–250.
- Birare N, Lwaleed BA, Cooper AJ: **Multidrug resistance in a urothelial cancer cell line after 1-hour mitomycin C exposure.** *J Urol* 2009, **182**:2472–2476.
- Leslie EM, Deeley RG, Cole SPC: **Multidrug resistance proteins: role of P-glycoprotein MRP1, MRP2, and BCRP (ABCG2) in tissue defense.** *Toxicol Appl Pharmacol* 2005, **204**:216–237.
- Borst P, Evers R, Kool M, Wijnholds J: **A family of drug transporters: the multidrug resistance-associated proteins.** *J Natl Cancer Inst* 2000, **92**:1295–1302.
- Tan B, Piwnicka-Worms D, Ratner L: **Multidrug resistance transporters and modulation.** *Curr Opin Oncol* 2000, **12**:450–458.
- Jedlitschky G, Leier I, Buchholz U, Barnouin K, Kurz G, Keppler D: **Transport of glutathione, glucuronate, and sulfate conjugates by the MRP gene-encoded conjugate export pump.** *Cancer Res* 1996, **56**:988–994.
- Nakagawa M, Emoto A, Nasu N, Hanada T, Kuwano M, Cole SP, Nomura Y: **Clinical significance of multi-drug resistance associated protein and P-glycoprotein in patients with bladder cancer.** *J Urol* 1996, **157**:1260–1265.
- Diestra JE, Condom E, Del Muro XG, Scheffer GL, Pérez J, Zurita AJ, Muñoz-Seguí J, Vigués F, Scheper RJ, Capellá G, *et al*: **Expression of multidrug resistance proteins P-glycoprotein, multidrug resistance protein 1, breast cancer resistance protein and lung resistance related protein in locally advanced bladder cancer treated with neoadjuvant chemotherapy: biological and clinical implications.** *J Urol* 2003, **170**:1383–1387.
- Smithand WL, Dewitt DL: **Prostaglandin endoperoxide H synthases-1 and -2.** *Adv Immunol* 1996, **62**:167–215.
- Mohammed SI, Dhawan D, Abraham S, Snyder PW, Waters DJ, Craig BA, Lu M, Wu L, Zheng R, Stewart J, *et al*: **Cyclooxygenase inhibitors in urinary bladder cancer: in vitro and in vivo effects.** *Mol Cancer Ther* 2006, **5**:329–336.
- Altorki NK, Keresztes RS, Port JL, Libby DM, Korst RJ, Flieder DB, Ferrara CA, Yankelevitz DF, Subbaramaiah K, Pasmantier MW, *et al*: **Celecoxib, a selective cyclooxygenase-2 inhibitor, enhances the response to preoperative paclitaxel and carboplatin in early-stage non-small-cell lung cancer.** *J Clin Oncol* 2003, **21**:2645–2650.
- Milella M, Gelibter A, Di Cosimo S, Bria E, Ruggeri EM, Carlini P, Malaguti P, Pellicciotta M, Terzoli E, Cognetti F: **Pilot study of celecoxib and infusional 5-fluorouracil as second-line treatment for advanced pancreatic carcinoma.** *Cancer* 2004, **101**:133–138.
- Liu B, Qu L, Tao H: **Cyclo-oxygenase 2 up-regulates the effect of multidrug resistance.** *Cell Biol Int* 2009, **34**:21–25.
- Kang HK, Lee E, Pyo H, Lim SJ: **Cyclooxygenase-independent down-regulation of multidrug resistance-associated protein-1 expression by celecoxib in human lung cancer cells.** *Mol Cancer Ther* 2005, **4**:1358–1363.
- Klein RD, Van Pelt CS, Sabichi AL, Dela Cerda J, Fischer SM, Fürstenberger G, Müller-Decker K: **Transitional cell hyperplasia and carcinomas in urinary bladders of transgenic mice with keratin 5 promoter-driven cyclooxygenase-2 overexpression.** *Cancer Res* 2005, **65**:1808–1813.
- Shariat SF, Kim JH, Ayala GE, Kho K, Wheeler TM, Lerner SP: **Cyclooxygenase-2 is highly expressed in carcinoma in situ and T1 transitional cell carcinoma of the bladder.** *J Urol* 2003, **169**:938–942.
- Miyata Y, Kanda S, Nomata K, Eguchi J, Kanetake H: **Expression of cyclooxygenase-2 and EP4 receptor in transitional cell carcinoma of the upper urinary tract.** *J Urol* 2005, **173**:56–60.
- Sun Y, Tang XM, Half E, Kuo MT, Sinicropo FA: **Cyclooxygenase-2 overexpression reduces apoptotic susceptibility by inhibiting the cytochrome c-dependent apoptotic pathway in human colon cancer cells.** *Cancer Res* 2002, **62**:6323–6328.
- Polli JW, Wring SA, Humphreys JE, Huang L, Morgan JB, Webster LO, Serabjit-Singh CS: **Rational use of in vitro P-glycoprotein assays in drug discovery.** *J Pharmacol Exp Ther* 2001, **299**:620–628.
- Taub ME, Podila L, Almeida I: **Functional assessment of multiple P-glycoprotein (P-gp) probe substrates: influence of cell line and modulator concentration on P-gp activity.** *Drug Metab Dispos* 2005, **33**:1679–1687.
- Kangas L, Grönroos M, Nieminen AL: **Bioluminescence of cellular ATP: a new method for evaluating cytotoxic agents in vitro.** *Med Biol* 1984, **62**:338–343.
- Korjamo T, Honkakoski P, Toppinen MR, Niva S, Reinisalo M, Palmgrén JJ, Mönkkönen J: **Absorption properties and P-glycoprotein activity of modified caco-2 cell lines.** *Eur J Pharm Sci* 2005, **26**:266–279.
- Eneroth A, Aström E, Hoogstraate J, Schrenk D, Conrad S, Kauffmann HM, Gjellan K: **Evaluation of a vincristine resistant caco-2 cell line for use in a calcein AM extrusion screening assay for P-glycoprotein interaction.** *Eur J Pharm Sci* 2001, **12**:205–214.
- Feng B, Mills JB, Davidson RE, Mireles RJ, Janiszewski JS, Troutman MD, de Moraes SM: **In vitro P-glycoprotein assays to predict the in vivo interactions of P-glycoprotein with drugs in the central nervous system.** *Drug Metab Dispos* 2008, **36**:268–275.
- Azzariti A, Colabufo NA, Berardi F, Snyder PW, Waters DJ, Craig BA, Lu M, Wu L, Zheng R, Stewart J, *et al*: **Cyclohexylpiperazine derivative PB28, a sigma-2 agonist and sigma-1 antagonist receptor, inhibits cell growth, modulates P-glycoprotein and synergizes with anthracyclines in breast cancer.** *Mol Cancer Ther* 2006, **5**:1807–1816.
- Kalalinia F, Elahian F, Behravan J: **Potential role of cyclooxygenase-2 on the regulation of the drug efflux transporter ABCG2 in breast cancer cell lines.** *J Cancer Res Clin Oncol* 2011, **137**:321–330.
- Fantappiè O, Masini E, Sardi I, Raimondi L, Bani D, Solazzo M, Vannacci A, Mazzanti R: **The MDR phenotype is associated with the expression of COX-2 and iNOS in a human hepatocellular carcinoma cell line.** *Hepatology* 2002, **35**:843–852.
- Patel VA, Dunn MJ, Sorokin A: **Regulation of MDR-1 (P-glycoprotein) by cyclooxygenase-2.** *J Biol Chem* 2002, **277**:38915–38920.
- Sorokin A: **Cyclooxygenase-2: potential role in regulation of drug efflux and multidrug resistance phenotype.** *Curr Pharm Des* 2004, **10**:647–657.

35. Raspollini MR, Amunni G, Villanuoci A, Boddi V, Taddei GL: **Increased cyclooxygenase-2 (COX-2) and P-glycoprotein-170 (MDR1) expression is associated with chemotherapy resistance and poor prognosis. Analysis in ovarian carcinoma patients with low and high survival.** *Int J Gynecol Cancer* 2005, **15**:255–260.
36. Ratnasinghe D, Daschner PJ, Anver MR, Kasprzak BH, Taylor PR, Yeh GC, Tangrea JA: **Cyclooxygenase-2, P-glycoprotein-170 and drug resistance; is chemoprevention against multidrug resistance possible?** *Anticancer Res* 2001, **21**:2141–2147.
37. Half E, Tang XM, Gwyn K, Sahin A, Wathen K, Sinicrope FA: **Cyclooxygenase-2 expression in human breast cancers and adjacent ductal carcinoma in situ.** *Cancer Res* 2002, **62**:1676–1681.
38. Liu B, Qu L, Tao H: **Cyclo-oxygenase 2 up-regulates the effect of multidrug resistance.** *Cell Biol Int* 2009, **34**:21–25.
39. Shinohara N, Liebert M, Wedemeyer G, Chang JH, Grossman HB: **Evaluation of multiple drug resistance in human bladder cancer cell lines.** *J Urol* 1993, **150**:505–509.
40. Hasegawa S, Abe T, Naito S, Kotoh S, Kumazawa J, Hipfner DR, Deeley RG, Cole SP, Kuwano M: **Expression of multidrug resistance-associated protein (MRP), MDR1 and DNA topoisomerase II in human multidrug-resistant bladder cancer cell lines.** *Br J Cancer* 1995, **71**:907–913.
41. Kimiya K, Naito S, Soejima T, Sakamoto N, Kotoh S, Kumazawa J, Tsuruo T: **Establishment and characterization of doxorubicin-resistant human bladder cancer cell line, KK47/ADM.** *J Urol* 1992, **148**:441–445.
42. Nakagawa M, Emoto A, Nasu N, Hanada T, Kuwano M, Cole SP, Nomura Y: **Clinical significance of multi-drug resistance associated protein and P-glycoprotein in patients with bladder cancer.** *J Urol* 1997, **157**:1260–1265.
43. Malmström PU, Sylvester RJ, Crawford DE, Friedrich M, Krega S, Rintala E, Solsona E, Di Stasi SM, Witjes JA: **An individual patient data meta-analysis of the long-term outcome of randomised studies comparing intravesical mitomycin C versus bacillus calmette-guérin for non-muscle-invasive bladder cancer.** *Eur Urol* 2009, **56**:247–256.
44. Erlichman C, Boerner SA, Hallgren CG, Spieker R, Wang XY, James CD, Scheffer GL, Maliepaard M, Ross DD, Bible KC, Kaufmann SH: **The HER tyrosine kinase inhibitor CI1033 enhances cytotoxicity of 7-ethyl-10-hydroxycamptothecin and topotecan by inhibiting breast cancer resistance protein-mediated drug efflux.** *Cancer Res* 2001, **61**:739–748.
45. Zrieki A, Farinotti R, Buyse M: **Cyclooxygenase inhibitors down regulate P-glycoprotein in human colorectal caco-2 cell line.** *Pharm Res* 2008, **25**:1991–2001.
46. van Wijngaarden J, van Beek E, van Rossum AC, van der Bent C, *et al*: **Celecoxib, enhances doxorubicin-induced cytotoxicity in MDA-MB231 cells by NF- κ B-mediated increase of intracellular doxorubicin accumulation.** *Eur J Cancer* 2007, **43**:433–442.
47. Ye CG, Wu WK, Yeung JH, Li HT, Li ZJ, Wong CC, Ren SX, Zhang L, Fung KP, Cho CH: **Indomethacin and SC236 enhance the cytotoxicity of doxorubicin in human hepatocellular carcinoma cells via inhibiting P-glycoprotein and MRP1 expression.** *Cancer Lett* 2011, **304**:90–96.
48. Grubbs CJ, Juliana MM, Eto I, *et al*: **Chemoprevention by indomethacin of N-butyl-N-(4-hydroxybutyl)-nitrosamine-induced urinary bladder tumors.** *Anticancer Res* 1993, **13**:33–36.
49. Rao KV, Detrisac CJ, Steele VE, *et al*: **Differential activity of aspirin, ketoprofen and sulindac as cancer chemopreventive agents in the mouse urinary bladder.** *Carcinogenesis* 1996, **17**:1435–1438.
50. Daugherty SE, Pfeiffer RM, Sigurdson AJ, Hayes RB, Leitzmann M, Schatzkin A, Hollenbeck AR, Silverman DT: **Nonsteroidal antiinflammatory drugs and bladder cancer: a pooled analysis.** *Am J Epidemiol* 2011, **173**:721–730.
51. Castela JE, Yuan JM, Gago-Dominguez M, Yu MC, Ross RK: **Non-steroidal anti-inflammatory drugs and bladder cancer prevention.** *Br J Cancer* 2000, **82**:1364–1369.
52. Sabichi AL, Lee JJ, Grossman HB, Liu S, Richmond E, Czerniak BA, De la Cerda J, Eagle C, Viner JL, Palmer JL, Lerner SP: **A randomized controlled trial of celecoxib to prevent recurrence of nonmuscle-invasive bladder cancer.** *Cancer Prev Res* 2011, **4**:1580–1589.

doi:10.1186/1476-4598-12-47

Cite this article as: Pagliarulo *et al*: The interaction of celecoxib with MDR transporters enhances the activity of mitomycin C in a bladder cancer cell line. *Molecular Cancer* 2013 **12**:47.

Submit your next manuscript to BioMed Central and take full advantage of:

- **Convenient online submission**
- **Thorough peer review**
- **No space constraints or color figure charges**
- **Immediate publication on acceptance**
- **Inclusion in PubMed, CAS, Scopus and Google Scholar**
- **Research which is freely available for redistribution**

Submit your manuscript at
www.biomedcentral.com/submit

