




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Contents lists available at ScienceDirect

# Comparative Immunology, Microbiology and Infectious Diseases

journal homepage: [www.elsevier.com/locate/cimid](http://www.elsevier.com/locate/cimid)

## Detection and molecular characterization of feline hemoplasmas in wild felid species in Iran in the Middle East



Fereshteh Ghazisaeeedi<sup>a,\*</sup>, Nahid Atyabi<sup>a</sup>, Taghi Zahraei Salehi<sup>b</sup>, Saeid Tabatabaei<sup>b</sup>, Iraj Ashrafi Tamai<sup>b</sup>, Iman Memarian<sup>c</sup>, Séverine Tasker<sup>d</sup>

<sup>a</sup> Department of Veterinary Clinical Pathology, Faculty of Veterinary Medicine, University of Tehran, Qareeb Street, Azadi Av. P.O. Box: 14155-6453, Tehran, Iran

<sup>b</sup> Department of Microbiology, Faculty of Veterinary Medicine, University of Tehran, Qareeb Street, Azadi Av. P.O. Box: 14155-6453, Tehran, Iran

<sup>c</sup> Chief Veterinarian at Tehran Zoo and Pardisan Rehabilitation Center Tehran Zoo, 4th Kilometer of Tehran Karaj Freeway, 1484613111, Tehran, Iran

<sup>d</sup> The Feline Centre, Langford Veterinary Services and School of Veterinary Sciences, University of Bristol, Langford, BS40 5DU, Bristol, UK

### ARTICLE INFO

#### Keywords:

Feline hemoplasma  
*Panthera pardus saxicolor*  
Persian leopard

### ABSTRACT

Three feline hemoplasma species exist in felids: *Mycoplasma haemofelis*, ‘*Candidatus Mycoplasma haemominutum*’, and ‘*Candidatus Mycoplasma turicensis*’.

The aims of the study were to determine the presence of, and molecularly characterize, any hemoplasmas in wild felids, including the endangered Persian leopard in Iran, the Middle East.

Blood samples were collected from 19 wild felids, including three Persian leopards. Using species-specific hemoplasma PCRs and ELISA serological testing for feline leukaemia virus and feline immunodeficiency virus (FIV), two Persian leopards were found to be infected with ‘*Ca. M. haemominutum*’ and were seropositive for FIV. Partial 16S rRNA gene sequences were generated for these ‘*Ca. M. haemominutum*’ species and subsequent phylogenetic analysis revealed 97.70% to 99.45% sequence identity with those found in domestic cats from Iran and other countries.

This study confirms the presence of ‘*Ca. M. haemominutum*’ and concurrent FIV antibody in wild felids in Iran. This represents the first report of hemoplasma in wild felids in the Middle East as well as the first report of infection in Persian leopards.

### 1. Introduction

Hemoplasmas are hemotropic mycoplasmal bacteria that infect a wide range of mammals [1,2]. At least three feline hemoplasma species have been described in domestic cats including *Mycoplasma haemofelis*, ‘*Candidatus Mycoplasma haemominutum*’, and ‘*Candidatus Mycoplasma turicensis*’ [1–4]. The most pathogenic species is *M. haemofelis*, which can cause hemolytic anemia [5,6] in immunocompetent cats. Coinfection of hemoplasmas with other pathogens such as feline leukemia virus (FeLV) and feline immunodeficiency virus (FIV) may worsen the severity of the hemoplasma-induced anemia and result in anemia following infection with less pathogenic hemoplasmas such as ‘*Ca. M. haemominutum*’, and ‘*Ca. M. turicensis*’ [7,8].

Hemoplasma infection with *M. haemofelis*, ‘*Ca. M. haemominutum*’ and/or ‘*Ca. M. turicensis*’ has been reported in around nine wild felid species worldwide [9–11], with wildlife isolates showing near identity to those found in domestic feline species [9]. There are, however, only limited studies of hemoplasma infections in wild felids, and no studies

have yet been performed in countries in the Middle East, such as Iran, and the natural transmission route for hemoplasmas is not known [9].

The Persian leopard is an endangered wild felid, native to Iran and some neighboring countries. Following the extinction of the lion *Panthera leo persica* and tiger *Panthera tigris virgate* in Iran, it is the only large wild felid now existing in Iran [12–15], and no studies have yet evaluated this species as a host for feline hemoplasma infection. We have recently reported the presence and molecular characterization of feline hemoplasma infections in domestic cats in Iran [16], and the aim of this study was to document the presence and molecularly characterize of feline hemoplasma species in wild felids in Iran in the Middle East.

### 2. Materials & methods

#### 2.1. Sample collection and processing

Nineteen EDTA-anticoagulated blood samples (FL Medical K3 EDTA

\* Corresponding author at: Institute of Microbiology and Epizootics School of Veterinary Medicine, Freie Universität Berlin, Germany.

E-mail addresses: [Fereshteh.Ghazisaeeedi@fu-berlin.de](mailto:Fereshteh.Ghazisaeeedi@fu-berlin.de) (F. Ghazisaeeedi), [natyabi@ut.ac.ir](mailto:natyabi@ut.ac.ir) (N. Atyabi), [tsalehi@ut.ac.ir](mailto:tsalehi@ut.ac.ir) (T. Zahraei Salehi), [tabatabaei.saeid@gmail.com](mailto:tabatabaei.saeid@gmail.com) (S. Tabatabaei), [iradjashrafi@gmail.com](mailto:iradjashrafi@gmail.com) (I. Ashrafi Tamai), [imanmemarianvet@gmail.com](mailto:imanmemarianvet@gmail.com) (I. Memarian), [S.Tasker@bristol.ac.uk](mailto:S.Tasker@bristol.ac.uk) (S. Tasker).

<http://dx.doi.org/10.1016/j.cimid.2017.07.004>

Received 27 September 2016; Received in revised form 6 July 2017; Accepted 9 July 2017  
0147-9571/ © 2017 Elsevier Ltd. All rights reserved.

K3E, Lot. F111332 2.5 mL tube, Torreglia, Italy) were obtained from the following cats; twelve African lions, four leopards (three Persian leopards and one African leopard), one Eurasian lynx, one Bengal tiger and one Caracal, using a blowpipe filled with a combination of drugs and dosage to each species; ketamine (3 mg/kg) and medetomidine (0.03 mg/kg) for Bengal tiger, Persian and African leopards, butorphanol (0.2 mg/kg), medetomidine (0.035 mg/kg) and midazolam (0.15 mg/kg) for Eurasian lynx and Caracal caracal and tiletamine/zolazepam (1.5 mg/kg) and medetomidine (0.015 mg/kg) for African lion. These are given intramuscularly to anesthetize the animals followed by femoral vein sampling. Approval was granted for the study from the Iran Veterinary Organization since samples were taken as part of a national and international cooperative project for conservation of Persian leopards, supported by the Iranian Department of the Environment, the International Union for Conservation of Nature, The Wildlife Conservation Society and Panthera. The sampled animals were kept either in Tehran zoo or in the Tandoureh National Park. Tandoureh National Park has been protected since 1968 and is located in north eastern Iran and is around 355 km<sup>2</sup> in size. Signalment data for these wild felids, as well as their origin and current residence are shown in Table 1.

Hematological parameters including white blood cell, red blood cell (RBC), Hematocrit (HCT), hemoglobin concentration (Hb), mean corpuscular volume, mean corpuscular hemoglobin, mean corpuscular hemoglobin concentration and platelets were measured using an automatic hemocytometer (Hema-screen 18, Hospitex diagnostic, Florence, Italy). Blood smears were prepared for differential white blood cells count and examination for hemoparasites. Plasma was submitted for serological retrovirus testing for FeLV and FIV using a commercially available rapid diagnostic ELISA kit (Quicking FIV Ab + FeLV Ag Combined Test, W81099, China), according to the manufacturer's instructions, and were confirmed by repeat ELISA testing using a different serological retrovirus test (ELISA kit for serodiagnosis of FeLV and FIV Ab, Biopronix, Agrolabo, Italy).

## 2.2. DNA extraction

DNA was extracted from 100 µL whole blood from each sample

**Table 1**  
Signalment data, origin and residence of sampled wild felids.

No.	Species	Scientific name	Gender	Age (years)	Origin	Residence at time of sampling
1	African lion	<i>Panthera leo</i>	Female	6	Born in zoos in Iran	Zoo of Pardisan, Tehran, Iran
2	African lion	<i>Panthera leo</i>	Male	3	Born in zoos in Iran	Zoo of Pardisan, Tehran, Iran
3	African lion	<i>Panthera leo</i>	Female	7	Born in zoos in Iran	Zoo of Pardisan, Tehran, Iran
4	African lion	<i>Panthera leo</i>	Female	1	Born in zoos in Iran	Zoo of Pardisan, Tehran, Iran
5	African lion	<i>Panthera leo</i>	Female	1	Born in zoos in Iran	Zoo of Pardisan, Tehran, Iran
6	African lion	<i>Panthera leo</i>	Female	7	Born in zoos in Iran	Zoo of Pardisan, Tehran, Iran
7	African lion	<i>Panthera leo</i>	Female	2	Born in zoos in Iran	Zoo of Pardisan, Tehran, Iran
8	African lion	<i>Panthera leo</i>	Male	3	Born in zoos in Iran	Zoo of Pardisan, Tehran, Iran
9	African lion	<i>Panthera leo</i>	Male	1	Born in zoos in Iran	Zoo of Pardisan, Tehran, Iran
10	African lion	<i>Panthera leo</i>	Male	2	Born in zoos in Iran	Zoo of Pardisan, Tehran, Iran
11	African lion	<i>Panthera leo</i>	Female	5	Born in zoos in Iran	Zoo of Pardisan, Tehran, Iran
12	African lion	<i>Panthera leo</i>	Female	3	Born in zoos in Iran	Zoo of Pardisan, Tehran, Iran
13	Persian leopard	<i>Panthera pardus</i> ssp. <i>saxicolor</i>	Male	15	Born in wild in Khorasan, Iran	In wild in Iran (transferred to National Park of Tandoureh, Iran at time of sampling)
14	Persian leopard	<i>Panthera pardus</i> ssp. <i>saxicolor</i>	Male	14	Born in wild in Mazandaran, Iran before being transferred to National Park of Tandoureh, Iran	National Park of Tandoureh, Iran
15	Persian leopard	<i>Panthera pardus</i> ssp. <i>saxicolor</i>	Female	4	Born in wild in Golestan, Iran before being transferred to National Park of Tandoureh, Iran	National Park of Tandoureh, Iran
16	African leopard	<i>Panthera pardus</i> ssp. <i>pardus</i>	Male	22	Born in wild in Kenya before being transferred to National Park of Tandoureh, Iran	National Park of Tandoureh, Iran
17	Eurasian lynx	<i>Lynx lynx</i>	Male	8	Born in wild in Iran before being transferred to National Park of Tandoureh, Iran	National Park of Tandoureh, Iran
18	Caracal	<i>Caracal caracal</i>	Male	6	Born in wild in Iran before being transferred to National Park of Tandoureh, Iran	National Park of Tandoureh, Iran
19	Bengal tiger	<i>Panthera tigris</i> ssp. <i>Tigris</i>	Male	1.5	Born in zoo in Denmark	Zoo of Pardisan, Tehran, Iran

using a commercial kit (QIAamp cadior pathogen Mini kit, Qiagen, Hilden, Germany), following the manufacturer's instructions, and stored at –20 °C until further use.

Distilled water and known positive blood samples for each of the three feline hemoplasma species, obtained from the School of Veterinary Sciences, University of Bristol, Bristol, UK and Bologna University, Bologna, Italy, were used as negative and positive controls respectively during each run of DNA extractions.


## 2.3. Diagnostic polymerase chain reaction (PCR) assays

A control conventional PCR to amplify a fragment of feline glyceraldehyde-3-phosphate dehydrogenase (GAPDH) gene was performed to detect possible PCR inhibitors in DNA samples [17]. Screening hemoplasma PCR analysis was performed using a previously described generic universal hemoplasma conventional PCR assay using 5'-ATACGG-CCCATATTCCTACG-3' and 5'-TGCTCCACCCTGTGTTCA-3' as forward and reverse primers, respectively [18].

All samples were then subjected to species-specific conventional PCRs for each of the three feline hemoplasma species using previously described conventional PCR assays [19,20]. Positive controls of *M. haemofelis*, '*Ca. M. haemominutum*' and '*Ca. M. turicensis*' were used for both the generic haemoplasma and species specific PCRs.

## 2.4. 16S rRNA gene sequencing

The 16S rRNA gene of positive samples on generic screening hemoplasma PCR was amplified using primers 8F: 5'-AGAGTTTGATCC TGGCTCAG-3' and 1492R: 5'-GGTTACCTGTGTTACGACTT-3', as previously described, with resulting PCR products then subjected to sequencing using the Sanger technique (ABI, 96-capillary XL) [21]. After evaluating the quality of sequence reading in Finch TV software (Geospiza), 5' and 3' ends of the forward and reverse sequence reading were trimmed. The forward and reverse sequences of each sample were then overlapped and aligned with available 16S rRNA sequences of '*Ca. M. haemominutum*' in Genbank. Finally, partial 16S rRNA sequences of 1086 bp (lacking about 200 bp from each 5' and 3' end of complete 16S rRNA sequence of '*Ca. M. haemominutum*') were obtained.



**Fig. 1.** Phylogenetic analysis of partial 16S rRNA gene sequences from “*Candidatus Mycoplasma haemominutum*” isolates from Persian leopards. Sequences from this study are shown all in bold. Bootstrap values are given at the nodes of the tree. The following sequences are shown: *Mycoplasma haemofelis* (Cat, South Africa AF548631; Cat, Iran, KU852584; Eurasian Lynx, Switzerland, DQ825458; European Wildcat, France, DQ825441; *Leopardus Weidii*, Brazil, DQ825438; Cat, Switzerland, DQ157160; Cat, United Kingdom, AY150985; Cat, Australia, AY150977; Cat, Australia, AY150976; Cat, France, AY150972; Iberian Lynx, Spain, DQ825447; Cat, Iran, KU852583; Cat, United States, U95297, Lion, Tanzania, DQ825451), *Mycoplasma haemocanis* (Dog, United States, AF407208; Dog Germany AY150973), “*Candidatus Mycoplasma haemomuris*” (*Apodemus argenteus*, Japan, AB758437; wild mouse, Japan, U82963; *Apodemus argenteus*, Japan, AB758436; *Rattus rattus*, Japan, AB758439) “*Candidatus Mycoplasma turicensis*” (Lion, Tanzania, DQ825454; *Leopardus Pardalis*, Brazil, DQ825448; Cat Switzerland, DQ157150; European Wildcat, France, DQ825450; *Mycoplasma haemolamae* AF306346, “*Candidatus Mycoplasma haematoparvum*” (Dog, United States, AY383241; Dog, United Kingdom, AY532390) “*Candidatus Mycoplasma haemominutum*” (*Leopardus tigrinus*, Brazil, DQ825439; Iberian Lynx, Spain, DQ825445; Cat, Israel, AY150974; *Leopardus Weidii*, Brazil, DQ825440; Iberian Lynx, Spain, DQ825446; Iberian Lynx, Spain, DQ825444; Euroasian Lynx, Switzerland, DQ825456; Persian Leopard, Iran, KU852587; Persian Leopard, Iran, KU852586; Lion, Tanzania, DQ825452; Cat, United Kingdom, AY150980; Cat United States, U88564; European Wildcat, France, DQ825442; Cat, United Kingdom, AY150981; Cat, Iran, KU852585), *Clostridium perfringens* NR 121697.

## 2.5. Statistical analysis

Data analysis, including descriptive statistics, was performed using SPSS software (16.0 IBM, New York, USA). African lion hematology reference intervals were calculated using the mean  $\pm$  SD data available in previously published work [22], using the formula mean  $\pm$  1.96SD. Sequence data analysis and phylogenetic tree construction were performed with MEGA6 software using the partial 16S rRNA sequences derived in this study as well as other wild and domestic cat hemoplasma sequences downloaded from Genbank (Accession numbers shown in Fig. 1). Bootstrap testing (1000 replicates) and out-grouping were used to validate the phylogenetic tree [23]. The evolutionary distances were computed using the Kimura 2-parameter method [24] and the Neighbor-Joining method [25] used for tree construction [26].

**Nucleotide Sequence Accession Numbers.** The partial 16S rRNA gene sequences derived from this study were submitted to Genbank with accession numbers KU852586 and KU852587.

## 3. Results

Of the 19 samples analyzed, all were PCR-positive for GAPDH, and two (10.5%) were PCR-positive by generic universal hemoplasma conventional PCR. Only the same two samples were positive on species-specific PCR; both for '*Ca. M. haemominutum*' only. All positive controls had expected amplified band in the generic universal and the species-specific conventional hemoplasma PCRs and distilled water as the negative control had none. Both of the positive samples were from old (14 and 15 years) male Persian leopards (Case numbers 13 and 14 in Table 1), from two different geographical areas of Iran. No hemoplasma organisms were observed on blood smear examination.

The two hemoplasma ('*Ca. M. haemominutum*') infected Persian leopards were both also FIV seropositive, and one African lion was also FeLV and FIV positive but not hemoplasma infected. No other samples were retrovirus positive.

To the authors' knowledge, no hematological reference ranges exist for Persian leopards, nor for any closely related species (e.g. African leopard, Arabian leopard), limiting interpretation of the hematology profiles of the Persian leopards in the current study. However, as shown in Table 2, the hematology profiles of the two '*Ca. M. haemominutum*' and FIV-seropositive Persian leopards (Case numbers 13 & 14) showed HCT, Hb, and RBC counts at the lower end of the reference range used for domestic cats, and HCT and RBC counts below the reference range

calculated for African lions based on Larsson et al. [22], and were lower than those recorded in the non-infected Persian leopard (Case number 15). Thus it is possible that '*Ca. M. haemominutum*' and FIV infection were associated with a reduction in erythrocyte indices in the infected cats, but further data from larger numbers of cats would be required to confirm this.

The hematology profiles of the 12 African lions were also compared to the reference range calculated for African lions based on Larsson et al. [22], and 11 of the 12 lions had HCT and Hb values within or above the reference range. The FeLV and FIV seropositive but hemoplasma PCR negative lion had a hypochromic normocytic anemia (HCT 20.7%). The four remaining cases (Case numbers 16, 17, 18 and 19) could not have their hematological profiles determined due to sample hemolysis.

The partial (1086 bp) '*Ca. M. haemominutum*' 16S rRNA gene sequences derived for the two hemoplasma infected Persian leopards in the current study (KU852586 and KU852587) showed high sequence identity (97.7–99.45%) with, and were closely related to the '*Ca. M. haemominutum*' sequences in Genbank derived from worldwide wild felids and domestic cats [9,27–29], including Iranian domestic cats. Data are shown in Fig. 1. The '*Ca. M. haemominutum*' Persian leopard sequence KU852586 was slightly more closely related to the Iranian domestic cat sequence KU852585 than the other Persian leopard sequence KU852587, with 99.26% sequence identity. The sequence identity between the two Persian leopards '*Ca. M. haemominutum*' (KU852586 and KU852587) was 98.43%. The highest sequence identities of 99.63% and 98.62% were between '*Ca. M. haemominutum*' sequences KU852586 and KU852587 derived from Persian leopards and DQ825452 from a lion in Tanzania.

## 4. Discussion

This study documents the presence of hemoplasmas in wild felids for the first time in Iran in the Middle East. It is also the first documentation of hemoplasma infection in the endangered Persian leopard species [12]. The prevalence of hemoplasma infection in wild felids has varied in different studies but is not frequently high. In a surveillance study in Brazil on neotropical and exotic felids, 9.2% of 109 felids were hemoplasma positive (all '*Ca. M. haemominutum*') [30], whilst in free-ranging cheetahs in Namibia, only one of 63 cheetahs was positive [31]. However higher prevalence was reported in another study evaluating a large sample size (275) from worldwide geographical areas where prevalence of 18%, 32% and 20% were found for *M. haemofelis*,

**Table 2**  
Hematological parameters for Persian leopards.

Species	Persian leopard Case no. 13 <sup>a</sup>	Persian leopard Case no. 14 <sup>a</sup>	Persian leopard Case no. 15	Domestic cat Reference range [49]	African lion Reference range [22](mean $\pm$ SD)	Derived African lion Reference range <sup>c</sup>	Unit
Hct <sup>b</sup>	29,4	28,2	34,15	29–45	42,38 $\pm$ 4,73	33,11–51,65	%
Hb	10,2	8,8	12,1	8–14	14,11 $\pm$ 1,63	10,92–17,30	g/dl
RBC	6,1	6,59	7,92	6–10	8,97 $\pm$ 1,43	6,17–11,77	10 <sup>6</sup> / $\mu$ l
MCV	48	50	48,5	41.0–54	47,70 $\pm$ 4,53	38,82–56,58	fl
MCH	12,8	15,7	14,53	13.3–17.5	15,48 $\pm$ 1,25	13,03–17,93	pg
MCHC	26,6	31	29,38	31–36	33,3 $\pm$ 2,02	29,34–37,26	%
Plt	139	102	98,5	2.3–6.8	–	–	10 <sup>5</sup> / $\mu$ l
WBC	9,53	4,06	10,75	5.5–19.5	9,73 $\pm$ 1,43	6,93–12,53	10 <sup>3</sup> / $\mu$ l
Seg.	7,1475	2,436	7,821	2.5–12.5	7,748 $\pm$ 1,209	5,38–10,12	10 <sup>3</sup> / $\mu$ l
Band	0,1906	0,0406	0,131	0–0.3	0	0,00–0,00	10 <sup>3</sup> / $\mu$ l
Lymph	1,906	1,421	2,281	1,5–7	894 $\pm$ 456	0,24–1787,76	10 <sup>3</sup> / $\mu$ l
Mono	0,0953	0,0406	0,067	0–0.85	365 $\pm$ 193	0–743,28	10 <sup>3</sup> / $\mu$ l
Eos	0,1906	0,0812	0,101	0–1.5	372 $\pm$ 364	0–1085,44	10 <sup>3</sup> / $\mu$ l
Baso	0	0	0	Rare	0	0,00–0,00	10 <sup>3</sup> / $\mu$ l

NB: No hematology reference range is available for Persian leopards.

<sup>a</sup> '*Ca. M. haemominutum*' FIV infected Persian leopards.

<sup>b</sup> Hematocrit (HCT), hemoglobin concentration (Hb), Red blood cell count (RBC), mean corpuscular volume (MCV), mean corpuscular hemoglobin (MCH) mean corpuscular hemoglobin concentration (MCHC), platelets (Plt), white blood cell count (WBC), segmented neutrophil (seg), band cell (Band), lymphocyte (Lymph), monocyte (Mono), eosinophil (Eos), basophil (Baso).

<sup>c</sup> Reference range for African lion derived using mean  $\pm$  1.96SD from Larsson et al. [23].

'*Ca. M. haemominutum*', and '*Ca. M. turicensis*', respectively [9]. In the current study, hemoplasma infection was confirmed in just two of 19 samples (10.5%), although the sample size was small since access to wild felid samples in the Middle East is very limited due to the difficulties in access to hosts and collection of blood. Both '*Ca. M. haemominutum*' infected wild felids in the current study were old male Persian leopards, an indigenous species in Iran. This is in agreement with other studies in domestic cats showing that being male and, older, are risk factors for '*Ca. M. haemominutum*' infection [27,32–37]. Fighting behavior is also regarded as a risk factor for hemoplasma infection [27,36,38], although the fighting behavior of the cats sampled was not completely known and leopards (*Panthera pardus*) are generally not regarded as an aggressive species [39]. However, in the literature there are cases of intraspecific killing among leopards over a kill, territory or cannibalism, so aggression is possible [40,41]. There are also two reports of intraspecific killing from Persian leopards in Tandoureh in 2007 and 2016 over food and territory (Memarian. I, personal communication).

Neither of the two '*Ca. M. haemominutum*' infected Persian leopards identified in the current study lived in zoos. As reported in an extensive study on feline hemoplasma infection in wild felid species worldwide [9], free-ranging felids had higher hemoplasma infection prevalence [9,42,43] than captive felids. This may be because free-ranging felids have more fighting and hunting habits and/or more exposure to vectors, than captive or zoo-based felids. In the same study described, a significant correlation between FeLV PCR positivity and hemoplasma infection was found in European wild cats [9]. There are several reports of retrovirus infections in free-ranging and captive wild felids [44–47], and multiple other concurrent infections such as feline calicivirus, feline herpesvirus, feline parvovirus, and feline coronavirus [42,43]. In the current study both '*Ca. M. haemominutum*' Persian leopards were FIV seropositive and this is, to the authors' knowledge, the first report for such a co-infection in a wild felid species.

It is not known if the '*Ca. M. haemominutum*' infection in the Persian leopards caused anemia. This was difficult to assess since no reference ranges exist for hematological parameters in this species, and it is known that greater anemia can occur in cats with concurrent '*Ca. M. haemominutum*' and retrovirus infection compared to '*Ca. M. haemominutum*' alone [7]. The very small sample size did not permit a statistical comparison between '*Ca. M. haemominutum*' FIV-seropositive and FIV-seronegative Persian leopards. Nevertheless, it was of note that the HCT, Hb and RBC counts of the two '*Ca. M. haemominutum*' FIV-seropositive leopards were lower than the Persian leopard free from hemoplasmas and retroviral infection, suggesting that coinfection of '*Ca. M. haemominutum*' and FIV could have been associated with reduced RBC parameters.

The partial 16S rRNA gene phylogenetic analysis found that the '*Ca. M. haemominutum*' isolates derived from this study were closely related to those from different geographical origins and from both domestic and wild felids. In a previous phylogenetic study of domestic feline hemoplasmas, using both 16S rRNA gene and RNaseP genes, almost 100% identity was reported between Europe, Asia, Africa and United States species [48]. In a Japanese study, the identities of the detected hemoplasma sequences was very high, such that it was not possible to assume the origin of *M. haemofelis* and '*Ca. M. turicensis*' from endangered Iriomote cats. In agreement with our findings, previous studies describing feline hemoplasma phylogenetic analysis based on the RNaseP gene revealed similar close relationships between the hemoplasma species of both domestic and wild felids [9,31,48].

A limitation of this study is the small sample size, but despite this, it is interesting to note that two of the three Persian leopards tested were '*Ca. M. haemominutum*' positive, suggesting that hemoplasma infection may be prevalent in this species, especially as the two positive Persian leopards were from geographically distinct areas.

In conclusion, we have documented that hemoplasma infections occur in wild felids and we have reported, for the first time,

hemoplasma infection in wild felids in the Middle East and hemoplasma infection in Persian leopards. Interestingly the two '*Ca. M. haemominutum*' infected Persian leopards were seropositive for FIV. The prevalence of infectious diseases in wild felids is difficult to assess and monitor but should be considered by those working to save endangered animal species such as the Persian leopard.

## Funding

This research did not receive any grants from any funding agencies.

## Conflict of interest

None.

## References

- [1] H. Neimark, K.E. Johansson, Y. Rikihisa, J.G. Tully, Proposal to transfer some members of the genera *Haemobartonella* and *Eperythrozoon* to the genus *Mycoplasma* with descriptions of '*Candidatus Mycoplasma haemofelis*' '*Candidatus Mycoplasma haemomuris*', '*Candidatus Mycoplasma haemostuis*' and '*Candidatus Mycoplasma wenyonii*', *Int. J. Syst. Evol. Microbiol.* 51 (2001) 891–899.
- [2] B. Willi, F.S. Boretti, S. Tasker, M.L. Meli, N. Wengi, C.E. Reusch, H. Lutz, R. Hofmann-Lehmann, From *Haemobartonella* to hemoplasma: molecular methods provide new insights, *Vet. Microbiol.* 125 (2007) 197–209.
- [3] B. Willi, F.S. Boretti, V. Cattori, S. Tasker, M.L. Meli, C. Reusch, H. Lutz, R. Hofmann-Lehmann, Identification, molecular characterization, and experimental transmission of a new hemoplasma isolate from a cat with hemolytic anemia in Switzerland, *J. Clin. Microbiol.* 43 (2005) 2581–2585.
- [4] B. Willi, K. Museux, M. Novacco, E.M. Schraner, P. Wild, K. Groebel, U. Ziegler, G.A. Wolf-Jäckel, Y. Kessler, C. Geret, S. Tasker, H. Lutz, R. Hofmann-Lehmann, First morphological characterization of '*Candidatus Mycoplasma turicensis*' using electron microscopy, *Vet. Microbiol.* 149 (2011) 367–373.
- [5] S. Tasker, Haemotropic mycoplasmas: what's their real significance in cats? *J. Feline Med. Surg.* 12 (2010) 369–381.
- [6] L.M. Berent, J.B. Messick, S.K. Cooper, Detection of *Haemobartonella felis* in cats with experimentally induced acute and chronic infections, using a polymerase chain reaction assay, *Am. J. Vet. Res.* 59 (1998) 1215–1220.
- [7] J.W. George, B.A. Rideout, S.M. Griffey, N.C. Pedersen, Effect of preexisting FeLV infection or FeLV and feline immunodeficiency virus coinfection on pathogenicity of the small variant of *Haemobartonella felis* in cats, *Am. J. Vet. Res.* 63 (2002) 1172–1178.
- [8] D.B. Macieira, C. de Menezes Rde, C.B. Damico, N.R. Almosny, H.L. McLane, J.K. Daggy, J.B. Messick, Prevalence and risk factors for hemoplasmas in domestic cats naturally infected with feline immunodeficiency virus and/or feline leukemia virus in Rio de Janeiro–Brazil, *J. Feline Med. Surg.* 10 (2008) 120–129.
- [9] B. Willi, C. Filoni, J.L. Catão-Dias, V. Cattori, M.L. Meli, A. Vargas, F. Martínez, M.E. Roelke, M.-P. Ryser-Degiorgis, C.M. Leutenegger, H. Lutz, R. Hofmann-Lehmann, Worldwide occurrence of feline hemoplasma infections in wild felid species, *J. Clin. Microbiol.* 45 (2007) 1159–1166.
- [10] M. Haefner, T.J. Burke, B.E. Kitchell, L.A. Lamont, D.J. Schaeffer, M. Behr, J.B. Messick, Identification of *Haemobartonella felis* (*Mycoplasma haemofelis*) in captive nondomestic cats, *J. Zoo Wildl. Med.* 34 (2003) 139–143.
- [11] M.R. André, C.H. Adania, S.M. Allegritti, R.Z. Machado, Hemoplasmas in wild canids and felids in Brazil, *J. Zoo. Wildl. Med.* 42 (2011) 342–347.
- [12] International Union for Conservation of Nature (IUCN), The IUCN Red List of Threatened Species, (2015) <http://www.iucnredlist.org>.
- [13] M.S. Farhadinia, H. Farahmand, A. Gavashelishvili, M. Kaboli, M. Karami, B. Khalili, S. Montazamy, Molecular and craniological analysis of leopard, *Panthera pardus* (Carnivora: Felidae) in Iran: support for a monophyletic clade in Western Asia, *Biol. J. Linn. Soc.* 114 (2015) 721–736.
- [14] A. Ghoddousi, A. Kh Hamidi, T. Ghadirian, D. Ashayeri, I. Khorozyan, The status of the endangered Persian leopard *Panthera pardus saxicolor* in Bamou National Park, Iran, *Oryx* 44 (2010) 551–557.
- [15] B.H. Kiabi, B.F. Dareshouri, R.A. Ghaemi, M. Jahanshahi, Population status of the Persian Leopard (*Panthera pardus saxicolor* Pocock, 1927) in Iran, *Zool. Middle East* 26 (2002) 41–47.
- [16] F. Ghazisaeedi, N. Atyabi, T. Zahrai Salehi, F. Gentilini, I. Ashrafi Tamai, H. Akbarain, S. Tasker, A molecular study of hemotropic mycoplasmas (hemoplasmas) in cats in Iran, *Vet. Clin. Pathol.* 43 (2014) 381–386.
- [17] A.J. Birkenheuer, M.G. Levy, E.B. Breitschwerdt, Development and evaluation of a seminested PCR for detection and differentiation of *Babesia gibsoni* (Asian genotype) and *B. canis* DNA in canine blood samples, *J. Clin. Microbiol.* 41 (2003) 4172–4177.
- [18] A. Criado-Fornelio, A. Martinez-Marcos, A. Buling-Saraña, J.C. Barba-Carretero, Presence of *Mycoplasma haemofelis*, *Mycoplasma haemominutum* and piroplasmids in cats from southern Europe: a molecular study, *Vet. Microbiol.* 93 (2003) 307–317.
- [19] W.A. Jensen, M.R. Lappin, S. Kamkar, W.J. Reagan, Use of a polymerase chain reaction assay to detect and differentiate two strains of *Haemobartonella felis* in naturally infected cats, *Am. J. Vet. Res.* 62 (2001) 604–608.
- [20] A.P. Santos, J.B. Messick, A.W. Biondo, S.T. Oliveira, V. Pedralli, C.S. Lasta,

- L.A. Lacerda, V.S. Esteves, R. Hofmann-Lehmann, B. Willi, F.H.D. González, Design optimization, and application of a conventional PCR assay with an internal control for detection of 'Candidatus Mycoplasma turicensis' 16S rDNA in domestic cats from Brazil, *Vet. Clin. Pathol.* 38 (2009) 443–452.
- [21] C. Pitulle, D.M. Citron, B. Bochner, R. Barbers, M.D. Appleman, Novel bacterium isolated from a lung transplant patient with cystic fibrosis, *J. Clin. Microbiol.* 37 (1999) 3851–3855.
- [22] M.H.M.A. Larsson, P.L. do Espírito Santo, R.M.S. Mirandola, J.D.L. Fedullo, F.H. Ito, P.H. Itikawa, R.B. Pessoa, Hematologic parameters of captive Lions (*Panthera leo*) and Siberian Tigers (*Panthera tigris altaica*), *Acta Sci. Vet.* 43 (2015) 1311.
- [23] J. Felsenstein, Confidence limits on phylogenies: an approach using the bootstrap, *Evolution* 39 (1985) 783–791.
- [24] M. Kimura, A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences, *J. Mol. Evol.* 16 (1980) 111–120.
- [25] N. Saitou, M. Nei, The neighbor-joining method: a new method for reconstructing phylogenetic trees, *Mol. Biol. Evol.* 4 (1987) 406–425.
- [26] K. Tamura, G. Stecher, D. Peterson, A. Filipski, S. Kumar, MEGA6: molecular evolutionary genetics analysis version 6.0, *Mol. Biol. Evol.* 30 (2013) 2725–2729.
- [27] B. Willi, F.S. Boretti, C. Baumgartner, S. Tasker, B. Wenger, V. Cattori, M.L. Meli, C.E. Reusch, H. Lutz, R. Hofmann-Lehmann, Prevalence, risk factor analysis, and follow-up of infections caused by three feline hemoplasma species in cats in Switzerland, *J. Clin. Microbiol.* 44 (2006) 961–969.
- [28] S. Tasker, C.R. Helps, M.J. Day, T.J. Gruffydd-Jones, D.A. Harbour, Use of real-time PCR to detect and quantify *Mycoplasma haemofelis* and *Candidatus Mycoplasma haemominutum* DNA, *J. Clin. Microbiol.* 41 (2003) 439–441.
- [29] Y. Rikihisa, M. Kawahara, B. Wen, G. Kociba, P. Fuerst, F. Kawamori, C. Suto, S. Shibata, M. Futohashi, Western immunoblot analysis of *Haemobartonella muris* and comparison of 16S rRNA gene sequences of *H. muris*, *H. felis*, and *Eperythrozoon suis*, *J. Clin. Microbiol.* 35 (1997) 823–829.
- [30] C. Filoni, J.L. Catão-Dias, V. Cattori, B. Willi, M.L. Meli, S.H.R. Corrêa, M.C. Marques, C.H. Adania, J.C.R. Silva, M.F.V. Marvulo, J.S.F. Neto, E.L. Durigon, V.M. de Carvalho, S.D.A. Coutinho, H. Lutz, R. Hofmann-Lehmann, Surveillance using serological and molecular methods for the detection of infectious agents in captive Brazilian neotropical and exotic felids, *J. Vet. Diagn. Invest.* 24 (2012) 166–173.
- [31] A. Krenzel, M.L. Meli, V. Cattori, B. Wachter, B. Willi, S. Thalwitzer, J. Melzheimer, H. Hofer, H. Lutz, R. Hofmann-Lehmann, First evidence of hemoplasma infection in free-ranging Namibian cheetahs (*Acinonyx jubatus*), *Vet. Microbiol.* 162 (2013) 972–976.
- [32] S. Tasker, S.H. Binns, M.J. Day, T.J. Gruffydd-Jones, D.A. Harbour, C.R. Helps, W.A. Jensen, C.S. Olver, M.R. Lappin, Use of a PCR assay to assess the prevalence and risk factors for *Mycoplasma haemofelis* and 'Candidatus Mycoplasma haemominutum' in cats in the United Kingdom, *Vet. Rec.* 152 (2003) 193–198.
- [33] C.B. Grindem, W.T. Corbett, M.T. Tomkins, Risk factors for *Haemobartonella felis* infection in cats, *J. Am. Vet. Med. Assoc.* 196 (1990) 96–99.
- [34] F. Gentilini, M. Novacco, M.E. Turba, B. Willi, M.L. Bacci, R. Hofmann-Lehmann, Use of combined conventional and real-time PCR to determine the epidemiology of feline hemoplasma infections in northern Italy, *J. Feline Med. Surg.* 11 (2009) 277–285.
- [35] B. Willi, S. Tasker, F.S. Boretti, M.G. Doherr, V. Cattori, M.L. Meli, R.G. Lobetti, R. Malik, C.E. Reusch, H. Lutz, R. Hofmann-Lehmann, Phylogenetic analysis of *Candidatus Mycoplasma turicensis* isolates from pet cats in the United Kingdom, Australia, and South Africa, with analysis of risk factors for infection, *J. Clin. Microbiol.* 44 (2006) 4430–4435.
- [36] L.C. Aquino, C.A. Hicks, M.C. Scaloni, M.G. Lima, S. Lemos Mdos, G.R. Paludo, C.R. Helps, S. Tasker, Prevalence and phylogenetic analysis of hemoplasmas from cats infected with multiple species, *J. Microbiol. Methods* 107 (2014) 189–196.
- [37] S. Tasker, J.A. Braddock, R. Baral, C.R. Helps, M.J. Day, T.J. Gruffydd-Jones, R. Malik, Diagnosis of feline hemoplasma infection in Australian cats using a real-time PCR assay, *J. Feline Med. Surg.* 6 (2004) 345–354.
- [38] M. Tanahara, S. Miyamoto, T. Nishio, Y. Yoshii, M. Sakuma, Y. Sakata, K. Nishigaki, H. Tsujimoto, A. Setoguchi, Y. Endo, An epidemiological survey of feline hemoplasma infection in Japan, *J. Vet. Med. Sci.* 72 (2010) 1575–1581.
- [39] A.B. Stein, V. Hayssen, *Panthera pardus* (Carnivora: felidae), *Mamm. Spec.* 45 (2013) 30–48.
- [40] G.A. Balme, R. Slotow, L.T.B. Hunter, Impact of conservation interventions on the dynamics and persistence of a persecuted leopard (*Panthera pardus*) population, *Biol. Conserv.* 142 (2009) 2681–2690.
- [41] V. Steyn, P.J. Funston, A case of cannibalism in leopards, *S. Afr. J. Wildl. Res.* 36 (2006) 189–190.
- [42] R. Hofmann-Lehmann, D. Fehr, M. Grob, M. Elgizoli, C. Packer, J.S. Martenson, S.J. O'Brien, H. Lutz, Prevalence of antibodies to feline parvovirus calicivirus, herpesvirus, coronavirus, and immunodeficiency virus and of feline leukemia virus antigen and the interrelationship of these viral infections in free-ranging lions in east Africa, *Clin. Diagn. Lab. Immunol.* 3 (1996) 554–562.
- [43] C.M. Leutenegger, R. Hofmann-Lehmann, C. Riols, M. Liberek, G. Worel, P. Lups, D. Fehr, M. Hartmann, P. Weilenmann, H. Lutz, Viral infections in free-living populations of the European wildcat, *J. Wildl. Dis.* 35 (1999) 678–686.
- [44] R.A. Olmsted, R. Langley, M.E. Roelke, R.M. Goeken, D. Adger-Johnson, J.P. Goff, J.P. Albert, C. Packer, M.K. Laurenson, T.M. Caro, Worldwide prevalence of lentivirus infection in wild feline species: epidemiologic and phylogenetic aspects, *J. Virol.* 66 (1992) 6008–6018.
- [45] M.C. Barr, P.P. Calle, M.E. Roelke, F.W. Scott, Feline immunodeficiency virus infection in nondomestic felids, *J. Zoo Wildl. Med.* 20 (1989) 265–272.
- [46] M.J. Daniels, M.C. Golder, O. Jarrett, D.W. MacDonald, Feline viruses in wildcats from Scotland, *J. Wildl. Dis.* 35 (1999) 121–124.
- [47] H. Lutz, E. Isenbügel, R. Lehmann, R.H. Sabapara, C. Wolfensberger, Retrovirus infections in non-domestic felids: serological studies and attempts to isolate a lentivirus, *Vet. Immunol. Immunopath.* 35 (1992) 215–224.
- [48] S. Tasker, C.R. Helps, M.J. Day, D.A. Harbour, S.E. Shaw, S. Harrus, G. Baneth, R.G. Lobetti, R. Malik, J.P. Beaufils, C.R. Belford, T.J. Gruffydd-Jones, Phylogenetic analysis of hemoplasma species: an international study, *J. Clin. Microbiol.* 41 (2003) 3877–3880.
- [49] D.J. Weiss, J.K. Wardrop, *Schalm's Veterinary Hematology*, 6th ed., Wiley-Blackwell, 2010.