

CASE REPORT

Open Access



Studies on *Trueperella pyogenes* isolated from an okapi (*Okapia johnstoni*) and a royal python (*Python regius*)

Marwa F. E. Ahmed¹, Mazen Alssahen², Christoph Lämmle², Tobias Eisenberg³, Madeleine Plötz⁴ and Amir Abdulmawjood^{4*}

Abstract

Background: The present study was designed to characterize phenotypically and genotypically two *Trueperella pyogenes* strains isolated from an okapi (*Okapia johnstoni*) and a royal python (*Python regius*).

Case presentation: The species identity could be confirmed by phenotypic properties, by MALDI-TOF MS analysis and by detection of *T. pyogenes* chaperonin-encoding gene *cpn60* with a previously developed loop-mediated isothermal amplification (LAMP) assay. Furthermore, sequencing of the 16S ribosomal RNA (rRNA) gene, the 16S-23S rDNA intergenic spacer region (ISR), the target genes *rpoB* encoding the β -subunit of bacterial RNA polymerase, *tuf* encoding elongation factor tu and *plo* encoding the putative virulence factor pyolysin allowed the identification of both *T. pyogenes* isolates at species level.

Conclusions: Both strains could be clearly identified as *T. pyogenes*. The *T. pyogenes* strain isolated in high number from the vaginal discharge of an okapi seems to be of importance for the infectious process; the *T. pyogenes* strain from the royal python could be isolated from an apparently non-infectious process. However, both strains represent the first isolation of *T. pyogenes* from these animal species.

Keywords: 16S rRNA gene, 16S-23S rDNA intergenic spacer region, LAMP, MALDI-TOF MS, pyolysin, *rpoB*, *tuf*

Background

Trueperella pyogenes is worldwide considered as part of the commensal biota of skin and mucous membranes of the upper respiratory and urogenital tract of animals [1]. However, *T. pyogenes* is also an important opportunistic pathogen that causes mastitis, abortion and a variety of diverse pyogenic infections in livestock, including cattle, sheep, goats, horses, and pigs [2–4]. In cattle, *T. pyogenes* appears to be responsible for infections of the reproductive tract [5] and the mammary gland [6], as well as cases of pneumonia and liver abscessation of large

and small ruminants [7]. In swine, *T. pyogenes* is well known as a causative agent of different types of inflammation in various organs including the lung, heart, joints, mammary glands, and in the reproductive tract [8, 9]. Furthermore, *T. pyogenes* could be found in companion animals [4]. One of the first reported cases in companion animals was an otitis externa detected in a cat and cystitis in a dog [10]. More recently, Wareth et al. [11] described a co-infection case of *T. pyogenes* with *Brucella abortus* in a cat and dog. Additionally, various wildlife animals could harbour *T. pyogenes* [3]. In 2010, Ülbeği-Mohyla et al. [12] characterized two *T. pyogenes* strains isolated from a bearded dragon and a gecko. Additionally, *T. pyogenes* infections were reported from a bison and from camels [13, 14], from goitered

* Correspondence: amir.abdulmawjood@tiho-hannover.de

⁴Institute of Food Quality and Food Safety, Research Center for Emerging Infections and Zoonoses (RIZ), University of Veterinary Medicine Hannover, Bischofsholer Damm 15, D-30173 Hannover, Germany
Full list of author information is available at the end of the article



© The Author(s). 2020 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

gazelles [15] and from a white-tailed deer [16]. Likewise, some other sporadic cases of infectious diseases associated with *T. pyogenes* were described in a galago [17], in gray slender lorises [18, 19] and in a eurasian lynx [20].

Besides conventional bacteriological methods for identifying *T. pyogenes* isolates, other new, fast and reliable techniques were described and utilized in this study: matrix-assisted laser desorption/ionization time-of-flight mass spectrometry (MALDI-TOF MS) [20–23], a loop-mediated isothermal amplification (LAMP) assay [24] and 16S rRNA gene sequencing [25, 26].

The present study was designed to identify and further characterize *T. pyogenes* isolated from wildlife animals phenotypically and genotypically. To the best of our knowledge, the present study provides a first detailed description of *T. pyogenes* recovered from an okapi and a royal python.

Case presentation

As part of routine examination and diagnostics performed on zoo animals at Frankfurt Zoo (Frankfurt am Main, Germany) in 2019, *T. pyogenes* 24398 was isolated

Table 1 Biochemical properties of *T. pyogenes* 24398 (okapi), *T. pyogenes* 171003246 (royal python) and type strain *T. pyogenes* DSM 20630^T


Biochemical properties	<i>T. pyogenes</i> 24398 (okapi)	<i>T. pyogenes</i> 171003246 (royal python)	<i>T. pyogenes</i> ^a DSM 20630 ^T
Haemolysis on SBA ^b	+	+	+
CAMP-like hemolytic reaction with ^c :			
<i>Staphylococcus aureus</i>	+	+	+
<i>Streptococcus agalactiae</i>	–	–	–
<i>Rhodococcus hoagii</i>	+	+	+
Reverse CAMP reaction	–	–	–
Nitrate reduction	–	–	–
Pyrazinamidase	–	–	–
Pyrrolidonyl Arylamidase	+	+	+
Alkaline phosphatase	+	+	+
β-Glucuronidase (β-GUR)	+	+	+
β-Galactosidase (β-GAL)	+	+	+
α-Glucosidase (α-GLU)	+	+	+
N-Acetyl-β-glucosaminidase (β-NAG)	+	+	+
Esculin (β-glucosidase)	–	–	–
Urease	–	–	–
Gelatine	+	+	+
Fermentation of:			
Glucose	+	+	+
Ribose	+	+	+
Xylose	+	+	+
Mannitol	–	–	–
Maltose	+	+	+
Lactose	+	+	+
Saccharose	–	+	–
Glycogen	+	+	+
Catalase	–	–	–
Serolysis on Loeffler agar	+	+	+
Identification % according to API-Coryne test System	99.9	99.9	99.9

^aResults taken from (Ülbeği-Mohyla et al. 2010 [12]; Hijazin et al. 2011 [27]; Eisenberg et al. 2012 [18]; Alssahen et al. 2020 [20])

^bSBA: Sheep Blood Agar

^csynergistic or reverse CAMP-like reaction with indicator strains

+: positive reaction, –: negative reaction, ^T:type strain



from a vaginal discharge of an okapi (*Okapia johnstoni*) in high numbers (+++), together with *Enterobacter cloacae* (+) and *Pasteurella* spp. (+). The initial bacteriology analysis for *T. pyogenes* 24398 was performed at Hessian State Laboratory (LHL) Gießen, Germany. As a result of post-mortem examination conducted in 2017, *T. pyogenes* 171003246 was recovered in low numbers (+) from a kidney of a seven-year-old female royal python (*Python regius*). The python was found dead in a bird park in Hesse (Germany) and was 107 cm in length and weighted 1.23 kg. In addition, *Escherichia coli* (+), α -hemolytic streptococci (+), *Corynebacterium* spp. (+) and *Clostridium sardiniense* (+) were cultured from the python specimen. The post-mortem analysis of the royal python revealed a good body condition and in the throat and head area a 15 cm lung edema and swelling, possibly caused by traumatic reasons. The post-mortem examination and the initial bacteriology analysis were also performed at Hessian State Laboratory. Both *T. pyogenes*

strains were further investigated phenotypically and genotypically.


Phenotypic characterization

A phenotypic characterization was performed using conventional cultural and biochemical assays as previously described [12, 18, 20, 27] and the API-Coryne test System (BioMérieux, Nürtingen, Germany) in accordance with the manufacturer’s instructions. Furthermore, the bacterial isolates were identified by MALDI-TOF MS using a Microflex LT (Bruker Daltonik GmbH, Bremen, Germany) instrument and MBT Compass Explorer 4.1 software (Bruker Daltonik GmbH). Sample preparation was carried out in accordance with the manufacturer’s instructions using the direct transfer method. Briefly, one microbial colony was first smeared in duplicate onto spots of the MALDI target MSP 96 target (MicroScout Target plate; Bruker Daltonik GmbH) with sterile toothpicks. The air-dried bacteria were overlaid with 1 μ l of an α -cyan 4-hydroxycinnamic acid matrix solution

Table 2 Results of LAMP including detection time and annealing temperature of the tested isolate, positive and negative control

Sample ID	<i>T. pyogenes</i> 24398	<i>T. pyogenes</i> 171003246	<i>T. pyogenes</i> DSM 20630 ^T	<i>T. abortus</i> DSM 19515 ^T	<i>T. bernardiae</i> DSM 9152 ^T	<i>T. bonasi</i> DSM 17163 ^T	HPLC water and Master mix
Result	+ve	+ve	+ve	-ve	-ve	-ve	-ve
Detection time hh:mm:ss	00:11:00	00:08:30	00:10:00	0	0	0	0
Annealing	89.6	89.4	89.9	0	0	0	0

+ve: Positive, -ve: Negative, ^T:type strain



(HCCA, in 50% acetonitrile and 2.5% trifluoroacetic acid in pure water) followed by drying and loading into the mass spectrometer.

Genotypic properties

The genomic DNA of both isolates and the type strains *T. pyogenes* DSM 20630^T (pig), *T. abortus* DSM 19515^T (placenta of sow after abortion), *T. bernardiae* DSM 9152^T (human blood) and *T. bonasi* DSM 17163^T (european bison) were extracted using the DNeasy blood and tissue kit (Qiagen GmbH, Hilden, Germany), in accordance with the manufacturer's instructions. The concentration and purity of DNA were measured by utilizing a Nano Drop spectrophotometer (ND1000; Thermo Fisher Scientific GmbH, Dreieich, Germany).

The detection of gene *cpn60* of *T. pyogenes* was performed using a previously designed loop-mediated isothermal amplification (LAMP) assay [24] with a portable real-time fluorometer (Genie II[®], OptiGene Ltd, Hershaw, UK) and the reference strains *T. pyogenes* DSM 20630^T, *T. abortus* DSM 19515^T, *T. bernardiae* DSM 9152^T and *T. bonasi* DSM 17163^T.

Both *T. pyogenes* isolates were also evaluated by PCR for the presence of five genomic targets: 16S rRNA gene (16S), 16S-23S rDNA intergenic spacer region (ISR), the β -subunit of bacterial RNA polymerase encoding gene *rpoB*, the elongation factor tu encoding gene *tuf*, and pyolysin encoding gene *plo*. The sequence of the oligonucleotide primers and PCR conditions were previously described by Hassan et al. [25], Ülbegi-Mohyla et al. [12], Hijazin et al. [27], Eisenberg et al. [18], Wickhorst et al. [28], Wickhorst et al. [23], Alssahen et al. [20].

The PCR products were purified and sequenced by Eurofins Umwelt Nord GmbH (Göttingen, Germany). The obtained sequences were analyzed via the cluster method of the MegAlign program (DNASTAR Inc., ver. 15, Madison, WI, USA) by comparing with the nucleotide sequences of 16S rRNA, ISR, *rpoB*, *tuf* and *plo* from different *Truiperella* reference strains. Moreover, the resulting amino acid sequences of pyolysin of both *T. pyogenes* isolates were compared with the respective sequences of pyolysin of *T. pyogenes* DSM 20630^T, closely related pore-forming toxins of genus *Arcanobacterium* and with other bacterial pore-forming toxins. All the nucleotide and amino acid sequences were obtained from the NCBI GenBank.

Discussion and Conclusion

Both *T. pyogenes* strains investigated in the present study showed a narrow zone of complete hemolysis on 5% sheep blood agar and CAMP-like reactions in the staphylococcal β -hemolysin zone with *Rhodococcus hoagii* as an indicator strain. The conventional biochemical properties and the results of the commercial identification system revealed almost identical results to previously investigated *T. pyogenes* of various origins and *T. pyogenes* DSM 20630^T [12, 18, 20, 27] (Table. 1). The *T. pyogenes* isolates yielded positive reactions for pyrrolidonyl arylamidase, alkaline phosphatase, β -glucuronidase, β -galactosidase, α -glucosidase and N-acetyl- β -glucosaminidase and negative reactions for nitrate reduction and pyrazinamidase. Additionally, the isolates hydrolyzed gelatine, but not esculin and urea. The isolates also fermented D-glucose, D-ribose, D-xylose, D-maltose, D-




Fig. 3 Phylogenetic analyses based on ISR (a), tuf (b) and ropB (c) nucleotide sequences of the investigated *T. pyogenes* 24398 and *T. pyogenes* 171003246 and *T. pyogenes* S 1276/1/18 isolated from a eurasian lynx and the control strains, *T. pyogenes* DSM 20630^T, *T. abortus* DSM 19515^T, *T. bernardiae* DSM 9152^T, *T. bialowiezensis* DSM 17162^T, and *T. bonasi* DSM 17163^T

lactose and glycogen, but not D-mannitol. *T. pyogenes* 24398 fermented D-saccharose; however, *T. pyogenes* 171003246 was D-saccharose negative. In addition, both isolates showed a negative catalase reaction and a

positive reaction on Löffler agar (Table 1). A positive reaction on Löffler agar is typical for *T. pyogenes* and widely used for phenotypic identification of this species [2, 18].




Fig. 4 Phylogenetic relationships among amino acid sequences of PLO of the investigated *T. pyogenes* 24398 and *T. pyogenes* 171003246, PLO of type strain *T. pyogenes* 20630^T, *T. pyogenes* S 1276/1/18 isolated from a eurasian lynx, ALN of *A. haemolyticum*, PHL of *A. phocae*, HLY of *L. monocytogenes*, PLY of *S. pneumoniae*

Moreover, MALDI-TOF MS identified *T. pyogenes* 24398 and *T. pyogenes* 171003246 with log score values of 2.35 and 2.29 for the first hit and log score values of 2.28 and 1.9 for the second hit, respectively (data not shown). These log score values confirmed, in accordance with the current decision rules of the manufacturer, the species designation. Comparable to the present results, MALDI-TOF MS had already been shown to be a rapid and reliable technique for identifying bacteria of genera *Arcanobacterium* and *Trueperella*, including *T. pyogenes* [20, 21].

The previously described *cpn60*-specific LAMP assay could successfully be used to identify the species-specific gene *cpn60* of *T. pyogenes* 24,398 and *T. pyogenes* 171,003,246 in the present investigation. This was comparable to the LAMP assay for detecting gene *cpn60* of the previously described *T. pyogenes* of various origins [24], a *T. pyogenes* strain isolated from an adult roebuck (*Capreolus capreolus*) [23], and a *T. pyogenes* strain isolated from a eurasian lynx (*Lynx lynx*) [20]. The results of the *cpn60* LAMP assay are shown in Fig. 1; Table 2.

The oligonucleotide primers, 16SUNI-L and 16SUNI-R, were used for amplifying of 16S rRNA gene of the investigated *T. pyogenes* isolates. The nucleotide sequence data of *T. pyogenes* 24398 (GenBank accession numbers: MN946520) and *T. pyogenes* 171003246 (MN712476) were compared with type strain *T. pyogenes* DSM 20630^T (AAC45754) and with the previously described strain *T. pyogenes* S 1276/1/18 isolated from a eurasian

lynx (MN135984), *T. abortus* DSM 19515^T (FN667628), *T. bernardiae* DSM 9152^T (X79224), *T. bielowiezensis* DSM 17162^T (EU194569), and *T. bonasi* DSM 17163^T (EU194570). The nucleotide sequence data of *T. pyogenes* 24398 and *T. pyogenes* 171003246 revealed a sequence homology of 98.9% among both strains, a sequence homology of 99.5% and 98.7% with *T. pyogenes* DSM 20,630^T, and a sequence homology of 99.9% and 99.1% with *T. pyogenes* S1276/1/18, respectively. The control strains of genus *Trueperella* yielded a sequence homology to both *T. pyogenes* isolates \leq 98.7% (Fig. 2).

Both strains *T. pyogenes* 24398 and *T. pyogenes* 171003246 were further identified by sequencing ISR, the genes *tuf* and *rpoB* and the putative virulence factor pyolysin encoding gene *plo*. *T. pyogenes* 24398 and *T. pyogenes* 171003246 showed sequence similarities of ISR (MN947249, MN724920) of 99.8% and 98.9% with *T. pyogenes* DSM 20630^T (EU194563) and 100% and 99.8% with *T. pyogenes* S 1276/1/18 (MN164031), respectively with 98.5% identity between both strains. The additionally investigated gene *tuf* (MN956808, MN741111) showed a sequence similarity of 99.6% and 99.7% with *T. pyogenes* DSM 20630^T (HG941716), and 99.6% and 99.7% with *T. pyogenes* S 1276/1/18 (MN163266), respectively; gene *rpoB* (MN956807, MN741109), a sequence similarity of 99.8% and 98.3% with *T. pyogenes* DSM 20630^T (FN550375), and 98.8% and 98.3% with *T. pyogenes* S 1276/1/18 (MN163265), respectively, and

gene *plo* (MN956806, MN741110), a sequence similarity of 99.5% with *T. pyogenes* DSM 20630^T (U84782) for both isolates, and 99.1% with *T. pyogenes* S 1276/1/18 (MN163264) for both isolates.

Dendrograms of the *ISR*, *tuf* and *rpoB* genes are presented in Fig. 3.

A phylogenetic analysis of the amino acid sequences of pyolysin (PLO) encoded by gene *plo* of *T. pyogenes* 24398 (MN956806), and *T. pyogenes* 171003246 (MN741110) PLO of type strain *T. pyogenes* DSM 20630^T (AAC45754), PLO of *T. pyogenes* S 1276/1/18 (MN163264), arcanolysin (ALN) of *Arcanobacterium haemolyticum* (ACV96715), phocaelysin (PHL) of *Arcanobacterium phocae* 10002^T (SMR98720), listeriolysin O (HLY) of *Listeria monocytogenes* (NP_463733), intermedilysin (ILY) of *Streptococcus intermedius* (BAA89790), pneumolysin (PLY) of *Streptococcus pneumoniae* (ADF28298) and streptolysin O (SLO) of *Streptococcus pyogenes* (BAB41212). The results showed an amino acid similarity of 99.5% for both *T. pyogenes* 24,398 and *T. pyogenes* 171003246 with PLO of *T. pyogenes* DSM 20630^T and 99.1% with PLO of *T. pyogenes* S 1276/1/18 (Fig. 4).

T. pyogenes 24398 was isolated in high numbers from vaginal discharge of an okapi and seems to be responsible for the infectious process; *T. pyogenes* 171003246 was isolated from a non-infectious process of a royal python suffering from a throat swelling, possibly caused by trauma. Both *T. pyogenes* isolates were identified by a biochemical test, LAMP and MALDI-TOF MS. The genomic targets of the two isolates, 16S rRNA gene, *ISR*, *tuf*, *rpoB* and *plo* were sequenced and compared to the respective targets of reference and other strains. Thus, the report is the first to provide a detailed characterization of *T. pyogenes* strains of these origin.

Abbreviations

MALDI-TOF MS: Matrix-assisted laser desorption/ionization time-of-flight mass spectrometry; LAMP: Loop-mediated isothermal amplification; rRNA: Ribosomal RNA; *ISR*: 16S-23S rDNA intergenic spacer region; CAMP test: Christie-Atkins-Munch-Peterson-test; DSM: German Collection of Microorganisms; ALN: Arcanolysin; PHL: Phocaelysin; HLY: Listeriolysin O; ILY: Intermedilysin; PLY: Pneumolysin; SLO: Streptolysin O

Acknowledgements

Not applicable.

Authors' contributions

MFEA, MA, CL, AA and MP contributed to the design of the study, collected and analysed the data. TE performed the initial examination of the isolates. MFEA, AA and MP drafted the manuscript. All authors read and approved the final manuscript.

Funding

Not applicable.

Availability of data and materials

The datasets generated during the current study are available in the NCBI GenBank repository, under the accession number: MN741109, MN741110, MN741111, MN956806, MN956807 and MN956808.

Ethics approval and consent to participate

This study did not require official or institutional ethical approval. The material was collected post mortem and/or during routine diagnosis. According to competent authorities, this kind of research does not require ethics approval or general approval with respect to German law.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Hygiene and Zoonoses Department, Faculty of Veterinary Medicine, Mansoura University, Elgomhoria Street 60, 35516 Mansoura, Egypt. ²Institut für Hygiene und Infektionskrankheiten der Tiere, Justus-Liebig-Universität Gießen, Frankfurterstraße 85-91, D-35392 Gießen, Germany. ³Landesbetrieb Hessisches Landeslabor (LHL), Schubertstraße 60, D-35392 Gießen, Germany. ⁴Institute of Food Quality and Food Safety, Research Center for Emerging Infections and Zoonoses (RIZ), University of Veterinary Medicine Hannover, Bischofsholer Damm 15, D-30173 Hannover, Germany.

Received: 9 June 2020 Accepted: 3 August 2020

Published online: 14 August 2020

References

- Rzewuska M, Kwiecien E, Chrobak-Chmiel D, Kizerwetter-Swida M, Stefanska I, Gierynska M. Pathogenicity and virulence of *Trueperella pyogenes*: a review. *Int J Mol Sci*. 2019;20(11):2737.
- Lämmle C, Hartwig H. *Actinomyces pyogenes* und *Arcanobacterium haemolyticum*. In: Handbuch der bakteriellen Infektionen bei Tieren. 2nd edn: Gustav Fischer: Jena 1995: 196–240.
- Jost BH, Billington SJ. *Arcanobacterium pyogenes*: molecular pathogenesis of an animal opportunist. *Antonie Van Leeuwenhoek*. 2005;88(2):87–102.
- Ribeiro MG, Rissetti RM, Bolanos CAD, Caffaro KA, de Morais ACB, Lara GHB, et al. *Trueperella pyogenes* multispecies infections in domestic animals: a retrospective study of 144 cases (2002 to 2012). *Vet Quart*. 2015;35(2):82–7.
- Tamai IA, Mohammadzadeh A, Salehi TZ, Mahmoodi P. Genomic characterisation, detection of genes encoding virulence factors and evaluation of antibiotic resistance of *Trueperella pyogenes* isolated from cattle with clinical metritis. *Antonie Van Leeuwenhoek*. 2018;111(12):2441–53.
- Ishiyama D, Mizomoto T, Ueda C, Takagi N, Shimizu N, Matsuura Y, et al. Factors affecting the incidence and outcome of *Trueperella pyogenes* mastitis in cows. *J Vet Med Sci*. 2017;79(3):626–31.
- Rogovskyy AS, Lawhon S, Kuczanski K, Gillis DC, Wu J, Hurley H, et al. Phenotypic and genotypic characteristics of *Trueperella pyogenes* isolated from ruminants. *J Vet Diagn Invest*. 2018;30(3):348–53.
- Jarosz LS, Gradzki Z, Kalinowski M. *Trueperella pyogenes* infections in swine: clinical course and pathology. *Pol J Vet Sci*. 2014;17(2):395–404.
- Dong WL, Liu L, Odah KA, Atiah LA, Gao YH, Kong LC, et al. Antimicrobial resistance and presence of virulence factor genes in *Trueperella pyogenes* isolated from pig lungs with pneumonia. *Trop Anim Health Pro*. 2019;51(7): 2099–103.
- Billington SJ, Post KW, Jost BH. Isolation of *Arcanobacterium (Actinomyces) pyogenes* from cases of feline otitis externa and canine cystitis. *J Vet Diagn Invest*. 2002;14(2):159–62.
- Wareth G, El-Diasty M, Melzer F, Murugaiyan J, Abdulmawjood A, Sprague LD, et al. *Trueperella pyogenes* and *Brucella abortus* coinfection in a dog and a cat on a dairy farm in Egypt with recurrent cases of mastitis and abortion. *Vet Med Int*. 2018;2018:2056436.
- Ülbegi-Mohyla H, Hijazin M, Alber J, Lämmle C, Hassan AA, Abdulmawjood A, et al. Identification of *Arcanobacterium pyogenes* isolated by post mortem examinations of a bearded dragon and a gecko by phenotypic and genotypic properties. *J Vet Sci*. 2010;11(3):265–7.
- Wareth G, Murugaiyan J, Khater DF, Moustafa SA. Subclinical pulmonary pathogenic infection in camels slaughtered in Cairo, Egypt. *J Infect Dev Countr*. 2014;8(7):909–13.
- Tarazi YH, Al-Ani FK. An outbreak of dermatophilosis and caseous lymphadenitis mixed infection in camels (*Camelus dromedaries*) in Jordan. *J Infect Dev Countr*. 2016;10(5):506–11.

15. Çeçen G, Buyukcangaz EK, Caliskan UG, Cangul TI, Akdesir E. Interdigital necrobacillosis associated with *Trueperella pyogenes* in Goitered Gazelles (*Gazella Subgutturosa*). J Zoo Wildlife Med. 2018;49(2):429–34.
16. Cohen BS, Belser EH, Keeler SP, Yabsley MJ, Miller KV. Isolation and genotypic characterization of *Trueperella (Arcanobacterium) pyogenes* recovered from active cranial abscess Infections of Male White-Tailed Deer (*Odocoileus Virginianus*). J Zoo Wildlife Med. 2015;46(1):62–7.
17. Salleng KJ, Burton BJ, Apple TM, Sanchez S. Isolation of *Trueperella pyogenes* in a case of thoracic and abdominal abscess in a galago (*Otolemur garnettii*). J Med Primatol. 2016;45(4):198–201.
18. Eisenberg T, Nagib S, Hijazin M, Alber J, Lämmmler C, Hassan AA, et al. *Trueperella pyogenes* as cause of a facial abscess in a grey slender loris (*Loris lydekkerianus nordicus*) - a case report. Berl Münch Tierärztl. 2012;125(9–10): 407–10.
19. Nagib S, Glaeser SP, Eisenberg T, Sammra O, Lämmmler C, Kämpfer P, et al. Fatal infection in three Grey Slender Lorises (*Loris lydekkerianus nordicus*) caused by clonally related *Trueperella pyogenes*. BMC Vet Res. 2017;13:273.
20. Alssahen M, Peters M, Rau J, Hassan AA, Sammra O, Lämmmler C, et al. Phenotypic and genotypic approach to characterize a *Trueperella pyogenes* strain isolated from an Eurasian Lynx (*Lynx lynx*). Berl Münch Tierärztl. 2020. <https://doi.org/10.2376/0005-9366-19037>.
21. Hijazin M, Hassan AA, Alber J, Lämmmler C, Timke M, Kostrzewa M, et al. Evaluation of matrix-assisted laser desorption ionization-time of flight mass spectrometry (MALDI-TOF MS) for species identification of bacteria of genera *Arcanobacterium* and *Trueperella*. Vet Microbiol. 2012;157(1–2):243–5.
22. Randall LP, Lemma F, Koylass M, Rogers J, Ayling RD, Worth D, et al. Evaluation of MALDI-TOF as a method for the identification of bacteria in the veterinary diagnostic laboratory. Res Vet Sci. 2015;101:42–9.
23. Wickhorst JP, Hassan AA, Sheet OH, Eisenberg T, Sammra O, Alssahen M, et al. *Trueperella pyogenes* isolated from a brain abscess of an adult roebuck (*Capreolus capreolus*). Folia Microbiol. 2018;63(1):17–22.
24. Abdulmawjood A, Wickhorst J, Hashim O, Sammra O, Hassan AA, Alssahen M, et al. Application of a loop-mediated isothermal amplification (LAMP) assay for molecular identification of *Trueperella pyogenes* isolated from various origins. Mol Cell Probe. 2016;30(4):205–10.
25. Hassan AA, Ülbegi-Mohyla H, Kanbar T, Alber J, Lämmmler C, Abdulmawjood A, et al. Phenotypic and genotypic characterization of *Arcanobacterium haemolyticum* isolates from infections of horses. J Clin Microbiol. 2009;47(1): 124–8.
26. Moreno LZ, Matajira CEC, da Costa BLP, Ferreira TSP, Silva GFR, Dutra MC, et al. Characterization of porcine *Trueperella pyogenes* by matrix-assisted laser desorption ionization-time of flight mass spectrometry (MALDI-TOF MS), molecular typing and antimicrobial susceptibility profiling in Sao Paulo State. Comp Immunol Microb. 2017;51:49–53.
27. Hijazin M, Ülbegi-Mohyla H, Alber J, Lämmmler C, Hassan AA, Abdulmawjood A, et al. Molecular identification and further characterization of *Arcanobacterium pyogenes* isolated from bovine mastitis and from various other origins. J Dairy Sci. 2011;94(4):1813–9.
28. Wickhorst JP, Sammra O, Hassan AA, Alssashen M, Lämmmler C, Prenger-Berninghoff E, et al. Identification of *Arcanobacterium hippocoleae* by MALDI-TOF MS analysis and by various genotypical properties. Res Vet Sci. 2017;115:10–2.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

