



Bleaching efficacy and quality of life of different bleaching techniques — randomized controlled trial

Ruben Pereira¹ · João Silveira² · Susana Dias¹ · Ana Cardoso¹ · António Mata² · Duarte Marques²

Received: 21 March 2022 / Accepted: 10 August 2022

© The Author(s), under exclusive licence to Springer-Verlag GmbH Germany, part of Springer Nature 2022

Abstract

Objectives To evaluate bleaching efficacy and oral health-related quality of life (ORHQoL) of three bleaching systems with similar hydrogen peroxide (HP) concentration for up to 6 months post-treatment.

Materials and methods A randomized controlled trial was designed with three parallel groups: group A — in-office 6% HP paint-on varnish; group B — at-home 6% HP with adaptable tray; group C — at-home 16% carbamide peroxide with custom tray. At three different stages (baseline, after bleaching, and 6-month follow-up), ORHQoL was evaluated by the OHIP-14 questionnaire and tooth color of the upper canines and central incisors were measured by two shade guides and a spectrophotometer (measuring CIE L*a*b* with respective color/whiteness differences — $\Delta E_{00}/\Delta WI_D$). Results were presented as mean and 95% confidence intervals and statistical tests were performed appropriately, considering a significance level of $\alpha=0.05$.

Results All groups presented significant color differences ($P<0.05$) between all stages, with $\Delta E_{00}/\Delta WI_D$ surpassing the perceptibility threshold in 98% cases, with group C's results being significantly ($P<0.05$) higher when compared to other groups, although with significantly ($P<0.05$) higher values of color relapse. Significant ORHQoL improvements ($P<0.05$) were detected after bleaching in a global analysis with no differences between techniques.

Conclusions All techniques presented bleaching efficacy, color stability, and improvements in ORHQoL up to 6 months post-treatment.

Clinical significance Clinicians may consider both at-home and in-office bleaching techniques with 6% HP to attain long-lasting satisfactory clinical results while producing positive changes in ORHQoL.

Keywords Tooth bleaching · Tooth bleaching agents · Color · Aesthetics · Quality of life

Introduction

Patients seek care from dental professionals for preventive treatments or to address their current oral health problems. Common concerns regard the appearance and color of their teeth, as dissatisfaction with tooth color is widely reported in several adult populations, ranging from 19.6 to 65.9% [1–3]. This dissatisfaction has led to an increased desire for

treatments that improve dental aesthetics, including tooth bleaching, which is a conservative and viable option for attaining a patient's desired smile when tooth integrity is acceptable [1, 4, 5].

Tooth bleaching can be performed at home or in the dental office by a wide range of techniques [4, 6]. At-home bleaching has become increasingly popular since the introduction of the nightguard vital bleaching in 1989, which is the most prescribed technique among dentists, mainly due to its high efficacy and safety profile [5, 7–13]. Although the described protocol for at-home bleaching is the overnight use of a custom tray with a 10% carbamide peroxide (CP) gel (which requires medical prescription), nowadays, several modifications and formulations can be found among manufacturers, with application times ranging between 1 and 8 hours a day [14, 15].

As an alternative to at-home bleaching, dentists can perform in-office techniques which are viable options typically associated

✉ Duarte Marques
duarte.marques@campus.ul.pt

¹ Oral Biology and Biochemistry Research Group, Faculdade de Medicina Dentária, Universidade de Lisboa, Lisbon, Portugal

² Oral Biology and Biochemistry Research Group - LIB-Phys FCT UID/FIS/04559/2013, Faculdade de Medicina Dentária, Universidade de Lisboa, Lisbon, Portugal

with higher hydrogen peroxide (HP) concentrations. Most of the products have 35% to 40% HP and are available in the form of a base and catalyst gel, either ready-mixed or supplied as a powder/liquid combination to be freshly mixed at the dental office [16–19]. The rationale for those higher HP concentrations lies in obtaining faster results, thus being indicated for situations when immediate whitening is required [16, 20]. However, HP's oxidative properties prompted manufacturers and clinicians to search for in-office techniques with lower HP concentrations to prevent hazardous effects on biological tissues [6, 21–23]. As a result, a wide range of bleaching products with lower peroxide concentrations have been developed over the years, and even an at-home paint-on varnish technique (VivaStyle Paint On Plus, Ivoclar Vivadent, Liechtenstein) was proposed for in-office use due to its fast-bleaching rate suggested by a fast HP release in approximately 10 min [18, 24–26]. Although evidence shows promising efficacy results (tooth color change), many issues have not yet been addressed as no studies have compared this proposed technique with at-home techniques [27, 28]. Concomitantly, studies lack evidence regarding tooth color stability, which is a great concern since tooth color relapses are common issues that may require touch-up bleaching [4, 29–33].

The quantitative analysis of tooth color changes and stability is important to evaluate the efficacy/effectiveness of a bleaching technique; however, patient-reported outcomes are also major aspects of a successful treatment and can be characterized by changes in oral health-related quality of life (OHRQoL) [34–37]. Currently, tooth bleaching is known to potentially influence OHRQoL by affecting the patient's self-esteem and social behaviors, such as smiling, laughing, or showing teeth without embarrassment [34, 35]. Therefore, the long-term effects of tooth bleaching are not only related to tooth color stability but may also impact the patient's everyday life.

This study aimed to compare the bleaching efficacy and OHRQoL of three different bleaching systems with a similar HP concentration of 6% or its CP equivalent while assessing the outcomes for up to 6 months. The following null hypotheses were established: (1) there were no differences in bleaching efficacy between the three tested bleaching systems; (2) there were no differences in tooth color stability, at the 6-month follow-up, between the three tested bleaching systems; (3) there were no differences in OHRQoL, at the end of treatment, between the three tested bleaching systems; (4) there were no differences in OHRQoL, at the 6-month follow-up, between the three tested bleaching systems.

Materials and methods

This randomized clinical trial took place between November 2019 and October 2021 at the Faculty of Dental Medicine of the University of Lisbon and was conducted in full compliance

with the Helsinki World Medical Association Declaration's most recent amendments [38]. Additionally, the local ethics committee gave ethical approval, and the trial was registered at the U.S. National Library of Medicine ClinicalTrials.gov website under the reference number NCT03588871.

Study design and participants

A randomized clinical trial was designed with three parallel groups corresponding to different products and techniques: group A, in-office paint-on varnish 6% HP (VivaStyle Paint On Plus, Ivoclar Vivadent, Liechtenstein); group B, at-home 6% HP with a prefilled disposable tray (Opalescence GO, Ultradent, EUA); group C, at-home 16% CP with a customized tray (Opalescence PF 16% CP, Ultradent, EUA).

Participants attending the faculty clinic were screened according to the following inclusion criteria and consecutively recruited: being at least 18 years of age, having the upper canines darker than A3.5 in VITA Classical (VC) shade guide (assessed by spectrophotometry), accepting to interrupt smoking habits during the full duration of the study, and signing an informed consent form. The exclusion criteria were the presence of fixed orthodontic appliances, decayed teeth, pregnancy, poor oral hygiene, anterior teeth (16 anterior teeth, from the second premolar to the second premolar) with dental restorations, endodontic treatment, and severe anomalies of the dental structure or intrinsic stain. A flowchart of the study is summarized in Fig. 1.


Randomization process and blinding

Each bleaching system was coded from A to C using a randomization software (GraphPad QuickCals, <http://www.graphpad.com/quickcalcs/randomize1.cfm>), and the information was held by external personnel until the end of the study. A third party (blinded to the allocation results) analyzed the data in an SPSS worksheet (IBM Statistics, Inc., Chicago, IL, USA) where each bleaching system was referred to as groups A to C.

Participant and clinical operator blinding was not possible due to the three whitening systems' different formulations. However, the tooth color examiners were blinded, and spectrophotometric analysis is not susceptible to interpretation, thus reducing the potential bias.

Calibration of examiners for clinical analysis

Visual shade selection was performed by dentists with at least 5-year clinical experience and negative history of visual color deficiencies (confirmed using X-Rite Color Challenge by Pantone®) who were submitted to a calibration process. This

Fig. 1 Flowchart of the study, according to CONSORT

process was based on a consecutive determination of VITA Classical (VC) visual shade guides using two VC scales (one of which had a blinded identification; VITA Zahnfabrick, Germany). The dentist would be considered a valid operator with an intraclass correlation coefficient (ICC; model: two-way random; type: absolute agreement) higher than 0.80 (considered excellent agreement [39]). The same process was repeated for the VITA Bleachedguide 3D-Master shade guide (VB) (VITA Zahnfabrick, Germany). The calibrated operators' ICC ranged between 0.86 and 0.93. During the study, if disagreements occurred, the examiners reached a consensus. To standardize lighting conditions, the Smile Lite device (Smile Line AS, Switzerland; serial number 052015) with LED lights at 5500 K and a polarization filter was used.

An independent and blinded examiner performed objective tooth color measurements with a spectrophotometer, SpectroShade micro (SS) (MHT Optic Research, Niederhasli, Switzerland; serial number HDL3973), which is considered a diagnostic device for tooth color assessment [40–43]. The SS intradevice's calibration process was performed before each measuring round according to the manufacturer's instructions.

Oral health-related quality of life evaluation

The validated Portuguese version of the Oral Health Impact Profile 14 (OHIP-14) was applied at baseline, at the end of treatment (after bleaching), and after 6 months (6-month

follow-up) [44]. The questionnaire consisted of 14 questions with seven domains (2 questions per domain): functional limitation, physical pain, psychological discomfort, physical disability, psychological disability, social disability, and handicap. The answers were scored according to a Likert scale [45] from 0 to 4 (never = 0, rarely = 1, sometimes = 2, repeatedly = 3, always = 4), with higher scores representing a worse OHRQoL (OHIP-14 total score ranged from 0 to 56 and each domain score from 0 to 8). Effect size (ES; calculated by Kendall's W) and standardized response mean (SRM; calculated by dividing the mean score change by the standard deviation of the change) were calculated as previously recommended for health questionnaires (ES and SRM were described as small < 0.3, moderate 0.3–0.8, or large \geq 0.8 effect) [46–48]. A minimal important difference (MID) of five in the total OHIP-14 score change was also considered [46, 49].

The OHIP-14 was applied twice to each participant with a 1-week interval before bleaching treatment to evaluate the questionnaire's reliability and internal consistency. Test–retest reliability was evaluated for each question by ICC (model: two-way random; type: absolute agreement) with values ranging from 0.51 (moderate agreement) to 0.74 (substantial agreement) [39]. Internal consistency was evaluated by Cronbach's alpha with an obtained value of 0.81, which is desirable and considered good consistency [50].

Clinical procedures and measurements

In the first appointment, each patient was screened according to the previously described inclusion/exclusion criteria and submitted to professional dental prophylaxis with interproximal radiographs for diagnosis purposes. The professional dental prophylaxis was performed using an ultrasonic scaler and a nylon brush with prophylaxis paste (Cleanic, Kerr Orange, USA) in a low-rotation contra-angle handpiece by a dentist. Each patient was assigned to one group, according to the randomization process. One week after, the clinical bleaching protocol was performed according to the technique's description (Table 1) [4, 24, 27, 51].

To assess tooth sensitivity that could lead to treatment interruption, all patients were instructed to fill a daily visual analogic scale (VAS) form during the treatment (15 days),

numbered from 0 (no pain) to 10 (maximum extreme pain), while notifying medication intake and oral lesions occurrences. Additionally, instruction forms were delivered with information regarding at-home bleaching procedures, food intake (to avoid acidic and potential staining foods), and oral hygiene. Patients were instructed to use their regular toothpaste during the whole study to avoid any potential change in tooth sensitivity unless it was a whitening toothpaste, in which case they were instructed to change to a non-whitening 1450-ppm fluoride-containing toothpaste.

Tooth color measurements were performed at baseline, after bleaching treatment, and at the 6-month follow-up. The color of the upper central incisors and canines' buccal surfaces was assessed with the VC and VB shade guides with the patient seated in the high Fowler's position on the dental chair while the calibrated examiner used the Smile Lite device with LED lights at 5500 K and a polarization filter for standard lighting conditions. The shade tabs received a number to categorize each color: VC tabs were numbered from 1 to 16 according to the color's value order from the highest (B1) to lowest (C4), and VB tabs were also numbered according to the color's value order from 1 to 15 (highest: 0M1; lowest: 5M3). An interpolated guide corresponding to the American Dental Association's Eq. (1 $ccu = 1$; $SGU = 1 \Delta E$; ccu , color difference unit; SGU , shade guide unit; ΔE , overall color difference) was used to express results in shade guide units, and the differences were expressed by ΔSGU (SGU_{VC} and ΔSGU_{VC} for the VC shade guide; SGU_{VB} and ΔSGU_{VB} for the VB shade guide) [36, 52]. The SS performed three measuring replicates to obtain the CIE $L^*a^*b^*$ values of the upper central incisors and canines' buccal surfaces, following the manufacturer's instructions. The total tooth color difference (ΔE_{00}) and tooth whiteness index (WI_D), with the corresponding difference (ΔWI_D), both based on the CIE $L^*a^*b^*$ color notation system, were calculated to evaluate bleaching efficacy at the end of treatment and color relapse at the 6-month follow-up [53–56].

Sample size calculation

The sample size was calculated based on the upper canines' color difference (ΔE_{00} — a primary outcome) recorded in a previously performed pilot study with 30 randomized

Table 1 Simplified clinical protocol for each bleaching product. *HP*, hydrogen peroxide; *CP*, carbamide peroxide

Bleaching product	Clinical protocol
Group A VivaStyle Paint On Plus 6% HP	In-office applications: 2 sessions with 6 applications of 10 min with a 1-week interval (2 h application time)
Group B Opalescence GO 6% HP	At-home applications: 1 daily application of 90 min for 10 days (15 h application time)
Group C Opalescence PF 16% CP	At-home applications: 1 daily application of 6 h for 14 days (84 h application time)

participants (GraphPad QuickCals, <http://www.graphpad.com/quickcals/randomize1.cfm>), using the G*Power 3.1 software (Heinrich-Heine-Universität, Düsseldorf, Germany). The size effect was calculated based on the perceptibility threshold $\Delta E_{00} = 0.8$ with a standard deviation of 0.81 [57, 58]. Considering an *F* test (one-way ANOVA) with a significance level of 5% and a power of 80%, a minimum of 20 participants per group was required. To offset a possible attrition bias, 50% was added to each group, resulting in 30 patients' samples (a total of 90 patients).

Statistical analysis

All collected data were analyzed using IBM SPSS version 25 (IBM Statistics, Inc. Chicago, IL, USA). Results are presented as mean and 95% confidence interval (IC) of CIE L*a*b* color parameters, SGU, and OHIP-14 scores, with the respective WI_D , ΔE_{00} , ΔWI_D , and ΔSGU calculated. The CIEDE2000 formula from the Commission Internationale De l'Eclairage (CIE, International Commission on Illumination) was used to calculate ΔE_{00} . Computations with this color difference formula were performed according to the following equation [56]:

$$\Delta E_{00} = \sqrt{\left(\frac{L_2 - L_1}{K_L S_L}\right)^2 + \left(\frac{C_2 - C_1}{K_C S_C}\right)^2 + \left(\frac{H_2 - H_1}{K_H S_H}\right)^2 + R_T \left(\frac{C_2 - C_1}{K_C S_C}\right) \left(\frac{H_2 - H_1}{K_H S_H}\right)}$$

The variables of the CIEDE2000 formula were calculated from the CIE L*a*b* values using a free online code while setting the parametric factors to 1 (https://www.rit.edu/cos/colorscience/rc_useful_data.php).

The whiteness index was calculated before and after tooth bleaching with the following formula: $WI_D = 0.511L^* - 2.324a^* - 1.100b^*$ [54]. Color and whiteness difference perception was assessed according to two major thresholds: perceptibility threshold (PT for ΔE_{00} ; WPT for ΔWI_D) considered $\Delta E_{00} = 0.8$ and $\Delta WI_D = 0.72$; acceptability threshold (AT for ΔE_{00} ; WAT for ΔWI_D) considered $\Delta E_{00} = 1.8$ and $\Delta WI_D = 2.60$ [55, 58]. The percentages of cases in which ΔE_{00} and ΔWI_D were higher than the respective thresholds were calculated, and the bleaching efficacy was considered when both perceptibility thresholds were surpassed. When comparing post-treatment results with the 6-month follow-up, surpassing of both perceptibility thresholds was considered a tooth color relapse. The surpassing of both acceptability thresholds at the 6-month follow-up compared to the baseline values was considered undoubted color differences with no requirement for touch-up bleaching.

Since darker-colored teeth may have a different response to tooth bleaching, the upper central incisors and canines were analyzed in individual groups. Statistical analysis was performed with parametric tests whenever the minimal

sample of 30 was attained according to the central limit theorem [59]. The variables without a minimal sample of 30 were evaluated regarding their distribution with the Kolmogorov–Smirnov test, and non-parametric tests were used to analyze the OHIP-14 score (results were also presented with median values). Repeated-measures analysis of variance (ANOVA) with Tukey post hoc test was performed to analyze intragroup differences at different times (baseline, after bleaching, and 6-month follow-up) in CIE L*a*b*, WI_D , and SGU, while the one-way ANOVA with Tukey post hoc test analyzed intergroup differences in ΔE_{00} , ΔWI_D , and ΔSGU . Differences in the OHIP-14 score among different times were determined by the Friedman test, while the Kruskal–Wallis test was adopted for multiple group comparison. In all statistical tests, the level of significance considered was 5%.

Results

Ninety participants were included in the study after the recruitment procedures: 56 females and 24 males, aged between 18 and 40 years old with a mean of 23.0 [22.8:23.4] years. A total of 80 bleaching treatments were completed (group A: 27; group B: 26; group C: 27) with an overall 11.1% attrition bias due to COVID-19 quarantine measures, leading to an overall 32.2% attrition bias at the 6-month follow-up (group A: 20; group B: 20; group C: 21) (Fig. 1). Baseline CIE L*a*b*, WI_D , and SGU values are depicted in Table 2 and did not show significant ($P > 0.05$) differences between groups, resulting in tooth-color and whiteness homogeneity before bleaching treatment.

Bleaching efficacy analysis (depicted in Table 3) detected that the perceptibility thresholds (PT and WPT) in all techniques were surpassed in at least 98% of cases and attained 100% in the upper canines (98% for acceptability thresholds). Thus, all techniques showed bleaching efficacy even though the $\Delta E_{00}/\Delta WI_D/\Delta SGU$ were significantly higher ($P < 0.05$) in group C after bleaching. Concomitantly, the ANOVA also detected significant differences ($P < 0.05$) in all groups' CIE L*a*b*, WI_D , and SGU (both VC and VB) values after bleaching (depicted in Table 2). The L* color coordinate presented significantly ($P < 0.05$) higher mean values while a* and b* were lower when compared to baseline, indicating a lighter and less yellow tooth color post-treatment. The WI_D mean values were significantly ($P < 0.05$) higher after bleaching in all groups, thus indicating increased levels of whiteness in tooth color. The SGU_{VC} and SGU_{VB} mean values were significantly ($P < 0.05$) lower after bleaching, indicating that the examiners detected higher value color tabs.

At the 6-month follow-up, an inverse response was detected in all variables, with values becoming closer to

Table 2 Mean and 95% IC values for CIE L*a*b*, WI_D, SGU_{VC}, and SGU_{VB} at different times. In all groups, the repeated measures ANOVA with Tukey post hoc presented statistically significant ($P < 0.05$) intragroup differences between baseline, after bleaching, and 6-month follow-up evaluations

		Baseline 80 cases			After bleaching 80 cases			6-month follow-up 61 cases		
		Group A 27 cases n=54	Group B 26 cases n=52	Group C 27 cases n=54	Group A 27 cases n=54	Group B 26 cases n=52	Group C 27 cases n=54	Group A 20 cases n=40	Group B 20 cases n=40	Group C 21 cases n=42
L*	Canines	70.6 [70.1:71.1]	70.0 [69.6:70.3]	70.0 [69.1:70.2]	74.3 [73.8:74.7]	74.1 [73.6:74.6]	76.3 [75.6:77.0]	73.6 [73.1:74.2]	73.9 [73.4:74.4]	75.6 [75.0:76.2]
	Incisors	75.7 [75.3:76.1]	75.7 [75.3:76.2]	75.3 [74.7:75.8]	77.8 [77.4:78.2]	77.4 [77.0:77.9]	78.4 [77.8:78.9]	77.0 [76.4:77.6]	77.4 [77.0:77.8]	78.1 [77.5:78.7]
a*	Canines	5.1 [4.9:5.3]	5.4 [5.2:5.5]	5.3 [5.1:5.6]	3.6 [3.4:3.8]	3.3 [3.1:3.6]	2.4 [2.1:2.6]	3.5 [3.3:3.7]	3.5 [3.2:3.7]	2.5 [2.3:2.7]
	Incisors	2.1 [2.0:2.3]	2.2 [2.0:2.3]	2.3 [2.0:2.5]	1.6 [1.5:1.7]	1.5 [1.3:1.6]	1.3 [1.2:1.5]	1.6 [1.4:1.7]	1.5 [1.4:1.6]	1.2 [1.0:1.3]
b*	Canines	24.1 [23.5:24.6]	24.0 [23.5:24.5]	23.9 [23.2:24.6]	19.3 [18.7:19.8]	19.7 [18.9:20.4]	15.0 [14.3:15.8]	20.2 [19.6:20.8]	20.2 [19.5:20.9]	16.4 [15.8:16.9]
	Incisors	17.6 [17.1:18.2]	17.0 [16.4:17.7]	17.2 [16.6:17.9]	14.7 [14.1:15.1]	14.2 [13.5:14.8]	12.0 [11.4:12.6]	20.2 [19.6:20.8]	20.2 [19.5:20.9]	16.4 [15.8:16.9]
WI _D	Canines	-2.3 [-3.4:-1.2]	-3.1 [-3.9:-2.2]	-2.5 [-4.3:-0.6]	8.4 [7.4:9.5]	9.0 [7.8:10.2]	17.8 [16.7:18.9]	7.2 [6.1:8.3]	7.4 [6.1:8.8]	14.7 [13.6:15.8]
	Incisors	14.3 [13.3:15.3]	14.9 [13.7:16.1]	14.2 [12.8:15.6]	19.9 [19.2:20.6]	20.8 [19.0:21.8]	24.3 [23.4:25.1]	17.5 [15.9:19.1]	19.7 [18.7:20.8]	22.7 [21.8:23.5]
SGU _{VC}	Canines	12.1 [11.9:12.4]	11.9 [11.5:12.3]	12.0 [11.7:12.2]	4.5 [4.1:5.1]	5.1 [4.8:5.6]	2.9 [2.6:3.3]	7.2 [6.1:8.3]	7.4 [6.1:8.8]	14.7 [13.6:15.8]
	Incisors	4.9 [4.4:5.4]	4.2 [3.6:4.8]	4.1 [3.7:4.5]	1.3 [1.1:1.4]	1.6 [1.4:1.9]	1.0 [1.0:1.8]	17.5 [15.9:19.1]	19.7 [18.7:20.8]	22.7 [21.8:23.5]
SGU _{VB}	Canines	10.9 [10.7:11.1]	10.9 [10.6:11.1]	10.7 [10.5:10.9]	6.8 [6.5:7.1]	6.7 [6.3:7.1]	4.8 [4.5:5.1]	7.2 [6.1:8.3]	7.4 [6.1:8.8]	14.7 [13.6:15.8]
	Incisors	6.8 [6.5:7.1]	6.2 [5.7:6.7]	6.5 [6.2:6.8]	3.7 [3.5:3.9]	3.6 [3.3:3.9]	2.8 [2.5:3.0]	17.5 [15.9:19.1]	19.7 [18.7:20.8]	22.7 [21.8:23.5]

the respective baseline. However, significant differences ($P < 0.05$) were still detected in $\Delta E_{00}/\Delta WI_D/\Delta SGU$ (Table 3) and CIE L*a*b*, WI_D, and SGU (Table 2), with a maximum of 16.7% of cases needing touch-up bleaching (evaluated by the acceptability thresholds). Thus, all techniques showed color stability even though tooth color relapse cases were higher in group C (83.3% cases) with significantly superior ($P < 0.05$) $\Delta E_{00}/\Delta WI_D$ values compared to after bleaching.

There were no reports of treatment interruption due to tooth hypersensitivity or presence of oral lesions, with the following overall VAS mean values: 1.0 [0.6:1.5] in group A, 1.2 [0.7:1.7] in group B, and 1.6 [0.9:2.2] in group C, without significant ($P > 0.05$) differences in the ANOVA. Oral lesions were reported in eight out of 27 cases in group A, four out of 26 cases in group B, and four out of 27 cases in group C, with the following occasional intakes of paracetamol or ibuprofen: three cases in group A, two in group B, and two in group C.

There was a noticeable improvement in OHRQoL after tooth bleaching, represented by significantly lower ($P < 0.05$) OHIP-14 total score values when all treatments were considered (global analysis — Table 4), with an ES of 0.1 and

an SRM of 0.4 (low to moderate effect). That improvement compared to baseline was maintained up to 6 months, with an ES of 0.1 and an SRM of 0.2 (low effect). However, no significant differences ($P > 0.05$) in OHIP-14 scores were detected within or between groups, indicating that changes in OHRQoL are not related to the bleaching technique.

The percentage of cases in which the OHIP-14 total score difference was attained or surpassed the MID value of five was 18.8% after post-treatment and 13.8% at the 6-month follow-up compared to baseline values. Additionally in Table 4, are presented the scores for the seven domains of the OHIP-14 individually which assessment did not reveal any significant results ($P > 0.05$) between techniques or stages.

Discussion

The tested techniques presented bleaching efficacy with mean $\Delta E_{00}/\Delta WI_D$ above the respective AT and WAT values of 1.8 and 2.6, while the PT of 0.8 and the WPT of 0.72 were surpassed in at least 98% of treatments. Group C

Table 3 Mean and 95% IC values for ΔE_{00} , ΔWI_D , ΔSGU_{VC} , and ΔSGU_{VB} at different times with intergroup analysis. Also presented the percentages of cases, for each group, in which $\Delta E_{00}/\Delta WI_D$ sur-

passed the respective perceptibility (PT/WPT) and acceptability (WAT) thresholds. *Statistically significant difference ($P < 0.05$) by one-way ANOVA with Tukey post hoc

		Baseline — after bleaching 80 cases			After bleaching — 6-month follow-up 80 cases			Baseline — 6-month follow-up 61 cases		
		Group A 27 cases <i>n</i> = 54	Group B 26 cases <i>n</i> = 52	Group C 27 cases <i>n</i> = 54	Group A 20 cases <i>n</i> = 40	Group B 20 cases <i>n</i> = 40	Group C 21 cases <i>n</i> = 42	Group A 20 cases <i>n</i> = 40	Group B 20 cases <i>n</i> = 40	Group C 21 cases <i>n</i> = 42
ΔE_{00}	Canines	4.0 [3.7:4.3]	4.3 [4.0:4.7]	7.7* [7.0:8.3]	0.9 [0.8:1.1]	1.2 [0.7:1.7]	2.1* [1.7:2.4]	3.7 [3.4:4.0]	4.0 [3.7:4.3]	6.0* [5.3:6.7]
	Incisors	2.6 [2.3:2.8]	2.4 [2.1:2.7]	4.4* [4.0:4.8]	1.1 [0.9:1.3]	1.0 [0.7:1.2]	1.7* [1.4:2.1]	1.9 [1.6:2.2]	2.1 [1.8:2.4]	3.2* [2.7:3.7]
ΔWI_D	Canines	11.0 [10.0:11.0]	11.8 [10.7:12.9]	21.1* [19.6:23.4]	1.3 [1.0:1.6]	1.7 [1.1:2.2]	4.5* [3.6:5.3]	10.8 [9.9:11.6]	10.8 [9.8:11.8]	18.5* [16.5:20.4]
	Incisors	5.8 [4.9:6.6]	5.7 [4.9:6.4]	10.7* [9.5:12.0]	1.4 [1.1:1.8]	1.3 [0.9:1.7]	2.5* [1.9:3.1]	7.0 [5.8:8.3]	5.1* [4.3:5.9]	8.5 [7.3:10.2]
ΔSGU_{VC}	Canines	7.5 [6.9:8.0]	6.7 [6.2:7.1]	9.1* [8.6:9.5]	2.7 [1.9:3.5]	1.6 [1.0:2.3]	2.5 [1.9:3.1]	5.1 [4.4:5.9]	5.1 [4.4:5.8]	7.4* [6.7:8.2]
	Incisors	3.6 [3.1:4.1]	2.6* [2.1:3.1]	3.1 [2.7:3.4]	0.9* [0.7:1.1]	0.3 [0.1:0.6]	0.4 [0.2:0.5]	2.8 [2.0:3.6]	2.3 [1.7:2.9]	2.7 [2.3:3.0]
ΔSGU_{VB}	Canines	4.0 [3.7:4.4]	4.2 [3.8:4.6]	5.9* [5.6:6.3]	1.5 [1.2:1.9]	1.1* [0.8:1.3]	2.0 [1.7:2.4]	2.7 [2.3:3.1]	3.4 [2.9:3.9]	4.2* [3.7:4.7]
	Incisors	3.1 [2.8:3.4]	2.6 [2.2:3.0]	3.7* [3.4:4.0]	1.4 [1.1:1.6]	0.9 [0.7:1.2]	1.0 [0.8:1.3]	1.9 [1.5:2.2]	2.1 [1.7:2.5]	2.7* [2.4:3.0]
% cases $\Delta E_{00} > PT$	Canines	100	100	100	49.5	47.2	90.5	100	100	90.5
	Incisors	98.1	98.1	100	48.4	42.8	83.3	95.2	97.2	92.9
% cases $\Delta E_{00} > AT$	Canines	98.1	98.1	100	2.6	11.1	47.6	97.4	97.2	90.5
	Incisors	72.2	75.0	96.3	2.6	8.3	35.7	88.1	88.1	83.3
% cases $\Delta WI_D > WPT$	Canines	100	100	100	67.8	69.4	95.2	100	100	100
	Incisors	98.1	98.1	100	68.4	69.4	88.1	100	100	100
% cases $\Delta WI_D > WAT$	Canines	98.1	98.1	100	13.2	25.0	71.4	100	100	95.2
	Incisors	81.5	94.2	100	15.8	13.9	40.5	86.8	88.8	92.9

presented the higher bleaching efficacy and the highest tooth color relapse at the 6-month follow-up, thus rejecting both first and second null hypotheses. Additionally, tooth bleaching with 6% HP or its CP equivalent significantly improved patients' OHRQoL, although without detectable differences between techniques, thus accepting the third and fourth null hypothesis.

To the authors' knowledge, this is the first randomized controlled trial comparing multiple different bleaching systems (both in-office and at-home) that evaluates efficacy (tooth color change) and long-term outcomes (tooth color relapse) by objective methods while evaluating patient-reported outcomes (effects in OHRQoL). Visual shade analysis was also performed since shade guides are a commonly employed method in clinical practice; however, the subjectivity of its assessment reduces accuracy and reliability, requiring the supporting use of an objective method [60].

By testing systems with similar HP concentration, the authors intended to pragmatically assess different factors on treatment outcomes, such as the application time, the

delivery method, or the necessary patient's compliance. In fact, the results suggest that efficacy outcomes and treatment time efficiency depend on the clinical protocol since the group C, which presented the highest $\Delta E_{00}/\Delta WI_D/\Delta SGU$, requires approximately 30 to 40 h to attain the same bleaching effect as 2 h treatment in the group A. When applied as an in-office technique, this varnish can attain or even surpass tooth color/whitening acceptability thresholds in just 1 h, probably due to the proper soft-tissue isolation that reduces contact with crevicular fluids or the continuous evaporation of the varnish's solvent leading to a potential HP concentration increase in the tooth surface [25]. Therefore, the VivaStyle Paint On Plus (group A) has a higher efficacy in a shorter time than at-home techniques with the same HP concentration. However, further studies are required to evaluate if the product's full performance is attained with the current protocol, as increasing the number of in-office applications could potentially increase $\Delta E_{00}/\Delta WI_D/\Delta SGU$ values. A previous meta-analysis highlighted this problem stating that

Table 4 Mean, median, and 95% IC values for OHIP-14 total score and domain score at different times, divided by global and group analysis. No significant differences ($P > 0.05$) were detected between groups with Kruskal–Wallis test. *Statistically significant difference ($P < 0.05$) by Friedman test

	Baseline 80 cases $n = 80$			After bleaching 80 cases $n = 80$			6-month follow-up 61 cases $n = 61$		
OHIP-14 Total score	Mean 2.8* Median 1.0 [1.9:3.8]			Mean 1.7 Median 0 [1.0:2.1]			Mean 2.1 Median 0 [1.3:3.0]		
	Group A 27 cases $n = 27$	Group B 26 cases $n = 26$	Group C 27 cases $n = 27$	Group A 27 cases $n = 27$	Group B 26 cases $n = 26$	Group C 27 cases $n = 27$	Group A 20 cases $n = 20$	Group B 20 cases $n = 20$	Group C 21 cases $n = 21$
	Mean 3.0 Median 2.0 [1.2:5.1]	Mean 2.4 Median 1.0 [0.9:3.5]	Mean 3.0 Median 1.0 [1.0:5.4]	Mean 1.7 Median 1.0 [1.0:3.3]	Mean 1.1 Median 0 [0.2:1.9]	Mean 1.7 Median 0 [0.3:3.1]	Mean 2.9 Median 1.0 [0.:5.1]	Mean 1.6 Median 0 [0.4:2.8]	Mean 2.0 Median 0 [0.4:3.5]
OHIP-14 functional limitation score	Mean 0.2 Median 0 [0:0.6]	Mean 0.1 Median 0 [0:0.2]	Mean 0.1 Median 0 [0:0.3]	Mean 0.4 Median 0 [0.1:0.7]	Mean 0.1 Median 0 [0:0.2]	Mean 0.1 Median 0 [0:0.3]	Mean 0.1 Median 0 [0:0.3]	Mean 0.1 Median 0 [0:0.2]	Mean 0.1 Median 0 [0:0.2]
OHIP-14 physical pain score	Mean 0.7 Median 0 [0.2:1.1]	Mean 0.6 Median 0 [0.3:0.9]	Mean 0.7 Median 0 [0.1:1.2]	Mean 0.7 Median 0 [0.3:1.1]	Mean 0.3 Median 0 [0.0:0.6]	Mean 0.3 Median 0 [0:0.6]	Mean 0.6 Median 0 [0.2:1.0]	Mean 0.4 Median 0 [0:0.9]	Mean 0.1 Median 0 [0:0.3]
OHIP-14 psychological discomfort score	Mean 1.3 Median 0 [0.5:2.1]	Mean 0.8 Median 0 [0.1:1.2]	Mean 1.2 Median 0 [0.4:2.0]	Mean 0.9 Median 0 [0.3:1.5]	Mean 0.4 Median 0 [0.1:0.8]	Mean 0.6 Median 0 [0.1:1.2]	Mean 1.3 Median 0 [0.4:2.0]	Mean 0.8 Median 0 [0.1:1.3]	Mean 1.0 Median 0 [0.2:1.7]
OHIP-14 physical disability score	Mean 0.3 Median 0 [0:0.7]	Mean 0.2 Median 0 [0:0.4]	Mean 0.3 Median 0 [0:0.6]	Mean 0.2 Median 0 [0:0.5]	Mean 0.1 Median 0 [0:0.2]	Mean 0.1 Median 0 [0:0.3]	Mean 0.1 Median 0 [0:0.3]	Mean 0.1 Median 0 [0:0.2]	Mean 0.2 Median 0 [0:0.5]
OHIP-14 psychological disability score	Mean 0.6 Median 0 [0.3:1.0]	Mean 0.4 Median 0 [0.1:0.6]	Mean 0.4 Median 0 [0.0:0.7]	Mean 0.2 Median 0 [0:0.4]	Mean 0.2 Median 0 [0:0.4]	Mean 0.3 Median 0 [0:0.5]	Mean 0.5 Median 0 [0.1:0.9]	Mean 0.2 Median 0 [0:0.4]	Mean 0.5 Median 0 [0.1:0.9]
OHIP-14 social disability score	Mean 0.4 Median 0 [0.0:0.8]	Mean 0.1 Median 1.0 [0:0.1]	Mean 0.4 Median 0 [0.0:0.8]	Mean 0.3 Median 0 [0.1:0.5]	Mean 0.1 Median 0 [0:0.3]	Mean 0.2 Median 0 [0:0.4]	Mean 0.6 Median 0 [0.2:1.0]	Mean 0.2 Median 0 [0.0:0.5]	Mean 0.3 Median 0 [0.0:0.6]
OHIP-14 handicap score	Mean 0.1 Median 0 [0:0.2]	Mean 0.1 Median 0 [0:0.1]	Mean 0.1 Median 0 [0:0.1]	Mean 0.1 Median 0 [0:0.2]	Mean 0.1 Median 0 [0:0.1]	Mean 0.1 Median 0 [0:0.1]	Mean 0.2 Median 0 [0:0.3]	Mean 0.1 Median 0 [0:0.1]	Mean 0.1 Median 0 [0:0.1]

the application time or the number of applications may be inadequate for a full HP release in some bleaching products, thus undervaluing their efficacy/effectiveness [61].

Because patients' concerns are not exclusively related to the treatment result but also the long-term outcomes, it is important to follow up on every possible case. All tested systems presented color stability up to 6 months, regardless of the percentage of tooth color relapses, since most differences were still superior to the AT thresholds compared to baseline. Additionally, only a small percentage of cases needed touch-up (between 11.9 and 16.7%), in agreement with previous studies that tested similar concentration products [9, 29–33, 36]. However, color stability values differed among techniques, and the bleaching system that attained a higher tooth color change was the same with a higher tooth color relapse — Opalescence PF (group C). This higher tooth color relapse may explain the similar whiteness results in the upper central incisors at the 6-month follow-up between groups A and C (considering group A's lower tooth color relapse values).

Patients' reports revealed improvements in OHRQoL after bleaching treatment that were maintained up to 6 months, even though that effect seems to decrease over time, as suggested by lower SRM values. This finding supports the idea that whiter and lighter tooth color may be related to patient satisfaction, consequently impacting OHRQoL, since tooth color is a major factor of an aesthetic smile and can have a psychological influence by positively changing a patient's self-esteem and social behaviors [2, 3, 34]. However, a meta-analysis concluded that the impact of tooth bleaching improvements on OHRQoL is hardly detected clinically, especially in heterogeneous populations [34]. Accordingly, our results suggest that only a small percentage of cases attained the OHIP-14 MID value, indicating that the improvements in overall OHRQoL may not be clinically relevant. Nevertheless, this could be related to the fact that the OHIP-14 MID value was established for general oral treatments and not specifically for tooth bleaching, thus not providing a reliable threshold [46].

Our results also suggest that the OHRQoL improvement is related to the bleaching treatment itself and not to any specific

technique, as significant differences in the OHIP-14 total score were detected when evaluating all cases globally. When evaluating each tested system individually, a similar effect on OHRQoL was detected despite minor adverse effects mostly related to mild and transient tooth sensitivity and gingival irritation, which were reported during the whole treatment in at-home techniques (constant low level of tooth sensitivity) and on treatment days in the in-office technique. These higher tooth sensitivity levels and some transitory oral lesion occurrences, as reported in other studies, were expected for the in-office technique due to the lower viscosity and faster HP release of the varnish, increasing the difficulty of soft tissues' isolation and tooth sensitivity during application [24, 27, 28].

As in most tooth bleaching clinical studies, limitations include participants being mostly between 20 and 30 years of age, probably due to the inclusion/exclusion criteria requiring anterior teeth free of decays/restorations and the higher demand for aesthetic treatments by younger individuals [62]. Although evidence suggests a significant relationship between the subject's age and the magnitude of the whitening response (younger subjects experience greater effects), in older populations, tooth bleaching could be effective considering the reported positive correlation between yellow hues and bleaching effects [63].

The generality of this study's results may be applied to clinical practice, although a patient-centered approach should always be considered. An in-office technique, such as the VivaStyle Paint On Plus (group A), would be suitable for faster treatments or low compliance patients, while at-home techniques could be a treatment option when it is not suitable to perform several clinical bleaching sessions. Additionally, the choice between wearing a custom (Opalescence PF — group C) or an adaptable/disposable tray (Opalescence GO — group B) could be made based on the inability to be submitted to dental impressions (e.g., vomit reflex) or difficulties in positioning the adaptable tray causing frequent displacements, as detected in our study patients' reports. Since the tested systems have their own advantages, a combined approach, where the treatment is performed in the dental office along with an at-home protocol, could be proposed to achieve a balance between tooth color change, time efficiency, and color stability. Therefore, further studies should address assistant bleaching protocols to evaluate the treatment potential of the tested bleaching systems in a combine approach.

Conclusions

All techniques presented bleaching efficacy and color stability up to the 6-month follow-up, even though a small percentage of color relapses must be expected. Tooth bleaching produces positive changes in OHRQoL, with low to moderate effects that are still detectable at a 6-month follow-up; however, these improvements are not associated with any of the tested systems.

Acknowledgements The authors thank the members of Oral Biology and Biochemistry Research Group for their support and valuable help in conducting this research. The authors also thank the Faculty of Dental Medicine of the University of Lisbon for supporting this research providing the facilities and resources to conduct the study.

Author contribution Conceptualization: Ruben Pereira; methodology: Duarte Marques; formal analysis and investigation: Ruben Pereira, João Silveira, Susana Dias, Ana Cardoso; writing — original draft preparation: Ruben Pereira; writing — review and editing: Duarte Marques; funding acquisition: António Mata; resources: Ana Cardoso; supervision: António Mata, Duarte Marques.

Funding The main funding entity was the Faculty of Dental Medicine of the University of Lisbon who financially supported this study while providing the facilities and resources.

Declarations

Ethics approval This study was performed in line with the principles of the Declaration of Helsinki. Approval was granted by the Ethics Committee of Faculty of Dental Medicine of the University of Lisbon.

Consent to participate and consent for publication It was obtained from all individual participants included in the study an informed consent regarding study procedures/information and publishing data.

Conflict of interest The authors declare no competing interests.

References

1. Joiner A, Luo W (2017) Tooth colour and whiteness: a review. *J Dent* 67S:S3–S10. <https://doi.org/10.1016/j.jdent.2017.09.006>
2. Tin-Oo MM, Saddki N, Hassan N (2011) Factors influencing patient satisfaction with dental appearance and treatments they desire to improve aesthetics. *BMC Oral Health* 11:6. <https://doi.org/10.1186/1472-6831-11-6>
3. Al-Zarea BK (2013) Satisfaction with appearance and the desired treatment to improve aesthetics. *Int J Dent* 2013:912368. <https://doi.org/10.1155/2013/912368>
4. Perdigão J (2016) *Tooth whitening: an evidence-based prespective*. Springer, Minneapolis
5. Boushell LW, Ritter AV, Garland GE, Tiwana KK, Smith LR, Broome A, Leonard RH (2012) Nightguard vital bleaching: side effects and patient satisfaction 10 to 17 years post-treatment. *J Esthet Restor Dent* 24(3):211–219. <https://doi.org/10.1111/j.1708-8240.2011.00479.x>
6. Alkahtani R, Stone S, German M, Waterhouse P (2020) A review on dental whitening. *J Dent* 100:103423. <https://doi.org/10.1016/j.jdent.2020.103423>
7. Tredwin CJ, Naik S, Lewis NJ, Scully C (2006) Hydrogen peroxide tooth-whitening (bleaching) products: review of adverse effects and safety issues. *Br Dent J* 200(7):371–376. <https://doi.org/10.1038/sj.bdj.4813423>
8. Kihn PW (2007) Vital tooth whitening. *Dent Clin North Am* 51(2):319–31, viii. <https://doi.org/10.1016/j.cden.2006.12.001>
9. Llana C, Villanueva A, Mejias E, Forner L (2020) Bleaching efficacy of at home 16% carbamide peroxide. A long-term clinical follow-up study. *J Esthet Restor Dent* 32(1):12–18. <https://doi.org/10.1111/jerd.12560>
10. Meireles SS, Heckmann SS, Leida FL, dos Santos IS, Della Bona A, Demarco FF (2008) Efficacy and safety of 10% and

- 16% carbamide peroxide tooth-whitening gels: a randomized clinical trial. *Oper Dent* 33(6):606–612. <https://doi.org/10.2341/07-150>
11. Majeed A, Farooq I, Grobler SR, Rossouw RJ (2015) Tooth-bleaching: a review of the efficacy and adverse effects of various tooth whitening products. *J Coll Physicians Surg Pak* 25(12):891–896
 12. Haywood VB (2000) Current status of nightguard vital bleaching. *Compend Contin Educ Dent Suppl* (28):S10–7; quiz S48.
 13. Haywood VB, Heymann HO (1989) Nightguard vital bleaching. *Quintessence Int* 20(3):173–176
 14. Demarco FF, Meireles SS, Masotti AS (2009) Over-the-counter whitening agents: a concise review. *Braz Oral Res* 23(Suppl 1):64–70. <https://doi.org/10.1590/S1806-83242009000500010>
 15. Naidu AS, Bennani V, Brunton J, Brunton P (2020) Over-the-counter tooth whitening agents: a review of literature. *Braz Dent J* 31(3):221–235. <https://doi.org/10.1590/0103-6440202003227>
 16. Basson RA, Grobler SR, Kotze TJ, Osman Y (2013) Guidelines for the selection of tooth whitening products amongst those available on the market. *SADJ* 68(3):122–129
 17. Goldstein RE (1997) In-office bleaching: where we came from, where we are today. *J Am Dent Assoc* 128(Suppl):11S–15S
 18. Maran BM, Matos TP, de Castro ADS, Vochikovski L, Amadori AL, Loguercio AD, Reis A, Berger SB (2020) In-office bleaching with low/medium vs. high concentrate hydrogen peroxide: a systematic review and meta-analysis. *J Dent* 103:103499. <https://doi.org/10.1016/j.jdent.2020.103499>
 19. de Geus JL, Wambier LM, Kossatz S, Loguercio AD, Reis A (2016) At-home vs in-office bleaching: a systematic review and meta-analysis. *Oper Dent* 41(4):341–356. <https://doi.org/10.2341/15-287-LIT>
 20. Haywood VB, Sword RJ (2017) Tooth bleaching questions answered. *Br Dent J* 223(5):369–380. <https://doi.org/10.1038/sj.bdj.2017.767>
 21. Carey CM (2014) Tooth whitening: what we now know. *J Evid Based Dent Pract* 14(Suppl):70–76. <https://doi.org/10.1016/j.jebdp.2014.02.006>
 22. Alqahtani MQ (2014) Tooth-bleaching procedures and their controversial effects: a literature review. *Saudi Dent J* 26(2):33–46. <https://doi.org/10.1016/j.sdentj.2014.02.002>
 23. Pontes M, Gomes J, Lemos C, Leao RS, Moraes S, Vasconcelos B, Pellizzer EP (2020) Effect of bleaching gel concentration on tooth color and sensitivity: a systematic review and meta-analysis. *Oper Dent* 45(3):265–275. <https://doi.org/10.2341/17-376-L>
 24. da Mata AD, Marques DN (2006) A novel technique for in-office bleaching with a 6% hydrogen peroxide paint-on varnish. *Eur J Esthet Dent* 1(1):70–77
 25. da Silva Marques DN, Silveira JM, Marques JR, Amaral JA, Guilherme NM, da Mata AD (2012) Kinetic release of hydrogen peroxide from different whitening products. *Eur J Esthet Dent* 7(3):344–352
 26. Eachempati P, KumbargereNagraj S, Kiran Kumar Krishanappa S, Gupta P, Yaylali IE (2018) Home-based chemically-induced whitening (bleaching) of teeth in adults. *Cochrane Database Syst Rev* 12(12):CD006202. <https://doi.org/10.1002/14651858.CD006202.pub2>
 27. Cardoso A, Pereira R, Silveira J, Dias S, Mata A, Marques D (2021) In-office tooth bleaching effectiveness with different soft-tissue barriers – randomized controlled trial. *Rev Port Estomatol Med Dent Cir Maxilofac* 62(3):141–149. <https://doi.org/10.24873/j.rpemd.2021.07.840>
 28. Benbachir N, Ardu S, Krejci I (2008) Spectrophotometric evaluation of the efficacy of a new in-office bleaching technique. *Quintessence Int* 39(4):299–306
 29. Ferraz NKL, Nogueira LC, Neiva IM, Ferreira RC, Moreira AN, Magalhaes CS (2018) Longevity, effectiveness, safety, and impact on quality of life of low-concentration hydrogen peroxides in-office bleaching: a randomized clinical trial. *Clin Oral Investig* 23(5):2061–2070. <https://doi.org/10.1007/s00784-018-2607-7>
 30. Bersezio C, Ledezma P, Estay J, Mayer C, Rivera O, Fernandez E (2019) Color regression and maintenance effect of intracoronal whitening on the quality of life: RCT-A one-year follow-up study. *Oper Dent* 44(1):24–33. <https://doi.org/10.2341/17-288-C>
 31. Angel P, Bersezio C, Estay J, Werner A, Retamal H, Araya C, Martin J, Fernandez E (2018) Color stability, psychosocial impact, and effect on self-perception of esthetics of tooth whitening using low-concentration (6%) hydrogen peroxide. *Quintessence Int* 49(7):557–566. <https://doi.org/10.3290/j.qi.a40468>
 32. Bersezio C, Martin J, Mayer C, Rivera O, Estay J, Vernal R, Haidar ZS, Angel P, Oliveira OB Jr, Fernandez E (2018) Quality of life and stability of tooth color change at three months after dental bleaching. *Qual Life Res* 27(12):3199–3207. <https://doi.org/10.1007/s11136-018-1972-7>
 33. Fernandez E, Bersezio C, Bottner J, Avalos F, Godoy I, Inda D, Vildosola P, Saad J, Oliveira OB Jr, Martin J (2017) Longevity, esthetic perception, and psychosocial impact of teeth bleaching by low (6%) hydrogen peroxide concentration for in-office treatment: a randomized clinical trial. *Oper Dent* 42(1):41–52. <https://doi.org/10.2341/15-335-C>
 34. Kothari S, Gray AR, Lyons K, Tan XW, Brunton PA (2019) Vital bleaching and oral-health-related quality of life in adults: a systematic review and meta-analysis. *J Dent* 84:22–29
 35. Klaric Sever E, Budimir Z, Cerovac M, Stambuk M, Par M, Negovetic Vranic D, Tarle Z (2018) Clinical and patient reported outcomes of bleaching effectiveness. *Acta Odontol Scand* 76(1):30–38. <https://doi.org/10.1080/00016357.2017.1376111>
 36. Mailart MC, Sakasagawa PA, Santos KC, Torres CRG, Palo RM, Borges AB One-year follow-up comparing at-home bleaching systems outcomes and the impact on patient's satisfaction: Randomized clinical trial. *J Esthet Restor Dent* 33(8):1175–1185. <https://doi.org/10.1111/jerd.12814>
 37. John MT, Rener-Sitar K, Baba K, Celebic A, Larsson P, Szabo G, Norton WE, Reissmann DR (2016) Patterns of impaired oral health-related quality of life dimensions. *J Oral Rehabil* 43(7):519–527. <https://doi.org/10.1111/joor.12396>
 38. A. General Assembly of the World Medical (2014) World Medical Association Declaration of Helsinki: ethical principles for medical research involving human subjects. *J Am Coll Dent* 81(3):14–18
 39. Fleiss J (1986) The design and analysis of clinical experiments. Wiley
 40. Klotz AL, Habibi Y, Corcodel N, Rammelsberg P, Hassel AJ, Zenthofer A (2018) Laboratory and clinical reliability of two spectrophotometers. *J EsthetRestor Dent* 28. <https://doi.org/10.1111/jerd.12452>
 41. Klotz AL, Habibi Y, Hassel AJ, Rammelsberg P, Zenthofer A How reliable and accurate is the shade determination of premolars by spectrophotometry?. *Clin Oral Investig* 24(4):1439–1444. <https://doi.org/10.1007/s00784-019-03162-x>
 42. Mahn E, Tortora SC, Olate B, Cacciuttolo F, Kernitsky J, Jorquera G (2021) Comparison of visual analog shade matching, a digital visual method with a cross-polarized light filter, and a spectrophotometer for dental color matching. *J Prosthet Dent* 125(3):511–516. <https://doi.org/10.1016/j.prosdent.2020.02.002>
 43. Ferreira Dias SB, Lourenco Silveira JM, Nunes Pereira RM, Cardoso AB, Duarte Sola Pereira da Mata A, da Silva Marques DN (2021) CIEL*a*b* values in vita classical and vita 3d master by two dental spectrophotometers. *Int J Prosthodont* 18. <https://doi.org/10.11607/ijp.7235>
 44. Amaral J, Sanches C, Marques D, Vaz Patto J, Barcelos F, Mata A (2018) Validation of Oral Health Impact Profile-14 and its

- association with Hyposialia in a Sjogren Syndrome Portuguese Population. *Acta Reumatol Port* 43(2):137–145
45. Slade GD (1997) Measuring oral health and quality of life: the oral health impact profile. University of North Carolina.
 46. Tsakos G, Allen PF, Steele JG, Locker D (2012) Interpreting oral health-related quality of life data. *Community Dent Oral Epidemiol* 40(3):193–200. <https://doi.org/10.1111/j.1600-0528.2011.00651.x>
 47. Ialongo C (2016) Understanding the effect size and its measures. *Biochem Med (Zagreb)* 26(2):150–63. <https://doi.org/10.11613/BM.2016.015>
 48. Middel B, van Sonderen E (2002) Statistical significant change versus relevant or important change in (quasi) experimental design: some conceptual and methodological problems in estimating magnitude of intervention-related change in health services research. *Int J Integr Care* 2:e15. <https://doi.org/10.5334/ijic.65> (Epub 2002 Dec 17)
 49. Locker D, Jokovic A, Clarke M (2004) Assessing the responsiveness of measures of oral health-related quality of life. *Community Dent Oral Epidemiol* 32(1):10–18. <https://doi.org/10.1111/j.1600-0528.2004.00114.x>
 50. Bland J, Altman D (1997) Cronbach's alpha. *BMJ* 314(7080):572
 51. A. Mata, D. Marques, J. Silveira, J. Marques, Branqueamento Dentário, in: J. Ramos (Ed.), *Estética em Medicina Dnetária*, Edição de autor, Coimbra, 2009, pp. 16–22.
 52. Paravina RD, Johnston WM, Powers JM (2007) New shade guide for evaluation of tooth whitening—colorimetric study. *J Esthet Restor Dent* 19(5):276–283. https://doi.org/10.1111/j.1708-8240.2007.00118_1.x
 53. International Organization for Standardization (2016) ISO/TR 28642 dentistry—guidance on color measurement. International Organization for Standardization, Geneva
 54. Perez Mdel M, Ghinea R, Rivas MJ, Yebra A, Ionescu AM, Paravina RD, Herrera LJ (2016) Development of a customized whiteness index for dentistry based on CIELAB color space. *Dent Mater* 32(3):461–467. <https://doi.org/10.1016/j.dental.2015.12.008>
 55. Perez MM, Herrera LJ, Carrillo F, Pecho OE, Dudea D, Gasparik C, Ghinea R, Bona AD (2019) Whiteness difference thresholds in dentistry. *Dent Mater* 35(2):292–297. <https://doi.org/10.1016/j.dental.2018.11.022>
 56. Commission Internationale de l'Eclairage (2004) CIE 15:2004 Technical Report. Colorimetry, 3rd Edition. Vienna, Austria.
 57. Paravina RD, Ghinea R, Herrera LJ, Bona AD, Igiel C, Linninger M, Sakai M, Takahashi H, Tashkandi E, Perez Mdel M (2015) Color difference thresholds in dentistry. *J Esthet Restor Dent* 27(Suppl 1):S1-9. <https://doi.org/10.1111/jerd.12149>
 58. Paravina RD, Perez MM, Ghinea R (2019) Acceptability and perceptibility thresholds in dentistry: a comprehensive review of clinical and research applications. *J Esthet Restor Dent* 31(2):103–112. <https://doi.org/10.1111/jerd.12465>
 59. Hogg R, Tanis E, Zimmerman D (2015) Probability and statistical inference. Pearson 9th Edition
 60. Borse S, Chaware SH (2020) Tooth shade analysis and selection in prosthodontics: a systematic review and meta-analysis. *J Indian Prosthodont* 20(2):131–140. https://doi.org/10.4103/jips.jips_399_19 (Epub 2020 Apr 7)
 61. Luque-Martinez I, Reis A, Schroeder M, Munoz MA, Loguercio AD, Masterson D, Maia LC (2016) Comparison of efficacy of tray-delivered carbamide and hydrogen peroxide for at-home bleaching: a systematic review and meta-analysis. *Clin Oral Investig* 20(7):1419–1433. <https://doi.org/10.1007/s00784-016-1863-7>
 62. Silva FBD, Chisini LA, Demarco FF, Horta BL, Correa MB (2018) Desire for tooth bleaching and treatment performed in Brazilian adults: findings from a birth cohort. *Braz Oral Res* 32:e12. <https://doi.org/10.1590/1807-3107bor-2018.vol32.0012>
 63. Joiner A (2006) The bleaching of teeth: a review of the literature. *J Dent* 34(7):412–419. <https://doi.org/10.1016/j.jdent.2006.02.002>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.