

# Nutritional Screening Tools among Hospitalized Children: from Past and to Present

Yeoun Joo Lee

*Department of Pediatrics, Pusan National University Children's Hospital, Yangsan, Korea*

Increased awareness of the importance of nutrition among hospitalized children has increased the use of nutrition screening tool (NST). However, it is not well known the NST for hospitalized children. Therefore, the purpose of this study is to understand the past and present state of adult and child NST and discuss the pros and cons of each NST.

**Key Words:** Nutrition, Screening, Malnutrition, Child

## INTRODUCTION

Nutrition management is essential for the outcomes of patients receiving advanced medical care. Nutritional support prevents complications from infections and shortens the length of hospital stay [1,2]. A recent study about medical cost related to malnourished hospitalized patients has proven that a comprehensive nutrition-focused quality improvement program reduced the per-patient healthcare cost [3].

Although there is a growing interest in preventing malnutrition in hospitalized patients, the recent study has also shown that the prevalence rate of malnutrition among hospitalized children ranged from 7.5% to 17% in Europe [4,5]. In a study in Korea, the prevalence of malnutrition among hospitalized children and adults were 12.5% and 22%, respectively

[6,7].

To systematically manage the nutrition of hospitalized patients, the European Society for Parenteral and Enteral Nutrition (ESPEN) and American Society for Parenteral and Enteral Nutrition (ASPEN) recommended the use of the guidelines on nutritional screening to identify hospitalized patients who are at risk for malnutrition [8,9]. The guidelines suggested the identification of patients at risk for malnutrition and provision of interventions and treatments to these patients with the help from a multidisciplinary team of doctors, dietitians, nurses, and pharmacists [10]. Furthermore, the Joint Commission in the United States proposed the nutritional screening of all patients within 24 hours after admission.

The use of appropriate nutrition screening tools (NSTs) is important. Overly complex NSTs are difficult to access, whereas the simplicity but poorly vali-

Received : March 21, 2018, Accepted : March 26, 2018

**Corresponding author:** Yeoun Joo Lee, Department of Pediatrics, Pusan National University Children's Hospital, 20 Geumo-ro, Mulgeum-eup, Yangsan 50612, Korea. Tel: +82-55-360-2180, Fax: +82-55-360-2181, E-mail: moonmissing@gmail.com

Copyright © 2018 by The Korean Society of Pediatric Gastroenterology, Hepatology and Nutrition

This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

dated still has limit. NSTs should be practical, reliable, valid, and evidence based [11]. This study aimed at validating the characteristics, application, and validation results of the available NSTs that are used for hospitalized children.

## NSTs FOR HOSPITALIZED ADULTS

A variety of NSTs that can be used to screen hospitalized adult patients were developed. The main concept for nutrition screening had started from improving surgical outcomes. The prognostic nutritional index [12] and prognostic inflammatory and nutritional index [13] published in the 1980s were used as the primary index to screen for nutrition based on triceps skin fold, skin sensitivity, and albumin, prealbumin, or transferrin level. With the emphasis of the importance of nutritional screening, NSTs have been developed to improve its accessibility and validity. In the 1990s, several NSTs were developed such as the nutritional risk index (NRI) [14], Birmingham nutrition risk score [15], nutrition risk classification (NRC) [16], and malnutrition screening tool (MST) [17]. They had changed their focus on history taking and physical examination findings. Appetite, dietary intake, weight loss, or body mass index was used as an index of NSTs.

Since 2000, representative screening tools, such as simple screening tool [18], malnutrition universal screening tool (MUST) [11], nutritional risk screening (NRS) 2002 [19], short nutritional assessment questionnaire (SNAQ) [20], and recent Canadian nutrition screening tool [21], have been developed and are still used. The presence of acute disease and severity of the diagnosis along with anthropometric measurements and dietary factors are considered important in the use of MUST and NRS 2002, which are among the NSTs developed since the 2000s. Subjective global assessment (SGA) [22] and mini nutritional assessment (MNA) [23] are convenient assessment tools not only assessing but also screening of the nutritional status. SGA is a method of nutritional assessment based on medical history (weight and diet changes, primary diagnosis, and stress lev-

el) and physical symptoms (presence of subcutaneous fat, muscle wasting, ankle edema, ascites, functional capacity, and gastrointestinal symptoms).

A validation study on the individual screening tool, medical environment, and population and age groups has been published in a variety of countries. Moreover, several researches compared each screening tool that was used in a limited setting [24-27].

Aside from the universal NSTs, NST for a specific disease has also been developed, considering the specific nature of the disease. Several NSTs were developed for specific diseases, such as cirrhosis [28], cancers [29], and cerebral palsy [30], and critically-ill patient groups [31].

## SCREENING TOOLS FOR HOSPITALIZED CHILDREN

Studies on NSTs for children are limited compared to those of adults. Secker and Jeejeebhoy [32] have used the SGA for hospitalized children who underwent major thoracic or abdominal surgery. A correlation was observed between malnourished children and a higher risk of developing nutrition-associated complications and prolonged hospitalizations. A Brazilian study that used SGA for children with acute illness had found an association between the SGA score and anthropometric measurement. However, no association was observed between the SGA score and length of hospital stay [33].

In 2000, Sermet-Gaudelus et al. [34] have published the pediatric nutritional risk score (PNRS) by developing a unique equation with the study endpoint at >2% weight loss during the first week of admission. The factors for PNRS are food intake <50%, pain, and the presence of pathologic conditions. Each component can be assigned a score from 1 to 5. Unlike other NSTs, PNRS did not use anthropometric data.

In the UK, the screening tool for the assessment of malnutrition in paediatrics (STAMP) [35] was established in 2004 and evaluated in 2007. To evaluate the quick and easy-to-use NSTs for hospitalized children, three factors were considered: diagnosis, nutri-

tional intake, as well as weight and height. After evaluating these factors, the sum was classified into low, medium, and high risk, and the TAMP also suggested that a care plan in the last step.

In a large tertiary children's hospital in the UK, the pediatric Yorkhill malnutrition score (PYMS) [36] was developed for nutritional screening, and the PYMS used four factors for the screening: body mass index, history of recent weight loss, changes in nutritional intake, and the predicted effect of the current medical condition on nutritional status. PYMS used anthropometric data for two of the four factors by placing weight on the anthropometric data, and subjective data were also used for the effect of the current medical condition.

In 2009, the Dutch Society published the national survey results to test their own NST for the screening tool for risk on nutritional status and growth (STRONG<sub>kids</sub>) [37]. They also focused on developing STRONG<sub>kids</sub> to improve its applicability. They used four factors for the subjective assessment, high-risk disease, nutritional intake/losses and weight loss/poor weight gain, and a score of 0 to 2 was provided. Like PYMS, STRONG<sub>kids</sub> also recommended a nutritional intervention for each risk.

Recently, the pediatric digital scaled malnutrition risk screening tool (PeDiSMART) [38] was introduced for hospitalized children by using computer-based information systems in Greece. One of the significant advantages of PeDiSMART is its high reproducibility. Moreover, it can help professionals save time. The factors are weight-for-age z score, nutritional intake level, overall disease impact, and disease symptoms affecting intake. Weight loss might be significantly associated with nutrition support during hospitalization in the PeDiSMART malnutrition risk group after adjusting anthropometric data.

Pediatric nutrition screening tool (PNST) [39] consists of four simple questions that a child's caregiver can answer with Yes or No. Among the four questions about recent weight loss, poor weight gain over the last few months, poor oral intake within the last few weeks, and obvious weight loss or gain based on the PNST, two affirmative responses can identify

patients who are at risk. Moreover, a correlation between nutrition risk identified using the PNST and pediatric SGA was observed. PNST also correlated with nutritional status using z-score. PNST may be the easiest NST that can be used. However, interrater reliability or reproducibility data are limited.

Except for PNST, most of the NSTs established the nutritional risk based on three categories: low (mild or grade 1), medium (moderate or grade 2), and severe (grade 3). All the NSTs used to monitor intake evaluate at least one factor for nutritional screening. Although PNST did not focus on anthropometric data, all the other NSTs focused on weight and height or recent weight loss. NSTs for hospitalized children are summarized in Table 1 [34-39].

## VALIDATION STUDY ABOUT NSTs FOR HOSPITALIZED CHILDREN

A limited number of studies about NSTs for hospitalized children compared with those for adults are available. There are studies that compared several representative NSTs [40-46]. Although some studies considered that the PNST is the most suitable for clinical practice since the results of high sensitivity and specificity in PNST compared with SGA [40,43]. However, in New Zealand, a study has shown that STRONG<sub>kids</sub> was the most reliable NST in their clinical setting [45]. Moreover, there are other reports that have reported that the STRONG<sub>kids</sub> was the most suitable for clinical use [42,46]. Even in studies about acute burn injuries, it is difficult to conclude that one NST is superior than the other [47]. A study that used PeDiSMART has shown a correlation among PYMS, STRONG<sub>kids</sub>, and STAMP. In addition, the area under the curve for weight loss/nutrition support and the length of hospital stay (>7 days) was superior to that of the other three indicators [38]. However, newly developed NSTs, such as PeDiSMART and PNST, have not been fully validated.

**Table 1.** Nutrition Screening Tools Made for Hospitalized Children and Representative Validation Study of Each NST

Name of NST	Country	Number of subjects	Factors used in the NST				Score	
			Number of factors	Anthropometry	Appetite or food intake	Diagnosis or pathologic conditions		Others
PNRS [34]	France	296	3		Food intake <50%	Pathologic condition	Pain	1 to 5
STAMP [35]	UK	122 (developing) 238 (evaluation)	3	Weight and height	Nutritional intake	Diagnosis		0 to 9
PYMS [36]	UK	247	4	Body mass index, history of recent weight loss	Changes in nutritional intake	Predicted effect of the current medical condition on the nutritional status		0 to 7
STRONG <sub>kids</sub> [37]	Netherlands	424	4	Weight loss or poor weight increase	Nutritional intake and losses	High risk disease	Subjective global assessment	0 to 5
PeDiSMART (computer software) [38]	Greece	500	4	Weight-for-age z score	Nutrition intake level, symptoms affecting intake	Overall disease impact		0 to 18
PNST [39]	Australia	295	4	Recent weight loss, poor weight gains over the last few months, obviously underweight/significantly overweight	Eating/feeding less in the last few weeks			0 to 4

NST: nutrition screening tool, PNRS: pediatric nutritional risk score, STAMP: screening tool for the assessment of malnutrition in paediatrics, PYMS: pediatric Yorkhill malnutrition score, STRONG<sub>kids</sub>: screening tool for risk on nutritional status and growth, PeDiSMART: pediatric digital scaled malnutrition risk screening tool, PNST: pediatric nutrition screening tool.

## DISEASE-SPECIFIC SCREENING TOOLS

A study on paradigm shift was also conducted for the evaluation and treatment of children with disease-specific malnutrition [48]. Since malnutrition is a serious health problem in children with cancer, malnutrition in these children has been a topic of interest. The nutrition screening tool for childhood cancer (SCAN) was developed in Australia and had an excellent accuracy in term of pediatric SGA [49]. It is difficult to identify poor nutritional status with a simple method, and researchers attempted to assess and treat poor nutrition in pediatric patients with cancer [50,51]. Although this tool is not only for hospitalized children, a NST for children with cystic

fibrosis is also available [52].

## CAN THE NST OBTAIN SIMILAR RESULTS REGARDLESS OF PERFORMER?

Nutritional screening after hospitalization is usually performed by nurses. However, nutritional assessment in hospitalized children is usually conducted by a clinical dietitian. Several screening tools that can be used by nurses during the developmental stage have been developed [11,36]. In addition, a validation study is usually performed by research dietitians [36]. Good reproducibility without obtaining different results depending on the performers is

one of the important factors of a good screening tool. A study about the validity and reliability of nutritional screening in adults by two independence nurses within 24 hours of admission was conducted. The article reported that even within the same occupation, inter-observer agreement showed 78.3%, though 100% agreement in detecting severely malnourished patients [53]. To overcome these differences in the performer's distinction, PeDiSMART attempted to increase its reproducibility by using a computer software [38]. Whether NST can reproduce similar results regardless of the performer must be highly considered.

## CONCLUSION

To date, the performance accuracy of NSTs for children and adults are still being developed. Previous studies do not show that any screening tool is superior than the other. Screening tools that tailored for each hospital and diagnosis and those with excellent reproducibility regardless of performers must be developed. However, in actual settings, this cannot be easily performed. Therefore, health care professionals in hospitals must identify and use screening tools that are the most appropriate and suitable for their hospital setting.

Moreover, the sensitivity and specificity of the tools must be re-evaluated compared with the actual outcomes in hospital settings. Most importantly, patients who are at risk for malnutrition should be treated, and malnutrition must be prevented in these patients [54].

## REFERENCES

- Zhang H, Wang Y, Jiang ZM, Kondrup J, Fang H, Andrews M, et al. Impact of nutrition support on clinical outcome and cost-effectiveness analysis in patients at nutritional risk: a prospective cohort study with propensity score matching. *Nutrition* 2017;37:53-9.
- Hecht C, Weber M, Grote V, Daskalou E, Dell'Era L, Flynn D, et al. Disease associated malnutrition correlates with length of hospital stay in children. *Clin Nutr* 2015;34:53-9.
- Pimenta FS, Oliveira CM, Hattori WT, Teixeira KR. Agreement between the methods: Subjective Global Nutritional Assessment and the nutritional assessment of the World Health Organization. *J Pediatr (Rio J)* 2017. doi: 10.1016/j.jped.2017.09.007.
- Sissaoui S, De Luca A, Piloquet H, Guimber D, Colomb V, Peretti N, et al. Large scale nutritional status assessment in pediatric hospitals. *e-SPEN J* 2013;8:e68-72.
- Freijer K, van Puffelen E, Joosten KF, Hulst JM, Koopmanschap MA. The costs of disease related malnutrition in hospitalized children. *Clin Nutr ESPEN* 2018;23:228-33.
- Kang MC, Kim JH, Ryu SW, Moon JY, Park JH, Park JK, et al. Prevalence of malnutrition in hospitalized patients: a multicenter cross-sectional study. *J Korean Med Sci* 2018;33:e10.
- Hwang EH, Park JH, Chun P, Lee YJ. Prevalence and risk factors for the weight loss during hospitalization in children: a single Korean children's hospital experience. *Pediatr Gastroenterol Hepatol Nutr* 2016; 19:269-75.
- Kondrup J, Allison SP, Elia M, Vellas B, Plauth M. ESPEN guidelines for nutrition screening 2002. *Clin Nutr* 2003;22:415-21.
- Mehta NM, Compber C; A.S.P.E.N. Board of Directors. A.S.P.E.N. Clinical Guidelines: nutrition support of the critically ill child. *JPEN J Parenter Enteral Nutr* 2009;33:260-76.
- Agostoni C, Axelson I, Colomb V, Goulet O, Koletzko B, Michaelsen KF, et al. The need for nutrition support teams in pediatric units: a commentary by the ESPGHAN committee on nutrition. *J Pediatr Gastroenterol Nutr* 2005;41:8-11.
- Elia M. The "MUST" Report. Nutritional screening of adults: a multidisciplinary responsibility. Redditch, UK: British Association for Parenteral and Enteral Nutrition; 2003 [cited 2009 Mar]. Available from: [http://www.bapen.org.uk/pdfs/must/must\\_exec\\_sum.pdf](http://www.bapen.org.uk/pdfs/must/must_exec_sum.pdf).
- Buzby GP, Mullen JL, Matthews DC, Hobbs CL, Rosato EF. Prognostic nutritional index in gastrointestinal surgery. *Am J Surg* 1980;139:160-7.
- Ingenbleek Y, Carpentier YA. A prognostic inflammatory and nutritional index scoring critically ill patients. *Int J Vitam Nutr Res* 1985;55:91-101.
- Veterans Affairs Total Parenteral Nutrition Cooperative Study Group. Perioperative total parenteral nutrition in surgical patients. *N Engl J Med* 1991;325: 525-32.
- Reilly HM, Martineau JK, Moran A, Kennedy H. Nutritional screening--evaluation and implemen-

- tation of a simple Nutrition Risk Score. *Clin Nutr* 1995;14:269-73.
16. Kovacevich DS, Boney AR, Braunschweig CL, Perez A, Stevens M. Nutrition risk classification: a reproducible and valid tool for nurses. *Nutr Clin Pract* 1997;12:20-5.
  17. Ferguson M, Capra S, Bauer J, Banks M. Development of a valid and reliable malnutrition screening tool for adult acute hospital patients. *Nutrition* 1999;15:458-64.
  18. Laporte M, Villalon L, Thibodeau J, Payette H. Validity and reliability of simple nutrition screening tools adapted to the elderly population in healthcare facilities. *J Nutr Health Aging* 2001;5:292-4.
  19. Kondrup J, Rasmussen HH, Hamberg O, Stanga Z; Ad Hoc ESPEN Working Group. Nutritional risk screening (NRS 2002): a new method based on an analysis of controlled clinical trials. *Clin Nutr* 2003;22:321-36.
  20. Kruizenga HM, Van Tulder MW, Seidell JC, Thijs A, Ader HJ, Van Bokhorst-de van der Schueren MA. Effectiveness and cost-effectiveness of early screening and treatment of malnourished patients. *Am J Clin Nutr* 2005;82:1082-9.
  21. Laporte M, Keller HH, Payette H, Allard JP, Duerksen DR, Bernier P, et al. Validity and reliability of the new Canadian Nutrition Screening Tool in the 'real-world' hospital setting. *Eur J Clin Nutr* 2015;69:558-64.
  22. Detsky AS, McLaughlin JR, Baker JP, Johnston N, Whittaker S, Mendelson RA, et al. What is subjective global assessment of nutritional status? *JPEN J Parenter Enteral Nutr* 1987;11:8-13.
  23. Guigoz Y. The Mini Nutritional Assessment (MNA) review of the literature--What does it tell us? *J Nutr Health Aging* 2006;10:466-85; discussion 485-7.
  24. Rabito EI, Marcadenti A, da Silva Fink J, Figueira L, Silva FM. Nutritional risk screening 20, short nutritional assessment questionnaire, malnutrition screening tool, and malnutrition universal screening tool are good predictors of nutrition risk in an emergency service. *Nutr Clin Pract* 2017;32:526-32.
  25. Chi J, Yin S, Zhu Y, Gao F, Song X, Song Z, et al. A comparison of the nutritional risk screening 2002 tool with the subjective global assessment tool to detect nutritional status in Chinese patients undergoing surgery with gastrointestinal cancer. *Gastroenterol Nurs* 2017;40:19-25.
  26. Fiol-Martínez L, Calleja-Fernández A, Pintor de la Maza B, Vidal-Casariago A, Villar-Taibo R, Urioste-Fondo A, et al. Comparison of two nutritional screening tools to detect nutritional risk in hematologic inpatients. *Nutrition* 2017;34:97-100.
  27. Koren-Hakim T, Weiss A, HersHKovitz A, Otrzateni I, Anbar R, Gross Nevo RF, et al. Comparing the adequacy of the MNA-SF, NRS-2002 and MUST nutritional tools in assessing malnutrition in hip fracture operated elderly patients. *Clin Nutr* 2016;35:1053-8.
  28. Schneider AC, Pinto RB, Silveira TR. Nutritional risk and malnutrition determination by anthropometry in cirrhotic children and adolescents. *Arq Gastroenterol* 2007;44:345-9.
  29. Decoster L, Van Puyvelde K, Mohile S, Wedding U, Basso U, Colloca G, et al. Screening tools for multidimensional health problems warranting a geriatric assessment in older cancer patients: an update on SIOG recommendations †. *Ann Oncol* 2015;26:288-300.
  30. Benigni I, Devos P, Rofidal T, Seguy D. The CP-MST, a malnutrition screening tool for institutionalized adult cerebral palsy patients. *Clin Nutr* 2011;30:769-73.
  31. Rahman A, Hasan RM, Agarwala R, Martin C, Day AG, Heyland DK. Identifying critically-ill patients who will benefit most from nutritional therapy: further validation of the "modified NUTRIC" nutritional risk assessment tool. *Clin Nutr* 2016;35:158-62.
  32. Secker DJ, Jeejeebhoy KN. Subjective Global Nutritional Assessment for children. *Am J Clin Nutr* 2007; 85:1083-9.
  33. Vermilyea S, Slicker J, El-Chammas K, Sultan M, Dasgupta M, Hoffmann RG, et al. Subjective global nutritional assessment in critically ill children. *JPEN J Parenter Enteral Nutr* 2013;37:659-66.
  34. Sermet-Gaudelus I, Poisson-Salomon AS, Colomb V, Brusset MC, Mosser F, Berrier F, et al. Simple pediatric nutritional risk score to identify children at risk of malnutrition. *Am J Clin Nutr* 2000;72:64-70.
  35. McCarthy H, Dixon M, Crabtree I, Eaton-Evans MJ, McNulty H. The development and evaluation of the Screening Tool for the Assessment of Malnutrition in Paediatrics (STAMP©) for use by healthcare staff. *J Hum Nutr Diet* 2012;25:311-8.
  36. Gerasimidis K, Keane O, Macleod I, Flynn DM, Wright CM. A four-stage evaluation of the Paediatric Yorkhill Malnutrition Score in a tertiary paediatric hospital and a district general hospital. *Br J Nutr* 2010;104:751-6.
  37. Hulst JM, Zwart H, Hop WC, Joosten KF. Dutch national survey to test the STRONGkids nutritional risk screening tool in hospitalized children. *Clin Nutr* 2010;29:106-11.
  38. Karagiozoglou-Lampoudi T, Daskalou E, Lampoudis D, Apostolou A, Agakidis C. Computer-based malnutrition risk calculation may enhance the ability to identify pediatric patients at malnutrition-related risk for unfavorable outcome. *JPEN J Parenter Enteral Nutr*

- 2015;39:418-25.
39. White M, Lawson K, Ramsey R, Dennis N, Hutchinson Z, Soh XY, et al. Simple nutrition screening tool for pediatric inpatients. *JPEN J Parenter Enteral Nutr* 2016;40:392-8.
  40. Lestari NE, Nurhaeni N, Wanda D. The pediatric york-hill malnutrition score is a reliable malnutrition screening tool. *Compr Child Adolesc Nurs* 2017;40 (sup1):62-8.
  41. Chourdakis M, Hecht C, Gerasimidis K, Joosten KF, Karagiozoglou-Lampoudi T, Koetse HA, et al. Malnutrition risk in hospitalized children: use of 3 screening tools in a large European population. *Am J Clin Nutr* 2016;103:1301-10.
  42. Márquez Costa MV, Alberici Pastore C. Nutritional screening tool versus anthropometric assessment in hospitalized children: which method is better associated to clinical outcomes? *Arch Latinoam Nutr* 2015;65:12-20.
  43. Huysentruyt K, Devreker T, Dejonckheere J, De Schepper J, Vandenplas Y, Cools F. Accuracy of nutritional screening tools in assessing the risk of undernutrition in hospitalized children. *J Pediatr Gastroenterol Nutr* 2015;61:159-66.
  44. Wonoputri N, Djais JT, Rosalina I. Validity of nutritional screening tools for hospitalized children. *J Nutr Metab* 2014. doi: 10.1155/2014/143649.
  45. Moeeni V, Walls T, Day AS. Nutritional status and nutrition risk screening in hospitalized children in New Zealand. *Acta Paediatr* 2013;102:e419-23.
  46. Huysentruyt K, Alliet P, Muysfont L, Rossignol R, Devreker T, Bontems P, et al. The STRONG(kids) nutritional screening tool in hospitalized children: a validation study. *Nutrition* 2013;29:1356-61.
  47. Bang YK, Park MK, Ju YS, Cho KY. Clinical significance of nutritional risk screening tool for hospitalised children with acute burn injuries: a cross-sectional study. *J Hum Nutr Diet* 2017. doi: 10.1111/jhn.12518.
  48. Mehta NM, Corkins MR, Lyman B, Malone A, Goday PS, Carney LN, et al. Defining pediatric malnutrition: a paradigm shift toward etiology-related definitions. *JPEN J Parenter Enteral Nutr* 2013;37:460-81.
  49. Murphy AJ, White M, Viani K, Mosby TT. Evaluation of the nutrition screening tool for childhood cancer (SCAN). *Clin Nutr* 2016;35:219-24.
  50. Murphy AJ, White M, Davies PS. The validity of simple methods to detect poor nutritional status in paediatric oncology patients. *Br J Nutr* 2009;101:1388-92.
  51. Robinson DL, Loman DG, Balakas K, Flowers M. Nutritional screening and early intervention in children, adolescents, and young adults with cancer. *J Pediatr Oncol Nurs* 2012;29:346-55.
  52. McDonald CM. Validation of a nutrition risk screening tool for children and adolescents with cystic fibrosis ages 2-20 years. *J Pediatr Gastroenterol Nutr* 2008; 46:438-46.
  53. Lim SL, Ang E, Foo YL, Ng LY, Tong CY, Ferguson M, et al. Validity and reliability of nutrition screening administered by nurses. *Nutr Clin Pract* 2013;28:730-6.
  54. Weekes CE, Spiro A, Baldwin C, Whelan K, Thomas JE, Parkin D, et al. A review of the evidence for the impact of improving nutritional care on nutritional and clinical outcomes and cost. *J Hum Nutr Diet* 2009;22:324-35.