

Prevalence and Genetic Characterization of *Toxoplasma gondii* in House Sparrows (*Passer domesticus*) in Lanzhou, China

Wei Cong^{1,2,†}, Si-Yang Huang^{1,†,*}, Dong-Hui Zhou¹, Xiao-Xuan Zhang^{1,3}, Nian-Zhang Zhang¹, Quan Zhao² and Xing-Quan Zhu^{1,2}

¹State Key Laboratory of Veterinary Etiological Biology, Key Laboratory of Veterinary Parasitology of Gansu Province, Lanzhou Veterinary Research Institute, Chinese Academy of Agricultural Sciences, Lanzhou, Gansu Province 730046, P.R. China; ²College of Animal Science and Technology, Jilin Agricultural University, Changchun, Jilin Province 130118, P.R. China; ³Department of Veterinary Medicine, College of Agriculture, Yanbian University, Yanji, Jilin Province 133000, P.R. China

Abstract: The prevalence of *Toxoplasma gondii* infection in birds has epidemiological significance because birds are indeed considered as a good indicator of environmental contamination by *T. gondii* oocysts. In this study, the prevalence of *T. gondii* in 313 house sparrows in Lanzhou, northwestern China was assayed by the modified agglutination test (MAT). Antibodies to *T. gondii* were positive in 39 (12.46%) of 313 samples (MAT titer \geq 1:5). Tissues of heart, brain, and lung from the 39 seropositive house sparrows were tested for *T. gondii* DNA, 11 of which were found to be positive for the *T. gondii* B1 gene by PCR amplification. These positive DNA samples were typed at 9 genetic markers, including 8 nuclear loci, i.e., SAG1, 5'- and 3'-SAG2, alternative SAG2, SAG3, GRA6, L358, PK1, c22-8 and an apicoplast locus Apico. Of them, 4 isolates were genotyped with complete data for all loci, and 2 genotypes (Type II variants; ToxoDB #3 and a new genotype) were identified. These results showed that there is a potential risk for human infection with *T. gondii* in this region. To our knowledge, this is the first report of *T. gondii* seroprevalence in house sparrows in China.

Key words: *Toxoplasma gondii*, sparrow, seroprevalence, PCR-RFLP typing, northwestern China

Toxoplasma gondii is an important protozoan parasite that infects most species of warm-blooded animals worldwide, including humans and birds [1-3]. It has a complex life cycle in that sexual development occurs only in the intestine of felines, but asexual replication can occur extra-intestinally in many vertebrate hosts. Humans become infected postnatally via ingesting *T. gondii* tissue cysts from undercooked meat, consuming food or drinking water contaminated with *T. gondii* oocysts, or by accidentally ingesting oocysts from the environment [4]. Birds as intermediate wild hosts are very important in the circulation of the parasite in wildlife, and *T. gondii* infection in birds is considered to be epidemiologically significant because of their dietary habit. The house sparrow is frequently commensal with humans and easily adaptable to urban and rural areas [5]. Therefore, the investigation of *T. gondii* infec-

tion in house sparrows is an effective way to estimate environmental contamination with *T. gondii* oocysts.

The genetic diversity of *T. gondii* varies in different geographical regions and different hosts. Isolates of *T. gondii* in South America, for example, are genetically and biologically different from those in North America and Europe [6-13], where the *T. gondii* population structure is highly clonal and composed mainly of 4 distinct lineages, i.e., Types I, II, III, and 12 [13]. We have previously identified limited genotypes in isolates from humans, cats, pigs, and sheep in China [14-17], but there is little genetic data on *T. gondii* isolates from wild birds [18]. In the present paper, we report the seroprevalence and genetic characterization of *T. gondii* isolates from house sparrows in Lanzhou, northwest China.

A total of 313 blood samples were obtained from house sparrows between September and November 2011 in Lanzhou. Blood samples were incubated at 37°C for 2 hr and then centrifuged at 3,000 g for 5 min to separate the serum samples. The separated serum samples were stored at -20°C until further analysis.

Antibodies to *T. gondii* were determined in house sparrow

•Received 14 January 2013, revised 16 March 2013, accepted 20 March 2013.

*Corresponding author (siyang.huang@hotmail.com)

†These authors contributed equally to this work.

© 2013, Korean Society for Parasitology and Tropical Medicine

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

sera by the modified agglutination test (MAT) as described previously [19,20]. Briefly, sera were added to the “U” bottom of 96-well microtiter plates, and diluted 2-fold starting from 1:5 to 1:20. Birds sera with MAT titers of 1:5 or higher were considered positive for *T. gondii* antibodies based on a previous study [21], those sera with doubtful reactions were re-tested, and positive and negative controls were included in each test.

Heart, brain, and lung tissues from 39 seropositive house sparrows were used for DNA extraction. Genomic DNA was extracted from 5-10 g of heart, brain, or lung tissues by SDS/proteinase K treatment, column-purified (Tiangen™, Beijing, China) and eluted into 50 ml H₂O according to the manufacturer’s recommendations. A nested PCR targeting the *T. gondii* B1 gene was performed to detect possible infection with *T. gondii* [22]. DNA samples giving positive B1 amplification were then used for genetic characterization.

Genetic characterization of *T. gondii* isolates from sparrows was carried out using the multilocus PCR-RFLP method [16, 18,23]. Multiplex PCR-amplified products were 1:1 diluted in sterile, double-distilled water, and then used for nested PCR amplifications with internal primers for each marker, separately. The nested PCR products were digested with restriction enzymes. The restriction fragments were resolved in 2.5% agarose gel, stained by the GoldenView™, and photographed using a gel documentation system (UVP GelDoc-It™ Imaging System, Cambridge, U.K.).

Thirty nine (12.46%) out of 313 serum samples were detected positive for *T. gondii* antibodies by MAT (Table 1). However, most of them had low titers, with MAT titers 1:5 in 29 house sparrows, and with 1:10 or higher in 10 house sparrows. Of the 117 DNA samples, 11 were positive for the *T. gondii* B1 gene by PCR amplification (Fig. 1), including 6 from brains, 3 from

lungs, and 2 from hearts. Moreover, among the 11 positive DNA samples, 2 samples were come from the same house sparrow (sample LZS232) but different tissues (brain and heart).

Four DNA samples showed complete genotyping results, 2 from brain (sample LZS232, LZS253), 1 from heart (sample LZS232), and 1 from lung (sample LZS107) (Fig. 2). Due to low DNA concentration, 7 of the 11 positive samples could not be completely genotyped and was therefore not used. Two genotypes (ToxoDB #3 and a new genotype) were identified from the 11 positive samples (Table 2).

Little is known of the prevalence of *T. gondii* in sparrows in China. Sparrows could be one of the important species during the cycle of *T. gondii* transmission. Sparrows especially house sparrows are normally distributed around the human environment. Due to their habit of feeding close to the ground, sparrows, like free-range chickens, are indeed considered as a good indicator of environmental contamination by *T. gondii* oocysts [4]. More importantly, comparing to chickens, sparrows can transmit *T. gondii* faster and wider because of their flying ability, hence are more important intermediate hosts of *T. gondii* than chickens.

Any meat from warm-blooded animals and birds has been traditionally considered a major source of *Toxoplasma* infection in Western countries [4]. Some local people in this study region like to eat the sparrow meat very much. The risk associated with the type of meat varies among different countries according to local eating habits and according to the *T. gondii*

Table 1. Seroprevalence of *Toxoplasma gondii* in house sparrows in Lanzhou, northwest China using the modified agglutination test (MAT)

	No. tested	No. positive	Prevalence (%)	95% CI
House sparrow	313	39	12.46	11.25-13.67




Fig. 1. Representative PCR products of *Toxoplasma gondii* B1 gene amplified from tissue DNA samples of house sparrows. Lanes 1-12 represent negative amplification, and lanes 13-18 represent positive amplification, respectively.




Fig. 2. PCR-RFLP analysis of *Toxoplasma gondii* isolates from house sparrows in Lanzhou, China based on 9 different loci. M represents a DNA marker. Lanes 1-10 represent GT1, PTG, CTG, MAS, TgCatCal, TgCaBr5, LZS253, LZS232 (brain), LZS232 (heart), and LZS107, respectively.

Table 2. Summary of genotyping of *Toxoplasma gondii* from house sparrows in Lanzhou, northwest China

Isolate ID	Host	Tissue	Location	SAG1	5'+3'SAG2	Alternative SAG2	SAG3	GRA6	c22-8	L358	PK1	Apico	Genotype
GT1I	Goat		United States	I	I	I	I	I	I	I	I	I	Reference, Type I, ToxoDB #10
PTG	Sheep		United States	II/III	II	II	II	II	II	II	II	II	Reference, Type II, ToxoDB #1
CTG	Cat		United States	II/III	III	III	III	III	III	III	III	III	Reference, Type III, ToxoDB #2
MAS	Human		France	u-1 ^a	I	II	III	III	u-1 ^a	I	III	I	Reference, ToxoDB #17
TgCgCa1	Cougar		Canada	I	I	II	III	II	II	I	u-2 ^a	I	Reference, ToxoDB #66
TgCatBr5	Cat		Brazil	I	III	III	III	III	I	I	u-1 ^a	I	Reference, ToxoDB #19
LZS253	House Sparrow	Brain	Lanzhou, Gansu	II	II	II	II	II	I	II	II	I	Type II variant, new genotype
LZS232	House Sparrow	Brain	Lanzhou, Gansu	II	II	II	II	II	II	II	II	I	Type II variant, ToxoDB #3
LZS232	House Sparrow	Heart	Lanzhou, Gansu	II	II	II	II	II	II	II	II	I	Type II variant, ToxoDB #3
LZS107	House Sparrow	Lung	Lanzhou, Gansu	II	II	II	II	II	II	II	II	I	Type II variant, ToxoDB #3

^au-1 and u-2 represent unique RFLP genotypes, respectively.

prevalence in meat-producing animals. We previously reported that 4 of 178 bird DNA samples were positive for the *T. gondii* B1 gene by PCR amplification. Because serum samples of these 178 birds were not available for serological examination, the data could not indicate the seroprevalence of *T. gondii* infection in these wild birds in China [18]. Meanwhile, only breast muscle samples were available in that study [18]. Here, we collected heart, brain, and lung tissues for genetic characterization. In the present study, antibodies to *T. gondii* were examined by MAT, and found in 39 (12.46%) of 313 house sparrows at the cut of 1:5. Although with low titers, house sparrows are easily exposed to *T. gondii* through the ingestion of food or water contaminated with sporulated oocysts excreted by infected felids. Free-living animals such as wild birds and mammals were considered to be sentinels of environmental

spreading of *T. gondii* oocysts, because of the increasing probability of sharing the same environment with humans [24].

Two genotypes (both are Type II variants) were identified in this study, one of them had a Type I allele at the Apico locus and is considered the Type II variant (ToxoDB #3), which is widely distributed in the world [25-27]. This was the third time that this Type II variant was identified in China, as this genotype was previously reported from sheep in Qinghai Province [17] and wild birds in Xinjiang Uygur Autonomous Region [18]. However, the other strain isolated from sample LZS253 had the Type I allele at the c22-8 locus in addition to the Apico locus, this genotype has not been reported before, and is considered a Type II variant (a new genotype). This is different from previous studies that showed the genotype ToxoDB #9 was predominant in cats and other animals in southern, south-

western, and central parts of China [14,17,18,28]. Therefore, the results of the present study and other recent studies indicate that ToxoDB #3 is the major clonal *T. gondii* genotype circulating in northwestern China [17,18].

To our knowledge, this is the first report of *T. gondii* seroprevalence in house sparrows in China. The present work also provided new genetic information about *T. gondii* infection in house sparrows in northwestern China. The prevalence of *T. gondii* infection in birds has epidemiological significance because infection of the ground-foraging birds like sparrows suggests soil contamination with *T. gondii* oocysts. These birds are potential reservoirs and good indicators for *T. gondii* transmission.

ACKNOWLEDGMENTS

The project was supported, in part, by the National Natural Science Foundation of China (no. 31230073, 31172316, 31228022, and 31101812), the International Science and Technology Cooperation Project of Gansu Province (Grant no. 1204-WCGA023), the Science Fund for Creative Research Groups of Gansu Province (Grant no. 1210RJA006) and the National S & T Major Program (Grant no. 2012ZX10004220). The authors thank Dr. J. P. Dubey, Animal Parasitic Diseases Laboratory, Animal and Natural Resources Institute, Beltsville Agricultural Research Center, United States Department of Agriculture, USA for providing the *Toxoplasma gondii* MAT antigen. Associate Professor Chunlei Su at the Department of Microbiology, the University of Tennessee, Knoxville, USA is gratefully thanked for providing reference *T. gondii* strains and for constructive comments and suggestions on the draft manuscript.

REFERENCES

- Dubey JP. *Toxoplasmosis of Animals and Humans*, second Edition. CRC Press, Boca Raton, Florida. 2010; 46-48.
- Lee SE, Hong SH, Lee SH, Jeong YI, Lim SJ, Kwon OW, Kim SH, You YS, Cho SH, Lee WJ. Detection of ocular *Toxoplasma gondii* infection in chronic irregular recurrent uveitis by PCR. *Korean J Parasitol* 2012; 50: 229-231.
- Yang Z, Cho PY, Ahn SK, Ahn HJ, Kim TS, Chong CK, Hong SJ, Cha SH, Nam HW. A surge in the seroprevalence of toxoplasmosis among the residents of islands in Gangwha-gun, Incheon, Korea. *Korean J Parasitol* 2012; 50: 191-197.
- Robert-Gangneux F, Darde ML. Epidemiology of and diagnostic strategies for toxoplasmosis. *Clin Microbiol Rev* 2012; 25: 264-296.
- Gondim LS, Abe-Sandes K, Uzeda RS, Silva MS, Santos SL, Mota RA, Vilela SM, Gondim LF. *Toxoplasma gondii* and *Neospora caninum* in sparrows (*Passer domesticus*) in the Northeast of Brazil. *Vet Parasitol* 2010; 168: 121-124.
- Dubey JP, Graham DH, Blackston CR, Lehmann T, Gennari SM, Ragozo AM, Nishi SM, Shen SK, Kwok OC, Hill DE, Thulliez P. Biological and genetic characterisation of *Toxoplasma gondii* isolates from chickens (*Gallus domesticus*) from Sao Paulo, Brazil: unexpected findings. *Int J Parasitol* 2002; 32: 99-105.
- Darde ML, Bouteille B, Pestre-Alexandre M. Isoenzyme analysis of 35 *Toxoplasma gondii* isolates and the biological and epidemiological implications. *J Parasitol* 1992; 78: 786-794.
- Howe DK, Sibley LD. *Toxoplasma gondii* comprises three clonal lineages: correlation of parasite genotype with human disease. *J Infect Dis* 1995; 172: 1561-1566.
- Ajzenberg D, Banuls AL, Tibayrenc M, Darde ML. Microsatellite analysis of *Toxoplasma gondii* shows considerable polymorphism structured into two main clonal groups. *Int J Parasitol* 2002; 32: 27-38.
- Dubey JP, Sundar N, Gennari SM, Minervino AH, Farias NA, Ruas JL, dos Santos TR, Cavalcante GT, Kwok OC, Su C. Biologic and genetic comparison of *Toxoplasma gondii* isolates in free-range chickens from the northern Para state and the southern state Rio Grande do Sul, Brazil revealed highly diverse and distinct parasite populations. *Vet Parasitol* 2007; 143: 182-188.
- Dubey JP, Cortes-Vecino JA, Vargas-Duarte JJ, Sundar N, Velmurugan GV, Bandini LM, Polo LJ, Zambrano L, Mora LE, Kwok OC, Smith T, Su C. Prevalence of *Toxoplasma gondii* in dogs from Colombia, South America and genetic characterization of *T. gondii* isolates. *Vet Parasitol* 2007; 145: 45-50.
- Dubey JP, Sundar N, Hill D, Velmurugan GV, Bandini LA, Kwok OC, Majumdar D, Su C. High prevalence and abundant atypical genotypes of *Toxoplasma gondii* isolated from lambs destined for human consumption in the USA. *Int J Parasitol* 2008; 38: 999-1006.
- Khan A, Dubey JP, Su C, Ajioka JW, Rosenthal BM, Sibley LD. Genetic analyses of atypical *Toxoplasma gondii* strains reveal a fourth clonal lineage in North America. *Int J Parasitol* 2011; 41: 645-655.
- Dubey JP, Zhu XQ, Sundar N, Zhang H, Kwok OC, Su C. Genetic and biologic characterization of *Toxoplasma gondii* isolates of cats from China. *Vet Parasitol* 2007; 145: 352-356.
- Zhou P, Nie H, Zhang LX, Wang HY, Yin CC, Su C, Zhu XQ, Zhao JL. Genetic characterization of *Toxoplasma gondii* isolates from pigs in China. *J Parasitol* 2010; 96: 1027-1029.
- Zhou P, Sun XT, Yin CC, Yang JF, Yuan ZG, Yan HK, Zhu XQ, Zou FC. Genetic characterization of *Toxoplasma gondii* isolates from pigs in southwestern China. *J Parasitol* 2011; 97: 1193-1195.
- Zhou P, Zhang H, Lin RQ, Zhang DL, Song HQ, Su C, Zhu XQ. Genetic characterization of *Toxoplasma gondii* isolates from China. *Parasitol Int* 2009; 58: 193-195.
- Huang SY, Cong W, Zhou P, Zhou DH, Wu SM, Xu MJ, Zou FC, Song HQ, Zhu XQ. First report of genotyping of *Toxoplasma gon-*

- dii* isolates from wild birds in China. J Parasitol 2012; 98: 681-682.
19. Wu SM, Zhu XQ, Zhou DH, Fu BQ, Chen J, Yang JF, Song HQ, Weng YB, Ye DH. Seroprevalence of *Toxoplasma gondii* infection in household and stray cats in Lanzhou, northwest China. Parasit Vectors 2011; 4: 214.
 20. Wu SM, Huang SY, Fu BQ, Liu GY, Chen JX, Chen MX, Yuan ZG, Zhou DH, Weng YB, Zhu XQ, Ye DH. Seroprevalence of *Toxoplasma gondii* infection in pet dogs in Lanzhou, Northwest China. Parasit Vectors 2011; 4: 64.
 21. Dubey JP. *Toxoplasma gondii* infections in chickens (*Gallus domesticus*): prevalence, clinical disease, diagnosis and public health significance. Zoonoses Public Health 2010; 57: 60-73.
 22. Hill DE, Chirukandoth S, Dubey JP, Lunney JK, Gamble HR. Comparison of detection methods for *Toxoplasma gondii* in naturally and experimentally infected swine. Vet Parasitol 2006; 141: 9-17.
 23. Su C, Shwab EK, Zhou P, Zhu XQ, Dubey JP. Moving towards an integrated approach to molecular detection and identification of *Toxoplasma gondii*. Parasitology 2010; 137: 1-11.
 24. Lopes AP, Sargo R, Rodrigues M, Cardoso L. High seroprevalence of antibodies to *Toxoplasma gondii* in wild animals from Portugal. Parasitol Res 2011; 108: 1163-1169.
 25. Dubey JP, Patitucci AN, Su C, Sundar N, Kwok OC, Shen SK. Characterization of *Toxoplasma gondii* isolates in free-range chickens from Chile, South America. Vet Parasitol 2006; 140: 76-82.
 26. Dubey JP, Rajapakse RP, Wijesundera RR, Sundar N, Velmurugan GV, Kwok OC, Su C. Prevalence of *Toxoplasma gondii* in dogs from Sri Lanka and genetic characterization of the parasite isolates. Vet Parasitol 2007; 146: 341-346.
 27. Prestrud KW, Asbakk K, Mork T, Fuglei E, Tryland M, Su C. Direct high-resolution genotyping of *Toxoplasma gondii* in arctic foxes (*Vulpes lagopus*) in the remote arctic Svalbard archipelago reveals widespread clonal Type II lineage. Vet Parasitol 2008; 158: 121-128.
 28. Chen ZW, Gao JM, Huo XX, Wang L, Yu L, Halm-Lai F, Xu YH, Song WJ, Hide G, Shen JL, Lun ZR. Genotyping of *Toxoplasma gondii* isolates from cats in different geographic regions of China. Vet Parasitol 2011; 183: 166-170.

