



Molecular basis and evolutionary drivers of endosperm-based hybridization barriers

Heinrich Bente ¹ and Claudia Köhler ^{1,2,*}

¹ Department of Plant Reproductive Biology and Epigenetics, Max Planck Institute of Molecular Plant Physiology, Potsdam 14476, Germany

² Department of Plant Biology, Swedish University of Agricultural Sciences and Linnean Center for Plant Biology, Uppsala 75007, Sweden

*Author for correspondence: koehler@mpimp-golm.mpg.de

The author responsible for distribution of materials integral to the findings presented in this article in accordance with the policy described in the Instructions for Authors (<https://academic.oup.com/plphys/pages/General-Instructions>) is: Claudia Köhler (koehler@mpimp-golm.mpg.de).

Abstract

The endosperm, a transient seed tissue, plays a pivotal role in supporting embryo growth and germination. This unique feature sets flowering plants apart from gymnosperms, marking an evolutionary innovation in the world of seed-bearing plants. Nevertheless, the importance of the endosperm extends beyond its role in providing nutrients to the developing embryo by acting as a versatile protector, preventing hybridization events between distinct species and between individuals with different ploidy. This phenomenon centers on growth and differentiation of the endosperm and the speed at which both processes unfold. Emerging studies underscore the important role played by type I MADS-box transcription factors, including the paternally expressed gene *PHERES1*. These factors, along with downstream signaling pathways involving auxin and abscisic acid, are instrumental in regulating endosperm development and, consequently, the establishment of hybridization barriers. Moreover, mutations in various epigenetic regulators mitigate these barriers, unveiling a complex interplay of pathways involved in their formation. In this review, we discuss the molecular underpinnings of endosperm-based hybridization barriers and their evolutionary drivers.

Introduction

Angiosperms comprise approximately 300,000 extant species and represent the most species-rich plant clade, accounting for approximately 90% of all land plant species (Christenhusz and Byng 2016; Katz 2018). Their rapid rise to ecological dominance in the mid-Cretaceous remained the “abominable mystery” to Charles Darwin, as evident from his 1879 letter to Joseph Hooker (Darwin and Seward 1903). Understanding this enigmatic rise to prominence has remained a central objective in evolutionary biology, necessitating insights into the evolution of reproductive barriers that limit gene flow between populations. Much previous research has focused on comprehending prefertilization reproductive barriers, which prevent successful zygote formation through mechanisms like pollinator adaptations or pollen–pistil incompatibilities (Widmer et al. 2009; Hernández-Hernández et al. 2021). Recent advances

in the field of evolutionary biology have recognized postzygotic mechanisms as substantial drivers of plant speciation (Coughlan 2023). These mechanisms operate at various time points after fertilization, leading to impaired viability of hybrid seeds or seedlings or sterility of the resulting hybrids (He et al. 2023). In this review, we focus on the evolution of postzygotic barriers arising from the endosperm, particularly in the context of interspecies and interploidy hybridizations. We discuss the molecular mechanisms underlying these barriers and explore their impact on plant speciation.

Endosperm development

Seed development in flowering plants starts with double fertilization, whereby 1 of the 2 pollen-delivered sperm cells fertilizes the egg cell, initiating the formation of the embryo, and the other sperm cell fertilizes the central cell, giving rise to the

Received August 31, 2023. Accepted January 06, 2024. Advance access publication January 31, 2024

© The Author(s) 2024. Published by Oxford University Press on behalf of American Society of Plant Biologists.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

Open Access

ADVANCES

- Genetic data substantiate a model wherein AGLs function upstream of a regulatory cascade governing endosperm cellularization, making understanding of AGL regulation and their downstream targets of crucial relevance.
- Maternal Pol IV–dependent siRNAs play a pivotal role in suppressing AGL expression, thus fostering endosperm cellularization. Conversely, paternal Pol IV–dependent siRNAs exert a negative influence on endosperm cellularization, illustrating the antagonistic functions of maternal and paternal siRNAs in this process.
- The inhibitory effect of auxin on endosperm cellularization underscores the importance of unraveling auxin signaling within the endosperm.
- Endosperm cellularization serves to protect the embryo from desiccation, a process typically occurring during maturation. The precise role of ABA in this process remains in important open question.

endosperm (Dresselhaus et al. 2016). The endosperm is a unique feature of angiosperms, playing a crucial role as a nourishing tissue to support the growth and germination of the embryo (Doll and Ingram 2022). In most angiosperms, the endosperm is a triploid tissue with 2 maternal and 1 paternal genome copies. This triploid constitution arises since the central cell is a homodiploid cell (Dresselhaus et al. 2016). This increased maternal copy number may have evolved to regulate resource allocation to the embryo, possibly reducing the influence of paternally provided growth-promoting alleles and ensuring equal provisioning of seeds (Haig and Westoby 1989). The ancestral state of the endosperm, observed in sister lineages to all angiosperms such as Nymphaeales and Austrobaileyales, is diploid. Resource allocation in these species is typically mediated by the perisperm, a nutritive tissue derived from the maternal sporophyte (Floyd and Friedman 2000; Williams and Friedman 2002).

Endosperm development leads to the formation of a fully cellularized tissue surrounding the embryo. However, the process of endosperm cellularization is not uniform across all species. In most angiosperms, the endosperm does not become cellular from the start; instead, nuclear divisions occur initially without cellularization, resulting in the formation of a coenocyte. Subsequently, cellularization takes place after a defined number of mitotic cycles, followed by the differentiation of distinct cell types (Boisnard-Lorig et al. 2001; Sorensen et al. 2002; Berger 2003). This developmental transition is critical for embryo survival; impaired endosperm cellularization in *Arabidopsis thaliana* mutants is generally connected with embryo arrest (Grossniklaus et al. 1998; Kiyosue et al. 1999; Pignocchi et al. 2009; Hehenberger et al. 2012). In contrast to the nuclear

type of endosperm development, certain genera like *Solanum*, *Mimulus*, and *Nicotiana* exhibit the cellular endosperm type, where nuclear division and cellularization are coupled events (Lopes and Larkins 1993; Floyd and Friedman 2001). Additionally, an intermediate type of endosperm development exists, known as the helobial endosperm. After the first division of the fertilized central cell, 1 cell follows the nuclear type of development while the other follows the cellular type, resulting in a unique helobial endosperm. Helobial endosperm development is relatively rare but can be found in species of the Cabombaceae, Sabiaceae, and Saxifragaceae families (Geeta 2003).

The endosperm plays a crucial role as a nutrient reservoir for the embryo, and its reserves can be remobilized during embryo development and germination. While in monocots the endosperm is preserved in the mature seed, serving to support embryo germination, in most eudicots, the endosperm is consumed by the developing embryo as it grows (Li and Berger 2012).

The endosperm establishes reproductive isolation

Numerous studies conducted in the last century have reported the failure of producing viable crop hybrids when crossing related plant species due to the widespread occurrence of seed lethality (Brink and Cooper 1947; Williams and White 1976; Gill and Waines 1978; Johnston and Hanneman Jr 1982; Sukno et al. 1999). By histological observations and in vitro embryo rescue experiments, these studies identified abnormal endosperm development as the primary cause of hybrid seed lethality in response to interspecies crosses (Brink and Cooper 1947; Valentine and Woodell 1963; Sukno et al. 1999; Dinu et al. 2005; Roy et al. 2011). This phenomenon's prevalence across diverse plant taxa (Fig. 1; Coughlan 2023) suggests that endosperm-based hybrid seed lethality (later on referred to as endosperm-based barriers [EBBs]) represents a substantial mode of reproductive isolation in flowering plants.

Similar to interspecies crosses, crosses involving plants with differing numbers of chromosome sets (ploidy) also result in seed lethality, with the outcome dependent on the direction of the cross. To describe this phenomenon, we will employ the term “paternal excess seeds” for seeds resulting from crosses where the paternal parent has a higher ploidy, while “maternal excess seeds” refer to those resulting from the reciprocal cross direction. Both types of seeds, whether they exhibit maternal or paternal excess, show abnormalities in endosperm development. In species characterized by the nuclear mode of endosperm development, paternal excess leads to a delay in endosperm cellularization, whereas maternal excess has the opposite effect, causing precocious cellularization (Scott et al. 1998; Pennington et al. 2008; Sekine et al. 2013). In species with the cellular mode of endosperm development, paternal excess seeds are characterized by larger and more vacuolated endosperm cells, while maternal excess seeds have smaller and less vacuolated cells (Woodell and




Figure 1. Widespread occurrence of EBBs. The presence of endosperm corresponds closely with the emergence of robust hybridization barriers. In gymnosperms, embryo development is supported by the enlarged female gametophyte (denoted by parallel lines), whereas angiosperms have successfully integrated fertilization with the formation of a nourishing endosperm tissue. The endosperm may initiate as a cellular structure, as seen in ANA (Amborella, Nymphaeales, and Austrobaileyales) species that are sister lineages to all other angiosperms (indicated by the checkerboard pattern), or as a coenocyte where initial nuclear divisions are not synchronized with cellularization (depicted by the dotted pattern). References for each category of barriers are provided; for a more comprehensive exploration of reported barriers, we direct readers to Coughlan (2023). References in light green refer to barriers occurring in nuclear-type endosperm. Dark green references refer to cellular-type barriers. Checkmarks symbolize presence of EBBs. Brackets symbolize that insufficient data are available to determine whether the endosperm is affected in hybrid seeds. Question marks symbolize that no data are available yet.

Valentine 1961; von Wangenheim and Peterson 2004). Maternal and paternal excess-like phenotypes have also been observed in interspecies crosses where the ploidy of the parental plants is alike (Brink and Cooper 1947; Sukno et al. 1999; Dinu et al. 2005; Roy et al. 2011; Baek et al. 2016; Oneal et al. 2016; Roth et al. 2018a; Coughlan et al. 2020). These deviations in endosperm development observed in both interploidy and interspecies crosses suggest a common mechanistic basis for these phenomena. This notion gains substantial support from the observation that interspecies EBBs can be overcome by increasing the ploidy of 1 parent, highlighting the involvement of a dosage-sensitive component in both types of EBBs (Johnston and Hanneman Jr 1982; Lafon-Placette et al. 2017; Tonosaki et al. 2018).

The concept of the endosperm balance number (EBN) provides a theoretical framework for understanding the dosage sensitivity of the endosperm and offers a means to assess hybridization success between different species (Johnston et al. 1980). According to this concept, each species possesses an effective ploidy, which may differ from its actual (karyotypic) ploidy. The effective ploidy is determined through test crosses with a defined species, allowing the assessment of cross-compatibility with other species (Johnston et al. 1980; Städler et al. 2021). Only species with similar EBNs can produce viable seeds by maintaining an effective 2:1 maternal-to-paternal ratio in the endosperm, a characteristic feature in most angiosperms (Lin 1984; Scott et al. 1998; Leblanc et al. 2002). From now on, we will refer to maternal or paternal excess crosses when the maternal or paternal parent has the higher effective ploidy, respectively. The focus however will be on paternal excess crosses, as most molecular research has concentrated on unraveling the molecular mechanisms underlying EBBs originating from these specific types of crosses.

Genomic imprinting: An epigenetic phenomenon underpinning reproductive isolation

The nonreciprocal effects observed in interploidy and interspecies hybridizations defy Mendelian expectations, revealing that parental genomes exert varying influences on offspring. Genomic imprinting, an epigenetic phenomenon that induces alterations in gene expression based on parental origin, explains these nonreciprocal hybridization outcomes. In the context of flowering plants, genomic imprinting primarily occurs within the endosperm and substantially influences its growth (Rodrigues and Zilberman 2015; Satyaki and Gehring 2017; Batista and Köhler 2020).

Imprinted genes are epigenetically modified, a process that typically initiates within the gametes and persists following fertilization. This results in genes that are specifically expressed either maternally (MEGs) or paternally (PEGs). Imprinted genes are enriched for transposable elements (TEs) in their flanking regions (Gehring et al. 2009; Wolff et al. 2011; Rodrigues et al. 2013; Pignatta et al. 2014;

Hatorangan et al. 2016), with many of them subject to regulation by the DNA glycosylase DEMETER (DME). DME excises 5'-methylcytosine, and its activity in the central cell of the female gametophyte leads to the activation of maternal alleles for certain MEGs that remain hypomethylated after fertilization (Choi et al. 2002; Pignatta et al. 2014; Park et al. 2016). Conversely, for PEGs, DNA hypomethylation facilitates Polycomb Repressive Complex2 (PRC2)-directed trimethylation at lysine 27 of histone H3 (H3K27me3), resulting in the suppression of maternal alleles of PEGs (Moreno-Romero et al. 2016, 2019).

The relationship between imprinted genes and flanking TEs initially led to the hypothesis that imprinting primarily functions to repress TEs (Gehring et al. 2009; Hsieh et al. 2009). However, recent research challenges this notion, revealing that numerous TEs harbor transcription factor binding sites. This discovery implies that DME-mediated demethylation plays a vital role in kick-starting gene expression by granting access to transcription factors (Schmitz et al. 2013; Batista et al. 2019b; Khouider et al. 2021). Hence, it appears that endosperm imprinting results from a process enabling the transcriptional activation of MEGs through DME-mediated demethylation, with some of these MEGs being crucial for seed development. Notably, *MEDEA* and *FERTILIZATION INDEPENDENT SEED2* (*FIS2*), encoding subunits of the PRC2, are essential MEGs that subsequently contribute to the suppression of the maternal alleles of PEGs (Zhang et al. 2014; Moreno-Romero et al. 2019).

Aside from the molecular explanations of how genomic imprinting is established, there are many considerations on the possible evolutionary advantages of this phenomenon. Among those, the “parental conflict” or “kinship theory” (Trivers 1974; Haig and Westoby 1989, 1991) is the most widely accepted evolutionary theory that provides a theoretical framework for the selection of imprinted genes. According to this theory, conflict arises because siblings compete for limited maternal resources. Since the maternal parent is equally related to its offspring, balanced provisioning of the progeny maximizes maternal fitness (Haig and Westoby 1989). Multiple paternity increases this conflict, since each offspring is more closely related to itself than to its siblings (Trivers 1974; Haig and Westoby 1989, 1991). This conflict drives the evolution of paternally derived alleles that enhance resource acquisition, while maternally derived alleles are selected to equalize nutrient allocation (Queller 1983; Haig and Westoby 1989, 1991). Consequently, MEGs and PEGs are predicted to be frequently associated with opposing functions in the endosperm, either restricting or promoting endosperm growth (Haig and Westoby 1989, 1991; Costa et al. 2012). Supporting this theory, an increased maternal genome dosage leads to reduced endosperm growth and smaller seeds, whereas an increased paternal genome dosage produces the opposite effect (Brink and Cooper 1947; Scott et al. 1998; von Wangenheim and Peterson 2004; Stoute et al. 2012; Sekine et al. 2013; Rebernig et al. 2015; Roth et al. 2019). Interestingly, the chalazal region of the

endosperm is particularly affected by changes in parental genome dosage (Scott et al. 1998; Sandstedt and Sweigart 2022). This region of the endosperm serves as an interface between maternal and endosperm tissues, playing a pivotal role in determining the extent of nutrient flow from the maternal plant to the embryo and endosperm (Povilus and Gehring 2022). PEGs are most highly expressed in the chalazal endosperm, pointing at a functional role of PEGs in regulating nutrient transfer by affecting the growth of the chalazal endosperm (Picard et al. 2021).

The cross-direction–dependent seed phenotypes observed in interspecies and interploidy hybridizations have led to the hypothesis that imprinted genes play a causal role in establishing EBBs (Haig and Westoby 1989, 1991; Moore 2001; Gutierrez-Marcos et al. 2003). Genetic studies provide support for this notion, demonstrating that mutations in several PEGs can suppress triploid seed abortion (referred to as triploid block) in *Arabidopsis* (Kradolfer et al. 2013b; Wolff et al. 2015; Huang et al. 2017; Batista et al. 2019b). Suppression of the triploid block by mutations in PEGs consistently corresponds with the restoration of endosperm cellularization (Wolff et al. 2015; Batista et al. 2019b), underscoring the critical role of endosperm cellularization in embryo growth. Many PEGs associated with the triploid block are generally highly upregulated in triploid seeds. However, this upregulation is not attributed to a breakdown of imprinting but rather to enhanced expression of the active paternal alleles (Batista et al. 2019b). In contrast, seed arrest in interspecies hybrids is associated with disruption of imprinting in hybrid endosperm (Josefsson et al. 2006; Burkart-Waco et al. 2015; Florez-Rueda et al. 2016; Kinser et al. 2021). Nevertheless, while the cause for the deregulation of imprinted genes in interploidy and interspecies hybrids may differ, the consequences of their deregulation are similar and lead to shared endosperm phenotypes with lethal consequences.

Small RNAs and DNA methylation in the establishment of reproductive isolation

Paternally inherited mutations in the RNA-directed DNA methylation (RdDM) pathway suppress the triploid block (Erdmann et al. 2017; Martinez et al. 2018; Satyaki and Gehring 2019; Wang et al. 2021), similar to mutants in the maintenance methyltransferase MET1 or chemically induced hypomethylation (Schatlowski et al. 2014; Huc et al. 2021). The RdDM pathway initiates with the activity of RNA polymerase IV (Pol IV), which generates short transcripts that are transformed into double-stranded RNAs by RNA-dependent RNA POLYMERASE 2. These double-stranded RNAs are then cleaved into small interfering RNAs (siRNAs) of 21 to 24 nucleotides (nts) by DICER-like proteins that can be loaded into different ARGONAUTE proteins (Cuerda-Gil and Slotkin 2016). In pollen, Pol IV is required for generating 21- to 22-nt epigenetically activated siRNAs (easiRNAs) as well as 24-nt siRNAs. easiRNAs originate from TEs within the vegetative nucleus and are likely

transported to sperm cells afterwards (Slotkin et al. 2009; Martinez et al. 2016). Notably, the formation of easiRNAs relies on microRNA845a (miR845a) and miR845b, 2 abundant pollen miRNAs that target long terminal repeat retrotransposons at their reverse transcription primer binding site. In line with the importance of easiRNAs in the triploid block, a mutation in *MIR845b* can alleviate this block (Borges et al. 2018).

Pol IV–generated siRNAs recruit the de novo DOMAINS REARRANGED METHYLTRANSFERASE2, which establishes DNA methylation in all sequence contexts (Chow et al. 2020). Interestingly, inbreeding of RdDM mutants enhances their ability to suppress the triploid block and is linked with a gradual decrease in DNA methylation over generations. In contrast to other RdDM pathway mutants, *nrpd1* (affected in the large subunit of Pol IV, NUCLEAR RNA POLYMERASE D1) exhibits a suppressive effect in the first generation of inbreeding (Wang et al. 2021), suggesting that the loss of easiRNAs and paternal DNA methylation can independently suppress the triploid block. Recent findings further support this notion, revealing that the depletion of easiRNAs in the vegetative cell of pollen (but not in sperm cells) can suppress the triploid block (Pachamuthu et al. 2023). Whether paternally delivered easiRNAs are causally responsible for DNA methylation changes in the endosperm as previously proposed (Martinez et al. 2018) remains to be investigated.

Despite that paternally inherited *nrpd1* can suppress the triploid block, the transcriptome changes in triploid seeds inheriting *nrpd1* through pollen remain relatively mild. Among the genes that continue to be upregulated in triploid *nrpd1* seeds are several PEGs previously identified to suppress the triploid block when mutated (Martinez et al. 2018; Satyaki and Gehring 2019). There are 2 possible hypotheses to explain this data: either Pol IV operates in a parallel pathway to the known suppressor genes or it acts downstream of these genes. Further research is required to distinguish between these hypotheses.

A link between the dosage of siRNAs and gene deregulation in hybrid endosperm has also been established in *Solanum*, where reciprocal hybrid seeds exhibit reduced siRNA levels corresponding with increased gene expression at specific loci potentially targeted by siRNAs (Florez-Rueda et al. 2021). Similarly, *Capsella* paternal excess hybrid endosperm shows strongly depleted DNA methylation, which associates with chromatin decondensation (Dziasek et al. 2021) and loss of siRNAs that depend on maternal Pol IV function (Dziasek et al. 2023). The loss of chromatin condensation in *Capsella* hybrid endosperm is associated with increased gene expression, particularly near heterochromatic pericentromeric regions, including many targets of type I MADS-box transcription factors (referred to as AGAMOUS-LIKE TFs or AGLs). These AGLs are consistently upregulated in paternal excess interploidy and interspecies hybrids (Josefsson et al. 2006; Erilova et al. 2009; Ishikawa et al. 2010; Burkart-Waco et al. 2013; Sekine et al. 2013; Rebernig et al. 2015; Roth et al. 2019; Bjerkan et al. 2020; Dziasek et al. 2021). AGL

upregulation is likely causally relevant in establishing EBBs, since mutants in several AGLs show weakened interploidy and interspecies barriers (Josefsson et al. 2006; Walia et al. 2009; Hehenberger et al. 2012; Batista et al. 2019b). Maternal siRNAs inversely correspond with AGL expression (Lu et al. 2012; Kirkbride et al. 2019; Dziasek et al. 2023), suggesting a connection between the loss of maternal siRNAs, DNA methylation, and increased expression of AGLs. These Pol IV-dependent siRNAs exhibit a proportional increase corresponding to the ploidy of the maternal genome. Consequently, maternal excess in Arabidopsis leads to heightened accumulation of siRNAs, while paternal excess results in diminished siRNA abundance, corresponding with increased AGL expression (Lu et al. 2012; Kirkbride et al. 2019). Notably, maternal Pol IV-dependent siRNAs, originating from a few loci in high abundance, have been observed across various flowering plant species. The loci responsible for producing these siRNAs appear to evolve rapidly (Grover et al. 2020), hinting at a potential role in establishing reproductive barriers between species. Where exactly maternal Pol IV-dependent siRNAs are produced remains to be established; while they have been proposed to be predominantly maternally produced in *Brassica* (Grover et al. 2018, 2020), biparental production was found in Arabidopsis (Satyaki and Gehring 2022). Collectively, alterations in the dosage of maternal and paternal Pol IV-dependent siRNAs impact EBBs. However, dependent on their parental origin, these siRNAs are generated from distinct loci and influence different targets, in line with their antagonistic effects on gene expression and endosperm development (Satyaki and Gehring 2022).

Epigenetic regulators as key players in overcoming the triploid block

PHERES1 (PHE1) is a highly upregulated AGL in triploid Arabidopsis seeds that directly binds to the promoter region of PEGs, many of those also being highly upregulated in triploid seeds (Batista et al. 2019b). Among them, ADMETOS (ADM), SU(VAR)3-9 HOMOLOG 7 (SUVH7), PICKLE RELATED2 (PKR2), and PEG2 have been identified to suppress the triploid block when mutated (Kradolfer et al. 2013a; Wolff et al. 2015; Huang et al. 2017; Fig. 2). ADM physically interacts with SUVH9 and various AT HOOK-LIKE (AHL) proteins. Mutations in *suvh9* and *ahl10*, similar to *adm*, effectively suppress the triploid block, suggesting the 3 genes act in the same pathway (Jiang et al. 2017). SUVH9 has the capacity to bind to methylated DNA, facilitating the recruitment of DNA-dependent RNA polymerase V (Pol V) to chromatin (Johnson et al. 2014; Liu et al. 2014b). Simultaneously, AHLs bind to AT-rich chromosomal regions, inducing increased levels of H3K9me2 via unknown mechanisms (Yun et al. 2012; Xu et al. 2013). Notably, in triploid seeds, ADM-dependent mislocalization of H3K9me2 is targeted at AT-rich small TEs, inversely correlating with CHH methylation. Recent work has shown that the catalytic domain of SUVH9 is essential for its function (Parent et al. 2021), suggesting that

AHL10-ADM-SUVH9 may collectively recruit H3K9me2 to TEs in the endosperm, potentially leading to the ectopic recruitment of H3K9me2 in triploid seeds (Jiang et al. 2017). This ectopic recruitment of H3K9me2 inversely correlates with CHH methylation, suggesting that AHL10-ADM-SUVH9-mediated H3K9me2 at specific TEs opposes the RdDM pathway in triploid seeds, in line with previous reports on the inhibitory effect of H3K9me2 on the RdDM pathway (Schoft et al. 2009; Zemach et al. 2013; Gent et al. 2014).

The roles of SUVH7 and PKR2 in relation to the triploid block are yet to be fully elucidated. Given that PKR2 shares homology with PICKLE (PKL), which acts in the RdDM pathway by supporting the accumulation of Pol V-dependent transcripts (Yang et al. 2017), it is plausible that PKR2 serves a related function. Similar to other mutants in the RdDM pathway (Wang et al. 2021), the suppressive effect of *pk2* appears to require several generations of inbreeding (Huang et al. 2017). The *pk2* mutant was not initially identified as a suppressor when tested after introgression into the meiotic mutant *omission of second division 1* (*osd1*; Wolff et al. 2015), in contrast to its suppressive effect following inbreeding (Huang et al. 2017).

The suppressive effect of *peg2* on the triploid block has been linked to the ability of the PEG2 transcript to sequester the TE-derived siRNA854 (Wang et al. 2018). This suggests that a reduced abundance of siRNA854 has functional consequences for endosperm development. siRNA854 regulates the abundance of the stress granule-associated protein OLIGOURIDYLATE BINDING PROTEIN 1B (UBP1b) by translational repression (McCue et al. 2012), which is consistent with *ubp1b* partially suppressing the triploid block (Wang et al. 2018). UBP1b has been proposed to play a role in the translational repression of *Athila* retrotransposon transcripts (McCue et al. 2012). However, whether this mechanism is mechanistically connected to the triploid block requires further investigation.

Collectively, many of the triploid block suppressors identified in Arabidopsis encode for chromatin regulators with potential roles in TE silencing or the establishment of heterochromatin (Wolff et al. 2015; Jiang et al. 2017; Yang et al. 2017; Satyaki and Gehring 2019; Wang et al. 2021). Similarly, many genes responsible for hybrid incompatibility in *Drosophila* encode for dosage-sensitive heterochromatin-interacting proteins or components of the PIWI pathway that silences TEs (Brideau et al. 2006; Bayes and Malik 2009; Thomae et al. 2013; Parhad et al. 2017). In *Drosophila*, TEs have been implicated in causing intraspecies incompatibility (Castillo and Moyle 2022). Nevertheless, like in plants, the mechanisms through which TEs and deregulated chromatin regulators induce lethality remain to be fully established.

The role of auxin in antagonizing endosperm cellularization

Beyond its interactions with various epigenetic regulators, PHE1 also binds to auxin biosynthesis genes YUCCA10 (YUC10) and TRYPTOPHAN AMINOTRANSFERASE RELATED




Figure 2. Molecular pathways underlying endosperm-based interploidy and interspecies barriers. Maternal Pol IV–dependent siRNAs act as dosage-dependent repressors of AGL MADS-box genes. Increased dosage of siRNAs in maternal excess crosses corresponds with strong AGL repression (reflected by blunt-ended arrows; thickness of the arrows corresponds to intensity of repression) and early endosperm cellularization, while decreased siRNA dosage in paternal excess crosses corresponds with increased AGL abundance and late endosperm cellularization. High and low AGL dosage causes increased or decreased expression of downstream AGL targets, respectively, including auxin biosynthesis genes. Downstream direct or indirect targets of this signaling cascade are genes involved in pectin metabolism, including pectin methyltransferase (PME) genes. Demethylsterification exposes pectin to pectin-degrading enzymes, thereby establishing a connection between the expression level of AGLs and the timing of endosperm cellularization. Abundance of AGLs and their downstream targets in response to hybridization are depicted by red (low), gray (balanced), and blue (high) arrows. Light and dark green endosperm regions mark micropylar and chalazal domains, respectively. ADM, ADMETOS; AGLs, AGAMOUS-LIKE genes; EBN, endosperm balance number; IKU2, HAIKU2; MINI3, MINISEED3; NRPD1, NUCLEAR RNA POLYMERASE D1; PEG2, PATERNALLY EXPRESSED GENE2; UBP1, OLIGOURIDYLATE BINDING PROTEIN 1B; PKR2, PICKLE RELATED2; SUVH7, SU(VAR)3–9 HOMOLOG 7; TAR1, TAA1-RELATED1; TEs, transposable elements; YUC10, YUCCA10.

1 (TAR1). Activated by PHE1, YUC10 and TAR1 are highly overexpressed in triploid seeds, concomitant with increased auxin activity (Batista et al. 2019a, 2019b; Fig. 3).

Elevated auxin levels wield a negative impact on endosperm cellularization, establishing a direct link between the upregulation of PHE1, increased auxin production, and the failure of endosperm cellularization (Batista et al. 2019a). Interestingly, increasing auxin levels, specifically in the seed coat, also block endosperm cellularization, implicating the seed coat as a target of auxin action that indirectly impacts endosperm cellularization. Notably, in triploid seeds, increased auxin activity is primarily detected in the seed coat rather than the endosperm, hinting at auxin's production in the endosperm and its subsequent transportation to the seed coat, drives seed coat expansion (Batista et al. 2019a). AGL62 emerges as a crucial player in regulating auxin transport from the endosperm to the seed coat, potentially by modulating the expression of the ABCB-type transporter gene PGP10 (Figueiredo et al. 2016) and regulating auxin biosynthesis (Guo et al. 2022). Supporting this notion is the observation that the *agl62* mutant exhibits partial suppression of EBBs (Walia et al. 2009; Hehenberger et al. 2012). This intricate regulation of auxin transport between the endosperm and seed coat potentially underpins the mechanism behind the triploid block.

The suppressive effect of *transparent testa* (*tt*) mutants, affecting flavonoid biosynthesis in the innermost layer of the inner integument, offers another potential link between auxin and the triploid block. Flavonoids impact polar auxin transport by interacting with auxin transporters such as PGPs (Peer and Murphy 2007). Several mutants within the TT pathway, including *transparent testa glabra2* (*ttg2*), *tt4*, *tt7*, *tt8*, and *tt13*, exhibit the ability to suppress the triploid block (Dilkes et al. 2008; Scott et al. 2013; Doughty et al. 2014; Zumajo-Cardona et al. 2023), although the precise mechanisms remain elusive. Intriguingly, the efficiency of this suppression varies among the *tt* mutants, with *tt8* causing complete suppression, while others have a milder effect (Zumajo-Cardona et al. 2023), hinting at the involvement of multiple pathways regulated by TT genes.

A plausible model suggests that type I MADS-box transcription factors, such as PHE1 and AGL62, govern auxin homeostasis within the endosperm. This regulation includes binding to the promoter regions of auxin biosynthesis genes, resulting in increased auxin production in direct proportion to the increased dosage of AGLs (Figueiredo et al. 2016; Batista et al. 2019a). Auxin reduces tissue rigidity in the shoot apical meristem by initiating the demethylesterification of homogalacturonan, a component of pectin. This process renders it susceptible to enzymatic depolymerization, facilitating gel formation and degradation (Braybrook and Peaucelle 2013; Wormit and Usadel 2018). Genes linked to demethylesterification and degradation of pectin are significantly overexpressed in triploid hybrid seeds that fail to undergo endosperm cellularization, suggesting that auxin regulates pectin metabolism with consequences on endosperm cellularization (Rebernig et al. 2015; Wolff et al. 2015).




Figure 3. Regulation of endosperm cellularization by antagonizing activities of auxin and ABA. ABA activates the downstream transcription factor ABI5, which, in turn, is stabilized by mobile factors such as TFL1, which is transported from the chalazal region to the peripheral endosperm. The mobilization of TFL1 is facilitated by RAN1. ABI5, under the influence of ABA, exerts its regulatory function by downregulating SHB1. SHB1 is a positive regulator of 2 critical genes, MINI3 and IKU2, both of which are direct targets of PHE1. MINI3 and IKU2 act as positive regulators of endosperm proliferation (symbolized by red dots). The repression of MINI3 and IKU2 via the ABA pathway likely promotes endosperm cellularization. Failure of endosperm cellularization can trigger an osmotic stress response in the embryo. Red arrows in this network represent the auxin-PHE1 pathway, and blue arrows highlight the ABA signaling pathway. Upward-pointing arrows reflect increased expression or activity, while downward-pointing arrows reflect decreased expression or activity. Arrows with blunt ends reflect repressive activity. Light and dark green endosperm regions mark micropylar and chalazal domains, respectively. ABI5, ABSCISIC ACID-INSENSITIVE5; IKU2, HAIKU2; MINI3, MINISEED3; PHE1, PHERES1; RAN1, Ras-related nuclear GTPase1; SHB1, SHORT HYPOCOTYL UNDER BLUE1; TFL1, TERMINAL FLOWER1.

Absciscic acid facilitates endosperm cellularization and enhances triploid seed viability

The precise timing of endosperm cellularization represents a critical factor for embryo survival, a phenomenon consistently disrupted in interploidy and interspecies hybrids (Muntzing 1933; Brink and Cooper 1947; Woodell and Valentine 1961; Scott et al. 1998; Dilkes et al. 2008; Hehenberger et al. 2012). The reasons underpinning the necessity of endosperm cellularization for embryo survival have

long been a matter of conjecture. One proposed hypothesis suggests that the failure of endosperm cellularization results in an inadequate supply of sucrose to the embryo, as the central vacuole remains the primary resource sink in the seed (Lafon-Placette and Kohler 2014). However, recent research suggests an alternative perspective: embryos surrounded by noncellularized endosperm elicit an embryo-specific osmotic stress response akin to the seed maturation process. This implies that endosperm cellularization serves to protect the embryo from desiccation, a process typically occurring during maturation (Kozaki and Aoyanagi 2022; Xu et al. 2023). Supporting this view, mutations in abscisic acid (ABA) biosynthesis and signaling exacerbate the triploid block, while elevated endogenous ABA levels induce endosperm cellularization and suppress the arrest of embryo growth (Xu et al. 2023). Nevertheless, whether the suppressive effect of increased ABA is a consequence of ABA-induced endosperm cellularization or an enhanced tolerance of the embryo to desiccation remains to be established.

Interestingly, exposure to cooler temperatures during early seed maturation leads to ABA retention in the endosperm upon desiccation (Chen et al. 2021). Similarly, cold treatment enhances the viability of *Arabidopsis* hybrid seeds impaired in endosperm cellularization (Bjerkan et al. 2020), suggesting that the survival of hybrid seeds in response to cold is linked to elevated ABA levels.

The process of ABA-mediated endosperm cellularization hinges on the downregulation of *SHORT HYPOCOTYL UNDER BLUE1* (*SHB1*) by *ABSCISIC ACID-INSENSITIVE5* (*ABI5*). *ABI5*-deficient plants exhibit delayed endosperm cellularization and develop enlarged seeds (Cheng et al. 2014), which, notably, intensify the triploid block (Xu et al. 2023). *SHB1*, in turn, upregulates *MINISEED3* (*MINI3*) and the LRR receptor kinase gene *HAIKU2* (*IKU2*), both of which are direct targets of *PHE1* and exert a positive influence on endosperm proliferation (Garcia et al. 2003; Luo et al. 2005; Batista et al. 2019a).

Recent investigations have unveiled the interaction between *ABI5* and *TERMINAL FLOWER1* (*TFL1*), a mobile regulator generated in the chalazal endosperm that then relocates to the peripheral endosperm to stabilize *ABI5* (Zhang et al. 2020). The movement of *TFL1* is modulated by Ras-related nuclear GTPases (*RANs*) previously implicated in seed size regulation (Liu et al. 2014a). A plausible model posits that ABA activates *ABI5*, which is subsequently stabilized by *TFL1* and *RAN1*. *ABI5* then represses *SHB1*, which in turn restrains *IKU2* and *MINI3*, recognized regulators of endosperm proliferation (Fig. 3). This intricate network is juxtaposed with *PHE1*, which binds to *IKU2* and *MINI3*, likely promoting their transcription (Batista et al. 2019b).

These findings underscore the convergence of multiple pathways on *MINI3* and *IKU2*, with 1 pathway positively regulated by auxin and *PHE1* and the other negatively regulated by ABA, *ABI5*, and *SHB1*. While the loss of *MINI3* function does not impact the triploid block (Batista et al. 2019a), a mutation in the *MINI3* interaction partner *IKU1* enhances the survival of *Arabidopsis* interspecies hybrids

(Burkart-Waco et al. 2013), suggesting that the dosage of components within the *IKU1*-*IKU2*-*MINI3* pathway plays a pivotal role in erecting hybridization barriers by affecting endosperm cellularization.

Understanding the relevance and drivers of endosperm-mediated reproductive barriers

As elaborated in the preceding sections, the establishment of reproductive barriers mediated by the endosperm is intricately linked to the dysregulation of imprinted genes, particularly PEGs (Wolff et al. 2015; Florez-Rueda et al. 2016; Batista et al. 2019b; Kinser et al. 2021). In line with theoretical expectations, species characterized by outbreeding tendencies are presumed to experience heightened parental conflict, thus predicting an increased abundance or elevated expression of imprinted genes (Brandvain and Haig 2005). Indeed, empirical evidence supports this prediction, as outbreeding species tend to exhibit higher numbers or levels of PEGs compared to their inbreeding relatives (Klosinska et al. 2016; Lafon-Placette et al. 2018).

This augmented prevalence and expression of PEGs in outbreeders could account for the robust EBBs that often manifest between inbreeding and outbreeding species. This phenomenon has been theoretically framed as the “weak inbreeder/strong outbreeder” (WISO) hypothesis (Brandvain and Haig 2005). The validity of the WISO hypothesis is substantiated by extensive evidence, with documented EBBs observed between inbreeders and outbreeders in plant genera such as *Arabidopsis*, *Arabis*, and *Capsella* (Brandvain and Haig 2005; Lafon-Placette and Kohler 2016; Lafon-Placette et al. 2018; İltaş et al. 2021; Petrén et al. 2023). Similar to differences in breeding mode, also differences in effective population size can lead to strong reproductive barriers. A decrease in population size is anticipated to diminish conflicts among siblings as mating partners become highly related, akin to the impact of selfing. Correspondingly, within outbreeding species of *Solanum* and *Mimulus*, a smaller population size (reflected by lower nt diversity) is associated with a lower EBN (Roth et al. 2018a, 2019; Coughlan et al. 2020; Sandstedt et al. 2021; Sandstedt and Sweigart 2022). Interestingly, in *Solanum*, out of 3 tested species, the one with the highest EBN has the highest number of PEGs (Roth et al. 2018b, 2019). This finding underscores a potential connection between number of PEGs, intensity of parental conflict, and strength of EBBs.

Genomic imprinting is a dynamically evolving phenomenon, characterized by low conservation of imprinted loci across species (Waters et al. 2013; Pignatta et al. 2014; Hatorangan et al. 2016; Klosinska et al. 2016; Chen et al. 2018). In alignment with the hypothesized causal role of PEGs in the establishment of reproductive barriers, EBBs themselves exhibit a propensity for rapid evolution (Rebernik et al. 2015; Oneal et al. 2016; Roth et al. 2018a; Coughlan et al. 2020; İltaş et al. 2021). The exploration of the functional role of imprinted genes in the context of interspecies hybridization barriers promises to be an exciting avenue for future research.

OUTSTANDING QUESTIONS

- What is the functional role of paternal Pol IV–dependent siRNAs (easiRNAs) in the establishment of reproductive barriers?
- Are maternal Pol IV–dependent siRNAs transmitted from maternal sporophytic tissues to the endosperm, and what are the underlying mechanisms of this process?
- What are the epistatic relationships among the identified suppressor genes associated with hybridization barriers?
- At what pace do hybridization barriers evolve?

Another compelling research direction lies in the relationship between the strength of EBBs and the mode of endosperm development. Notably, robust barriers are encountered in species characterized by both nuclear and cellular endosperm development, suggesting that the mode of endosperm development itself does not substantially influence barrier strength. Furthermore, the ploidy level of the endosperm does not appear to be the decisive factor, as evidenced by the robust interploidy barriers observed in species like evening primrose (*Oenothera hookeri*), which have a diploid endosperm (von Wangenheim 1962). Instead, the key determinant seems to be the role of the endosperm in provisioning the embryo, a role that varies among species. In species where the endosperm plays a pivotal role in embryo provisioning, strong EBBs tend to be prevalent. Conversely, barriers are generally weaker in species where the endosperm has a relatively minor role in this context, such as in Nymphaeales, which rely on the maternal sporophytic nucellus for seed storage (Les and Philbrick 1993; Povilus et al. 2018). An intriguing question for future investigation pertains to whether the presence of weak EBBs is a recurring pattern among angiosperms with reduced endosperms.

In conclusion, our comprehension of the genetic underpinnings of EBBs has made substantial progress in recent years, shedding light on intriguing facets while simultaneously unveiling additional inquiries (see Outstanding Questions). Tackling these emerging questions collectively as a scientific community promises to be an exciting endeavor for the future.

Acknowledgments

We thank Gesa Hoffmann for critically reading this manuscript and support in generating the figures. We furthermore thank Yichun Qiu, Katarzyna Dziasek, and Varsha Vasudevan for helpful suggestions to improve the manuscript.

Author contributions

H.B. and C.K. conceived, wrote, and edited the article.

Funding

This work was financially supported by the Max Planck Society.

Conflict of interest statement. None declared.

References

- Baek YS, Royer SM, Broz AK, Covey PA, López-Casado G, Nuñez R, Kear PJ, Bonierbale M, Orillo M, van der Knaap E, et al. Interspecific reproductive barriers between sympatric populations of wild tomato species (*Solanum* section *Lycopersicon*). *Am J Bot.* 2016;**103**(11):1964–1978. <https://doi.org/10.3732/ajb.1600356>
- Batista RA, Figueiredo DD, Santos-Gonzalez J, Köhler C. Auxin regulates endosperm cellularization in *Arabidopsis*. *Genes Dev.* 2019a;**33**(7–8):466–476. <https://doi.org/10.1101/gad.316554.118>
- Batista RA, Köhler C. Genomic imprinting in plants-revisiting existing models. *Genes Dev.* 2020;**34**(1–2):24–36. <https://doi.org/10.1101/gad.332924.119>
- Batista RA, Moreno-Romero J, Qiu Y, van Boven J, Santos-Gonzalez J, Figueiredo DD, Köhler C. The MADS-box transcription factor PHERES1 controls imprinting in the endosperm by binding to domesticated transposons. *Elife* 2019b;**8**:e50541. <https://doi.org/10.7554/elife.50541>
- Bayes JJ, Malik HS. Altered heterochromatin binding by a hybrid sterility protein in *Drosophila* sibling species. *Science* 2009;**326**(5959):1538–1541. <https://doi.org/10.1126/science.1181756>
- Berger F. Endosperm: the crossroad of seed development. *Curr Opin Plant Biol.* 2003;**6**(1):42–50. <https://doi.org/10.1016/S1369526602000043>
- Bjerkkan KN, Hornslien KS, Johannessen IM, Krabberød AK, van Ekelburg YS, Kalantarian M, Shirzadi R, Comai L, Brysting AK, Bramsiepe J, et al. Genetic variation and temperature affects hybrid barriers during interspecific hybridization. *Plant J.* 2020;**101**(1):122–140. <https://doi.org/10.1111/tpj.14523>
- Boisnard-Lorig C, Colon-Carmona A, Bauch M, Hodge S, Doerner P, Bancharel E, Dumas C, Haseloff J, Berger F. Dynamic analyses of the expression of the histone::YFP fusion protein in *Arabidopsis* show that syncytial endosperm is divided in mitotic domains. *Plant Cell.* 2001;**13**(3):495–509. <https://doi.org/10.1105/tpc.13.3.495>
- Borges F, Parent JS, van Ex F, Wolff P, Martinez G, Köhler C, Martienssen RA. Transposon-derived small RNAs triggered by miR845 mediate genome dosage response in *Arabidopsis*. *Nat Genet.* 2018;**50**(2):186–192. <https://doi.org/10.1038/s41588-017-0032-5>
- Brandvain Y, Haig D. Divergent mating systems and parental conflict as a barrier to hybridization in flowering plants. *Am Nat.* 2005;**166**(3):330–338. <https://doi.org/10.1086/432036>
- Braybrook SA, Peaucelle A. Mechano-chemical aspects of organ formation in *Arabidopsis thaliana*: the relationship between auxin and pectin. *PLoS One* 2013;**8**(3):e57813. <https://doi.org/10.1371/journal.pone.0057813>
- Brideau NJ, Flores HA, Wang J, Maheshwari S, Wang X, Barbash DA. Two Dobzhansky-Muller genes interact to cause hybrid lethality in *Drosophila*. *Science* 2006;**314**(5803):1292–1295. <https://doi.org/10.1126/science.1133953>
- Briggs BG. Interspecific hybridization in the *Ranunculus lappaceus* group. *Evolution* 1962;**16**(3):372–390. <https://doi.org/10.2307/2406286>
- Brink RA, Cooper DC. The endosperm in seed development. *Bot Rev.* 1947;**132**(8):423–541. <https://doi.org/10.1007/BF02861548>
- Burkart-Waco D, Ngo K, Dilkes B, Josefsson C, Comai L. Early disruption of maternal-zygotic interaction and activation of defense-like responses in *Arabidopsis* interspecific crosses. *Plant Cell.* 2013;**25**(6):2037–2055. <https://doi.org/10.1105/tpc.112.108258>
- Burkart-Waco D, Ngo K, Lieberman M, Comai L. Perturbation of parentally biased gene expression during interspecific hybridization.

- PLoS One 2015;10(2):e0117293. <https://doi.org/10.1371/journal.pone.0117293>
- Castillo DM, Moyle LC.** Hybrid incompatibility between *Drosophila virilis* and *D. lummei* is stronger in the presence of transposable elements. *J Evol Biol.* 2022;35(10):1319–1334. <https://doi.org/10.1111/jeb.14079>
- Chen C, Li T, Zhu S, Liu Z, Shi Z, Zheng X, Chen R, Huang J, Shen Y, Luo S, et al.** Characterization of imprinted genes in rice reveals conservation of regulation and imprinting with other plant species. *Plant Physiol.* 2018;177(4):1754–1771. <https://doi.org/10.1104/pp.17.01621>
- Chen X, Yoong F-Y, O'Neill CM, Penfield S.** Temperature during seed maturation controls seed vigour through ABA breakdown in the endosperm and causes a passive effect on DOG1 mRNA levels during entry into quiescence. *New Phytol.* 2021;232(3):1311–1322. <https://doi.org/10.1111/nph.17646>
- Cheng ZJ, Zhao XY, Shao XX, Wang F, Zhou C, Liu YG, Zhang Y, Zhang XS.** Absciscic acid regulates early seed development in *Arabidopsis* by ABI5-mediated transcription of SHORT HYPOCOTYL UNDER BLUE1. *Plant Cell.* 2014;26(3):1053–1068. <https://doi.org/10.1105/tpc.113.121566>
- Choi Y, Gehring M, Johnson L, Hannon M, Harada JJ, Goldberg RB, Jacobsen SE, Fischer RL.** DEMETER, a DNA glycosylase domain protein, is required for endosperm gene imprinting and seed viability in *Arabidopsis*. *Cell* 2002;110(1):33–42. [https://doi.org/10.1016/S0092-8674\(02\)00807-3](https://doi.org/10.1016/S0092-8674(02)00807-3)
- Chow HT, Chakraborty T, Mosher RA.** RNA-directed DNA methylation and sexual reproduction: expanding beyond the seed. *Curr Opin Plant Biol.* 2020;54:11–17. <https://doi.org/10.1016/j.pbi.2019.11.006>
- Christenhusz MJM, Byng JW.** The number of known plants species in the world and its annual increase. *Phytotaxa* 2016;261(3):201–217. <https://doi.org/10.11646/phytotaxa.261.3.1>
- Costa LM, Yuan J, Rouster J, Paul W, Dickinson H, Gutierrez-Marcos JF.** Maternal control of nutrient allocation in plant seeds by genomic imprinting. *Curr Biol.* 2012;22(2):160–165. <https://doi.org/10.1016/j.cub.2011.11.059>
- Coughlan JM.** The role of hybrid seed inviability in angiosperm speciation. *Am J Bot.* 2023;110(3):1–14. <https://doi.org/10.1002/ajb2.16135>
- Coughlan JM, Wilson Brown M, Willis JH.** Patterns of hybrid seed inviability in the *Mimulus guttatus* sp. complex reveal a potential role of parental conflict in reproductive isolation. *Curr Biol.* 2020;30(1):83–93.e85. <https://doi.org/10.1016/j.cub.2019.11.023>
- Cuerda-Gil D, Slotkin RK.** Non-canonical RNA-directed DNA methylation. *Nat Plants.* 2016;2(11):16163. <https://doi.org/10.1038/nplants.2016.163>
- Darwin F, Seward AC.** More letters of Charles Darwin. A record of his work in a series of hitherto. London: John Murray; 1903. (unpublished letters).
- Dilkes BP, Spielman M, Weizbauer R, Watson B, Burkart-Waco D, Scott RJ, Comai L.** The maternally expressed WRKY transcription factor TTG2 controls lethality in interploidy crosses of *Arabidopsis*. *PLoS Biol.* 2008;6(12):e308. <https://doi.org/10.1371/journal.pbio.0060308>
- Dinu II, Hayes RJ, Kynast RG, Phillips RL, Thill CA.** Novel inter-series hybrids in *Solanum*, section *Petota*. *Theor Appl Genet.* 2005;110(3):403–415. <https://doi.org/10.1007/s00122-004-1782-x>
- Doll NM, Ingram GC.** Embryo-endosperm interactions. *Annu Rev Plant Biol.* 2022;73(1):293–321. <https://doi.org/10.1146/annurev-arplant-102820-091838>
- Doughty J, Aljabri M, Scott RJ.** Flavonoids and the regulation of seed size in *Arabidopsis*. *Biochem Soc Trans.* 2014;42(2):364–369. <https://doi.org/10.1042/BST20140040>
- Dresselhaus T, Sprunck S, Wessel GM.** Fertilization mechanisms in flowering plants. *Curr Biol.* 2016;26(3):R125–R139. <https://doi.org/10.1016/j.cub.2015.12.032>
- Dziasek K, Santos-González J, Qiu Y, Rigola D, Nijbroek K, Köhler C.** Dosage sensitive maternal siRNAs determine hybridization success in *Capsella*. *Res Square.* 2023. <https://doi.org/10.21203/rs.21203.rs-3705839/v3705831>
- Dziasek K, Simon L, Lafon-Placette C, Laenen B, Wärdig C, Santos-González J, Slotte T, Köhler C.** Hybrid seed incompatibility in *Capsella* is connected to chromatin condensation defects in the endosperm. *PLoS Genet.* 2021;17(2):e1009370. <https://doi.org/10.1371/journal.pgen.1009370>
- Erdmann RM, Satyaki PR, Klosinska M, Gehring M.** A small RNA pathway mediates allelic dosage in endosperm. *Cell Rep.* 2017;21(12):3364–3372. <https://doi.org/10.1016/j.celrep.2017.11.078>
- Erilova A, Brownfield L, Exner V, Rosa M, Twell D, Mittelsten Scheid O, Hennig L, Köhler C.** Imprinting of the Polycomb group gene MEDEA serves as a ploidy sensor in *Arabidopsis*. *PLoS Genet.* 2009;5(9):e1000663. <https://doi.org/10.1371/journal.pgen.1000663>
- Figueiredo DD, Batista RA, Roszak PJ, Hennig L, Köhler C.** Auxin production in the endosperm drives seed coat development in *Arabidopsis*. *Elife* 2016;5:e20542. <https://doi.org/10.7554/eLife.20542>
- Florez-Rueda AM, Fiscalini F, Roth M, Grossniklaus U, Städler T.** Endosperm and seed transcriptomes reveal possible roles for small RNA pathways in wild tomato hybrid seed failure. *Genome Biol Evol.* 2021;13(8):evab107. <https://doi.org/10.1093/gbe/evab107>
- Florez-Rueda AM, Paris M, Schmidt A, Widmer A, Grossniklaus U, Stadler T.** Genomic imprinting in the endosperm is systematically perturbed in abortive hybrid tomato seeds. *Mol Biol Evol.* 2016;33(11):2935–2946. <https://doi.org/10.1093/molbev/msw175>
- Floyd S, Friedman W.** Evolution of endosperm developmental patterns among basal flowering plants. *Int J Plant Sci.* 2000;161(S6):S57S81. <https://doi.org/10.1086/317579>
- Floyd SK, Friedman WE.** Developmental evolution of endosperm in basal angiosperms: evidence from *Amborella* (Amborellaceae), *Nuphar* (Nymphaeaceae), and *Illicium* (Illiciaceae). *Plant Syst Evol.* 2001;228(3–4):153–169. <https://doi.org/10.1007/s006060170026>
- García D, Saingery V, Chambrier P, Mayer U, Jürgens G, Berger F.** *Arabidopsis* haiku mutants reveal new controls of seed size by endosperm. *Plant Physiol.* 2003;131(4):1661–1670. <https://doi.org/10.1104/pp.102.018762>
- Geeta R.** The origin and maintenance of nuclear endosperms: viewing development through a phylogenetic lens. *Proc R Soc Lond B Biol Sci.* 2003;270(1510):29–35. <https://doi.org/10.1098/rspb.2002.2206>
- Gehring M, Bubbs KL, Henikoff S.** Extensive demethylation of repetitive elements during seed development underlies gene imprinting. *Science* 2009;324(5933):1447–1451. <https://doi.org/10.1126/science.1171609>
- Gent JJ, Madzima TF, Bader R, Kent MR, Zhang X, Stam M, McGinnis KM, Dawe RK.** Accessible DNA and relative depletion of H3K9me2 at maize loci undergoing RNA-directed DNA methylation. *Plant Cell.* 2014;26(12):4903–4917. <https://doi.org/10.1105/tpc.114.130427>
- Gill BS, Waines JG.** Paternal regulation of seed development in wheat hybrids. *Theor Appl Genet.* 1978;51(6):265–270. <https://doi.org/10.1007/BF00274813>
- Grossniklaus U, Vielle-Calzada JP, Hoepfner MA, Gagliano WB.** Maternal control of embryogenesis by MEDEA a Polycomb group gene in *Arabidopsis*. *Science* 1998;280(5362):446–450. <https://doi.org/10.1126/science.280.5362.446>
- Grover JW, Burgess D, Kendall T, Baten A, Pokhrel S, King GJ, Meyers BC, Freeling M, Mosher RA.** Abundant expression of maternal siRNAs is a conserved feature of seed development. *Proc Natl Acad Sci U S A.* 2020;117(26):15305–15315. <https://doi.org/10.1073/pnas.2001332117>
- Grover JW, Kendall T, Baten A, Burgess D, Freeling M, King GJ, Mosher RA.** Maternal components of RNA-directed DNA methylation are required for seed development in *Brassica rapa*. *Plant J.* 2018;94(4):575–582. <https://doi.org/10.1111/tpj.13910>
- Guo L, Luo X, Li M, Joldersma D, Plunkert M, Liu Z.** Mechanism of fertilization-induced auxin synthesis in the endosperm for seed and fruit development. *Nat Commun.* 2022;13(1):3985. <https://doi.org/10.1038/s41467-022-31656-y>
- Gutierrez-Marcos JF, Pennington PD, Costa LM, Dickinson HG.** Imprinting in the endosperm: a possible role in preventing wide

- hybridization. *Philos Trans R Soc Lond B Biol Sci.* 2003;**358**(1434): 1105–1111. <https://doi.org/10.1098/rstb.2003.1292>
- Haig D, Westoby M.** Genomic imprinting in endosperm: its effect on seed development in crosses between species, and between different ploidies of the same species, and its implications for the evolution of apomixes. *Phil Trans R Soc Lond.* 1991;**333**(1266):1–13. <https://doi.org/10.1098/rstb.1991.0057>
- Haig D, Westoby M.** Parent-specific gene expression and the triploid endosperm. *Am Nat.* 1989;**134**(1):147–155. <https://doi.org/10.1086/284971>
- Hatorangan MR, Laenen B, Steige KA, Slotte T, Kohler C.** Rapid evolution of genomic imprinting in two species of the Brassicaceae. *Plant Cell.* 2016;**28**(8):1815–1827. <https://doi.org/10.1105/tpc.16.00304>
- He H, Shiragaki K, Tezuka T.** Understanding and overcoming hybrid lethality in seed and seedling stages as barriers to hybridization and gene flow. *Front Plant Sci.* 2023;**14**:1219417. <https://doi.org/10.3389/fpls.2023.1219417>
- Hehenberger E, Kradolfer D, Köhler C.** Endosperm cellularization defines an important developmental transition for embryo development. *Development* 2012;**139**(11):2031–2039. <https://doi.org/10.1242/dev.077057>
- Hernández-Hernández T, Miller EC, Román-Palacios C, Wiens JJ.** Speciation across the tree of life. *Biol Rev.* 2021;**96**(4):1205–1242. <https://doi.org/10.1111/brv.12698>
- Hsieh TF, Ibarra CA, Silva P, Zemach A, Eshed-Williams L, Fischer RL, Zilberman D.** Genome-wide demethylation of *Arabidopsis* endosperm. *Science* 2009;**324**(5933):1451–1454. <https://doi.org/10.1126/science.1172417>
- Huang F, Zhu QH, Zhu A, Wu X, Xie L, Wu X, Helliwell C, Chaudhury A, Jean Finnegan E, Luo M.** Mutants in the imprinted PICKLE RELATED 2 gene, suppress seed abortion of fertilization independent seed class mutants and paternal excess interploidy crosses in *Arabidopsis*. *Plant J.* 2017;**90**(2):383–395. <https://doi.org/10.1111/tpj.13500>
- Huc J, Dziasek K, Pachamuthu K, Woh T, Köhler C, Borges F.** Bypassing reproductive barriers in hybrid seeds using chemically induced epimutagenesis. *Plant Cell.* 2021;**34**(3):989–1001. <https://doi.org/10.1093/plcell/koab284>
- İltaş Ö, Svitok M, Cornille A, Schmickl R, Lafon Placette C.** Early evolution of reproductive isolation: a case of weak inbreeder/strong outbreeder leads to an intraspecific hybridization barrier in *Arabidopsis lyrata*. *Evolution* 2021;**75**(6):1466–1476. <https://doi.org/10.1111/evo.14240>
- Ishikawa R, Ohnishi T, Kinoshita Y, Eiguchi M, Kurata N, Kinoshita T.** Rice interspecies hybrids show precocious or delayed developmental transitions in the endosperm without change to the rate of syncytial nuclear division. *Plant J.* 2010;**65**(5):798–806. <https://doi.org/10.1111/j.1365-313X.2010.04466.x>
- Jiang H, Moreno-Romero J, Santos-González J, De Jaeger G, Gevaert K, Van De Slijke E, Köhler C.** Ectopic application of the repressive histone modification H3K9me2 establishes post-zygotic reproductive isolation in *Arabidopsis thaliana*. *Genes Dev.* 2017;**31**(12):1272–1287. <https://doi.org/10.1101/gad.299347.117>
- Johnson LM, Du J, Hale CJ, Bischof S, Feng S, Chodavarapu RK, Zhong X, Marson G, Pellegrini M, Segal DJ, et al.** SRA- and SET-domain-containing proteins link RNA polymerase V occupancy to DNA methylation. *Nature* 2014;**507**(7490):124–128. <https://doi.org/10.1038/nature12931>
- Johnston SA, Hanneman RE Jr.** Manipulations of endosperm balance number overcome crossing barriers between diploid *Solanum* species. *Science* 1982;**217**(4558):446–448. <https://doi.org/10.1126/science.217.4558.446>
- Johnston SA, Nijs TPM, Peloquin SJ, Hanneman RE Jr.** The significance of genic balance to endosperm development in interspecific crosses. *Theor Appl Genet.* 1980;**57**(1):5–9. <https://doi.org/10.1007/BF00276002>
- Josefsson C, Dilkes B, Comai L.** Parent-dependent loss of gene silencing during interspecies hybridization. *Curr Biol.* 2006;**16**(13):1322–1328. <https://doi.org/10.1016/j.cub.2006.05.045>
- Katz O.** Extending the scope of Darwin's 'abominable mystery': integrative approaches to understanding angiosperm origins and species richness. *Ann Bot.* 2018;**121**(1):1–8. <https://doi.org/10.1093/aob/mcx109>
- Khouider S, Borges F, LeBlanc C, Ungu A, Schnittger A, Martienssen R, Colot V, Bouyer D.** Male fertility in *Arabidopsis* requires active DNA demethylation of genes that control pollen tube function. *Nat Commun.* 2021;**12**(1):410. <https://doi.org/10.1038/s41467-020-20606-1>
- Kinser TJ, Smith RD, Lawrence AH, Cooley AM, Vallejo-Marín M, Conradi Smith GD, Puzey JR.** Endosperm-based incompatibilities in hybrid monkeyflowers. *Plant Cell.* 2021;**33**(7):2235–2257. <https://doi.org/10.1093/plcell/koab117>
- Kirkbride RC, Lu J, Zhang C, Mosher RA, Baulcombe DC, Chen ZJ.** Maternal small RNAs mediate spatial-temporal regulation of gene expression, imprinting, and seed development in *Arabidopsis*. *Proc Natl Acad Sci U S A.* 2019;**116**(7):2761–2766. <https://doi.org/10.1073/pnas.1807621116>
- Kiyosue T, Ohad N, Yadegari R, Hannon M, Dinneny J, Wells D, Katz A, Margossian L, Harada JJ, Goldberg RB, et al.** Control of fertilization-independent endosperm development by the MEDEA Polycomb gene in *Arabidopsis*. *Proc Natl Acad Sci U S A.* 1999;**96**(7):4186–4191. <https://doi.org/10.1073/pnas.96.7.4186>
- Klosinska M, Picard CL, Gehring M.** Conserved imprinting associated with unique epigenetic signatures in the *Arabidopsis* genus. *Nat Plants.* 2016;**2**(10):16145. <https://doi.org/10.1038/nplants.2016.145>
- Kozaki A, Aoyanagi T.** Molecular aspects of seed development controlled by gibberellins and abscisic acids. *Int J Mol Sci.* 2022;**23**(3):1876. <https://doi.org/10.3390/ijms23031876>
- Kradolfer D, Hennig L, Köhler C.** Increased maternal genome dosage bypasses the requirement of the FIS Polycomb Repressive Complex 2 in *Arabidopsis* seed development. *PLoS Genet.* 2013a;**9**(1):e1003163. <https://doi.org/10.1371/journal.pgen.1003163>
- Kradolfer D, Wolff P, Jiang H, Siretskiy A, Köhler C.** An imprinted gene underlies postzygotic reproductive isolation in *Arabidopsis thaliana*. *Dev Cell.* 2013b;**26**(5):525–535. <https://doi.org/10.1016/j.devcel.2013.08.006>
- Lafon-Placette C, Hatorangan MR, Steige KA, Cornille A, Lascoux M, Slotte T, Kohler C.** Paternally expressed imprinted genes associate with hybridization barriers in *Capsella*. *Nat Plants.* 2018;**4**(6):352–357. <https://doi.org/10.1038/s41477-018-0161-6>
- Lafon-Placette C, Johannessen IM, Hornslien KS, Ali MF, Bjerkkan KN, Bramsiepe J, Glockle BM, Rebernig CA, Brysting AK, Grini PE, et al.** Endosperm-based hybridization barriers explain the pattern of gene flow between *Arabidopsis lyrata* and *Arabidopsis arenosa* in Central Europe. *Proc Natl Acad Sci U S A.* 2017;**114**(6):E1027–E1035. <https://doi.org/10.1073/pnas.1615123114>
- Lafon-Placette C, Kohler C.** Embryo and endosperm, partners in seed development. *Curr Opin Plant Biol.* 2014;**17**:64–69. <https://doi.org/10.1016/j.pbi.2013.11.008>
- Lafon-Placette C, Kohler C.** Endosperm-based postzygotic hybridization barriers: developmental mechanisms and evolutionary drivers. *Mol Ecol.* 2016;**25**(11):2620–2629. <https://doi.org/10.1111/mec.13552>
- Leblanc O, Pointe C, Hernandez M.** Cell cycle progression during endosperm development in ze mays depends on parental dosage effects. *Plant J.* 2002;**32**(6):1057–1066. <https://doi.org/10.1046/j.1365-313X.2002.01491.x>
- Les DH, Philbrick CT.** Studies of hybridization and chromosome number variation in aquatic angiosperms: evolutionary implications. *Aquatic Bot.* 1993;**44**(2–3):181–228. [https://doi.org/10.1016/0304-3770\(93\)90071-4](https://doi.org/10.1016/0304-3770(93)90071-4)
- Li J, Berger F.** Endosperm: food for humankind and fodder for scientific discoveries. *New Phytol.* 2012;**195**(2):290–305. <https://doi.org/10.1111/j.1469-8137.2012.04182.x>
- Lin BY.** Ploidy barrier to endosperm development in maize. *Genetics* 1984;**107**(1):103–115. <https://doi.org/10.1093/genetics/107.1.103>

- Liu P, Qi M, Wang Y, Chang M, Liu C, Sun M, Yang W, Ren H. *Arabidopsis* RAN1 mediates seed development through its parental ratio by affecting the onset of endosperm cellularization. *Mol Plant*. 2014a;7(8):1316–1328. <https://doi.org/10.1093/mp/ssu041>
- Liu ZW, Shao CR, Zhang CJ, Zhou JX, Zhang SW, Li L, Chen S, Huang HW, Cai T, He XJ. The SET domain proteins SUVH2 and SUVH9 are required for Pol V occupancy at RNA-directed DNA methylation loci. *PLoS Genet*. 2014b;10(1):e1003948. <https://doi.org/10.1371/journal.pgen.1003948>
- Lopes MA, Larkins BA. Endosperm origin, development and function. *Plant Cell*. 1993;5(10):1383–1399. <https://doi.org/10.1105/tpc.5.10.1383>
- Lu J, Zhang C, Baulcombe DC, Chen ZJ. Maternal siRNAs as regulators of parental genome imbalance and gene expression in endosperm of *Arabidopsis* seeds. *Proc Natl Acad Sci U S A*. 2012;109(14):5529–5534. <https://doi.org/10.1073/pnas.1203094109>
- Luo M, Dennis ES, Berger F, Peacock WJ, Chaudhury A. MINISEED3 (MINI3), a WRKY family gene, and HAIKU2 (IKU2), a leucine-rich repeat (LRR) KINASE gene, are regulators of seed size in *Arabidopsis*. *Proc Natl Acad Sci U S A*. 2005;102(48):17531–17536. <https://doi.org/10.1073/pnas.0508418102>
- Martinez G, Panda K, Köhler C, Slotkin RK. Silencing in sperm cells is directed by RNA movement from the surrounding nurse cell. *Nat Plants*. 2016;2(4):16030. <https://doi.org/10.1038/nplants.2016.30>
- Martinez G, Wolff P, Wang Z, Moreno-Romero J, Santos-Gonzalez J, Conze LL, DeFraia C, Slotkin RK, Köhler C. Paternal easiRNAs regulate parental genome dosage in *Arabidopsis*. *Nat Genet*. 2018;50(2):193–198. <https://doi.org/10.1038/s41588-017-0033-4>
- McCue AD, Nuthikattu S, Reeder SH, Slotkin RK. Gene expression and stress response mediated by the epigenetic regulation of a transposable element small RNA. *PLoS Genet*. 2012;8(2):e1002474. <https://doi.org/10.1371/journal.pgen.1002474>
- Moore T. Genetic conflict, genomic imprinting and establishment of the epigenotype in relation to growth. *Reproduction* 2001;122(2):185–193. <https://doi.org/10.1530/rep.0.1220185>
- Moreno-Romero J, Del Toro-De León G, Yadav VK, Santos-Gonzalez J, Kohler C. Epigenetic signatures associated with imprinted paternally expressed genes in the *Arabidopsis* endosperm. *Genome Biol*. 2019;20(1):41. <https://doi.org/10.1186/s13059-019-1652-0>
- Moreno-Romero J, Jiang H, Santos-Gonzalez J, Kohler C. Parental epigenetic asymmetry of PRC2-mediated histone modifications in the *Arabidopsis* endosperm. *EMBO J*. 2016;35(12):1298–1311. <https://doi.org/10.15252/embj.201593534>
- Muntzing A. Hybrid incompatibility and the origin of polyploidy. *Hereditas* 1933;18(1–2):33–55. <https://doi.org/10.1111/j.1601-5223.1933.tb02596.x>
- Oneal E, Willis JH, Franks RG. Disruption of endosperm development is a major cause of hybrid seed inviability between *Mimulus guttatus* and *Mimulus nudatus*. *New Phytol*. 2016;210(3):1107–1120. <https://doi.org/10.1111/nph.13842>
- Pachamuthu K, Simon M, Borges F. Targeted suppression of siRNA biogenesis in *Arabidopsis* pollen reveals distinct Pol IV activities in the sperm and vegetative cell lineages. *Res Square*. 2023. <https://doi.org/10.21203/rs.21203.rs-2947716/v2947711>
- Parent J-S, Cahn J, Herridge RP, Grimanelli D, Martienssen RA. Small RNAs guide histone methylation in *Arabidopsis* embryos. *Genes Dev*. 2021;35(11–12):841–846. <https://doi.org/10.1101/gad.343871.120>
- Parhad SS, Tu S, Weng Z, Theurkauf WE. Adaptive evolution leads to cross-species incompatibility in the piRNA transposon silencing machinery. *Dev Cell*. 2017;43(1):60–70.e65. <https://doi.org/10.1016/j.devcel.2017.08.012>
- Park K, Kim MY, Vickers M, Park JS, Hyun Y, Okamoto T, Zilberman D, Fischer RL, Feng X, Choi Y, et al. DNA demethylation is initiated in the central cells of *Arabidopsis* and rice. *Proc Natl Acad Sci U S A*. 2016;113(52):15138–15143. <https://doi.org/10.1073/pnas.1619047114>
- Peer WA, Murphy AS. Flavonoids and auxin transport: modulators or regulators? *Trends Plant Sci*. 2007;12(12):556–563. <https://doi.org/10.1016/j.tplants.2007.10.003>
- Pennington PD, Costa LM, Gutierrez-Marcos JF, Greenland AJ, Dickinson HG. When genomes collide: aberrant seed development following maize interploidy crosses. *Ann Bot*. 2008;101(6):833–843. <https://doi.org/10.1093/aob/mcn017>
- Petrén H, Thosteman H, Stift M, Toräng P, Ågren J, Friberg M. Differences in mating system and predicted parental conflict affect post-pollination reproductive isolation in a flowering plant. *Evolution* 2023;4(4):1019–1030. <https://doi.org/10.1093/evolut/qpado16>
- Picard CL, Povilus RA, Williams BP, Gehring M. Transcriptional and imprinting complexity in *Arabidopsis* seeds at single-nucleus resolution. *Nat Plants*. 2021;7(6):730–738. <https://doi.org/10.1038/s41477-021-00922-0>
- Pignatta D, Erdmann RM, Scheer E, Picard CL, Bell GW, Gehring M. Natural epigenetic polymorphisms lead to intraspecific variation in *Arabidopsis* gene imprinting. *Elife* 2014;3:e03198. <https://doi.org/10.7554/eLife.03198>
- Pignocchi C, Minns GE, Nesi N, Koumproglou R, Kitsios G, Benning C, Lloyd CW, Doonan JH, Hills MJ. ENDOSPERM DEFECTIVE1 is a novel microtubule-associated protein essential for seed development in *Arabidopsis*. *Plant Cell*. 2009;21(1):90–105. <https://doi.org/10.1105/tpc.108.061812>
- Povilus RA, Diggle PK, Friedman WE. Evidence for parent-of-origin effects and interparental conflict in seeds of an ancient flowering plant lineage. *Proc Biol Sci*. 2018;285(1872):20172491. <https://doi.org/10.1098/rspb.2017.2491>
- Povilus RA, Gehring M. Maternal-filial transfer structures in endosperm: a nexus of nutritional dynamics and seed development. *Curr Opin Plant Biol*. 2022;65:102121. <https://doi.org/10.1016/j.pbi.2021.102121>
- Queller DC. Kin selection and conflict in seed maturation. *J Theor Biol*. 1983;100(1):153–172. [https://doi.org/10.1016/0022-5193\(83\)90099-1](https://doi.org/10.1016/0022-5193(83)90099-1)
- Rebernick CA, Lafon-Placette C, Hatorangan MR, Slotte T, Kohler C. Non-reciprocal interspecies hybridization barriers in the *Capsella* genus are established in the endosperm. *PLoS Genet*. 2015;11(6):e1005295. <https://doi.org/10.1371/journal.pgen.1005295>
- Rodrigues JA, Ruan R, Nishimura T, Sharma MK, Sharma R, Ronald PC, Fischer RL, Zilberman D. Imprinted expression of genes and small RNA is associated with localized hypomethylation of the maternal genome in rice endosperm. *Proc Natl Acad Sci U S A*. 2013;110(19):7934–7939. <https://doi.org/10.1073/pnas.1306164110>
- Rodrigues JA, Zilberman D. Evolution and function of genomic imprinting in plants. *Genes Dev*. 2015;29(24):2517–2531. <https://doi.org/10.1101/gad.269902.115>
- Roth M, Florez-Rueda AM, Griesser S, Paris M, Städler T. Incidence and developmental timing of endosperm failure in post-zygotic isolation between wild tomato lineages. *Ann Bot*. 2018a;121(1):107–118. <https://doi.org/10.1093/aob/mcx133>
- Roth M, Florez-Rueda AM, Paris M, Städler T. Wild tomato endosperm transcriptomes reveal common roles of genomic imprinting in both nuclear and cellular endosperm. *Plant J*. 2018b;95(6):1084–1101. <https://doi.org/10.1111/tj.14012>
- Roth M, Florez-Rueda AM, Städler T. Differences in effective ploidy drive genome-wide endosperm expression polarization and seed failure in wild tomato hybrids. *Genetics* 2019;212(1):141–152. <https://doi.org/10.1534/genetics.119.302056>
- Roy AK, Malaviya DR, Kaushal P. Generation of interspecific hybrids of *Trifolium* using embryo rescue techniques. *Methods Mol Biol*. 2011;710:141–151. https://doi.org/10.1007/978-1-61737-988-8_12
- Sandstedt GD, Sweigart AL. Developmental evidence for parental conflict in driving *Mimulus* species barriers. *New Phytol*. 2022;236(4):1545–1557. <https://doi.org/10.1111/nph.18438>
- Sandstedt GD, Wu CA, Sweigart AL. Evolution of multiple postzygotic barriers between species of the *Mimulus tilingii* complex. *Evolution* 2021;75(3):600–613. <https://doi.org/10.1111/evo.14105>
- Satyaki PR, Gehring M. DNA methylation and imprinting in plants: machinery and mechanisms. *Crit Rev Biochem Mol Biol*. 2017;52(2):163–175. <https://doi.org/10.1080/10409238.2017.1279119>

- Satyaki PR, Gehring M.** Paternally acting canonical RNA-directed DNA methylation pathway genes sensitize Arabidopsis endosperm to paternal genome dosage. *Plant Cell*. 2019;**31**(7):1563–1578. <https://doi.org/10.1105/tpc.19.00047>
- Satyaki PRV, Gehring M.** RNA pol IV induces antagonistic parent-of-origin effects on *Arabidopsis* endosperm. *PLoS Biol*. 2022;**20**(4):e3001602. <https://doi.org/10.1371/journal.pbio.3001602>
- Schatlowski N, Wolff P, Santos-Gonzalez J, Schoft V, Siretskiy A, Scott R, Tamaru H, Köhler C.** Hypomethylated pollen bypasses the interploidy hybridization barrier in Arabidopsis. *Plant Cell*. 2014;**26**(9):3556–3568. <https://doi.org/10.1105/tpc.114.130120>
- Schmitz RJ, Schultz MD, Urich MA, Nery JR, Pelizzola M, Libiger O, Alix A, McCosh RB, Chen H, Schork NJ, et al.** Patterns of population epigenomic diversity. *Nature* 2013;**495**(7440):193–198. <https://doi.org/10.1038/nature11968>
- Schoft VK, Chumak N, Mosiolek M, Slusarz L, Komnenovic V, Brownfield L, Twell D, Kakutani T, Tamaru H.** Induction of RNA-directed DNA methylation upon decondensation of constitutive heterochromatin. *EMBO Rep*. 2009;**10**(9):1015–1021. <https://doi.org/10.1038/embor.2009.152>
- Scott RJ, Spielman M, Bailey J, Dickinson HG.** Parent-of-origin effects on seed development in *Arabidopsis thaliana*. *Development* 1998;**125**(17):3329–3341. <https://doi.org/10.1242/dev.125.17.3329>
- Scott RJ, Tratt JL, Bolbol A.** Seed development in interploidy hybrids. In: **Chen ZJ, Birchler JA**, editors. *Polyploid and hybrid genomics*. Oxford, UK: Wiley; 2013. p. 271–290.
- Sekine D, Ohnishi T, Furuumi H, Ono A, Yamada T, Kurata N, Kinoshita T.** Dissection of two major components of the postzygotic hybridization barrier in rice endosperm. *Plant J*. 2013;**76**(5):792–799. <https://doi.org/10.1111/tpj.12333>
- Slotkin RK, Vaughn M, Borges F, Tanurdzic M, Becker JD, Feijo JA, Martienssen RA.** Epigenetic reprogramming and small RNA silencing of transposable elements in pollen. *Cell* 2009;**136**(3):461–472. <https://doi.org/10.1016/j.cell.2008.12.038>
- Sorensen MB, Mayer U, Lukowitz W, Robert H, Chambrier P, Jürgens G, Somerville C, Lepiniec L, Berger F.** Cellularisation in the endosperm of *Arabidopsis thaliana* is coupled to mitosis and shares multiple components with cytokinesis. *Development* 2002;**129**(24):5567–5576. <https://doi.org/10.1242/dev.00152>
- Städler T, Florez-Rueda AM, Roth M.** A revival of effective ploidy: the asymmetry of parental roles in endosperm-based hybridization barriers. *Curr Opin Plant Biol*. 2021;**61**:102015. <https://doi.org/10.1016/j.pbi.2021.102015>
- Stoute AI, Varenko V, King GJ, Scott RJ, Kurup S.** Parental genome imbalance in *Brassica oleracea* causes asymmetric triploid block. *Plant J*. 2012;**71**(3):503–516. <https://doi.org/10.1111/j.1365-3113.2012.05015.x>
- Sukno S, Ruso J, Jan CC, Melero-Vara JM, Fernández-Martínez JM.** Interspecific hybridization between sunflower and wild perennial *Helianthus* species via embryo rescue. *Euphytica* 1999;**106**(1):69–78. <https://doi.org/10.1023/A:1003524822284>
- Thomae AW, Schade GO, Padeken J, Borath M, Vetter I, Kremmer E, Heun P, Imhof A.** A pair of centromeric proteins mediates reproductive isolation in *Drosophila* species. *Dev Cell*. 2013;**27**(4):412–424. <https://doi.org/10.1016/j.devcel.2013.10.001>
- Tonosaki K, Sekine D, Ohnishi T, Ono A, Furuumi H, Kurata N, Kinoshita T.** Overcoming the species hybridization barrier by ploidy manipulation in the genus *Oryza*. *Plant J*. 2018;**93**(3):534–544. <https://doi.org/10.1111/tpj.13803>
- Trivers RL.** Parent-offspring conflict. *Am Zool*. 1974;**14**(1):249–264. <https://doi.org/10.1093/icb/14.1.249>
- von Wangenheim KH.** Zur Ursache Der Abortion Von Samenanlagen In Diploid-Polyploid-Kreuzungen. *Zeitschrift für Vererbungslehre*. 1962;**93**:319–334. <https://doi.org/10.1007/BF00888790>
- von Wangenheim KH, Peterson HP.** Aberrant endosperm development in interploidy crosses reveals a timer of differentiation. *Dev Biol*. 2004;**270**(2):277–289. <https://doi.org/10.1016/j.ydbio.2004.03.014>
- Valentine DH, Woodell SRJ.** Studies in British *Primulas* X. Seed incompatibility in intraspecific and interspecific crosses at diploid and tetraploid levels. *New Phytol*. 1963;**62**(2):125–143. <https://doi.org/10.1111/j.1469-8137.1963.tb06321.x>
- Walia H, Josefsson C, Dilkes B, Kirkbride R, Harada J, Comai L.** Dosage-dependent deregulation of an AGAMOUS-LIKE gene cluster contributes to interspecific incompatibility. *Curr Biol*. 2009;**19**(13):1128–1132. <https://doi.org/10.1016/j.cub.2009.05.068>
- Wang G, Jiang H, Del Toro-De León G, Martinez G, Kohler C.** Sequestration of a transposon-derived siRNA by a target mimic imprinted gene induces postzygotic reproductive isolation in *Arabidopsis*. *Dev Cell*. 2018;**46**(6):696–705. <https://doi.org/10.1016/j.devcel.2018.07.014>
- Wang Z, Butel N, Santos-González J, Simon L, Wärdig C, Köhler C.** Transgenerational effect of mutants in the RNA-directed DNA methylation pathway on the triploid block in *Arabidopsis*. *Genome Biol*. 2021;**22**(1):141. <https://doi.org/10.1186/s13059-021-02359-2>
- Waters AJ, Bilinski P, Eichten SR, Vaughn MW, Ross-Ibarra J, Gehring M, Springer NM.** Comprehensive analysis of imprinted genes in maize reveals allelic variation for imprinting and limited conservation with other species. *Proc Natl Acad Sci U S A*. 2013;**110**(48):19639–19644. <https://doi.org/10.1073/pnas.1309182110>
- Widmer A, Lexer C, Cozzolino S.** Evolution of reproductive isolation in plants. *Heredity (Edinb)*. 2009;**102**(1):31–38. <https://doi.org/10.1038/hdy.2008.69>
- Williams E, White DWR.** Early seed development after crossing of *Trifolium ambiguum* and *T. repens*. *N Z J Bot*. 1976;**14**(4):307–314. <https://doi.org/10.1080/0028825X.1976.10428903>
- Williams JH, Friedman WE.** Identification of diploid endosperm in an early angiosperm lineage. *Nature* 2002;**415**(6871):522–526. <https://doi.org/10.1038/415522a>
- Wolff P, Jiang H, Wang G, Santos-Gonzalez J, Köhler C.** Paternally expressed imprinted genes establish postzygotic hybridization barriers in *Arabidopsis thaliana*. *Elife* 2015;**4**:e10074. <https://doi.org/10.7554/eLife.10074>
- Wolff P, Weinhofer I, Seguin J, Roszak P, Beisel C, Donoghue MT, Spillane C, Nordborg M, Rehmsmeier M, Köhler C.** High-resolution analysis of parent-of-origin allelic expression in the *Arabidopsis* endosperm. *PLoS Genet*. 2011;**7**(6):e1002126. <https://doi.org/10.1371/journal.pgen.1002126>
- Woodell SRJ, Valentine DH.** Studies in British *Primulas*. IX. Seed incompatibility in diploid-autotetraploid crosses. *New Phytol*. 1961;**60**(3):282–294. <https://doi.org/10.1111/j.1469-8137.1961.tb06256.x>
- Wormit A, Usadel B.** The multifaceted role of pectin methylsterase inhibitors (PMEIs). *Int J Mol Sci*. 2018;**19**(10):2878. <https://doi.org/10.3390/ijms19102878>
- Xu H, Li F, Pan Y, Gong X.** Interspecific hybridization processes between *Michelia yunnanensis* and *M. crassipes* and embryogenesis of the heterozygote. *HortScience* 2017;**52**(8):1043–1047. <https://doi.org/10.21273/HORTSCI12086-17>
- Xu W, Sato H, Bente H, Santos-González J, Köhler C.** Endosperm cellularization failure induces a dehydration-stress response leading to embryo arrest. *Plant Cell*. 2023;**35**(2):874–888. <https://doi.org/10.1093/plcell/koac337>
- Xu Y, Wang Y, Stroud H, Gu X, Sun B, Gan ES, Ng KH, Jacobsen SE, He Y, Ito T.** A matrix protein silences transposons and repeats through interaction with retinoblastoma-associated proteins. *Curr Biol*. 2013;**23**(4):345–350. <https://doi.org/10.1016/j.cub.2013.01.030>
- Yang R, Zheng Z, Chen Q, Yang L, Huang H, Miki D, Wu W, Zeng L, Liu J, Zhou JX, et al.** The developmental regulator PKL is required to maintain correct DNA methylation patterns at RNA-directed DNA methylation loci. *Genome Biol*. 2017;**18**(1):103. <https://doi.org/10.1186/s13059-017-1226-y>
- Yun J, Kim YS, Jung JH, Seo PJ, Park CM.** The AT-hook motif-containing protein AHL22 regulates flowering initiation by modifying FLOWERING LOCUS T chromatin in *Arabidopsis*. *J Biol Chem*.

- 2012;**287**(19):15307–15316. <https://doi.org/10.1074/jbc.M111.318477>
- Zemach A, Kim MY, Hsieh PH, Coleman-Derr D, Eshed-Williams L, Thao K, Harmer SL, Zilberman D.** The *Arabidopsis* nucleosome remodeler DDM1 allows DNA methyltransferases to access H1-containing heterochromatin. *Cell* 2013;**153**(1):193–205. <https://doi.org/10.1016/j.cell.2013.02.033>
- Zhang B, Li C, Li Y, Yu H.** Mobile TERMINAL FLOWER1 determines seed size in *Arabidopsis*. *Nat Plants*. 2020;**6**(9):1146–1157. <https://doi.org/10.1038/s41477-020-0749-5>
- Zhang M, Xie S, Dong X, Zhao X, Zeng B, Chen J, Li H, Yang W, Zhao H, Wang G, et al.** Genome-wide high resolution parental-specific DNA and histone methylation maps uncover patterns of imprinting regulation in maize. *Genome Res*. 2014;**24**(1):167–176. <https://doi.org/10.1101/gr.155879.113>
- Zumajo-Cardona C, Aguirre M, Castillo-Bravo R, Mizzotti C, Di Marzo M, Banfi C, Mendes MA, Spillane C, Colombo L, Ezquer I.** Maternal control of triploid seed development by the TRANSPARENT TESTA 8 (TT8) transcription factor in *Arabidopsis thaliana*. *Sci Rep*. 2023;**13**(1): 1316. <https://doi.org/10.1038/s41598-023-28252-5>