


Barriers and facilitators to the adoption and implementation of evidence-based injury prevention training programmes: a narrative review

Mary Catherine Minnig ¹, Lauren Hawkinson,² Hayley J Root,³ Jeffrey Driban,⁴ Lindsay J DiStefano,^{5,6} Leigh Callahan,^{7,8} Kirsten R Ambrose,⁷ Jeffrey T Spang,⁹ Yvonne M Golightly^{7,10}

To cite: Minnig MC, Hawkinson L, Root HJ, *et al.* Barriers and facilitators to the adoption and implementation of evidence-based injury prevention training programmes: a narrative review. *BMJ Open Sport & Exercise Medicine* 2022;**8**:e001374. doi:10.1136/bmjsem-2022-001374

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bmjsem-2022-001374>).

JTS and YMG are joint senior authors.

Accepted 5 September 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to
Mary Catherine Minnig;
mminnig@email.unc.edu

ABSTRACT

While there is a multitude of evidence supporting the efficacy of injury prevention training programmes, the literature investigating the implementation of these programmes is, in contrast, rather limited. This narrative review sought to describe the commonly reported barriers and facilitators of the implementation of injury prevention training programmes among athletes in organised sport. We also aimed to identify necessary steps to promote the uptake and sustainable use of these programmes in non-elite athletic communities. We identified 24 publications that discussed implementing evidence-based injury prevention training programmes. Frequently reported barriers to implementation include the perceived time and financial cost of the programme, coaches lacking confidence in their ability to implement it, and the programme including exercises that were difficult or confusing to follow. Frequently reported facilitators to implementation include the coach being aware of programme efficacy, shared motivation to complete the programme from both coaches and athletes, and the ability to easily integrate the programme into practice schedules. The current literature is focused on high-income, high-resource settings. We recommend that future studies focus on understanding the best practices of programme dissemination in culturally and economically diverse regions. Programmes ought to be of no financial burden to the user, be simply adaptable to different sports and individual athletes and be available for use in easily accessible forms, such as in a mobile smartphone application.

INTRODUCTION

Osteoarthritis (OA) is one of the most common chronic health conditions in the USA, substantially contributing to the burden of chronic pain and disability.¹ The prevalence of knee OA has nearly doubled since the mid-20th century, even after adjusting for age and body mass index.² Recent studies demonstrate that knee OA makes up roughly 80% of the total burden from all forms of OA.² About 19% of Americans aged 45 years and older

WHAT IS ALREADY KNOWN ON THIS TOPIC

- ⇒ The previous literature has described the efficacy of evidence-based injury prevention training programmes.
- ⇒ Effective programmes are comprised of neuromuscular training, plyometric exercises, balance training and stretching.
- ⇒ Despite generally high efficacy, program implementation rates remain low.

WHAT THIS STUDY ADDS

- ⇒ Frequently reported barriers to injury prevention program implementation include the perceived time and financial burden, coaches lacking confidence in their knowledge to implement the program properly and the complexity of the programme's exercises.
- ⇒ Commonly reported facilitators to injury prevention program implementation include the coach's knowledge of the programme's efficacy, motivation to complete the program from both the coach and the players, and the programme's ease of integration into the usual practice schedule.
- ⇒ Current literature on this topic is focused in high-resource and high-income countries.
- ⇒ We provide specific recommendations for researchers, clinicians, and administrators to encourage the wider and more sustained implementation injury prevention exercise programs.

are affected by knee OA; however, increasing evidence shows that OA also should be of concern for certain young athletes, especially those with a history of knee injury.^{2,3}

Lower-limb injuries account for over 65% of all sports injuries.⁴ Arguably of greatest concern, anterior cruciate ligament (ACL) injuries are associated with long-term economic and health burdens. Surgical reconstruction of the ACL and its long-term costs are estimated to be as high as US\$38 000 per patient.⁴ There is a known association between ACL injury and the later development of OA.^{3,5-7} This is of particular concern



HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

- ⇒ We suggest that future studies on injury prevention exercise programmes use implementation-specific frameworks to guide the assessment of implementation success, as well as qualitative methods to better understand the attitudes and beliefs of key stakeholder groups.
- ⇒ Future studies should also investigate the barriers and facilitators to injury prevention exercise program implementation in a wider range of athletic communities.
- ⇒ Parents of adolescent athletes are important stakeholders in this realm and seem to be currently underutilised. Parental buy-in could be an invaluable tool to encourage the broader use of injury prevention exercise programs in youth sports.
- ⇒ We suggest that injury prevention exercise programmes be made available in free mobile smartphone applications to increase ease of access and sustained use.

for young athletes, who, after sustaining an ACL injury, are at significantly increased risk of developing OA as early as 5–15 years after an ACL injury.⁵ Working to prevent initial knee injury is thus necessary to reduce the additional burden of living over half their life with OA.

Numerous randomised controlled trials have demonstrated the efficacy of injury prevention training programmes in reducing ankle and knee injuries in athletes at youth, high school and elite levels.^{4 8–18} These programmes generally include neuromuscular training, plyometric exercises, balance training and stretching.⁴ Exercise regimens are recommended to be performed 2–3 times per week through the pre-competitive season, in-season and during out-of-season maintenance. However, they do not need to be overly time-intensive and can be integrated into a pre-existing warm-up or conditioning routine.⁴ Several injury and athletic organisations, including the National Athletic Trainers' Association, the Academy of Orthopaedic Physical Therapy, the American Academy of Sports Physical Therapy, the Osteoarthritis Action Alliance, the American College of Sports Medicine and the International Olympic Committee, support and offer guidance on the use of injury prevention programmes to prevent knee injuries.^{4 19–22}

While the literature investigating the implementation of these programmes is limited, some research indicates usage rates are as low as 20%–29% in the youth soccer community, despite the multitude of evidence supporting their effectiveness.^{23 24} Further information about the current state of implementation of injury prevention programmes is critical to promote widespread dissemination and adoption. This paper aims to describe the frequently reported barriers and facilitators of injury prevention training programme implementation among athletes in organised sports. We aim to identify steps needed to promote the uptake and continued use of these programmes in community-level sports environments.

METHODS

An electronic database search of the literature was conducted between February and March 2021 by two authors (MCM and LH). We did not restrict the search to publication date, but the results were limited to articles printed in English. The search took place in PubMed, Google Scholar, the Cochrane Library and the Cumulative Index of Nursing and Allied Health Literature. A combination of the following terms was used during this search: 'implementation', 'exercise programme', 'injury prevention', and 'sports'. We conducted a comprehensive narrative review rather than a traditional systematic review because our focus was to gain a more general, larger-scale view of the current literature investigating injury prevention training programme implementation barriers and facilitators within a broad range of youth and adult non-elite athletic communities. Thus, data extraction and synthesis were not feasible for all eligible articles because of the variability in the types of articles, study designs and research areas (eg, effectiveness of implementing a specific injury prevention training programme, barriers/facilitators to injury prevention training programmes, implementation science) on this topic. We did not use strict criteria to exclude articles found through our search to preserve our goal of gaining a broader scope on this topic. Titles and abstracts were reviewed by two authors (MCM and YMG), and the full text was obtained for articles that included the implementation of injury prevention training programmes. We performed quality assessments of the retrieved articles. The two authors reviewed the identified articles using a spreadsheet that included quality assessment items from Walsh *et al*²⁵ for qualitative studies and Shea *et al* for systematic reviews.²⁶

RESULTS

Overview of the current implementation literature

We identified 23 publications that discussed topics related to the implementation of evidence-based injury prevention training programmes: 16 analysed barriers and facilitators regarding specific injury prevention training programmes, while 7 discussed more generally how to approach implementation science research on this topic. Of the 16 studies that investigated specific barriers and facilitators, 4 were based in Australia,^{27 28} 2 were based in Scandinavia,^{29 30} 3 in the USA,^{23 24 31 32} 2 in Canada,^{33 34} 1 in Switzerland³⁵ and 4 were systematic reviews studying injury prevention training programmes based in various regions.^{31 36–38} A representative sample of these studies is described in detail in online supplemental table 1.

Summary of the current literature

We found that the majority of the studies included in this review met all identified quality assessment measures. Tables summarising these results are included in online supplemental tables 2 and 3. We found that a qualitative study by McGlashan *et al* did not fully describe the methods used to conduct semistructured interviews and

PERCEIVED BARRIERS		PERCEIVED FACILITATORS	
<ul style="list-style-type: none"> • Potential cost of implementation, including the need to hire additional staff to administer program • Coaches lack confidence in knowledge/ability to implement program • Perceived time burden of program • Lack of importance placed on injury prevention 	<ul style="list-style-type: none"> • Difficult/confusing exercises • Lack of motivation from the athletes • Lack of flexibility of program, especially for athletes of different skill levels and athletes who miss practice 	<ul style="list-style-type: none"> • Knowledge of injury risk and need for proactive action to prevent injuries • Knowledge of efficacy of injury prevention programs • Motivation to complete program from both coaches and athletes 	<ul style="list-style-type: none"> • Simple integration of program into practice schedule • Coaches have access to educational materials on the program • The program allows for individualization and exercise progression throughout the competitive season

Figure 1 Summary of perceived barriers and facilitators of injury prevention exercise program implementation.

a systematic review by O'Brien and Finch did not discuss the likelihood of publication bias from the articles they described.^{27 39}

While synthesising the results from the current literature, we found common themes in the reported barriers and facilitators to implementing injury prevention training programmes (figure 1). Many studies reported that the extra time, ranging from 5 to 20 min required to complete the programme, was the biggest barrier noted by coaches.^{28 30 31 33 40–42} Even coaches who reported they had not used an injury prevention training programme indicated that the perceived time burden of the programme was too high. Data on coaches' knowledge about injuries and injury prevention techniques varied greatly. Little knowledge in this area and viewing injury prevention as a low priority were common barriers to implementation.^{24 27 28 30 35 37 42} On the other hand, knowledge surrounding the burden of injury did not always facilitate the adoption and implementation of an injury prevention training programme.⁴³

Some studies found that coaches felt they lacked the necessary physical equipment and support staff resources to properly implement injury prevention training programmes.^{41 42} It is important to note, however, that additional physical equipment is not a requirement of many injury prevention programmes, thus this barrier can be easily overcome with certain programmes. Unsurprisingly, support from key stakeholders, including coaches, administrators, athletic trainers and injury prevention training programme experts, was found to be an important facilitator in injury prevention training programme implementation.²⁷ Additionally, the complexity and difficulty of the exercises included in injury prevention training programmes were reported as barriers to implementation.^{34 44} Conversely, studies reported that programmes easily adaptable to individual teams, sports and athletes were more likely to be implemented.^{34 42 45} This result could relate to the finding that players' lack of motivation also affected whether or not a programme was implemented.^{28 31 40 41}

Looking specifically at the US-based literature, some studies have investigated barriers and facilitators in injury prevention training programme implementation in high school and collegiate sports teams. A 2015 study of soccer and basketball coaches in Oregon high schools found that the coaches believed there was little need for an injury prevention training programme given low injury rates. The perception that injury prevention training programmes were complex and difficult to administer prevented the initial adoption of programmes.²⁴ In a 2020 study by Dix *et al*, NCAA Women's soccer coaches in the NCAA East Region reported that the perceived need to hire additional coaching staff who could administer an injury prevention training programme was the biggest barrier to implementing a programme with their teams.³² A study focusing on club-level soccer coaches indicated that even high levels of intent to incorporate an injury prevention training programme into team practices did not necessarily translate to high levels of implementation.⁴³ Even an educational workshop intervention aimed to increase the adoption of injury prevention training programmes in this population did not drastically increase adherence rates.⁴³ Implementation success with these types of educational workshops has been found in another study, however.⁴⁶

DISCUSSION

Directions for researchers

Many of the studies found through this review were set within randomised clinical trials. While testing the efficacy of new injury prevention training programmes cannot be disputed as essential, it is also necessary to use pragmatic or hybrid designs to better understand the implementation of injury prevention training programmes in real-world settings outside of randomised clinical trials.^{47–49} True effectiveness trials are limited.⁵⁰ The current literature is set on the qualities that make an effective injury prevention training programme when implemented appropriately. Still, the evidence is lacking on the optimal strategies to best disseminate these

programmes to ensure widespread uptake, adoption and long-term use. Similar to suggestions made previously by Finch,⁵¹ we support future studies' use of the seven-step Translating Research into Injury Prevention Practice (TRIPP) Framework as well as Padua's seven steps for developing and implementing a preventive programme as roadmaps to drive intervention planning, and the use of both the Reach, Effectiveness, Adoption, Implementation, and Maintenance Framework and the TRIPP Framework to direct the evaluation of programme implementation and adoption.^{39 52 53} Long-term studies are also needed to examine how injury prevention training programmes endure over time.

Furthermore, we believe that the inclusion of qualitative interviews with key stakeholders will be a valuable addition to the current body of the literature and provide essential feedback on how to improve their involvement in injury prevention training programme implementation, dissemination and maintenance. The use of qualitative methods in sports medicine is growing. Understanding stakeholder motivation and their knowledge, attitudes and beliefs surrounding the implementation of injury prevention training programmes will help identify education initiatives and improve implementation techniques. Researchers in the area of concussion prevention and management have successfully used qualitative methods to identify gaps in knowledge and factors related to the attitudes and beliefs of athletes, parents and coaches.^{54–56} Valuable insights have been gained using qualitative methodology to examine the implementation and use of an injury prevention training programme through research in Europe.^{57–59} Applying similar techniques in other geographic regions will improve injury prevention training programme implementation strategies.

Currently, the literature on injury prevention training programme efficacy, implementation and adoption are primarily set in homogenous, high-resource communities in high-income countries, such as Australia, Canada the USA and European nations. There is a significant gap in the literature investigating the use of injury prevention training programmes in culturally and economically diverse settings globally. Conducting this type of research in underserved areas requires careful attention to address an array of ethical concerns. For example, investigators should ensure communities can sustain the tested injury prevention training programme after the study. We suggest targeting implementation strategies to middle and high school sports programmes in underrepresented areas as an initial step in expanding the use of these programmes within these communities. Many evidence-based injury prevention training programmes benefit from not requiring additional equipment resources. We see this factor as a facilitator to implementation and a way to ensure programme accessibility no matter the user's socioeconomic status. It will, however, be important to study how well current injury prevention training programmes and new programmes integrate into diverse settings where sports culture and injury needs may differ.

Drawing from key ideas within the implementation and dissemination science realm, the use of systematic evaluation of the social, cultural and environmental context is necessary to encourage the spread, scalability and sustained use of evidence-based injury prevention programmes.⁶⁰

Directions for clinicians and administrators

The current literature provides a good base to describe the principles of effective and implementable injury prevention training programmes. We found that injury prevention training programmes that are flexible and easily incorporated into the existing practice schedule for a team are more likely to be adopted. We also found evidence supporting the importance of involving key stakeholders, including coaches and players. Studies show that buy-in from coaches and players is a necessary first step to injury prevention training programme implementation.^{53 61 62} Education on the importance of injury prevention and the efficacy of injury prevention training programmes can help coaches feel adequately equipped to administer the injury prevention training programme. However, buy-in from coaches alone is not enough—players also need to be involved and feel motivated to participate in the injury prevention training programme.^{63 64} To achieve this goal, it may be beneficial to promote the positive effects injury prevention exercise programmes can have on overall biomechanics and athletic performance. Some studies have found evidence associating injury prevention training programme completion with increased lower extremity strength and flexibility,^{65 66} vertical jump height,^{66 67} balance ability⁶⁷ and maximum aerobic power,⁶⁶ all of which can, in turn, advance athletic performance.^{65–67}

Additionally, it would be beneficial to encourage buy-in for these programmes from the parents of adolescent athletes. While the literature on this particular aspect of stakeholder involvement is lacking, we suggest that educational efforts on the importance of these programmes be expanded to include parents. Parent involvement—and influence—in youth sports continues to increase, highlighting the invaluable need to include this previously untapped group in efforts to expand the implementation of injury prevention programmes in community athletics.^{68 69} Parents' investments in terms of time and money into their children's sports teams can be powerful tools that assist in changing the overall culture around injury prevention in these types of community settings.

We also note that it will be important to encourage injury prevention training programmes in school settings and recreational non-elite athletic environments. Previous studies have investigated the efficacy of injury prevention training programmes in school physical education classes.^{70 71} While results from these studies indicate that these programmes can successfully reduce injury occurrence, further research is needed to understand the specific barriers and facilitators to implementation in adolescent physical education classes as they likely differ

EASE OF USE	COACH/ATHLETE INVOLVEMENT	COST
<ul style="list-style-type: none"> • Easily adapt team's warm-up routine to include key elements of the program • The program does not require staff with specialized training • Exercises are easy to understand and perform in a variety of weather conditions • Exercises do not require the use of any additional physical equipment to complete • The program has the flexibility to individualize to specific sports, athletes, and abilities 	<ul style="list-style-type: none"> • The program should be available in a format where both coaches and athletes have access to the program (e.g., mobile app) • Coaches ought to be provided with educational materials on the program and injury • Future policy measures ought to encourage and monitor use of the program 	<ul style="list-style-type: none"> • The program must be of no financial cost • The program must not be time-intensive (15 minutes or less) • Stakeholder training in the program and injury should be time efficient and available in a variety of user-centered methods

Figure 2 Recommendations to increase injury prevention exercise program implementation.

from those associated with outside-of-school non-elite athletic settings.

As previously detailed, the current literature describes the time required to complete an injury prevention programme as a key barrier to implementation. A number of the studies found through this review are limited by the detail of information around the comprehensive time burden of the studied injury prevention training programmes—in other words, questions remain around the number of times per week a programme was performed, the number of weeks during in-season competition the programme was performed, the number of weeks during out-of-season competition the programme was completed, and whether the programme was completed all at once or separated throughout practice. These are all important factors to consider in terms of how they contribute to programme implementation and long-term compliance.^{37 72} The current best practice guidelines from the National Association of Athletic Trainers indicate that evidence-based injury prevention programmes can be performed in as little as 10–15 min, 2–3 times per week to be effective.⁴ Future research will benefit from ensuring that data on the amount of time required to complete a programme is reported. This type of information is necessary to understand the optimal balance of programme time that will ensure high levels of efficacy while limiting the overall time burden in different non-elite athletic settings. As noted in Padua's seven-step guidelines for programme implementation, a 'one-size-fits-all' approach to injury prevention training programmes does not seem feasible.⁵³ Instead, programmes must be flexible to accommodate the setting-specific barriers and facilitators to implementation.

To increase ease of access to and the sustained use of injury prevention training programmes, we suggest that

programmes include a free smartphone mobile application version. An estimated 80% of the world's population has a mobile phone, suggesting that mobile applications would greatly expand access to evidence-based injury prevention exercises to communities that previously may have been excluded from this content due to geographic and monetary factors.⁷³ A literature review conducted in 2014 found that only four evidence-based injury prevention applications were available for mobile download.⁷³ A free mobile application could address a number of the perceived barriers in implementation found through this comprehensive review, including barriers related to cost, coaches lacking educational materials on injury prevention, a need to individualise the programme for different athletes and a lack of explanation for confusing or difficult exercises.

CONCLUSIONS

Recent literature agree on the qualities that make an injury prevention training programme effective at preventing injuries. Despite the known positive benefits of these programmes, uptake and adoption rates are still quite low. Even more surprising, there is a major gap in the literature investigating strategies aimed at improving implementation rates across a wide range of athletic communities. Based on the findings of this review, we propose recommendations to facilitate the implementation of future injury prevention training programmes (figure 2). Additionally, we recommend engaging professional sport and coaches' associations to promote the use of and disseminate educational information on evidence-based injury prevention programmes. Perhaps most novel, we recommend that injury prevention training programmes be available in free smartphone mobile applications. We see the convenience and ease of access of a mobile application as key factors that will facilitate

the initial uptake and long-term use of injury prevention training programmes.

Author affiliations

¹Department of Epidemiology, Gillings School of Global Public Health, University of North Carolina, Chapel Hill, North Carolina, USA

²Department of Exercise and Sport Science, University of North Carolina at Chapel Hill College of Arts and Sciences, Chapel Hill, North Carolina, USA

³Department of Physical Therapy and Athletic Training, Northern Arizona University, Flagstaff, Arizona, USA

⁴Division of Rheumatology, Allergy, and Immunology, Tufts Medical Center, Boston, Massachusetts, USA

⁵Department of Kinesiology, University of Connecticut, Storrs, Connecticut, USA

⁶Institute of Sports Medicine, University of Connecticut, Storrs, Connecticut, USA

⁷Thurston Arthritis Research Center, University of North Carolina at Chapel Hill School of Medicine, Chapel Hill, North Carolina, USA

⁸Department of Medicine, University of North Carolina at Chapel Hill School of Medicine, Chapel Hill, North Carolina, USA

⁹Department of Orthopaedics, University of North Carolina at Chapel Hill School of Medicine, Chapel Hill, North Carolina, USA

¹⁰College of Allied Health Professions, University of Nebraska Medical Center, Omaha, Nebraska, USA

Contributors Substantial contributions to study conception and design: MCM, JTS and YMG. Substantial contributions to acquisition of data: MCM, LH, HJR, JD, LJD and YMG. Substantial contributions to analysis and interpretation of data: MCM, LH, HJR, JD, LC, KRA, JTS and YMG. Manuscript preparation: MCM and YMG. Critical revisions to the manuscript for intellectual content: MCM, LH, HJR, JD, LJD, LC, KRA, JTS and YMG. Final approval of the submitted manuscript: MCM, LH, HJR, JD, LJD, LC, KRA, JTS and YMG.

Funding This study was funded by Centers for Disease Control and Prevention (Grant number: #NU58DP006980).

Competing interests None declared.

Patient consent for publication Not applicable.

Ethics approval Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Mary Catherine Minnig <http://orcid.org/0000-0002-1218-4965>

REFERENCES

- 1 US Burden of Disease Collaborators, Mokdad AH, Ballesteros K, *et al*. The state of US health, 1990-2016: burden of diseases, injuries, and risk factors among US states. *JAMA* 2018;319:1444-72.
- 2 Wallace IJ, Worthington S, Felson DT, *et al*. Knee osteoarthritis has doubled in prevalence since the mid-20th century. *Proc Natl Acad Sci U S A* 2017;114:9332-6.
- 3 Caine DJ, Golightly YM. Osteoarthritis as an outcome of paediatric sport: an epidemiological perspective. *Br J Sports Med* 2011;45:298-303.
- 4 Padua DA, DiStefano LJ, Hewett TE, *et al*. National athletic trainers' association position statement: prevention of anterior cruciate ligament injury. *J Athl Train* 2018;53:5-19.
- 5 Simon D, Mascarenhas R, Saltzman BM, *et al*. The relationship between anterior cruciate ligament injury and osteoarthritis of the knee. *Adv Orthop* 2015;2015:928301.
- 6 Harkey MS, Luc BA, Golightly YM, *et al*. Osteoarthritis-related biomarkers following anterior cruciate ligament injury and reconstruction: a systematic review. *Osteoarthritis Cartilage* 2015;23:1-12.
- 7 Allen KD, Golightly YM. State of the evidence. *Curr Opin Rheumatol* 2015;27:276-83.
- 8 Coppack RJ, Etherington J, Wills AK. The effects of exercise for the prevention of overuse anterior knee pain: a randomized controlled trial. *Am J Sports Med* 2011;39:940-8.
- 9 Crossley KM, Patterson BE, Culvenor AG, *et al*. Making football safer for women: a systematic review and meta-analysis of injury prevention programmes in 11 773 female football (soccer) players. *Br J Sports Med* 2020;54:1089-98.
- 10 Emery CA, Meeuwisse WH. The effectiveness of a neuromuscular prevention strategy to reduce injuries in youth soccer: a cluster-randomised controlled trial. *Br J Sports Med* 2010;44:555-62.
- 11 Foss KDB, Thomas S, Khoury JC, *et al*. A school-based neuromuscular training program and sport-related injury incidence: a prospective randomized controlled clinical trial. *J Athl Train* 2018;53:20-8.
- 12 Heidt RS, Sweeterman LM, Carlonas RL, *et al*. Avoidance of soccer injuries with preseason conditioning. *Am J Sports Med* 2000;28:659-62.
- 13 LaBella CR, Huxford MR, Grissom J, *et al*. Effect of neuromuscular warm-up on injuries in female soccer and basketball athletes in urban public high schools: cluster randomized controlled trial. *Arch Pediatr Adolesc Med* 2011;165:1033-40.
- 14 Lauersen JB, Andersen TE, Andersen LB. Strength training as superior, dose-dependent and safe prevention of acute and overuse sports injuries: a systematic review, qualitative analysis and meta-analysis. *Br J Sports Med* 2018;52:1557-63.
- 15 Petersen J, Thorborg K, Nielsen MB, *et al*. Preventive effect of eccentric training on acute hamstring injuries in men's soccer: a cluster-randomized controlled trial. *Am J Sports Med* 2011;39:2296-303.
- 16 Soligard T, Myklebust G, Steffen K, *et al*. Comprehensive warm-up programme to prevent injuries in young female footballers: cluster randomised controlled trial. *BMJ* 2008;337:a2469.
- 17 van der Horst N, Smits D-W, Petersen J, *et al*. The preventive effect of the Nordic hamstring exercise on hamstring injuries in amateur soccer players: a randomized controlled trial. *Am J Sports Med* 2015;43:1316-23.
- 18 Waldén M, Atroshi I, Magnusson H, *et al*. Prevention of acute knee injuries in adolescent female football players: cluster randomised controlled trial. *BMJ* 2012;344:e3042.
- 19 Selected issues in injury and illness prevention and the team physician: a consensus statement. *Med Sci Sports Exerc* 2016;48:159-71.
- 20 Ardern CL, Ekås GR, Grindem H, *et al*. 2018 international Olympic Committee consensus statement on prevention, diagnosis and management of paediatric anterior cruciate ligament (ACL) injuries. *Br J Sports Med* 2018;52:422-38.
- 21 Arundale AJH, Bizzini M, Giordano A, *et al*. Exercise-Based knee and anterior cruciate ligament injury prevention. *J Orthop Sports Phys Ther* 2018;48:A1-42.
- 22 Trojian T, Driban J, Nuti R, *et al*. Osteoarthritis action alliance consensus opinion - best practice features of anterior cruciate ligament and lower limb injury prevention programs. *World J Orthop* 2017;8:726-34.
- 23 Joy EA, Taylor JR, Novak MA, *et al*. Factors influencing the implementation of anterior cruciate ligament injury prevention strategies by girls soccer coaches. *J Strength Cond Res* 2013;27:2263-9.
- 24 Norcross MF, Johnson ST, Bovbjerg VE, *et al*. Factors influencing high school coaches' adoption of injury prevention programs. *J Sci Med Sport* 2016;19:299-304.
- 25 Walsh D, Downe S. Appraising the quality of qualitative research. *Midwifery* 2006;22:108-19.
- 26 Shea BJ, Grimshaw JM, Wells GA, *et al*. Development of AMSTAR: a measurement tool to assess the methodological quality of systematic reviews. *BMC Med Res Methodol* 2007;7:10.
- 27 McGlashan A, Verrinder G, Verhagen E. Working towards more effective implementation, dissemination and scale-up of lower-limb Injury-Prevention programs: insights from community Australian football coaches. *Int J Environ Res Public Health* 2018;15:351.
- 28 Saunders N, Otago L, Romiti M, *et al*. Coaches' perspectives on implementing an evidence-informed injury prevention programme in junior community netball. *Br J Sports Med* 2010;44:1128-32.

- 29 Dahlström Ö, Jacobsson J, Timpka T. Overcoming the organization-practice barrier in sports injury prevention: a nonhierarchical organizational model. *Scand J Med Sci Sports* 2015;25:e414–22.
- 30 Harøy J, Wiger EG, Bahr R, et al. Implementation of the Adductor Strengthening Programme: Players primed for adoption but reluctant to maintain - A cross-sectional study. *Scand J Med Sci Sports* 2019;29:1092–100.
- 31 Bogardus RL, Martin RJ, Richman AR, et al. Applying the socio-ecological model to barriers to implementation of ACL injury prevention programs: a systematic review. *J Sport Health Sci* 2019;8:8–16.
- 32 Dix C, Logerstedt D, Arundale A, et al. Perceived barriers to implementation of injury prevention programs among collegiate women's soccer coaches. *J Sci Med Sport* 2021;24:352–6.
- 33 Mawson R, Creech MJ, Peterson DC, et al. Lower limb injury prevention programs in youth soccer: a survey of coach knowledge, usage, and barriers. *J Exp Orthop* 2018;5:43.
- 34 Richmond SA, Donaldson A, Macpherson A, et al. Facilitators and barriers to the implementation of iSPRINT: a sport injury prevention program in junior high schools. *Clin J Sport Med* 2020;30:231–8.
- 35 Gebert A, Gerber M, Pühse U, et al. Injury prevention in amateur soccer: a nation-wide study on implementation and associations with injury incidence. *Int J Environ Res Public Health* 2019;16:1593.
- 36 DiStefano LJ, Frank BS, Root HJ, et al. Dissemination and implementation strategies of lower extremity preventive training programs in youth: a clinical review. *Sports Health* 2017;9:524–31.
- 37 Sugimoto D, Myer GD, Bush HM, et al. Compliance with neuromuscular training and anterior cruciate ligament injury risk reduction in female athletes: a meta-analysis. *J Athl Train* 2012;47:714–23.
- 38 Voskanian N. Acl injury prevention in female athletes: review of the literature and practical considerations in implementing an ACL prevention program. *Curr Rev Musculoskelet Med* 2013;6:158–63.
- 39 O'Brien J, Finch CF. A systematic review of core implementation components in team ball sport injury prevention trials. *Inj Prev* 2014;20:357–62.
- 40 Andersson SH, Bahr R, Olsen MJ, et al. Attitudes, beliefs, and behavior toward shoulder injury prevention in elite handball: fertile ground for implementation. *Scand J Med Sci Sports* 2019;29:1996–2009.
- 41 Donaldson A, Callaghan A, Bizzini M, et al. A concept mapping approach to identifying the barriers to implementing an evidence-based sports injury prevention programme. *Inj Prev* 2019;25:244–51.
- 42 O'Brien J, Finch CF. The implementation of musculoskeletal injury-prevention exercise programmes in team ball sports: a systematic review employing the RE-AIM framework. *Sports Med* 2014;44:1305–18.
- 43 Frank BS, Register-Mihalik J, Padua DA. High levels of coach intent to integrate a ACL injury prevention program into training does not translate to effective implementation. *J Sci Med Sport* 2015;18:400–6.
- 44 Finch CF, White P, Twomey D, et al. Implementing an exercise-training programme to prevent lower-limb injuries: considerations for the development of a randomised controlled trial intervention delivery plan. *Br J Sports Med* 2011;45:791–6.
- 45 Donaldson A, Finch CF. Applying implementation science to sports injury prevention. *Br J Sports Med* 2013;47:473–5.
- 46 Pryor JL, Root HJ, Vandermark LW, et al. Coach-led preventive training program in youth soccer players improves movement technique. *J Sci Med Sport* 2017;20:861–6.
- 47 Curran GM, Bauer M, Mittman B, et al. Effectiveness-implementation hybrid designs: combining elements of clinical effectiveness and implementation research to enhance public health impact. *Med Care* 2012;50:217–26.
- 48 Owøye OBA, Rauvola RS, Brownson RC. Dissemination and implementation research in sports and exercise medicine and sports physical therapy: translating evidence to practice and policy. *BMJ Open Sport Exerc Med* 2020;6:e000974.
- 49 Proctor EK, Powell BJ, McMillen JC. Implementation strategies: recommendations for specifying and reporting. *Implement Sci* 2013;8:139.
- 50 Myklebust G, Skjølberg A, Bahr R. Acl injury incidence in female handball 10 years after the Norwegian ACL prevention study: important lessons learned. *Br J Sports Med* 2013;47:476–9.
- 51 Finch CF. No longer lost in translation: the art and science of sports injury prevention implementation research. *Br J Sports Med* 2011;45:1253–7.
- 52 Finch C. A new framework for research leading to sports injury prevention. *J Sci Med Sport* 2006;9:3–9. discussion 10.
- 53 Padua DA, Frank B, Donaldson A, et al. Seven steps for developing and implementing a preventive training program: lessons learned from JUMP-ACL and beyond. *Clin Sports Med* 2014;33:615–32.
- 54 White PE, Register-Mihalik J, Donaldson A, et al. Concussion guideline implementation perceptions and experiences among parents of community-level Australian football junior players. *BMJ Open Sport Exerc Med* 2017;3:e000215.
- 55 Register-Mihalik JK, Williams RM, Marshall SW, et al. Demographic, parental, and personal factors and youth athletes' Concussion-Related knowledge and beliefs. *J Athl Train* 2018;53:768–75.
- 56 Kerr ZY, Nedimyer AK, Kay MC, et al. Factors associated with concussion-symptom knowledge and attitudes toward concussion care seeking in a national survey of parents of middle-school children in the US. *J Sport Health Sci* 2021;10:113–21.
- 57 Orr B, Brown C, Hemsing J, et al. Female soccer knee injury: observed knowledge gaps in injury prevention among players/parents/coaches and current evidence (the know study). *Scand J Med Sci Sports* 2013;23:271–80.
- 58 Møller M, Zebis MK, Myklebust G, et al. "Is it fun and does it enhance my performance?" - Key implementation considerations for injury prevention programs in youth handball. *J Sci Med Sport* 2021;24:1136–42.
- 59 Lindblom H, Carlffjord S, Häggglund M. Adoption and use of an injury prevention exercise program in female football: a qualitative study among coaches. *Scand J Med Sci Sports* 2018;28:1295–303.
- 60 Benjaminse A, Verhagen E. Implementing ACL injury prevention in daily sports Practice-It's not just the program: let's build together, involve the context, and improve the content. *Sports Med* 2021;51:2461–7.
- 61 Bizzini M, Junge A, Dvorak J. Implementation of the FIFA 11+ football warm up program: how to approach and convince the football associations to invest in prevention. *Br J Sports Med* 2013;47:803–6.
- 62 Root HJ, Frank BS, Denegar CR, et al. Application of a preventive training program implementation framework to youth soccer and Basketball organizations. *J Athl Train* 2019;54:182–91.
- 63 Owøye OBA, VanderWey MJ, Pike I. Reducing injuries in soccer (football): an umbrella review of best evidence across the epidemiological framework for prevention. *Sports Med Open* 2020;6:46.
- 64 Wilke J, Niederer D, Vogt L, et al. Head coaches' attitudes towards injury prevention and use of related methods in professional basketball: a survey. *Phys Ther Sport* 2018;32:133–9.
- 65 Lim B-O, Lee YS, Kim JG, et al. Effects of sports injury prevention training on the biomechanical risk factors of anterior cruciate ligament injury in high school female basketball players. *Am J Sports Med* 2009;37:1728–34.
- 66 Noyes FR, Barber Westin SD. Anterior cruciate ligament injury prevention training in female athletes: a systematic review of injury reduction and results of athletic performance tests. *Sports Health* 2012;4:36–46.
- 67 DiStefano LJ, Padua DA, Blackburn JT, et al. Integrated injury prevention program improves balance and vertical jump height in children. *J Strength Cond Res* 2010;24:332–42.
- 68 Stefansen K, Smette I, Strandbu Åse. Understanding the increase in parents' involvement in organized youth sports. *Sport Educ Soc* 2018;23:162–72.
- 69 Wuerth S, Lee MJ, Alfermann D. Parental involvement and athletes' career in youth sport. *Psychol Sport Exerc* 2004;5:21–33.
- 70 Collard DCM, Verhagen EALM, Chinapaw MJM, et al. Effectiveness of a school-based physical activity injury prevention program: a cluster randomized controlled trial. *Arch Pediatr Adolesc Med* 2010;164:145–50.
- 71 Richmond SA, Kang J, Doyle-Baker PK, et al. A school-based injury prevention program to reduce sport injury risk and improve healthy outcomes in youth: a pilot cluster-randomized controlled trial. *Clin J Sport Med* 2016;26:291–8.
- 72 Padua DA, DiStefano LJ, Marshall SW, et al. Retention of movement pattern changes after a lower extremity injury prevention program is affected by program duration. *Am J Sports Med* 2012;40:300–6.
- 73 van Mechelen DM, van Mechelen W, Verhagen EALM. Sports injury prevention in your pocket?! Prevention apps assessed against the available scientific evidence: a review. *Br J Sports Med* 2014;48:878–82.