

# Modulation of the Expression of the GABAA Receptor $\beta_1$ and $\beta_3$ Subunits by Pretreatment with Quercetin in the KA Model of Epilepsy in Mice

## -The Effect of Quercetin on GABA<sub>A</sub> Receptor Beta Subunits-

Sahar Moghbelinejad<sup>1</sup>, Zahra Rashvand<sup>1</sup>, Fatemeh Khodabandehloo<sup>2</sup>, Ghazaleh Mohammadi<sup>2</sup>, Marjan Nassiri-Asl<sup>1\*</sup>

<sup>1</sup>Cellular and Molecular Research Center, Qazvin University of Medical Sciences, Qazvin, Iran

<sup>2</sup>Department of Molecular Medicine, Qazvin University of Medical Sciences, Qazvin, Iran

### Key Words

beta subunits, GABAA, gene expression, kainic acid, quercetin

### Abstract

**Objectives:** Quercetin is a flavonoid and an important dietary constituent of fruits and vegetables. In recent years, several pharmacological activities of quercetin, such as its neuroprotective activity and, more specifically, its anti-convulsant effects in animal models of epilepsy, have been reported. This study evaluated the role of quercetin pretreatment on gene expression of  $\gamma$ -amino butyric acid type A (GABA<sub>A</sub>) receptor beta subunits in kainic acid (KA)-induced seizures in mice.

**Methods:** The animals were divided into four groups: one saline group, one group in which seizures were induced by using KA (10 mg/kg) without quercetin pretreatment and two groups pretreated with quercetin (50 and 100 mg/kg) prior to seizures being induced by using KA. Next, the messenger ribonucleic acid (mRNA) levels of the GABA<sub>A</sub> receptor  $\beta$  subunits in the hippocampus of each animal were assessed at 2 hours and 7 days after KA administration. Quantitative real-time polymerase chain reaction (RT-PCR) assay was

used to detect mRNA content in hippocampal tissues.

**Results:** Pretreatments with quercetin at doses of 50 and 100 mg/kg prevented significant increases in the mRNA levels of the  $\beta_1$  and the  $\beta_3$  subunits of the GABA<sub>A</sub> receptor at 2 hours after KA injection. Pretreatment with quercetin (100 mg/kg) significantly inhibited  $\beta_1$  and  $\beta_3$  gene expression in the hippocampus at 7 days after KA injection. But, this inhibitory effect of quercetin at 50 mg/kg on the mRNA levels of the  $\beta_3$  subunit of the GABA<sub>A</sub> receptor was not observed at 7 days after KA administration.

**Conclusion:** These results suggest that quercetin (100 mg/kg) modulates the expression of the GABA<sub>A</sub> receptor  $\beta_1$  and  $\beta_3$  subunits in the KA model of epilepsy, most likely to prevent compensatory responses. This may be related to the narrow therapeutic dose range for the anticonvulsant activities of quercetin.

## 1. Introduction

Systemic administration of kainic acid (KA) induces seizures and brain damage associated with epilepsy [1].  $\gamma$ -amino butyric acid (GABA) is the main inhibitory neurotransmitter in the mammalian brain, and it is composed of five subunits that form a chloride channel. The messenger ribonucleic acid (mRNA)

Received: Apr 07, 2016 Reviewed: May 10, 2016 Accepted: May 15, 2016

© This is an Open-Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted noncommercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

© This paper meets the requirements of KS X ISO 9706, ISO 9706-1994 and ANSI/NISO Z39.48-1992 (Permanence of Paper).

\*Corresponding Author

Marjan Nassiri-Asl. Cellular and Molecular Research Center, Department of Pharmacology, Qazvin University of Medical Sciences, Qazvin, P.O. Box: 341197-5981, Iran.  
Tel: +98-283-333-6001 Fax: +98-283-332-4971  
E-mail: mnassiriasl@qums.ac.ir

molecules of the three  $\beta$  subunits have been shown to be distributed throughout the principal cell layers of the hippocampal and the parahippocampal areas in rats [2]. In addition, an association between mutations in the GABRB3 gene, encoding the  $\beta_3$  subunit of the GABA receptor (GABR), and families from Honduras and Mexico with no incidents of childhood epilepsy has been established [3]. The association between mutations in the GABRB3 gene, encoding the  $\beta_3$  subunit of the GABAR, has been established in families with childhood absence epilepsy (CAE) from Honduras and Mexico [3].

Quercetin (3,3',4',5,7-pentahydroxyflavone) is a flavonoid and an important dietary constituent of fruits and vegetables [4]. It has several biological effects [5], including anti-oxidative [6] anti-inflammatory [7], and neuroprotective [8-10] activities. Quercetin at a concentration of 30  $\mu$ M has been shown to have an inhibitory effect on the  $\alpha_1\beta_1\gamma_2$  GABA<sub>A</sub> and the  $P1$  GABA<sub>C</sub> receptors in *Xenopus laevis* oocytes [11]. GABA<sub>A<sub>AP</sub></sub> receptors are known to have low affinity for allosteric modulators of GABA such as benzodiazepine and barbiturates, and quercetin has been shown to antagonize GABA<sub>A<sub>AP</sub></sub> receptors in a dose-dependent way via a redox-independent allosteric mechanism [12].

Our previous studies demonstrated that quercetin had an anticonvulsant activity in acute and chronic models of pentylenetetrazole (PTZ) in rats [13, 14]. Furthermore, the expressions of the GABA<sub>A</sub> receptor  $\beta$  subunits were reported to have been altered at early time points during or soon after acute seizures induced by KA injection, [15]. Thus, this study aimed to assess the effects of quercetin by determining the variations in the mRNA levels in the  $\beta$  subunits of GABA<sub>A</sub> receptors in the KA model of epilepsy in mice.

## 2. Materials and Methods

A total of 70 male BALB/c mice (20 — 25 g) were maintained at constant room temperature (21°C  $\pm$  2°C) under a 12:12 hour light/dark cycle and had free access to food and water. All animal experiments were performed in accordance with the European Communities Council Directive of November 24, 1986 (86/609/EEC).

The 70 mice were divided into 4 groups as follows: Each mouse in the saline group (group 1, n = 10) received a daily intraperitoneal (i.p.) injection of saline and Tween 80 (10 mL/kg) for 6 days and two i.p. injections of saline 30 minutes apart on the 7<sup>th</sup> day. Each mouse in the KA group (group 2, n = 20) received a daily i.p. injection of saline for 6 days and an i.p. injection of saline 30 minutes prior to the i.p. injection of KA (10 mg/kg) on the 7<sup>th</sup> day. The mice in groups 3 (n = 20) and 4 (n = 20) received daily i.p. injections with quercetin at doses of 50 and 100 mg/kg, respectively, for 6 days and, on the 7<sup>th</sup> day i.p. injections of KA (10 mg/kg, i.p.) were administered 30 minutes after the quercetin injections. Fifty percent of mice in each group were anesthetized with i.p. injections of ketamine (60 mg/mL)/xylazine (6 mg/kg), and then sacrificed at 2 hours after KA administration, with the other fifty percent undergoing the same procedure at 7 days after KA administration.

The  $\beta$ -actin gene was used as an internal control for quantifying the expressions of the target genes. Primers for the

Gabrb1 and the Gabrb3 genes of subunits of the GABA<sub>A</sub> receptors were designed by using Gene Runner software (version 3.05). Primer sets were Gabrb1 F: 5' CCTGGCTCTACTGGAATATG3', R: 5' ATTGGCACTCTGGTCTTG3'; Gabrb3 F: 5' CTCCTACGCTGGTGTCTGAA3', R: 5' CACTGTGTTCCCACATGACA3';  $\beta$ -actin F: TTACTGAGCTGCGTTTTACAC, R: ACAAAGCCATGCCAATGTTG.


Quantitative real-time polymerase chain reaction (RT-PCR) assays were conducted with SYBR Green I Master Mix (Bioneer, Korea). Thermal cycling was performed using the ABI-7500 Sequence Detection System (Applied Biosystems, Foster, CA, USA) as follows: 10 minutes at 95°C for the first denaturation step, followed by 40 cycles at 95°C for a total of 20 seconds and 40 cycles at 60°C for a total of 45 seconds. The values of the cycle threshold (Ct) were used to quantify mRNA fold changes. In this regard, we used the  $2^{-\Delta\Delta Ct}$  calculating method by normalizing the Ct of the target mRNAs to the Ct of housekeeping gene  $\beta$ -actin in the treated and the control samples.

Data were expressed as means  $\pm$  standard errors of the mean (SEMs). Statistical analyses were performed using the one-way analysis of variance (ANOVA), followed by the post-hoc Turkey test for multiple comparisons. For the analyses, Prism software was used.  $P < 0.05$  was considered statistically significant.

## 3. Results

Compared to the mRNA levels of the GABA<sub>A</sub> receptor  $\beta_1$  subunit in the saline group, those levels in the KA group were significantly increased in the hippocampi of the mice at 2 hours and 7 days after KA administration (Figs. 1A, 1B). In contrast, no statistically significant changes were observed in the quercetin groups compared to the saline at those times. However, statistically significant differences were observed when comparing the KA group and the 100-mg/kg quercetin group at 7 days after KA administration. The mRNA levels of the GABA<sub>A</sub> receptor  $\beta_3$  subunit were significantly higher in the KA group at 2 hours and 7 days after KA administration than they were in the saline group (Figs. 2A, 2B). In contrast, no significant differences were observed in the mRNA levels of the GABA<sub>A</sub> receptor  $\beta_3$  subunit between the 50-mg/kg quercetin group at 2 hours after KA administration and the saline group (Fig. 2A). However, mRNA levels were increased in the quercetin 50-mg/kg group at 7 days after KA administration compared to the saline group (Fig. 2B).


Furthermore, the mRNA levels of the GABA<sub>A</sub> receptor  $\beta_3$  subunit observed in the 100-mg/kg quercetin group at 2 hours and 7 days after KA administration were similar to those observed in the saline group at those times (Figs. 2A, 2B). However, significant differences were found when comparing the KA group and both quercetin groups at 2 hours after KA administration (Fig. 2A). Moreover, significant differences were observed between the KA group and the quercetin groups at 7 days after KA administration (Fig. 2B).



**Figure 1** Effect of quercetin on the mRNA ratio of the expression of the  $\beta_1$  subunit of the GABAA receptor in KA-induced seizure in mice at (A) 2 and (B) 7 hours after KA administration.

\* $P < 0.01$  compared to the saline, † $P < 0.001$  compared to the KA group at D7,  $n = 10$ .

mRNA, messenger ribonucleic acid; GABAA,  $\gamma$ -amino butyric acid type A; KA, kainic acid.



**Figure 2** Effect of quercetin on the mRNA ratio of expression of the  $\beta_3$  subunit of the GABAA receptor in the KA model in mice at (A) 2 and (B) 7 hours after KA administration. † $P < 0.001$  compared to the KA at 2 hours, \* $P < 0.01$  compared to the KA group at 2 hours, \* $P < 0.001$  compared to the saline at 2 hours, \* $P < 0.01$  compared to the saline at D7,  $n = 10$ , \* $P < 0.001$  compared to the KA group at D7, † $P < 0.001$  compared to the saline and  $P < 0.05$  compared to the KA group at D7,  $n = 10$ .

mRNA, messenger ribonucleic acid; GABAA,  $\gamma$ -amino butyric acid type A; KA, kainic acid.

#### 4. Discussion

This study aimed at determining the effects of quercetin pretreatment on the expressions of the  $\beta_1$  and the  $\beta_3$  subunits of the GABAA receptor in a mouse KA-induced seizure model. The doses of quercetin used in this study were the same as those employed in a previous study assessing quercetin's anticonvulsant activity [13, 14]. Compared to the saline group, pretreatments with quercetin at doses of 50 and 100 mg/kg prevented significant increases in the mRNA levels of the  $\beta_1$  and the  $\beta_3$  subunits of the GABAA receptor at 2 hours after KA injection. In addition, pretreat-

ment with 100 mg/kg of quercetin inhibited increases in the mRNA levels of the  $\beta_3$  subunit of the GABAA receptor at 7 days after KA administration. Our results showed similar mRNA levels of the  $\beta_1$  and the  $\beta_3$  subunits of the GABAA receptor both in the 100-mg/kg quercetin group at 2 hours and day 7 after KA administration and the saline group, suggesting that high doses of quercetin have inhibitory effects on the gene expression in the  $\beta_1$  and the  $\beta_3$  subunits of the GABAA receptor. However, this inhibitory effect of quercetin at 50 mg/kg on the mRNA levels of the  $\beta_3$  subunit of the GABAA receptor was not observed at 7 days after KA administration. This may be related to the narrow therapeutic dose range for the anticonvulsant activities of

quercetin in different modes of seizure.

Of the three  $\beta$  subunits of the GABA<sub>A</sub> receptor, the mRNA level in the  $\beta_3$  subunit is the one that is increased in animal models and human temporal lobe epilepsy (TLE), as well as in the KA-induced-seizure model [16]. In addition, significant increases in the mRNA levels in the  $\beta_1$  and the  $\beta_3$  subunits have been observed in the granule cells of the dentate gyrus at 12 hours to 30 days and at 7 to 30 days after KA injection. These changes are consistent with the presumed compensatory up-regulation of GABA<sub>A</sub> receptors observed in the dentate molecular layer in epileptic rats [17].

## 5. Conclusion

In conclusion, pretreatment with quercetin significantly inhibited the gene expression of the GABA<sub>A</sub> receptor  $\beta$  subunits in the KA model of seizure. Quercetin (100 mg/kg) seems to modulate the expression of the GABA<sub>A</sub> receptor  $\beta$  subunits to prevent any progress in neurodegenerative and compensatory responses.

## Acknowledgment

This research was supported by Qazvin University of Medical Sciences.

## Conflict of interest

The authors declare that there are no conflict of interest.

## ORCID

Marjan Nassiri-Asl. <http://orcid.org/0000-0003-3701-0758>.

## References

1. Sperk G. Kainic acid seizures in the rat. *Prog Neurobiol.* 1994;42(1):1-32.
2. Drexel M, Kirchmair E, Sperk G. Changes in the expression of GABA<sub>A</sub> receptor subunit mRNAs in parahippocampal areas after kainic acid induced seizures. *Front Neural Circuits.* 2013;18:7-142.
3. Lachance-Touchette P, Martin C, Poulin C, Gravel M, Carmant L, Cossette P. Screening of GABRB3 in French-Canadian families with idiopathic generalized epilepsy. *Epilepsia.* 2010;51(9):1894-7.
4. Huang J, Zhu M, Tao Y, Wang S, Chen J, Sun W, *et al.* Therapeutic properties of quercetin on monosodium urate crystal-induced inflammation in rat. *J Pharm Pharmacol.* 2013;64(8):1119-27.
5. Kawabata K, Mukai R, Ishisaka A. Quercetin and related polyphenols: new insights and implications for their bioactivity and bioavailability. *Food Funct.* 2015;6(5):1399-417.
6. Boots AW, Drent M, de Boer VC, Bast A, Haenen GR. Quercetin reduces markers of oxidative stress and inflammation in sarcoidosis. *Clin Nutr.* 2011;30(4):506-12.
7. Loke WM, Proudfoot JM, Stewart S, McKinley AJ, Needs PW, Kroon PA, *et al.* Metabolic transformation has a profound effect on anti-inflammatory activity of flavonoids such as quercetin: lack of association between antioxidant and lipoxygenase inhibitory activity. *Biochem Pharmacol.* 2008;75(5):1045-53.
8. Yang T, Kong B, Gu JW, Kuang YQ, Cheng L, Yang WT, *et al.* Anti-apoptotic and anti-oxidative roles of quercetin after traumatic brain injury. *Cell Mol Neurobiol.* 2014;34(6):797-804.
9. Beckmann DV, Carvalho FB, Mazzanti CM, Dos Santos RP, Andrades AO, Aiello G, *et al.* Neuroprotective role of quercetin in locomotor activities and cholinergic neurotransmission in rats experimentally demyelinated with ethidium bromide. *Life Sci.* 2014;103(2):79-87.
10. Qu X, Qi D, Dong F, Wang B, Guo R, Luo M, *et al.* Quercetin improves hypoxia-ischemia induced cognitive deficits via promoting remyelination in neonatal rat. *Brain Res.* 2014;1553:31-40.
11. Goutman JD, Waxemberg MD, Doñate-Oliver F, Pomata PE, Calvo DJ. Flavonoid modulation of ionic currents mediated by GABA(A) and GABA(C) receptors. *Eur J Pharmacol.* 2003;461(2-3):79-87.
12. Calero CI, Beltrán González AN, Gasulla J, Alvarez S, Evelson P, Calvo DJ. Quercetin antagonism of GABA<sub>AA</sub> $\rho_1$  receptors is prevented by ascorbic acid through a redox-independent mechanism. *Eur J Pharmacol.* 2013;714(1-3):274-80.
13. Nassiri-Asl M, Moghbelinejad S, Abbasi E, Yonesi F, Haghghi MR, Lotfizadeh M, *et al.* Effects of quercetin on oxidative stress and memory retrieval in kindled rats. *Epilepsy Behav.* 2013;28(2):151-5.
14. Nassiri-Asl M, Hajiali F, Taghilo M, Abbasi E, Mohseni F, Yousefi F. Comparison between the effects of quercetin on seizure threshold in acute and chronic seizure models. *Toxicol Ind Health.* 2016;32(5):936-44.
15. Schwarzer C, Tsunashima K, Wanzenböck C, Fuchs K, Sieghart W, Sperk G. GABA(A) receptor subunits in the rat hippocampus II: altered distribution in kainic acid-induced temporal lobe epilepsy. *Neuroscience.* 1997;80(4):1001-17.
16. Sperk G, Schwarzer C, Tsunashima K, Kandlhofer S. Expression of GABA(A) receptor subunits in the hippocampus of the rat after kainic acid-induced seizures. *Epilepsy Res.* 1998;32(1-2):129-39.
17. Tsunashima K, Schwarzer C, Kirchmair E, Sieghart W, Sperk G. GABA(A) receptor subunits in the rat hippocampus III: altered messenger RNA expression in kainic acid-induced epilepsy. *Neuroscience.* 1997;80(4):1019-32.