

(2Z,3Z)-3,4-Dihydro-2H-1,4-benzothiazine-2,3-dione dioxime dihydrate

Ali Kakanejadifard^{a*} and Vahid Amani^b

^aDepartment of Chemistry, Faculty of Science, Lorestan University, Khorramabad, Iran, and ^bDepartment of Chemistry, Islamic Azad University, Shahr-e-Rey Branch, Tehran, Iran

Correspondence e-mail: alikakanejadifard@yahoo.com


Received 19 July 2008; accepted 23 July 2008

Key indicators: single-crystal X-ray study; $T = 120\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.046; wR factor = 0.108; data-to-parameter ratio = 16.1.

In the molecule of the title compound, $\text{C}_8\text{H}_{11}\text{N}_3\text{O}_4\text{S}$, the thiazine ring adopts an envelope conformation. In the crystal structure, intermolecular $\text{N}-\text{H}\cdots\text{O}$, $\text{O}-\text{H}\cdots\text{O}$ and $\text{O}-\text{H}\cdots\text{N}$ hydrogen bonds link the molecules.

Related literature

For related literature, see: Kakanejadifard, Niknam *et al.* (2007); Kakanejadifard, Saniei *et al.* (2007); Kakanejadifard & Niknam (2006); Kakanejadifard & Amani (2008). For general background, see: Jones *et al.* (1961); Schrauzer & Kohnle (1964); Yari *et al.* (2006); Hashemi *et al.* (2006); Ghiasvand *et al.* (2004, 2005); Kakanejadifard, Niknam & Zabardasti (2007); Gok & Kantekin (1997); Hughes (1981).



Experimental

Crystal data

$\text{C}_8\text{H}_{11}\text{N}_3\text{O}_2\text{S}\cdot 2\text{H}_2\text{O}$
 $M_r = 245.26$
 Orthorhombic, $Pbca$
 $a = 9.1636 (18)\text{ \AA}$
 $b = 9.8195 (18)\text{ \AA}$
 $c = 24.165 (4)\text{ \AA}$

$V = 2174.4 (7)\text{ \AA}^3$
 $Z = 8$
 Mo $K\alpha$ radiation
 $\mu = 0.30\text{ mm}^{-1}$
 $T = 120 (2)\text{ K}$
 $0.5 \times 0.5 \times 0.1\text{ mm}$

Data collection

Bruker SMART 1000 CCD area-detector diffractometer

Absorption correction: multi-scan (*SADABS*; Sheldrick, 1998)
 $T_{\min} = 0.859$, $T_{\max} = 0.974$

15018 measured reflections
 2337 independent reflections

1482 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.082$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.046$
 $wR(F^2) = 0.107$
 $S = 1.02$
 2337 reflections

145 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.43\text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.25\text{ e \AA}^{-3}$

Table 1
 Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1W-H1WA \cdots N11 ⁱ	0.85	2.11	2.931 (2)	162
O1W-H1WA \cdots N13 ⁱ	0.85	2.62	3.225 (2)	129
O1W-H1WB \cdots O2W ⁱ	0.85	2.00	2.845 (2)	171
O2W-H2WA \cdots N13 ⁱⁱ	0.85	2.02	2.853 (3)	168
N4-H4 \cdots O14	0.87	2.08	2.493 (2)	108
O2W-H2WB \cdots O1W	0.85	1.94	2.780 (2)	171
O12-H12 \cdots O2W ⁱⁱⁱ	0.82	1.89	2.699 (2)	171
O14-H14 \cdots O1W	0.82	1.85	2.673 (2)	179

Symmetry codes: (i) $x + \frac{1}{2}, y, -z + \frac{1}{2}$; (ii) $-x + 1, y + \frac{1}{2}, -z + \frac{1}{2}$; (iii) $-x + \frac{1}{2}, y - \frac{1}{2}, z$.

Data collection: *SMART* (Bruker, 1998); cell refinement: *SAINT-Plus* (Bruker, 1998); data reduction: *SAINT-Plus*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL*.

The authors acknowledge the Research Grant Council of Lorestan University for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HK2502).

References

- Bruker (1998). *SMART* and *SAINT-Plus*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Ghiasvand, A. R., Ghaderi, R. & Kakanejadifard, A. (2004). *Talanta*, **62**, 287–292.
- Ghiasvand, A. R., Shadabi, S., Kakanejadifard, A. & Khajehkolaki, A. (2005). *Bull. Korean Chem. Soc.* **26**, 781–785.
- Gok, Y. & Kantekin, H. (1997). *Polyhedron*, **16**, 2413–2420.
- Hashemi, P., Rahmani, Z., Kakanejadifard, A. & Niknam, E. (2006). *Anal. Sci.* **21**, 1297–1301.
- Hughes, M. N. (1981). *The Inorganic Chemistry of Biological Processes*. New York: Wiley.
- Jones, M. E. B., Thornton, D. A. & Webb, R. F. (1961). *Makromol. Chem.* **49**, 62–66.
- Kakanejadifard, A. & Amani, V. (2008). *Acta Cryst. E* **64**, o1512.
- Kakanejadifard, A. & Niknam, E. (2006). *Pol. J. Chem.* **80**, 1645–1649.
- Kakanejadifard, A., Niknam, E., Ranjbar, B. & Naderi-Manesh, H. (2007). *Synth. Commun.* **37**, 2753–2756.
- Kakanejadifard, A., Niknam, E. & Zabardasti, A. (2007). *J. Coord. Chem.* **60**, 677–681.
- Kakanejadifard, A., Saniei, A., Delfani, F., Farnia, M. & Najafi, G. R. (2007). *J. Heterocycl. Chem.* **44**, 717–718.
- Schrauzer, G. N. & Kohnle, J. (1964). *Chem. Ber.* **97**, 3056–3063.
- Sheldrick, G. M. (1998). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Yari, A., Azizi, S. & Kakanejadifard, A. (2006). *Sens. Actuators B*, **119**, 167–173.

supplementary materials

Acta Cryst. (2008). E64, o1628 [doi:10.1107/S1600536808023301]

(2Z,3Z)-3,4-Dihydro-2H-1,4-benzothiazine-2,3-dione dioxime dihydrate

A. Kakanejadifard and V. Amani

Comment

We have been interested in the synthesis and chemical behavior of vic-dioximes, in the past decade. In our investigations, the reaction of amines with dichloroglyoxime or cyanogendi-N-oxide resulted in various symmetrically substituted diaminoglyoxime derivatives, in which some of them were quite suitable to act, as donor species, towards some transition metal ions (Kakanejadifard, Niknam *et al.*, 2007; Kakanejadifard, Saniei *et al.*, 2007; Kakanejadifard & Niknam, 2006; Kakanejadifard & Amani, 2008). Some oximes are widely used for various purposes in organic, inorganic, bioinorganic, pigment, analytical, dyes and medical chemistry (Jones *et al.*, 1961; Schrauzer & Kohnle, 1964; Yari *et al.*, 2006; Hashemi *et al.*, 2006; Ghiasvand *et al.*, 2004; Ghiasvand *et al.*, 2005; Kakanejadifard, Niknam & Zabardasti, 2007). vic-Dioximes, containing mildly acidic hydroxyl groups and slightly basic nitrogen atoms, are amphoteric and their transition metal complexes have been widely investigated as analytical reagents (Gok & Kantekin, 1997), and models for biological systems such as vitamin B₁₂ (Hughes, 1981). We report herein the synthesis and crystal structure of the title compound.

In the molecule of the title compound (Fig. 1), the bond lengths and angles are within normal ranges. Ring A (S1/N4/C2/C3/C5/C6) adopts envelope conformation, with C2 atom displaced by 0.248 (3) Å from the plane of the other ring atoms. Ring B (C5–C10) is, of course, planar.

In the crystal structure, intermolecular N—H···O, O—H···O and O—H···N hydrogen bonds (Table 1) link the molecules (Fig. 2), in which they may be effective in the stabilization of the structure.

Experimental

For the preparation of the title compound, a solution of NaHCO₃ (0.05 g, 0.6 mmol) in aqueous EtOH (10 ml) was added to a magnetically stirred solution of dichloroglyoxime (1.57 g, 10 mmol) in aqueous EtOH (15 ml) and a solution of 2-aminothiophenole (1.25 g, 10 mmol) in EtOH (15 ml) at room temperature. The solution was stirred for 4 h and then the mixture was filtered. The filtrate was placed at room temperature for 24 h. The gray precipitate was removed by filtration and precipitate was washed with cold THF. It was recrystallized from 2-propanol in one week (yield; 0.31 g, 74.2%, m.p. 492 K).

Refinement

H atoms were positioned geometrically, with O—H = 0.82 Å (for OH), N—H = 0.8691 Å (for NH), O—H = 0.8499–0.8501 Å (for H₂O) and C—H = 0.93 Å for aromatic H, and constrained to ride on their parent atoms with U_{iso}(H) = xU_{eq}(C,N,O), where x = 1.5 for OH H, x = 0.95 for NH H, and x = 1.2 for all other H atoms.

supplementary materials

Figures




Fig. 1. The molecular structure of the title molecule, with the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level.




Fig. 2. A packing diagram of the title molecule. Hydrogen bonds are shown as dashed lines.

(2Z,3Z)-3,4-Dihydro-2H-1,4-benzothiazine-2,3-dione dioxime dihydrate

Crystal data

C ₈ H ₇ N ₃ O ₂ S·2H ₂ O	$F_{000} = 1024$
$M_r = 245.26$	$D_x = 1.498 \text{ Mg m}^{-3}$
Orthorhombic, <i>Pbca</i>	Mo $K\alpha$ radiation
Hall symbol: -P 2ac 2ab	$\lambda = 0.71073 \text{ \AA}$
$a = 9.1636 (18) \text{ \AA}$	Cell parameters from 910 reflections
$b = 9.8195 (18) \text{ \AA}$	$\theta = 2.5\text{--}26.3^\circ$
$c = 24.165 (4) \text{ \AA}$	$\mu = 0.30 \text{ mm}^{-1}$
$V = 2174.4 (7) \text{ \AA}^3$	$T = 120 (2) \text{ K}$
$Z = 8$	Plate, yellow
	$0.5 \times 0.5 \times 0.1 \text{ mm}$

Data collection

Bruker SMART 1000 CCD area-detector diffractometer	2337 independent reflections
Radiation source: fine-focus sealed tube	1482 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.082$
$T = 120(2) \text{ K}$	$\theta_{\max} = 27.0^\circ$
ϕ and ω scans	$\theta_{\min} = 1.7^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 1998)	$h = -11 \rightarrow 11$
$T_{\min} = 0.859, T_{\max} = 0.974$	$k = -12 \rightarrow 12$
15018 measured reflections	$l = -30 \rightarrow 28$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: difference Fourier map

$R[F^2 > 2\sigma(F^2)] = 0.046$	H-atom parameters constrained
$wR(F^2) = 0.107$	$w = 1/[\sigma^2(F_o^2) + (0.0534P)^2 + 0.01P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.02$	$(\Delta/\sigma)_{\max} < 0.001$
2337 reflections	$\Delta\rho_{\max} = 0.43 \text{ e \AA}^{-3}$
145 parameters	$\Delta\rho_{\min} = -0.25 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\text{sigma}(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R- factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.18557 (6)	0.97098 (6)	0.44569 (2)	0.02751 (18)
O1W	0.74924 (16)	1.21073 (15)	0.24961 (6)	0.0294 (4)
H1WA	0.7567	1.1544	0.2230	0.035*
H1WB	0.8298	1.2523	0.2539	0.035*
O2W	0.50348 (16)	1.37490 (17)	0.23976 (6)	0.0302 (4)
H2WA	0.5201	1.4420	0.2185	0.036*
H2WB	0.5838	1.3317	0.2411	0.036*
O12	0.07401 (16)	0.93917 (17)	0.34431 (6)	0.0326 (4)
H12	0.0423	0.9158	0.3140	0.049*
O14	0.61786 (17)	1.16110 (16)	0.34627 (6)	0.0316 (4)
H14	0.6570	1.1757	0.3163	0.047*
N4	0.4660 (2)	1.14780 (19)	0.43170 (7)	0.0231 (4)
H4	0.5489	1.1895	0.4276	0.022*
N11	0.20659 (19)	1.00628 (19)	0.33728 (8)	0.0268 (5)
N13	0.48394 (19)	1.09374 (19)	0.33772 (8)	0.0236 (4)
C2	0.2680 (2)	1.0265 (2)	0.38436 (9)	0.0223 (5)
C3	0.4132 (2)	1.0922 (2)	0.38433 (9)	0.0211 (5)
C5	0.4075 (2)	1.1389 (2)	0.48497 (9)	0.0212 (5)
C6	0.2847 (2)	1.0617 (2)	0.49653 (9)	0.0228 (5)
C7	0.2320 (2)	1.0540 (2)	0.55071 (9)	0.0269 (5)
H7	0.1500	1.0014	0.5584	0.032*
C8	0.3006 (3)	1.1236 (2)	0.59269 (9)	0.0289 (6)
H8	0.2651	1.1183	0.6287	0.035*
C9	0.4222 (3)	1.2015 (2)	0.58121 (10)	0.0302 (6)

supplementary materials

H9	0.4683	1.2493	0.6094	0.036*
C10	0.4756 (3)	1.2084 (2)	0.52783 (9)	0.0269 (6)
H10	0.5583	1.2602	0.5205	0.032*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0211 (3)	0.0357 (3)	0.0257 (3)	-0.0062 (3)	0.0008 (2)	0.0020 (3)
O1W	0.0243 (9)	0.0344 (9)	0.0295 (9)	-0.0021 (7)	0.0050 (7)	-0.0041 (8)
O2W	0.0251 (8)	0.0325 (9)	0.0329 (10)	0.0035 (7)	0.0048 (7)	0.0067 (8)
O12	0.0237 (9)	0.0466 (11)	0.0275 (9)	-0.0140 (8)	-0.0027 (7)	-0.0017 (8)
O14	0.0269 (9)	0.0398 (10)	0.0282 (9)	-0.0107 (8)	0.0022 (7)	0.0030 (8)
N4	0.0217 (9)	0.0262 (11)	0.0213 (10)	-0.0026 (8)	0.0009 (8)	-0.0002 (8)
N11	0.0201 (10)	0.0316 (11)	0.0287 (11)	-0.0038 (8)	-0.0016 (8)	0.0025 (9)
N13	0.0209 (10)	0.0249 (10)	0.0251 (10)	-0.0049 (8)	-0.0011 (8)	0.0008 (8)
C2	0.0200 (11)	0.0249 (11)	0.0220 (12)	0.0024 (10)	0.0004 (9)	0.0010 (10)
C3	0.0215 (11)	0.0200 (11)	0.0219 (12)	0.0014 (10)	-0.0025 (9)	0.0018 (10)
C5	0.0220 (11)	0.0198 (11)	0.0218 (12)	0.0067 (10)	0.0003 (9)	0.0017 (10)
C6	0.0210 (12)	0.0239 (12)	0.0236 (13)	0.0061 (9)	-0.0015 (9)	0.0033 (9)
C7	0.0224 (11)	0.0316 (14)	0.0268 (13)	0.0050 (10)	0.0023 (10)	0.0060 (11)
C8	0.0320 (13)	0.0327 (13)	0.0219 (13)	0.0095 (11)	0.0042 (11)	0.0021 (11)
C9	0.0413 (15)	0.0251 (13)	0.0243 (14)	0.0050 (11)	-0.0033 (11)	-0.0024 (10)
C10	0.0310 (13)	0.0209 (12)	0.0289 (14)	0.0006 (10)	-0.0003 (11)	0.0019 (10)

Geometric parameters (\AA , $^\circ$)

S1—C2	1.750 (2)	C2—C3	1.479 (3)
S1—C6	1.769 (2)	C5—C6	1.385 (3)
S1—C5	2.785 (2)	C5—C10	1.388 (3)
S1—C3	2.823 (2)	C6—C7	1.398 (3)
O12—N11	1.393 (2)	C7—C8	1.376 (3)
O12—H12	0.8200	C7—H7	0.9300
O14—N13	1.409 (2)	C8—C9	1.380 (3)
O14—C3	2.195 (3)	C8—H8	0.9300
O14—H14	0.8200	C9—C10	1.381 (3)
N4—C3	1.357 (3)	C9—H9	0.9300
N4—C5	1.397 (3)	C10—H10	0.9300
N4—H4	0.8691	O1W—H1WA	0.8501
N11—C2	1.285 (3)	O1W—H1WB	0.8499
N11—C3	2.364 (3)	O2W—H2WA	0.8500
N13—C3	1.299 (3)	O2W—H2WB	0.8500
C2—S1—C6	102.11 (11)	C10—C5—S1	150.86 (16)
C2—S1—C5	77.83 (9)	N4—C5—S1	90.18 (13)
C6—S1—C3	76.88 (9)	C9—C5—S1	121.39 (10)
C5—S1—C3	52.39 (6)	C7—C5—S1	62.11 (8)
N11—O12—H12	109.5	C3—C5—S1	64.59 (7)
N13—O14—H14	109.5	C6—C5—C5 ⁱ	85.42 (14)
C3—O14—H14	142.8	C10—C5—C5 ⁱ	90.62 (14)

C3—N4—C5	127.96 (19)	N4—C5—C5 ⁱ	93.28 (14)
C3—N4—H4	113.8	C9—C5—C5 ⁱ	88.25 (10)
C5—N4—H4	118.1	C7—C5—C5 ⁱ	84.92 (10)
C2—N11—O12	110.30 (18)	C3—C5—C5 ⁱ	92.99 (10)
O12—N11—C3	143.99 (15)	S1—C5—C5 ⁱ	87.74 (9)
C3—N13—O14	108.23 (17)	C5—C6—C7	120.0 (2)
N11—C2—C3	117.44 (19)	C5—C6—S1	123.56 (17)
N11—C2—S1	120.86 (17)	C7—C6—S1	116.47 (17)
C3—C2—S1	121.64 (16)	C5—C6—C9	60.51 (13)
N13—C3—N4	123.3 (2)	C7—C6—C9	59.46 (14)
N13—C3—C2	117.02 (19)	S1—C6—C9	175.77 (13)
N4—C3—C2	119.71 (19)	C8—C7—C6	120.4 (2)
N4—C3—O14	85.70 (14)	C6—C7—C9	90.21 (16)
C2—C3—O14	154.59 (16)	C8—C7—C5	90.50 (16)
N13—C3—N11	89.24 (14)	C9—C7—C5	60.34 (10)
N4—C3—N11	146.56 (17)	C8—C7—H7	119.8
O14—C3—N11	126.34 (11)	C6—C7—H7	119.8
N13—C3—C5	149.31 (17)	C9—C7—H7	150.0
C2—C3—C5	93.51 (14)	C5—C7—H7	149.7
O14—C3—C5	111.88 (11)	C7—C8—C9	119.8 (2)
N11—C3—C5	121.44 (11)	C7—C8—H8	120.1
N13—C3—S1	145.97 (16)	C9—C8—H8	120.1
N4—C3—S1	89.42 (13)	C8—C9—C10	120.1 (2)
O14—C3—S1	168.89 (11)	C10—C9—C7	90.00 (16)
N11—C3—S1	60.67 (7)	C8—C9—C5	90.44 (16)
C5—C3—S1	63.02 (7)	C7—C9—C5	60.37 (9)
N13—C3—C9 ⁱ	90.05 (14)	C8—C9—C6	60.40 (14)
N4—C3—C9 ⁱ	88.63 (13)	C10—C9—C6	59.67 (14)
C2—C3—C9 ⁱ	91.67 (14)	C8—C9—C3 ⁱ	80.63 (14)
O14—C3—C9 ⁱ	89.14 (9)	C10—C9—C3 ⁱ	96.59 (15)
N11—C3—C9 ⁱ	99.92 (10)	C7—C9—C3 ⁱ	83.30 (9)
C5—C3—C9 ⁱ	85.70 (8)	C5—C9—C3 ⁱ	92.62 (9)
S1—C3—C9 ⁱ	80.76 (7)	C6—C9—C3 ⁱ	87.82 (8)
C6—C5—C10	118.9 (2)	C8—C9—H9	120.0
C6—C5—N4	122.1 (2)	C10—C9—H9	120.0
C10—C5—N4	119.0 (2)	C7—C9—H9	150.0
C6—C5—C9	89.45 (15)	C5—C9—H9	149.6
N4—C5—C9	148.43 (17)	C6—C9—H9	179.4
C10—C5—C7	88.76 (15)	C3 ⁱ —C9—H9	92.8
N4—C5—C7	152.27 (17)	C9—C10—C5	120.9 (2)
C9—C5—C7	59.29 (10)	C9—C10—H10	119.5
C6—C5—C3	96.54 (15)	C5—C10—H10	119.5
C10—C5—C3	144.53 (17)	H1WA—O1W—H1WB	109.6
C9—C5—C3	173.96 (13)	H2WA—O2W—H2WB	104.7
C7—C5—C3	126.70 (12)		
O12—N11—C2—C3	177.69 (18)	C2—S1—C5—C9	170.77 (13)

supplementary materials

O12—N11—C2—S1	0.5 (3)	C6—S1—C5—C9	-1.59 (17)
C3—N11—C2—S1	-177.2 (3)	C3—S1—C5—C9	179.04 (14)
C6—S1—C2—N11	-164.88 (19)	C2—S1—C5—C7	171.89 (11)
C5—S1—C2—N11	-168.1 (2)	C6—S1—C5—C7	-0.47 (18)
C3—S1—C2—N11	177.1 (3)	C3—S1—C5—C7	-179.84 (10)
C6—S1—C2—C3	18.0 (2)	C2—S1—C5—C3	-8.27 (10)
C5—S1—C2—C3	14.80 (18)	C6—S1—C5—C3	179.4 (2)
O14—N13—C3—N4	-0.1 (3)	C2—S1—C5—C5 ⁱ	-102.67 (11)
O14—N13—C3—C2	179.59 (18)	C6—S1—C5—C5 ⁱ	85.0 (2)
O14—N13—C3—N11	171.56 (13)	C3—S1—C5—C5 ⁱ	-94.39 (10)
O14—N13—C3—C5	-6.9 (4)	C10—C5—C6—C7	-0.4 (3)
O14—N13—C3—S1	-161.9 (2)	N4—C5—C6—C7	178.8 (2)
O14—N13—C3—C9 ⁱ	-88.52 (14)	C9—C5—C6—C7	-0.5 (2)
C5—N4—C3—N13	-172.2 (2)	C3—C5—C6—C7	-179.75 (18)
C5—N4—C3—C2	8.1 (3)	S1—C5—C6—C7	-179.2 (3)
C5—N4—C3—O14	-172.3 (2)	C5 ⁱ —C5—C6—C7	87.8 (2)
C5—N4—C3—N11	23.0 (4)	C10—C5—C6—S1	178.75 (16)
C5—N4—C3—S1	-2.3 (2)	N4—C5—C6—S1	-2.1 (3)
C5—N4—C3—C9 ⁱ	-83.1 (2)	C9—C5—C6—S1	178.65 (15)
N11—C2—C3—N13	-16.8 (3)	C7—C5—C6—S1	179.2 (3)
S1—C2—C3—N13	160.36 (17)	C3—C5—C6—S1	-0.57 (19)
N11—C2—C3—N4	162.9 (2)	C5 ⁱ —C5—C6—S1	-93.07 (16)
S1—C2—C3—N4	-19.9 (3)	C10—C5—C6—C9	0.10 (16)
N11—C2—C3—O14	-16.2 (5)	N4—C5—C6—C9	179.3 (2)
S1—C2—C3—O14	160.9 (3)	C7—C5—C6—C9	0.5 (2)
S1—C2—C3—N11	177.2 (3)	C3—C5—C6—C9	-179.22 (13)
N11—C2—C3—C5	166.47 (19)	S1—C5—C6—C9	-178.65 (15)
S1—C2—C3—C5	-16.35 (19)	C5 ⁱ —C5—C6—C9	88.29 (10)
N11—C2—C3—S1	-177.2 (3)	C2—S1—C6—C5	-7.6 (2)
N11—C2—C3—C9 ⁱ	-107.7 (2)	C3—S1—C6—C5	0.51 (17)
S1—C2—C3—C9 ⁱ	69.45 (17)	C2—S1—C6—C7	171.56 (17)
N13—O14—C3—N4	179.9 (2)	C5—S1—C6—C7	179.2 (3)
N13—O14—C3—C2	-0.9 (4)	C3—S1—C6—C7	179.71 (18)
N13—O14—C3—N11	-10.50 (17)	C5—C6—C7—C8	0.6 (3)
N13—O14—C3—C5	176.2 (2)	S1—C6—C7—C8	-178.67 (17)
N13—O14—C3—S1	115.7 (6)	C9—C6—C7—C8	0.03 (17)
N13—O14—C3—C9 ⁱ	91.21 (18)	C5—C6—C7—C9	0.5 (2)
C2—N11—C3—N13	165.1 (3)	S1—C6—C7—C9	-178.70 (13)
O12—N11—C3—N13	161.4 (2)	S1—C6—C7—C5	-179.2 (3)
C2—N11—C3—N4	-27.6 (3)	C9—C6—C7—C5	-0.5 (2)
O12—N11—C3—N4	-31.3 (4)	C6—C5—C7—C8	-179.5 (3)
O12—N11—C3—C2	-3.7 (3)	C10—C5—C7—C8	0.11 (18)
C2—N11—C3—O14	171.4 (3)	N4—C5—C7—C8	178.2 (3)
O12—N11—C3—O14	167.7 (2)	C9—C5—C7—C8	-0.13 (13)
C2—N11—C3—C5	-15.9 (2)	C3—C5—C7—C8	-179.20 (15)
O12—N11—C3—C5	-19.6 (3)	S1—C5—C7—C8	-179.02 (15)

C2—N11—C3—S1	1.7 (2)	C5 ⁱ —C5—C7—C8	90.85 (15)
O12—N11—C3—S1	−2.0 (2)	C10—C5—C7—C6	179.6 (3)
C2—N11—C3—C9 ⁱ	75.1 (2)	N4—C5—C7—C6	−2.3 (4)
O12—N11—C3—C9 ⁱ	71.4 (3)	C9—C5—C7—C6	179.4 (2)
C2—S1—C3—N13	−32.4 (3)	C3—C5—C7—C6	0.3 (2)
C6—S1—C3—N13	165.8 (3)	S1—C5—C7—C6	0.50 (19)
C5—S1—C3—N13	166.0 (3)	C5 ⁱ —C5—C7—C6	−89.6 (2)
C2—S1—C3—N4	162.8 (3)	C6—C5—C7—C9	−179.4 (2)
C6—S1—C3—N4	0.88 (13)	C10—C5—C7—C9	0.24 (13)
C5—S1—C3—N4	1.15 (11)	N4—C5—C7—C9	178.4 (4)
C6—S1—C3—C2	−161.9 (2)	C3—C5—C7—C9	−179.07 (15)
C5—S1—C3—C2	−161.6 (2)	S1—C5—C7—C9	−178.89 (10)
C2—S1—C3—O14	−133.4 (7)	C5 ⁱ —C5—C7—C9	90.98 (10)
C6—S1—C3—O14	64.8 (6)	C6—C7—C8—C9	−0.1 (3)
C5—S1—C3—O14	65.0 (6)	C5—C7—C8—C9	0.2 (2)
C2—S1—C3—N11	−1.56 (18)	C7—C8—C9—C10	−0.6 (3)
C6—S1—C3—N11	−163.45 (10)	C7—C8—C9—C5	−0.2 (2)
C5—S1—C3—N11	−163.18 (10)	C7—C8—C9—C6	0.03 (17)
C2—S1—C3—C5	161.6 (2)	C7—C8—C9—C3 ⁱ	−92.8 (2)
C6—S1—C3—C5	−0.27 (9)	C6—C7—C9—C8	180.0 (3)
C2—S1—C3—C9 ⁱ	−108.5 (2)	C5—C7—C9—C8	−179.7 (3)
C6—S1—C3—C9 ⁱ	89.59 (9)	C8—C7—C9—C10	179.5 (3)
C5—S1—C3—C9 ⁱ	89.86 (8)	C6—C7—C9—C10	−0.55 (18)
C3—N4—C5—C6	3.4 (3)	C5—C7—C9—C10	−0.24 (13)
C3—N4—C5—C10	−177.4 (2)	C8—C7—C9—C5	179.7 (3)
C3—N4—C5—C9	−177.9 (2)	C6—C7—C9—C5	−0.31 (12)
C3—N4—C5—C7	4.8 (5)	C8—C7—C9—C6	−180.0 (3)
C3—N4—C5—S1	2.3 (2)	C5—C7—C9—C6	0.31 (12)
C3—N4—C5—C5 ⁱ	90.1 (2)	C8—C7—C9—C3 ⁱ	82.9 (2)
N13—C3—C5—C6	−164.3 (3)	C6—C7—C9—C3 ⁱ	−97.18 (14)
N4—C3—C5—C6	−177.1 (3)	C5—C7—C9—C3 ⁱ	−96.88 (9)
C2—C3—C5—C6	9.93 (18)	C6—C5—C9—C8	0.44 (18)
O14—C3—C5—C6	−168.81 (14)	C10—C5—C9—C8	−179.4 (3)
N11—C3—C5—C6	17.53 (18)	N4—C5—C9—C8	−178.4 (3)
S1—C3—C5—C6	0.34 (11)	C7—C5—C9—C8	0.13 (13)
C9 ⁱ —C3—C5—C6	−81.48 (13)	S1—C5—C9—C8	1.28 (18)
N13—C3—C5—C10	16.7 (5)	C5 ⁱ —C5—C9—C8	−85.00 (15)
N4—C3—C5—C10	3.9 (3)	C6—C5—C9—C10	179.8 (3)
C2—C3—C5—C10	−169.0 (3)	N4—C5—C9—C10	1.0 (3)
O14—C3—C5—C10	12.2 (3)	C7—C5—C9—C10	179.5 (3)
N11—C3—C5—C10	−161.4 (2)	S1—C5—C9—C10	−179.3 (3)
S1—C3—C5—C10	−178.6 (3)	C5 ⁱ —C5—C9—C10	94.4 (2)
C9 ⁱ —C3—C5—C10	99.5 (3)	C6—C5—C9—C7	0.31 (12)
N13—C3—C5—N4	12.8 (3)	C10—C5—C9—C7	−179.5 (3)
C2—C3—C5—N4	−173.0 (3)	N4—C5—C9—C7	−178.5 (3)
O14—C3—C5—N4	8.3 (2)	S1—C5—C9—C7	1.15 (10)

supplementary materials

N11—C3—C5—N4	-165.4 (3)	C5 ⁱ —C5—C9—C7	-85.13 (10)
S1—C3—C5—N4	177.4 (3)	C10—C5—C9—C6	-179.8 (3)
C9 ⁱ —C3—C5—N4	95.6 (2)	N4—C5—C9—C6	-178.8 (4)
N13—C3—C5—C7	-164.5 (3)	C7—C5—C9—C6	-0.31 (12)
N4—C3—C5—C7	-177.2 (3)	S1—C5—C9—C6	0.84 (9)
C2—C3—C5—C7	9.77 (19)	C5 ⁱ —C5—C9—C6	-85.44 (14)
O14—C3—C5—C7	-168.97 (12)	C6—C5—C9—C3 ⁱ	81.08 (13)
N11—C3—C5—C7	17.37 (19)	C10—C5—C9—C3 ⁱ	-98.7 (2)
S1—C3—C5—C7	0.18 (11)	N4—C5—C9—C3 ⁱ	-97.8 (3)
C9 ⁱ —C3—C5—C7	-81.64 (13)	C7—C5—C9—C3 ⁱ	80.77 (9)
N13—C3—C5—S1	-164.6 (3)	S1—C5—C9—C3 ⁱ	81.92 (11)
N4—C3—C5—S1	-177.4 (3)	C5 ⁱ —C5—C9—C3 ⁱ	-4.36 (9)
C2—C3—C5—S1	9.60 (11)	C5—C6—C9—C8	-179.5 (2)
O14—C3—C5—S1	-169.15 (11)	C7—C6—C9—C8	-0.03 (17)
N11—C3—C5—S1	17.20 (10)	C5—C6—C9—C10	-0.11 (16)
C9 ⁱ —C3—C5—S1	-81.81 (7)	C7—C6—C9—C10	179.4 (2)
N13—C3—C5—C5 ⁱ	-78.6 (3)	C5—C6—C9—C7	-179.5 (2)
N4—C3—C5—C5 ⁱ	-91.3 (2)	C7—C6—C9—C5	179.5 (2)
C2—C3—C5—C5 ⁱ	95.66 (15)	C5—C6—C9—C3 ⁱ	-99.04 (13)
O14—C3—C5—C5 ⁱ	-83.09 (12)	C7—C6—C9—C3 ⁱ	80.43 (14)
N11—C3—C5—C5 ⁱ	103.26 (13)	C8—C9—C10—C5	0.7 (3)
S1—C3—C5—C5 ⁱ	86.06 (9)	C7—C9—C10—C5	0.4 (2)
C9 ⁱ —C3—C5—C5 ⁱ	4.25 (9)	C6—C9—C10—C5	0.11 (16)
C2—S1—C5—C6	172.4 (2)	C3 ⁱ —C9—C10—C5	83.7 (2)
C3—S1—C5—C6	-179.4 (2)	C6—C5—C10—C9	-0.2 (3)
C2—S1—C5—C10	170.1 (3)	N4—C5—C10—C9	-179.4 (2)
C6—S1—C5—C10	-2.2 (3)	C7—C5—C10—C9	-0.4 (2)
C3—S1—C5—C10	178.4 (3)	C3—C5—C10—C9	178.6 (2)
C2—S1—C5—N4	-9.39 (13)	S1—C5—C10—C9	1.1 (4)
C6—S1—C5—N4	178.2 (3)	C5 ⁱ —C5—C10—C9	-85.3 (2)
C3—S1—C5—N4	-1.12 (11)		


Symmetry codes: (i) $-x+1, -y+2, -z+1$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D\cdots H\cdots A$	$D—H$	$H\cdots A$	$D\cdots A$	$D—H\cdots A$
O1W—H1WA…N11 ⁱⁱ	0.85	2.11	2.931 (2)	162
O1W—H1WA…N13 ⁱⁱ	0.85	2.62	3.225 (2)	129
O1W—H1WB…O2W ⁱⁱ	0.85	2.00	2.845 (2)	171
O2W—H2WA…N13 ⁱⁱⁱ	0.85	2.02	2.853 (3)	168
N4—H4…O14	0.87	2.08	2.493 (2)	108
O2W—H2WB…O1W	0.85	1.94	2.780 (2)	171
O12—H12…O2W ^{iv}	0.82	1.89	2.699 (2)	171
O14—H14…O1W	0.82	1.85	2.673 (2)	179

Symmetry codes: (ii) $x+1/2, y, -z+1/2$; (iii) $-x+1, y+1/2, -z+1/2$; (iv) $-x+1/2, y-1/2, z$.

Fig. 1



supplementary materials

Fig. 2

