



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Original Article

Psychological Distress and Heavy Alcohol Consumption Among U.S. Young Women During the COVID-19 Pandemic

Hyeouk Chris Hahm, PhD^{a,*}, Kelly C. Hsu, BS^b, Sunah Hyun, PhD^c,
Kaiden Kane, BS^d, Cindy H. Liu, PhD^e

^a Boston University, School of Social Work, Boston, Massachusetts

^b Tufts University, Medford, Massachusetts

^c Department of Pediatric Newborn Medicine, Brigham and Women's Hospital, Boston, Massachusetts

^d Boston Children's Hospital, Boston, Massachusetts

^e Harvard Medical School, Brigham and Women's Hospital, Boston, Massachusetts

Article history: Received 16 March 2022; Received in revised form 24 August 2022; Accepted 15 September 2022

A B S T R A C T

Introduction: The predictors of heavy drinking among U.S. young women during the coronavirus disease 2019 (COVID-19) pandemic are not well-examined. This study aims to determine the prospective relationship between COVID-19 psychological distress and heavy alcohol use among U.S. young adult women.

Design: This study used the COVID-19 Adult Resilience Experiences Study (CARES) data collected from April 13, 2020, to August 31, 2020 (T₁) and September 21, 2020, to March 15, 2021 (T₂). Among the young adults (ages 18–30) who participated in both surveys, a total of 684 identified as women (including transgender women) were included in the analysis. Three or more drinks as a typical intake were defined as heavy drinking. Psychological distress in T₁ was measured using scores for financial stress, COVID-19–related worry, and COVID-19–related grief. A series of logistic regression analyses were performed to identify risk factors associated with young women's heavy drinking during the pandemic.

Results: Twenty-two and one-half percent of young women reported heavy drinking in a typical intake. After controlling for covariates, women who reported high levels of COVID-19–related grief were more likely to report heavy drinking (odds ratio, 1.06; 95% confidence interval [CI], 1.01–1.13; $p < .05$). Those with high levels of COVID-19–related worry were less likely to report drinking heavily (odds ratio, 0.90; 95% confidence interval, 0.85–0.97; $p < .01$).

Conclusions: Those who suffer from a deeper sense of COVID-19–related grief are particularly at risk of heavy drinking and should be targeted for outreach and clinical intervention. Further research is necessary to determine the long-term impacts of the pandemic on heavy drinking among young women and should include a more comprehensive assessment of psychological distress.

© 2022 Jacobs Institute of Women's Health, George Washington University. Published by Elsevier Inc. All rights reserved.

Heavy drinking is considered the riskiest form of alcohol consumption and is defined as the consumption of four or more drinks on any day for men and three or more drinks on any day for women (National Institute on Alcohol Abuse and Alcoholism, 2021). According to the National Survey on Drug Use and Health, the highest rate of alcohol use between 2019 and 2020 was found among individuals, both men and women, between the ages of

18 and 25. (Substance Abuse and Mental Health Services Administration, 2022).

Heavy drinking is a known risk for problematic behaviors and poor health outcomes, including interpersonal violence, property crimes, motor vehicle crashes, suicide, heart disease, sexually transmitted infections, liver disease, and cancer (Centers for Disease Control and Prevention [CDC], 2021). The financial cost of heavy alcohol use was estimated at approximately \$249 billion in 2010 (Sacks, Gonzales, Bouchery, Tomedi, & Brewer, 2015). Thus, from legal, financial, and public health standpoints, it is critical to understand the factors associated with heavy drinking among young adults.

Women with heavy drinking problems are particularly vulnerable to multiple dimensions of suffering. They are more

Funding Statement: Supported by the National Science Foundation (2027553) award (H.C.H., C.H.L.), a Mary A. Tynan Faculty Fellowship, and an NIMH K23 MH 107714-01 A1 award (to C.H.L.).

* Correspondence to: Hyeouk Chris Hahm, PhD, Professor Boston University, School of Social Work, 264 Bay State Road, #222, Boston, MA 02215. Phone: (617) 353-3925.

E-mail address: hahm@bu.edu (H.C. Hahm).

likely than their male counterparts to experience depression (King, Bernardy, & Hauner, 2003; Boykoff et al., 2010), develop a higher risk of psychiatric and physical comorbidities (Erol & Karpyak, 2015; Peltier et al., 2019), attempt suicide (Glasheen, Pemberton, Lipari, Copello, & Mattson, 2015), experience greater physiological consequences (Erol & Karpyak, 2015; Peltier et al., 2019), suffer from a higher level of sexual victimization and revictimization (Bryan et al., 2016), and report stigma as an alcohol treatment barrier (Khan et al., 2013). Women's alcohol consumption is associated with an increased risk of having liver disease, breast and other cancers, and negative impacts on the brain and heart (CDC, 2020). Furthermore, heavy alcohol use in pregnant women can lead to a higher risk of stillbirth, miscarriage, and sudden infant death syndrome after birth (CDC, 2020). Despite documented gender differences in alcohol use, few studies have examined the specific factors associated with heavy drinking among women during the COVID-19 pandemic (CDC, 2020). Given the evidence that women are more susceptible to alcohol-related health consequences (Peltier et al., 2020), it is essential to investigate young women's risk factors for heavy drinking during the COVID-19 pandemic.

The COVID-19 pandemic led to an enormous amount of stress owing to the rapid transmission of the disease and its associated morbidity and mortality (Sharif Nia et al., 2022). Combined with the unpredictable nature of the disease, a number of COVID-19-specific distresses have emerged, including worries about the health of self and loved ones, increased fear of transmission, feeling a sense of loss and grief, and financial stress (Bertuccio & Runion, 2020; Lechner et al., 2021). The stress and coping hypothesis indicate that individuals often cope with stress with either adaptive or maladaptive behaviors (Cohen & Wills, 1985). Alcohol use may be a behavior that mitigates COVID-19-related distress. In fact, the COVID-19 pandemic led to a dramatic increase in alcohol consumption; the average number of drinks between February and April of 2020 increased by 29% (Barbosa, Cowell, & Dowd, 2021). Studies demonstrate that compared with men, women are more likely to drink in response to stress (Becker & Koob, 2016; Peltier et al., 2019), which is a concerning trend given that women are more vulnerable to experiencing pandemic-related psychological distress (Brooks et al., 2020; Taylor, 2019).

Evidence showed that the pandemic exacerbated the gender imbalance in the division of labor such that women disproportionately absorbed the caregiving obligations for homeschooling their children as well as handled more domestic responsibilities. As a result, women were less able to contribute to the paid workforce (Stefanova, Farrell, & Latu, 2021). In May 2020, 11% of women faced underemployment compared with 7% of men (Golden & Kim, 2020). Additionally, women have endured more severe stress owing to health and childcare issues (Connor et al., 2020). As intimate partner violence increased during the pandemic, women were more likely to be victimized, placing their mental and physical well-being at risk (Lyons & Brewer, 2021; McNeil, Hicks, Yalcinoz-Ucan, & Browne, 2022). Studies suggest that, during the initial months of the COVID-19 pandemic, a greater proportion of women compared with men exceeded their drinking limits through engaging in heavy drinking (Barbosa et al., 2021) and that women drank alcohol more frequently than men in general (Pollard, Tucker, & Green, 2020). Thus, a gender-specific approach to better understand the role of COVID-19 distress on young women's heavy alcohol use is warranted.

In this study, we hypothesized that three specific types of psychological distress during the COVID-19 pandemic would predict women's heavy drinking. First, COVID-19-related grief may be linked to heavy drinking. During the pandemic, the loss of life, opportunities, and vital resources coupled with general uncertainties, all contributed to a collective sense of grief (Bertuccio & Runion, 2020). Drinking may be one coping mechanism used by individuals against unexpected losses and grief. Second, COVID-19-related worry (e.g., fear of contracting COVID, being unable to maintain contact with loved ones) may also be associated with increased drinking. Rogers, Shepherd, Garey, and Zvolensky (2020) found that those who initiated using substances during the pandemic had greater COVID-19-related worry than those who never used substances. Finally, COVID-19-related financial stress may be a critical factor when considering that financial stress was also associated with alcohol-related problems during the COVID-19 pandemic among U.S. adults (Rodriguez, Litt, & Stewart, 2021). Another study found that financial-related distress was the sole factor associated with an increase in risky drinking patterns over time among U.S. adults (Lechner et al., 2021).

In addition, women's poor mental health status—including the risk factors of depression and anxiety—may be associated with heavy drinking. Several studies found that preexisting mental health diagnoses and mental health problems (e.g., substance use disorders, anxiety, depression) were associated with elevated alcohol consumption during the pandemic (Capasso et al., 2021; Hansel, Saltzman, Melton, Clark, & Bordnick, 2022; Holmes et al., 2020; Pfefferbaum & North, 2020; Yazdi, Fuchs-Leitner, Rosenleitner, & Gerstgrasser, 2020). During the COVID-19 pandemic, high anxiety and depression were associated with increased drinking (Devoto, Himelein-Wachowiak, Liu, & Curtis, 2022). Overall, those with a preexisting mental health diagnosis may be at greater risk for increased alcohol use. Last, loneliness may be another risk factor among young women during the pandemic. One multicountry study found that loneliness is strongly associated with high alcohol use among women during the pandemic, suggesting that alcohol is used to cope with loneliness (O'Sullivan et al., 2021).

Studies have generally demonstrated an association between these risk factors and heavy drinking. However, no published research is based on longitudinal data to understand the extent to which COVID-19-related distress (grief, worry, and financial stress) is associated with heavy alcohol intake after controlling for other risk factors among young women during the pandemic. This study uses the longitudinal data from the COVID-19 Adult Resilience Experiences Study (CARES 2020 Project) that tracked the cohort panel of young adults in the U.S. in 2020 and 2021 to document changes in their health and well-being over time. The objectives of the present study are to 1) estimate the prevalence of heavy alcohol use among U.S. young women and 2) determine the relationship between COVID-19-related distresses, mental health status, and loneliness during T₁ and heavy alcohol consumption during T₂ among U.S. young women. Understanding these potential factors will help to design strategies and interventions to decrease excessive alcohol use for young women in the United States.

Methods

Recruitment and Procedure

The present study draws from CARES (see previous papers for details; Liu, Stevens, Conrad, & Hahm, 2020a; Liu, Zhang, Wong,

Hyun, & Hahm, 2020b; Hahm et al., 2021a; Hahm et al., 2021b), which follows young adults across two time points after their initial enrollment in 2020 when they were 18–30 years of age. Participants were recruited online via listservs and Facebook groups for the school organizations or various clubs, alumni groups, classes, and churches. Although the initial recruitment took place in the New England area, additional listservs in the Midwest, South, and West were used to make the sample more geographically diverse. During the study, participants were asked to complete a 30-minute online survey about their COVID-19-related experiences, risk and resilience, and physical and mental health outcomes. All participants completed an electronic consent form. For the T₁ survey, one out of every 10 participants received a \$25 gift card. The first invitation to the second survey was sent approximately 5.5 months after the T₁ completion date. Participants were given \$10 gift cards when they completed the T₂ survey. Three different attention checks were embedded in the survey to ensure data quality. Study approval was obtained from the Institutional Review Board of Boston University.

A total of 1,221 young adults took part in the T₁ survey (T₁ defined as April 13, 2020, to August 31, 2020). Participants who agreed to be contacted again during CARES T₁ were invited to participate in T₂ (T₂ defined as September 21, 2020, to March 15, 2021). Among those who were contacted, 65.9% completed surveys for T₂, for a total of 805 participants. For this study, those who identified themselves as women (including transgender women) were selected for the analysis ($n = 684$). Between T₁ and T₂, no significant differences in demographic characteristics were observed except for participants' student status, because many of them had graduated (students at T₁, 70.2% vs. students at T₂, 56.9%; $p < .001$).

Measures

Covariates

Demographics. Covariates were age, race/ethnicity, current student status (college/graduate students), and income from the previous year, including all sources such as social security/disability, stocks, alimony, and child support.

Duration of the pandemic. The pandemic duration was calculated by counting the number of days from the date when COVID-19 was declared as a global pandemic (March 11, 2020) by the World Health Organization. The pandemic duration was included as one of the covariates to consider possible correlations between the date when the survey was conducted and the outcome variables.

Psychological distress measures

COVID-19-related grief. A newly developed six-item measure adapted from the Inventory of Complicated Grief (Prigerson et al., 1995) assessed COVID-19-related feelings of grief and loss. After viewing six grief- or loss-related statements, participants rated their sense of agreement on a scale of 1 (strongly disagree) to 5 (strongly agree). Statements included missing out on significant life events, losing vital resources (e.g., housing, mentorship, food access), feeling stunned over COVID-19, and feeling bitter over loss in daily routines and activities (Liu et al., 2020a,b). Sum scores of measures ranged from 6 to 30. Cronbach's α for measure items was 0.75, indicating good reliability.

COVID-19-related worry. We chose four items from a new measure developed by Liu et al. (2020a,b). Participants were asked to

indicate their level of worry for each item on a scale of 1–5, with 1 being not worried at all and 5 being very worried. Items include, "Having enough groceries during city lockdowns/social distancing protocols," "obtaining a COVID-19 test if I become sick," "getting treated for COVID-19 if I contract it," and "keeping in touch with loved ones during social distancing protocols" (Liu et al., 2020a,b). Two items related to financial worry (worry about maintaining employment during the subsequent economic downturn; worry about having enough money to pay for rent and buy necessities) were omitted because these items were similar to the items on COVID-19-related financial stress. Sum scores were calculated with a possible range of 6–30 and recoded into a dichotomous variable with a cutoff score of 24 or greater as highly worried. Cronbach's α for measure items was 0.72, indicating good reliability.

COVID-19-related financial stress. Two questions were used to measure participants feeling stressed about personal finances and worrying about being able to pay monthly expenses on a scale ranging from 1 (strongly disagree) to 4 (strongly agree). Two items were summed to create the total score. The observed values ranged from 2 to 8. The correlation between these two items was 0.67.

Mental health status measures

Preexisting diagnosis. Respondents were asked whether they had ever been diagnosed with generalized anxiety disorder, depression, or post-traumatic stress disorder. For each disorder, respondents were asked to indicate yes, diagnosed but not treated, yes, diagnosed and treated, suspected, but not diagnosed, or no. We developed a dichotomous variable for each disorder. If respondents indicated either yes, diagnosed and treated or yes, diagnosed and untreated, they were coded as 1, indicating that they have a preexisting diagnosis for each of the disorders. If they responded as suspected, but not diagnosed or no, they were coded as 0, indicating not having a preexisting diagnosis for each disorder.

Depressive symptoms. The prevalence of depression and its severity in the past two weeks was measured using the 8-item Patient Health Questionnaire depression scale (Kroenke et al., 2009), and participants rated each item on a scale of 0 (not at all) to 3 (nearly every day). Items include, "How often in the past 2 weeks were you bothered by little interest or pleasure in doing things," "feeling down, depressed, or hopeless," and "poor appetite or overeating." The scores for each item were summed and ranged from 0 to 24 points. The scores were recoded dichotomously, with a cutoff of 10 or higher indicating major depression. The internal reliability of the eight-item Patient Health Questionnaire depression scale was excellent, with a Cronbach's α of 0.87.

Anxiety symptoms. Using the widely used Generalized Anxiety Disorder Scale (Spitzer, Kroenke, Williams, & Löwe, 2006), participants' frequency of anxiety symptoms in the past 2 weeks was assessed. The frequency of each symptom was rated on a scale of 0 (not at all) to 3 (nearly every day). Sum scores ranged from 0 to 21 and were recoded dichotomously to determine elevated anxiety based on a cutoff score of 10 or higher. This was based on previous cutoff conventions for this scale (Plummer et al., 2016). The internal consistency of the Generalized Anxiety Disorder Scale was excellent (Cronbach $\alpha = 0.90$).

Loneliness. Participants' sense of loneliness was assessed with the adapted three-item UCLA Loneliness Scale Short Form

(Hughes, Waite, Hawkey, & Cacioppo, 2004). Questions included, “How often do you feel that you lack companionship,” “feel left out,” and “feel isolated from others?” Participants responded on a scale of 1 (hardly ever) to 3 (often). The total score for loneliness ranged from 3 to 9 and recoded dichotomously with a cutoff score of 6 or greater, indicating high loneliness (Cronbach $\alpha = 0.79$).

Heavy drinking. Participants were asked, “How many drinks containing alcohol do you have on a typical day when you are drinking?” Participants selected a value from the following options: 1, or 2, 3; or 4 or 5; or 6 or 7; or 8, 9, or 10; or 10 or more drinks per day. We dichotomized scores ranging from 1 to 2 as no heavy drinking and scores ranging from 3 or 4 and beyond as heavy drinking.

Data Analytic Plan

Those who identified as women (including transgender women) were included in the analysis. Variables were checked for normal distribution and acceptable levels of collinearity (Variance Inflation Factor of <5). We performed t tests and Pearson χ^2 tests to compare the prevalence of demographics, mental health status, and COVID-19–related distress based on the participants' heavy alcohol use. Next, we fit a multiple logistic regression model to estimate the relative contribution of COVID-19–related distress to heavy alcohol use, controlling for covariates. The unadjusted model only included COVID-19–related distress (financial stress, COVID-19–related worry, and COVID-19–related grief at T₁). In addition to these predictors, the adjusted model included demographics, pandemic duration, preexisting diagnosis, depressive symptoms, anxiety symptoms, and loneliness at T₁. Analyses were conducted using SPSS 28.0.

Results

Table 1 displays the frequency of the sample characteristics. Among the total analytic sample of women, approximately 62% were White, followed by 22% Asians, 5% Black, and 5% Latinx. Between T₁ and T₂, no significant differences in demographic characteristics were observed, except for participants' student status. The mean of COVID-19–related worry scores was 9.9 (range, 6–30), and the mean of COVID-19–related grief scores was 19.2 (range, 6–30). Approximately 42% of the total sample had at least one preexisting mental health diagnosis. Among all participants, 27.5% of women at T₁ and 22.5% of women at T₂ reported heavy drinking.

Paired sample t tests were conducted to determine whether there is a significant difference in key variables between those in the no heavy drinking and heavy drinking groups (**Table 2**). Young women who reported heavy drinking, on average, reported statistically significantly higher levels of depressive and anxiety symptoms, loneliness, and COVID-19–related grief than those who did not.

Table 3 shows results from the unadjusted and adjusted regression models. We present the detailed findings from the adjusted model. After controlling for covariates (demographic variables, pandemic duration, preexisting diagnoses, depressive and anxiety symptoms, and loneliness), COVID-19–related grief was significantly associated with greater odds of heavy drinking (odds ratio, 1.06; 95% confidence interval, 1.01–1.13; $p < .05$), whereas COVID-19–related worry was associated with lower

Table 1
Characteristics of Respondents ($N = 684$) With Data from Times 1 and 2 of CARES 2020

Variables	Means \pm SD or % (n)
Age (mean years) at T ₁	24.2 \pm 3.3
Race	
White	62.3 (426)
Asian	21.5 (147)
Latinx	5.3 (36)
Black	4.8 (33)
Mixed	6.1 (42)
Student status at T ₁	
No	29.8 (204)
Yes	70.2 (480)
Income at T ₁	
<\$25,000	62.7 (428)
\$25,000–\$49,999	23.3 (159)
\$50,000–\$250,000+	14.1 (96)
Pandemic duration*	63.0 (27.0–171.0)
Mental health status at T ₁	
Preexisting diagnosis	
No	57.9 (396)
Yes	42.1 (288)
Depression (PHQ-8)	8.7 \pm 5.5
Anxiety (GAD-7)	9.4 \pm 5.5
Loneliness	6.1 \pm 1.9
COVID-19–related distress at T ₁	
COVID-19–related financial stress	5.0 \pm 1.7
COVID-19–related worry	9.9 \pm 3.6
COVID-19–related grief	19.2 \pm 4.6
Heavy drinking	
Heavy drinking at T ₁	
No	72.5 (471)
Yes	27.5 (179)
Heavy drinking at T ₂	
No	77.5 (507)
Yes	22.5 (147)

Abbreviations: COVID-19, coronavirus disease 2019; GAD, Generalized Anxiety Disorder Scale; PHQ-8, 8-item Patient Health Questionnaire depression scale; SD, standard deviation; T₁, April 13, 2020, to August 31, 2020; T₂, September 21, 2020, to March 15.

Note: May not add up to the total owing to missing.

* Pandemic duration indicates days from March 11, 2020, to T₁ survey completion.

odds of heavy drinking (odds ratio, 0.90; 95% confidence interval, 0.85–0.97; $p < .01$). Financial stress was not significantly associated with heavy drinking.

Discussion

This study represents one of the few targeted epidemiological studies of heavy alcohol use among U.S. young women during the COVID-19 pandemic. Our results show that 27.5% of women (T₁ defined as April to August 2020) and 22.5% (T₂ defined as September 2020 to March 2021) reported three drinks or more as their typical intake, the definition of heavy drinking. This difference in rates between the two time points may reflect a higher prevalence of heavy drinking among women during T₁, the earlier period of the pandemic that included the enactment of stay-at-home orders, compared with T₂. Although no known longitudinal study observed and measured heavy alcohol use consistent with our two time periods, one study showed a similar prevalence of heavy drinking among U.S. adults, at 27.7% during April 2020 (Barbosa et al., 2021).

Guided by prior findings showing a link between COVID-19–related psychological distress and women's levels of drinking

Table 2
Characteristics by Young Women's Alcohol Consumption (N = 684)

Variables	No Heavy Drinking at T ₂ (<2 drinks)	Yes Heavy Drinking at T ₂ (≥3 drinks)	χ ² or t
Age at T ₁	24.7 ± 3.19	22.7 ± 3.14	t = 6.69 [‡]
Race			χ ² = 3.06
White	61.3 (311)	68.7 (101)	
Asian	22.3 (113)	16.3 (24)	
Other (Latinx, Black, mixed)	16.4 (83)	15.0 (22)	
Student status at T ₁			χ ² = 12.74 [‡]
No	33.7 (171)	18.4 (27)	
Yes	66.3 (336)	81.6 (120)	
Income at T ₁			χ ² = 13.65 [‡]
<\$25,000	58.8 (298)	75.5 (111)	
\$25,000–\$49,999	25.6 (130)	15.6 (23)	
\$50,000–\$250,000+	15.6 (79)	8.8 (13)	
Mental health status at T ₁			
Preexisting diagnosis			χ ² = 2.13
No	55.8 (283)	62.6 (92)	
Yes	44.2 (224)	37.4 (55)	
Depression (PHQ-8)	8.43 ± 5.49	9.60 ± 5.47	t = -2.27 [*]
Anxiety (GAD-7)	9.16 ± 5.37	10.35 ± 5.77	t = -2.32 [*]
Loneliness	5.94 ± 1.87	6.49 ± 1.91	t = -3.13 [†]
COVID-19–related distress at T ₁			
COVID-19–related financial stress	4.93 ± 1.73	5.16 ± 1.72	t = -1.42
COVID-19–related worry	9.99 ± 3.56	9.73 ± 3.74	t = 0.78
COVID-19–related grief	18.79 ± 4.64	20.56 ± 4.18	t = -4.17 [‡]

Abbreviations: COVID-19, coronavirus disease 2019; GAD, Generalized Anxiety Disorder Scale; PHQ-8, 8-item Patient Health Questionnaire depression scale; T₁, April 13, 2020, to August 31, 2020; T₂, September 21, 2020, to March 15.

Note: May not add up to the total owing to missing. Values are% (n) or mean (standard deviation).

* p < .05.

† p < .01.

‡ p < .001.

Table 3
Odds Ratios and Confidence Intervals for Young Women's Heavy Drinking at T₂ (N = 684)

Variables	Unadjusted OR	Adjusted OR [§]
Age at T ₁	–	0.82 (0.75–0.89) [‡]
Race		
White	–	1.0
Asian	–	0.60 (0.35–1.04)
Other (Latinx, Black, mixed)	–	0.68 (0.39–1.20)
Income at T ₁		
<\$25,000	–	1.0
\$25,000–\$49,999	–	1.18 (0.63–2.22)
\$50,000–\$250,000+	–	1.69 (0.71–4.00)
Student status at T ₁		
No	–	1.0
Yes	–	1.43 (0.78–2.61)
Pandemic duration	–	1.00 (1.00–1.01)
Mental health status at T ₁		
Preexisting diagnosis		
No	–	1.0
Yes	–	0.74 (0.48–1.14)
Depression (PHQ-8)	–	1.01 (0.96–1.07)
Anxiety (GAD-7)	–	1.02 (0.97–1.08)
Loneliness	–	1.02 (0.90–1.16)
COVID-19–related distress at T ₁		
COVID-19–related financial stress	1.08 (0.96–1.22)	1.13 (1.00–1.28)
COVID-19–related worry	0.91 (0.85–0.97) [†]	0.90 (0.85–0.97) [†]
COVID-19–related grief	1.12 (1.07–1.18) [‡]	1.06 (1.01–1.13) [*]

Abbreviations: COVID-19, coronavirus disease 2019; GAD, Generalized Anxiety Disorder Scale; PHQ-8, 8-item Patient Health Questionnaire depression scale; T₁, April 13, 2020, to August 31, 2020; T₂, September 21, 2020, to March 15.

* p < .05.

† p < .01.

‡ p < .001.

§ Adjusted for age, race, income, student status, pandemic duration, preexisting mental health diagnosis, depressive and anxiety symptoms, and loneliness at T₁.

|| Pandemic duration indicates days from March 11, 2020, to T₁ survey completion.

(Rodriguez, Litt, & Stewart, 2020), our main research objectives were to test whether distinctive types of COVID-19–related distress—grief, worry, and financial stress—were associated with women's heavy drinking behavior. In our study, COVID-19–related grief was significantly associated with heavy drinking. This relationship remained significant after controlling for pre-existing mental health diagnoses, depressive and anxiety symptoms, loneliness, and demographic variables. Our measure of COVID-19–related grief captured degrees of reactions reflecting a sense of loss and bitter sentiments related to the pandemic, such as missing out on significant life events or experiencing the loss of vital resources. With sudden disruptions to various celebrations, networking opportunities, relationships, or academic engagements likely leading to such grief, alcohol may have been used as a temporary remedy to compensate for lost pleasures and to regulate feelings of vulnerability related to the unknownness stemming from the pandemic. Our finding supports the self-medication hypothesis in that an individual may engage in substance use as a coping device to help relieve or numb negative emotions, including one's own pain and suffering (Khantzian, 1997).

However, contrary to our hypothesis, we found that participants with higher levels of COVID-19–related worry were more likely to report nonheavy drinking than heavy drinking. Our measure of COVID-19–related worry is composed of concerns related to inaccessibility of COVID-19 tests or treatments, not having enough groceries, or an inability to keep in touch with loved ones. For many individuals, these are concerns that threaten their safety and well-being, but it does not seem that participants tried to cope with them by drinking heavily.

Similarly, Lechner et al. (2021) also found a negative relationship, where a greater level of COVID-19–related fear (i.e., fear of catching COVID-19) was associated with lower drinking rates among college students. A study in the UK also found that being worried about becoming seriously ill from COVID-19, being diagnosed with COVID-19, and having suspected COVID-19 were all associated with lower levels of alcohol use (Garnett et al., 2021). It is plausible that individuals with greater COVID-19–related worry or fear were more vigilant about their health and overall well-being, thereby refraining from heavy drinking. Additional research that examines the mechanisms of COVID-19–related worry in relation to heavy drinking is needed to identify intervention targets for reducing heavy drinking among young women.

Contrary to our hypothesis, there was no evidence that COVID-19–related financial stress measured in T_1 is associated with heavy drinking in T_2 . One plausible explanation is that the U.S. government responded to the economic crisis with several stimulus checks and policies for individuals during our T_1 period (April to August 2020). As a result, financial stress perceived by these young adults during this time might not have had a significant impact on heavy drinking behaviors. An alternative interpretation is that those who are financially stressed or stretched were living at their parents' home and, therefore, not necessarily in an environment where they would engage in binge drinking. Future research should explore the role of financial stress measured in T_2 in heavy drinking among women, especially as it captures stress related to the prolonged nature of the pandemic.

It is worthwhile to mention that we conducted an exploratory analysis to test whether COVID-19–related psychological distress was associated with heavy drinking in young men ($n = 121$). We

found that none of the psychological distress variables were significantly associated with men's heavy alcohol use during the pandemic. This finding may add to the growing evidence that women are more vulnerable to experiencing pandemic-related psychological distress. However, given the small sample size of young men ($n = 121$), future studies should further explore the factors associated with young men's heavy drinking during the COVID-19 pandemic.

Study limitations are the following. First, our measures were based on self-reports that may have been subject to biases, including social desirability bias and recall bias that may underestimate alcohol consumption. Second, considering that White women comprised the majority of our study sample, caution must be taken in the generalization of our study findings to the broader U.S. young adult population. Third, owing to the limitations of CARES data on alcohol use measures, we could not explore the range of alcohol-related problem behaviors. For future research, a comprehensive measure of alcohol use that includes the frequency and quantity of regular consumption, frequency and quantity of the largest amount consumed, and frequency of binge drinking should be used. Fourth, our COVID-19–related grief measure generally captures explicit and self-evident (or definite) types of grief. Future COVID-19–related grief measures should examine specific features of grief, including anticipatory or ambiguous loss (i.e., uncertainty about the current situation) and complicated grief (i.e., a persistent sense of loss) (Bertuccio & Runion, 2020). This process could help to distinguish the contributing role of each unique grief feature versus overall COVID-19–related grief in health risk behaviors during a pandemic. Finally, in terms of drinking behavior, CARES data are limited by a lack of measures exploring motivation and attitudes on drinking or genetic predispositions. Going forward, researchers should include a more comprehensive assessment to develop a greater nuanced understanding of coping-motivated drinking behaviors.

Implications for Practice and/or Policy

Despite these limitations, this study has some methodological strengths. Using two time points allowed us to predict heavy alcohol consumption because all predictors preceded the heavy drinking variable. With approximately one in four women reporting heavy drinking during the pandemic, this finding points to the need for alcohol reduction support for women. In addition, this study sheds light on the differential effects of various COVID-19–related distresses on alcohol-related behaviors. The finding that young women reporting higher levels of COVID-19–related grief were more likely to report heavy drinking suggests that mental health clinicians should address this form of COVID-19–related distress as the pandemic continues to claim lives and cause prolonged economic and mental health consequences. Women should be informed about the negative effect of pandemic-related grief on alcohol intake. In particular, clinicians and researchers should pay greater attention to screening COVID-19–related distress in women and design subsequent clinical interventions focusing on COVID-19–related grief to ensure an appropriate response to alcohol problems. Ongoing observation of drinking behavior trends across time is warranted to determine the long-term mental health and alcohol-related impacts of the COVID-19 pandemic.

References

- Barbosa, C., Cowell, A. J., & Dowd, W. N. (2021). Alcohol consumption in response to the COVID-19 pandemic in the United States. *Journal of Addiction Medicine*, 15, 341–344.
- Becker, J. B., & Koob, G. F. (2016). Sex differences in animal models: Focus on addiction. *Pharmacological Reviews*, 68, 242–263.
- Bertuccio, R. F., & Runion, M. C. (2020). Considering grief in mental health outcomes of COVID-19. *Psychological Trauma: Theory, Research, Practice, and Policy*, 12, S87–S89.
- Boykoff, N., Schneekloth, T. D., Hall-Flavin, D., Loukianova, L., Karpyak, V. M., Stevens, S. R., ... Frye, M. A. (2010). Gender differences in the relationship between depressive symptoms and cravings in alcoholism. *American Journal on Addictions*, 19, 352–356.
- Brooks, S. K., Webster, R. K., Smith, L. E., Woodland, L., Wessely, S., Greenberg, N., & Rubin, G. J. (2020). The psychological impact of quarantine and how to reduce it: Rapid review of the evidence. *Lancet*, 395, 912–920.
- Bryan, A. E., Norris, J., Abdallah, D. A., Stappenbeck, C. A., Morrison, D. M., Davis, K. C., ... Zawacki, T. (2016). Longitudinal change in women's sexual victimization experiences as a function of alcohol consumption and sexual victimization history: A latent transition analysis. *Psychology of Violence*, 6, 271.
- Capasso, A., Jones, A. M., Ali, S. H., Foreman, J., Tozan, Y., & DiClemente, R. J. (2021). Increased alcohol use during the COVID-19 pandemic: The effect of mental health and age in a cross-sectional sample of social media users in the U.S. *Preventive Medicine*, 145, 106422.
- Centers for Disease Control and Prevention (CDC). (2020). Excessive alcohol use and risks to women's health. Available: www.cdc.gov/alcohol/fact-sheets/womens-health.htm. Accessed: October 6, 2022.
- Centers for Disease Control and Prevention (CDC). (2021). Excessive alcohol use. Available: www.cdc.gov/chronicdisease/resources/publications/factsheets/alcohol.htm. Accessed: October 6, 2022.
- Cohen, S., & Wills, T. A. (1985). Stress, social support, and the buffering hypothesis. *Psychological Bulletin*, 98, 310–357.
- Connor, J., Madhavan, S., Mokashi, M., Amanuel, H., Johnson, N. R., Pace, L. E., & Bartz, D. (2020). Health risks and outcomes that disproportionately affect women during the Covid-19 pandemic: A review. *Social Science & Medicine*, 266, 113364.
- Devoto, A., Himelein-Wachowiak, M., Liu, T., & Curtis, B. (2022). Women's substance use and mental health during the COVID-19 pandemic. *Women's Health Issues*, 32, 235–240.
- Erol, A., & Karpyak, V. M. (2015). Sex and gender-related differences in alcohol use and its consequences: Contemporary knowledge and future research considerations. *Drug and Alcohol Dependence*, 156, 1–13.
- Garnett, C., Jackson, S., Oldham, M., Brown, J., Steptoe, A., & Fancourt, D. (2021). Factors associated with drinking behavior during COVID-19 social distancing and lockdown among adults in the UK. *Drug and Alcohol Dependence*, 219, 108461.
- Glasheen, C., Pemberton, M. R., Lipari, R., Copello, E. A., & Mattson, M. E. (2015). Binge drinking and the risk of suicidal thoughts, plans, and attempts. *Addictive Behaviors*, 43, 42–49.
- Golden, L., & Kim, J. (2020). Underemployment just Isn't working for US part-time workers. The Center for Law and Social Policy. Available: www.clasp.org/publications/report/brief/underemployment-just-isnt-working-us-part-time-workers. Accessed: October 6, 2022.
- Hahm, H. C., Hall, C. D. X., Garcia, K. T., Cavallino, A., Ha, Y., Cozier, Y. C., & Liu, C. H. (2021a). Experiences of COVID-19-related anti-Asian discrimination and affective reactions in a multiple race sample of U.S. young adults. *BMC Public Health*, 21, 1563.
- Hahm, H. C., Ha, Y., Scott, J. C., Wongchai, V., Chen, J. A., & Liu, C. H. (2021b). Perceived COVID-19-related anti-Asian discrimination predicts post-traumatic stress disorder symptoms among Asian American young adults. *Psychiatry Research*, 303, 114084.
- Hansel, T. C., Saltzman, L. Y., Melton, P. A., Clark, T. L., & Bordnick, P. S. (2022). COVID-19 behavioral health and quality of life. *Scientific Reports*, 12, 961.
- Holmes, E. A., O'Connor, R. C., Perry, V. H., Tracey, I., Wessely, S., Arseneault, L., ... Bullmore, E. (2020). Multidisciplinary research priorities for the COVID-19 pandemic: A call for action for mental health science. *Lancet Psychiatry*, 7, 547–560.
- Hughes, M. E., Waite, L. J., Hawkey, L. C., & Cacioppo, J. T. (2004). A short scale for measuring loneliness in large surveys: Results from two population-based studies. *Research on Aging*, 26, 655–672.
- King, A. C., Bernardy, N. C., & Hauner, K. (2003). Stressful events, personality, and mood disturbance: Gender differences in alcoholics and problem drinkers. *Addictive Behaviors*, 28, 171–187.
- Khan, S., Okuda, M., Hasin, D. S., Secades-Villa, R., Keyes, K., Lin, K.-H., ... Blanco, C. (2013). Gender differences in lifetime alcohol dependence: Results from the National Epidemiologic Survey on Alcohol and Related Conditions. *Alcoholism, Clinical and Experimental Research*, 37, 1696–1705.
- Khantzian, E. J. (1997). The self-medication hypothesis of substance use disorders: A reconsideration and recent applications. *Harvard Review of Psychiatry*, 4, 231–244.
- Kroenke, K., Strine, T. W., Spitzer, R. L., Williams, J. B., Berry, J. T., & Mokdad, A. H. (2009). The PHQ-8 as a measure of current depression in the general population. *Journal of Affective Disorders*, 114, 163–173.
- Lechner, W. V., Sidhu, N. K., Jin, J. T., Kittaneh, A. A., Laurene, K. R., & Kenne, D. R. (2021). Increases in risky drinking during the COVID-19 pandemic assessed via longitudinal cohort design: Associations with racial tensions, financial distress, psychological distress and virus-related fears. *Alcohol and Alcoholism (Oxford, Oxfordshire)*, 56, 702–707.
- Liu, C. H., Stevens, C., Conrad, R. C., & Hahm, H. C. (2020a). Evidence for elevated psychiatric distress, poor sleep, and quality of life concerns during the COVID-19 pandemic among U.S. young adults with suspected and reported psychiatric diagnoses. *Psychiatry Research*, 292, 113345.
- Liu, C. H., Zhang, E., Wong, G., Hyun, S., & Hahm, H. C. (2020b). Factors associated with depression, anxiety, and PTSD symptomatology during the COVID-19 pandemic: Clinical implications for U.S. young adult mental health. *Psychiatry Research*, 290, 113172.
- Lyons, M., & Brewer, G. (2021). Experiences of intimate partner violence during lockdown and the COVID-19 pandemic. *Journal of Family Violence*, 37, 969–977.
- McNeil, A., Hicks, L., Yalcinoz-Ucan, B., & Browne, D. T. (2022). Prevalence & correlates of intimate partner violence during COVID-19: A rapid review. *Journal of Family Violence*, 1–21. <https://doi.org/10.1007/s10896-022-00386-6>.
- National Institute on Alcohol Abuse and Alcoholism. (2021). Alcohol facts and statistics. Available: www.niaaa.nih.gov/publications/brochures-and-factsheets/alcohol-facts-and-statistics. Accessed: October 6, 2022.
- O'Sullivan, R., Burns, A., Leavey, G., Leroi, I., Burholt, V., Lubben, J., ... Prohaska, T. R. (2021). Impact of the COVID-19 pandemic on loneliness and social isolation: A multi-country study. *International Journal of Environmental Research and Public Health*, 18, 9982.
- Peltier, M. K. R., Verplaetse, T. L., Mineur, Y. S., Petrakis, I. L., Cosgrove, K. P., Picciotto, M. R., & McKee, S. A. (2019). Sex differences in stress-related alcohol use. *Neurobiology of Stress*, 10, 100149.
- Peltier, M. R., Verplaetse, T. L., Roberts, W., Moore, K., Burke, C., Marotta, P. L., ... McKee, S. A. (2020). Changes in excessive alcohol use among older women across the menopausal transition: A longitudinal analysis of the Study of Women's Health Across the Nation. *Biology of Sex Differences*, 11, 37.
- Pfefferbaum, B., & North, C. S. (2020). Mental health and the Covid-19 pandemic. *New England Journal of Medicine*, 383, 510–512.
- Plummer, F., Manea, L., Trepel, D., & McMillan, D. (2016). Screening for anxiety disorders with the GAD-7 and GAD-2: a systematic review and diagnostic meta analysis. *General Hospital Psychiatry*, 39, 24–31.
- Pollard, M. S., Tucker, J. S., & Green, H. D., Jr. (2020). Changes in adult alcohol use and consequences during the COVID-19 pandemic in the US. *JAMA Network Open*, 3, e2022942.
- Prigerson, H. G., Maciejewski, P. K., Reynolds, C. F., Bierhals, A. J., Newsom, J. T., Fasiczka, A., ... Miller, M. (1995). Inventory of complicated grief: A scale to measure maladaptive symptoms of loss. *Psychiatry Research*, 59, 65–79.
- Rodriguez, L. M., Litt, D. M., & Stewart, S. H. (2020). Drinking to cope with the pandemic: The unique associations of COVID-19-related perceived threat and psychological distress to drinking behaviors in American men and women. *Addictive Behaviors*, 110, 106532.
- Rodriguez, L. M., Litt, D. M., & Stewart, S. H. (2021). COVID-19 psychological and financial stress and their links to drinking: A dyadic analysis in romantic couples. *Psychology of Addictive Behaviors*, 35, 377–390.
- Rogers, A. H., Shepherd, J. M., Garey, L., & Zvolensky, M. J. (2020). Psychological factors associated with substance use initiation during the COVID-19 pandemic. *Psychiatry Research*, 293, 113407.
- Sacks, J. J., Gonzales, K. R., Bouchery, E. E., Tomedi, L. E., & Brewer, R. D. (2015). 2010 National and state costs of excessive alcohol consumption. *American Journal of Preventive Medicine*, 49, e73–e79.
- Sharif Nia, H., She, L., Kaur, H., Boyle, C., Khoshnavay Fomani, F., Hoseinzadeh, E., ... Rahmatpour, P. (2022). A predictive study between anxiety and fear of COVID-19 with psychological behavior response: The mediation role of perceived stress. *Frontiers in Psychiatry*, 13, 851212.
- Stefanova, V., Farrell, L., & Latu, I. (2021). Gender and the pandemic: Associations between caregiving, working from home, personal and career outcomes for women and men. *Current Psychology*, 30, 1–17.
- Substance Abuse and Mental Health Services Administration. (2022). 2020 NSDUH detailed tables. U.S. Department of Health and Human Services. Available: www.samhsa.gov/data/report/2020-nsduh-detailed-tables. Accessed: October 6, 2022.
- Spitzer, R. L., Kroenke, K., Williams, J. B. W., & Löwe, B. (2006). A brief measure for assessing generalized anxiety disorder: The GAD-7. *Archives of Internal Medicine*, 166, 1092–1097.
- Taylor, S. (2019). *The psychology of pandemics: Preparing for the next global outbreak of infectious disease*. Cambridge: Cambridge Scholars Publishing.
- Yazdi, K., Fuchs-Leitner, I., Rosenleitner, J., & Gerstgrasser, N. W. (2020). Impact of the COVID-19 pandemic on patients with alcohol use disorder and associated risk factors for relapse. *Frontiers in Psychiatry*, 11, 620612.

Author Descriptions

Hyeouk Chris Hahm is a Professor and Associate Dean of Research at Boston University, School of Social Work. She is dedicated to reducing health disparities

among vulnerable populations with a particular emphasis on building empirical evidence of health risk behaviors.

Kelly Hsu is a research assistant at the Boston University School of Social Work and graduated from Tufts University this year with a BS in Biology and Child Study and Human Development. She will begin her Medical Degree at Tufts University School of Medicine in the fall of 2022. In the future, she hopes to be able to better serve communities, including Asian American women, by tackling their existing physical and mental health disparities.

Sunah Hyun is a senior researcher for the student programs team at Tisch College, Tufts University. She currently focuses on creating new evaluation strategies and effective assessment systems to better support Tisch College's mission and values. She is a developmental psychologist and community-based researcher whose chief

aim is to develop culturally responsive interventions for marginalized families and conduct research to foster healthy development and school success for children and youth.

Kaiden Kane graduated from Boston University with a Bachelor of Science in Human Physiology with an emphasis on LGBTQ populations. He is currently at Boston Children's Hospital, working to improve young people's access to gender-affirming care.

Cindy H. Liu is the Director of the Developmental Risk and Cultural Resilience Laboratory at Brigham and Women's Hospital and an assistant professor of pediatrics at Harvard Medical School. She has a joint appointment within the departments of pediatric newborn medicine and psychiatry. Liu's research focuses on culture and socioemotional development and race and ethnic disparities as they relate to maternal and child mental health.