

Ginsenoside Rb1 reduces H₂O₂-induced HUVEC dysfunction by stimulating the sirtuin-1/AMP-activated protein kinase pathway

ZHENDA ZHENG^{1*}, MIN WANG^{1*}, CAILIAN CHENG², DINGHUI LIU¹,
LIN WU¹, JIEMING ZHU¹ and XIAOXIAN QIAN^{1,3}

Departments of ¹Cardiology and ²Nephrology; ³Institute of Integrated Traditional Chinese and Western Medicine, The Third Affiliated Hospital of Sun Yat-sen University, Sun Yat-sen University, Guangzhou, Guangdong 510630, P.R. China

Received July 23, 2019; Accepted April 1, 2020

DOI: 10.3892/mmr.2020.11096

Abstract. Endothelial dysfunction and senescence are closely associated with cardiovascular diseases including atherosclerosis and hypertension. Ginsenoside Rb1 (Rb1), the major active constituent of ginseng, has been investigated intensively because of its anti-obesity and anti-inflammatory effects. In a previous study, hydrogen peroxide (H₂O₂) was applied to induce human umbilical vein endothelial cell (HUVEC) aging. It was demonstrated that Sirtuin-1 (SIRT1) was activated by Rb1 to protect HUVECs from H₂O₂-induced senescence. However, the mechanisms are not fully understood. The present study examined the role of AMP-activated protein kinase (AMPK), an energy sensor of cellular metabolism, in the signaling pathway of SIRT1 during H₂O₂-stimulated HUVEC aging. It was identified that Rb1 restored the H₂O₂-induced reduction of SIRT1 expression, which was consistent with our previous study, together with the activation of AMPK phosphorylation. Using compound C, an AMPK inhibitor, the role of AMPK in the protective effect of Rb1 against H₂O₂-induced HUVEC senescence was examined. It was identified that the induction of phosphorylated AMPK by Rb1 markedly increased endothelial nitric oxide synthase expression and nitric oxide production, and suppressed PAI-1 expression, which were abrogated in HUVECs pretreated with compound C. Further experiments demonstrated that nicotinamide, a SIRT1 inhibitor, downregulated the phosphorylation of AMPK and reduced the protective effects of Rb1 against H₂O₂-induced endothelial aging. Taken together, these results provide new

insights into the possible molecular mechanisms by which Rb1 protects against H₂O₂-induced HUVEC senescence via the SIRT1/AMPK pathway.

Introduction

Endothelial cell aging is a major risk factor for cardiovascular disease (CVD) development (1-3). Numerous studies have shown that hydrogen peroxide (H₂O₂)-induced vascular endothelial dysfunction is partially responsible for the development of aging (4-6). H₂O₂-induced endothelial dysfunction reflects a loss of the balance between pro- and anti-oxidant, pro- and anti-inflammatory, and pro- and anti-thrombotic signals, all of which contribute to increased release of proinflammatory cytokines, including plasminogen activator inhibitor-1 (PAI-1), and a decline in antioxidants including endothelial nitric oxide synthase (eNOS) and nitric oxide (NO) (7-11). Thus, the prevention of endothelial dysfunction is essential to treat endothelial aging and CVD.

Ginseng, the root of *Panax ginseng* CA Meyer, is one of the most popular herbs in traditional Asian medicine. A growing body of evidence suggests that ginsenoside Rb1 (Rb1), a major component of ginsenosides extracted from ginseng, has various biological activities including antioxidative stress relief, anti-obesity and anti-inflammation (12-14). One of our previous studies also suggested that Rb1 at the concentration of 10-40 μM inhibits free fatty acid-induced inflammation partially through the blockade of nuclear factor (NF)-κB phosphorylation in 3T3-L1 adipocytes (15). Additionally, another study by our group demonstrated that Rb1 at the concentration of 20 μM attenuates human umbilical vein endothelial cell (HUVEC) senescence by improving the redox status (16). However, the range of effective concentrations and further modulated mechanisms of Rb1 in the endothelium are not fully elucidated.

AMP-activated protein kinase (AMPK) is a heterotrimeric member of an evolutionarily conserved protein kinase family that is sensitive to changes in oxygen tension and ATP consumption (17). Accumulating evidence has revealed that AMPK participates in the regulation of lipid metabolism, inflammation and angiogenesis in various animal models and cell types (18-22). Ido *et al* (23) reported that AMPK protects endothelial cells from the adverse effects of sustained

Correspondence to: Dr Jieming Zhu or Dr Xiaoxian Qian, Department of Cardiology, The Third Affiliated Hospital of Sun Yat-sen University, Sun Yat-sen University, 600 Tianhe Road, Guangzhou, Guangdong 510630, P.R. China
E-mail: zhujieming2011@126.com
E-mail: yyqianxiaoxian@163.com

*Contributed equally

Key words: ginsenoside Rb1, human umbilical vein endothelial cells, Sirtuin-1, AMP-activated protein kinase

hyperglycemia. Nagata *et al* (22) demonstrated that endothelial AMPK signaling plays a critical role in blood vessel recruitment to tissues responding to ischemic stress. In addition, studies have shown that AMPK exerts its beneficial role through multiple signaling pathways, including activation of eNOS and production of NO (24,25). However, it is still unclear whether endothelial senescence, eNOS activation and NO synthesis in HUVECs in response to Rb1 are related to the activation of AMPK.

Complementing our previous studies, the present study was undertaken to investigate the protective effects of Rb1 against H₂O₂-induced HUVEC dysfunction mediated by AMPK and the underlying mechanisms.

Materials and methods

Cell culture and treatments. Primary HUVECs were isolated from different six neonatal umbilical cords as previously described (26). Briefly, HUVECs at passages 2-4 were maintained in M199 medium (Invitrogen, Thermo Fisher Scientific, Inc.) supplemented with 20% fetal bovine serum (Hyclone, GE Healthcare Life Sciences) and 60 µg/ml endothelial cell growth supplement (BD Biosciences) at 37°C in a 5% CO₂ incubator and then exposed to the desired treatment in triplicate. The isolation procedure for HUVECs was approved by the Research Committee at the Third Affiliated Hospital of Sun Yat-sen University (approval nos. 2010-2-48 and 2018-02-057-01). The donors were negative for human immunodeficiency virus and hepatitis B virus and provided written informed consent to donate the umbilical cords.

To induce senescence, isolated HUVECs were treated with 60 µM H₂O₂ (Sigma-Aldrich, Merck KGaA) for 1 h and then cultured for another 24 h at 37°C. Rb1 (16071307, Chengdu Pufei De Biotech Co., Ltd.) used in the present study was extracted from Panax ginseng by HPLC according to the manufacturer's instructions and the purity of Rb1 used in the present study was 98.85%. To evaluate the effect of Rb1 on senescence, the cells were pretreated with 10 or 20 µM Rb1 for 30 min prior to H₂O₂ treatment. To measure the effect of the SIRT1 inhibitor nicotinamide (NAM; Sigma-Aldrich, Merck KGaA) and AMPK inhibitor compound C (Sigma-Aldrich, Merck KGaA), the cells were incubated with 20 mM NAM (27) or 8 µM compound C for 30 min as reported previously (18,28,29) and then treated with or without Rb1 at concentrations of 10 or 20 µM for 30 min before H₂O₂ treatment. At the end of each experiment, the cultured supernatants and monolayered cells were harvested for analyses.

Measurement of NO production. NO production was evaluated by measuring the accumulation of nitrites. The Griess method (30) was used to detect NO using a NO assay kit (Beyotime Institute of Biotechnology), following the manufacturer's instructions. Briefly, after the cells were cultured and treated as described above, 50 µl culture supernatant was incubated with 50 µl Griess reagent I and 50 µl Griess reagent II in a 96-well microplate at room temperature for 30 min. The optical density was measured with a Victor microplate reader (PerkinElmer, Inc.) at 540 nm. Nitrite concentrations in the medium were calculated according to a standard curve.

Senescence-associated β-galactosidase activity assay. Senescence was detected using a senescence-associated β-galactosidase (SA-β-gal)-positive approach according to a published protocol (31). After HUVECs were washed twice with prechilled PBS, the cells were fixed with 2% formaldehyde plus 1% glutaraldehyde for 5-10 min at room temperature. The cells were then washed twice with prechilled PBS for 3 min and stained with a staining solution [40 mmol/l citric acid/sodium phosphate buffer, 5 mmol/l potassium ferrocyanide (K₄[Fe(CN)₆]·3H₂O), 5 mmol/l potassium ferricyanide (K₃[Fe(CN)₆]), 150 mmol/l sodium chloride, 2 mmol/l magnesium chloride and 1 mg/ml X-gal] overnight at 37°C without CO₂. Senescent cells were identified as blue-stained cells under a TS100 inverted microscope (Nikon Corporation) at x100 magnification. At least 400 cells were examined to determine the percentage of SA-β-gal-positive cells in each group.

NAD⁺/NADH assay. The cellular NADP⁺/NADPH ratio was determined using a NAD⁺/NADH Quantification Kit (Beyotime Institute of Biotechnology), according to the manufacturer's instructions. In brief, HUVECs (1x10⁶/well) were seeded in six-well plates and exposed to the experimental conditions. To measure the NADP⁺/NADPH ratio, the cells were harvested, lysed with 200 µl NAD⁺/NADH buffer and gently pipetted to promote cell lysis. Then, 50-100 µl of the samples were collected and incubated for 30 min at 60°C. Then, 20 µl of the reacted samples were added to a 96-well plate and analyzed at 450 nm as the reference wavelength in the Victor microplate reader. The NAD⁺/NADH ratio was calculated according to a standard curve and normalized to the cell number, following the manufacturer's protocol.

Measurement of intracellular ATP. Determination of intracellular ATP in the indicated groups of HUVECs was performed by a bioluminescence assay (ATP Assay kit; Beyotime Institute of Biotechnology), according to the manufacturer's instructions. Briefly, the cells were washed twice with prechilled PBS and then lysed in lysis buffer on ice. Then, the samples were harvested and centrifuged at 12,000 x g for 5 min at 4°C and the supernatants were collected for subsequent analysis. After the reaction solutions containing luciferase and luciferin were added and background luminescence was measured, the ATP standard solution and samples were added and luminescence was measured. Then, the background luminescence was subtracted and the standard curve was constructed. The ATP concentrations were calculated from the standard curve and normalized to total protein content.

Western blot analysis. HUVECs were grown in six-well plates. After the indicated treatments, the cells were washed twice with prechilled PBS and then lysed in radioimmunoprecipitation buffer with a protease inhibitor cocktail, phenylmethylsulfonyl fluoride and sodium orthovanadate (Santa Cruz Biotechnology, Inc.). The protein concentration was measured by the Bradford method. Proteins (30 µg) in 30 µl reducing sample buffer were boiled for 5 min at 100°C and then resolved by SDS-PAGE (8 or 12% gels) for 2 h at 100 V. The proteins were transferred onto a polyvinylidene difluoride membrane for 90 min at 100 V. After transfer, the membrane was incubated in 25 ml blocking

buffer [1X Tris-buffered saline (TBS) and 0.1% Tween-20 with 5% non-fat dry milk] for 1 h at room temperature. The primary antibodies were rabbit polyclonal anti-eNOS (1:500; Cell Signaling Technology, Inc.), Rabbit polyclonal anti-PAI-1 (1:500; Cell Signaling Technology, Inc.), Rabbit polyclonal anti-AMPK (1:2,000; Cell Signaling Technology, Inc.), rabbit polyclonal anti-phosphorylated AMPK (Thr172; 1:1,000; Cell Signaling Technology, Inc.), rabbit polyclonal anti-SIRT1 (1:1,000; Cell Signaling Technology, Inc.), and rabbit polyclonal anti-GAPDH (control, 1:5,000; Cell Signaling Technology, Inc.). The membrane was incubated with the primary antibody in 10 ml primary antibody dilution buffer with gentle agitation overnight at 4°C. After washing three times for 10 min each with 15 ml of 10X TBS/0.1% TBST, the membrane was incubated with a horseradish peroxidase-conjugated secondary antibody (1:3,000; cat. no. 7074; Cell Signaling Technology, Inc.) in 10 ml blocking buffer with gentle agitation for 1 h at room temperature, followed by 3 washes for 10 min each. Membranes were developed using the enhanced chemiluminescence detection method (EMD Millipore). The signals were quantified using Quantity One software (version 4.6.9; Bio-Rad Laboratories, Inc.).

Statistical analysis. Results were obtained from at least three independent experiments ($n \geq 3$) if not otherwise stated. Data are expressed as the mean \pm standard deviation ($n \geq 3$). Statistical significance was calculated by one-way or two-way analysis of variance with Bonferroni's post-hoc test using GraphPad Prism software (version 5.0; GraphPad Software, Inc.). $P < 0.05$ was considered to indicate a statistically significant difference.

Results

Effects of H_2O_2 and Rb1 treatment in HUVECs. A number of studies have reported that H_2O_2 is a major inducer of cell senescence and oxidative stress in HUVECs and other different cell lines (32-35), which plays a central role in cell dysfunction. H_2O_2 was used at a concentration of 60 μM in the present study based on our previous studies (16,36,37) in which 60 μM H_2O_2 was reported to induce cell senescence and oxidative stress sufficiently. Senescence-associated β -galactosidase (SA- β -gal) staining was used to determine the degree of cell senescence. Aging cells stained blue indicated the senescent phenotype of HUVECs. In line with our previous studies (16,36,37), it was demonstrated that 60 μM H_2O_2 increased the SA- β -gal⁺ cell number (blue staining). In addition, the H_2O_2 treated group had more cells with a clear outline, enlarged cell body and difficulty in adhering to the bottom of the culture plate, all characteristics of stressed senescent cells (38,39). By contrast, pretreatment with Rb1 decreased the SA- β -gal⁺ cell number and attenuated cell senescence induced by H_2O_2 in a dose-dependent manner (Fig. 1A and B). PAI-1 and eNOS expression were then measured after the same treatment. The results demonstrated that 60 μM H_2O_2 increased PAI-1 expression (Fig. 1C) and decreased eNOS expression (Fig. 1D) in HUVECs, which were restored by Rb1 in a dose-dependent manner.

Effects of H_2O_2 and Rb1 on SIRT1 expression, AMPK phosphorylation and the $NAD^+/NADH$ ratio. A number

of studies have shown that SIRT1 exhibits anti-inflammatory (40-42) and antioxidant effects (43-45) in the endothelium. The effect of Rb1 on SIRT1 and phosphorylation of the catalytic subunit of AMPK (Thr172) in the presence or absence of H_2O_2 was examined. The results demonstrated that 60 μM H_2O_2 inhibited SIRT1 expression (Fig. 2A and B) as well as phosphorylation of AMPK (Fig. 2A and C) and that treatment with Rb1 at 10 and 20 μM restored Sirt-1 expression (Fig. 2A and B) and AMPK phosphorylation (Fig. 2A and C) in a dose-dependent manner.

The intracellular $NAD^+/NADH$ ratio in the presence or absence of Rb1 was further examined and it was identified that treatment with 60 μM H_2O_2 reduced the $NAD^+/NADH$ ratio, which was restored with Rb1 pretreatment (Fig. 2D). The data confirmed that H_2O_2 -induced decrease in the $NAD^+/NADH$ ratio were associated with reduced SIRT1 expression and it was demonstrated for the first time, to the best of the authors' knowledge, that treatment with Rb1 at 10 and 20 μM prevented the reductions in SIRT1 and the $NAD^+/NADH$ ratio, and inhibited phosphorylated AMPK, in HUVECs exposed to H_2O_2 .

Involvement of AMPK in Rb1-mediated inhibition of the H_2O_2 -induced oxidative response. Next, it was determined whether AMPK was involved in the inhibitory effects of Rb1 on the H_2O_2 -induced oxidative response. HUVECs were treated with 10 or 20 μM Rb1 for 24 h in the presence or absence of compound C, a specific inhibitor of AMPK. As shown in Fig. 3A, compound C clearly downregulated the phosphorylation of AMPK at 6, 8 and 10 μM in a dose-dependent manner. According to the results, 8 μM compound C was chosen to inhibit the phosphorylation of AMPK in the following experiment. Pretreatment of HUVECs with compound C significantly abolished the inhibitory effects of Rb1 on H_2O_2 -induced PAI-1 expression (Fig. 3C). Rb1 also significantly restored eNOS expression (Fig. 3D) and NO production (Fig. 3E) and this beneficial effect was markedly reversed by pretreatment of HUVECs with compound C (Fig. 3). The results suggest that AMPK activation is essential for the inhibitory effect of Rb1 on H_2O_2 -induced senescence and the oxidative response in HUVECs.

Involvement of SIRT1 in Rb1-mediated inhibition of the H_2O_2 -induced oxidative response. To further understand the role of SIRT1 in the AMPK pathway, it was investigated whether inactivation of SIRT1 by nicotinamide affected H_2O_2 -induced cell injury. As Guo *et al* (27) reported that 20 mM NAM decreases SIRT1 gene expression significantly, HUVECs were treated with H_2O_2 in the presence or absence of 20 mM NAM, followed by observation of the effects on H_2O_2 -induced senescence and the expression of antioxidant genes. As shown in Fig. 4, NAM reduced the phosphorylation of AMPK (Fig. 4A and B), indicating that AMPK is the downstream protein modulated by SIRT1.

As AMPK is one of the most important proteins in modulating the cellular energy metabolism and ATP is a major downstream product of mitochondrial energy coordinators, the effects of NAM with or without Rb1 on the ATP level were next investigated. Rb1 protected HUVECs from H_2O_2 -induced PAI-1 expression (Fig. 4C) and rescued the downregulation of




Figure 1. Rb1 inhibits H₂O₂-induced senescence and represses PAI-1 as well as eNOS expression in HUVECs. The HUVECs were pretreated with 10-20 μM Rb1, treated with or without 60 μM H₂O₂ for 1 h and then cultured for 24 h. (A) Representative phase contrast images of SA-β-gal⁺ cells (blue staining; magnification, x100). (B) Ratio of SA-β-gal-positive cells calculated from 400 cells per group. The protein levels of PAI-1 (C) and eNOS (D) were detected by western blot analysis and semi-quantified by densitometry. Representative blots of three independent experiments are shown. **P<0.01 vs. Ctrl group; #P<0.05 vs. 60 μM H₂O₂ group. Rb1, ginsenoside Rb1; H₂O₂, hydrogen peroxide; PAI-1, plasminogen activator inhibitor-1; eNOS, endothelial nitric oxide synthase; HUVEC, human umbilical vein endothelial cell; SA-β-gal, senescence-associated β-galactosidase; Ctrl, control.

eNOS expression (Fig. 4D) and NO production (Fig. 4E). As shown in Fig. 4F, treatment with Rb1 increased the ATP level under H₂O₂-induced oxidative stress. However, the promotional effect on the ATP level by Rb1 was not observed in the NAM-treated group (Fig. 4F). The beneficial effects of Rb1 on H₂O₂-induced injuries were reversed by treatment with NAM (Fig. 4). These *in vitro* data confirmed that inhibition of SIRT1 by NAM blocks Rb1 to increase phosphorylation of AMPK, ATP production, eNOS expression and NO production, which collectively indicate that Rb1 plays a protective role against H₂O₂-induced injury of HUVEC via the SIRT1/AMPK pathway.

Discussion

Our previous studies (16,36,37) demonstrated that endothelial senescence and dysfunction in HUVEC were characterized by enhanced H₂O₂-induced β-galactosidase activity and impaired anti-oxidant capacity. This was associated with an increased protein expression of PAI-1, decreased NO production and eNOS expression in HUVEC. The present study

identified that Ginsenoside Rb1 protects against cell senescence and dysfunction through activation of Sirt1. However, the involved molecular mechanisms have yet to be elucidated. In line with our previous studies (16,36,37), the present study demonstrated that Rb1 significantly increased NO content, eNOS expression and Sirt1 expression in H₂O₂-induced HUVEC senescence. In addition, it was identified for the first time, to the best of the authors' knowledge, that these changes were associated with upregulated phosphorylation of AMPK. Compound C, an inhibitor of AMPK, was used to clarify the relation of AMPK and Sirt1. It was identified that treatment with compound C markedly attenuated the protective effects of Rb1 in HUVECs. Furthermore, the role of SIRT1/AMPK pathway was further confirmed by administration of nicotinamide, which enhanced H₂O₂-induced senescence and dysfunction and decreased the protective effect of Rb1 on HUVEC. Collectively, the results provide novel mechanisms for Rb1 protected H₂O₂-induced HUVEC senescence involving SIRT1/AMPK pathway.

A number of studies have reported that Rb1, a main constituent of the root of *P. ginseng*, has various pharmacological




Figure 2. Rb1 restores the H₂O₂-induced reduction of SIRT1, promotes phosphorylation of AMPK and increases the NAD⁺/NADH ratio. The HUVECs were pretreated with 10-20 μM Rb1, treated with or without 60 μM H₂O₂ for 1 h and then cultured for 24 h. (A) The levels of SIRT1, phosphorylated AMPK, AMPK and GAPDH were determined by western blot analysis. (B) Quantitative analysis of SIRT1 expression ratio and SIRT1 expression levels. (C) Quantitative analysis of phosphorylated AMPK ratio and phosphorylated AMPK expression levels. (D) NAD⁺/NADH ratio measured using a colorimetric assay. **P<0.01 vs. control group; #P<0.05 vs. 60 μM H₂O₂ group. Rb1, ginsenoside Rb1; H₂O₂, hydrogen peroxide; SIRT1, Sirtuin-1; AMPK, AMP-activated protein kinase; HUVEC, human umbilical vein endothelial cell; p, phosphorylated.

effects that include mitigating endothelial inflammation and obesity *in vitro* and *in vivo*. Zhou *et al* (46) reported that Rb1 protects HUVECs from TNF-α-induced oxidative stress and inflammation by inhibiting NF-κB, JNK and p38 signaling pathways. Yuan *et al* (47) reported that Rb1 protects pulmonary microvascular endothelial cells from lipopolysaccharide-induced expression of inflammatory cytokines. Our previous studies demonstrated that treatment with Rb1 at the concentration of 20 μM significantly reduces H₂O₂-induced cell senescence and inhibits PAI-1 expression (37) partly through stimulation of SIRT1 (36). The present study demonstrated that Rb1 at the concentration of 10-20 μM played a beneficial role on H₂O₂-induced endothelial dysfunction, which complements our previous research. However, the molecular mechanisms need further elucidation.

Previous studies have suggested a critical role of AMPK in stabilizing endothelial functions through regulation of eNOS signaling to suppress inflammation and oxidative stress (48-51). In addition, there are some studies on the effects of AMPK activators on the expression of eNOS, suggesting that AMPK acts as an eNOS activator (52-54). Thus far, there has been a lack of studies from the present authors concerning the role of AMPK in the regulation of H₂O₂-induced oxidative stress and inflammatory responses

by Rb1 in HUVECs. However, the present study explored the use of compound C to inhibit AMPK and determined whether Rb1 upregulated eNOS expression and NO production through AMPK. As a result, it was identified that AMPK activity was enhanced by Rb1. The present study not only confirmed that the expression and activity of eNOS suppressed by H₂O₂ stimulation was enhanced by Rb1 treatment, which is in accordance with previous findings, but also identified that the beneficial effects of Rb1 were abolished by compound C, indicating a novel role of AMPK in regulating Rb1-dependent eNOS activity and NO synthesis. However, the mechanism of Rb1 in AMPK-dependent eNOS activation and NO production in HUVECs requires further investigation.

SIRT1 is highly sensitive to cellular redox states and considered to have a cardioprotective effect (55,56) that maintains endothelial functions by counteracting the effects of reactive oxygen species as a NAD⁺-dependent class III histone deacetylase (57). SIRT1 has been reported to regulate the functions of several important transcription factors with anti-inflammatory effects (58-60). It antagonizes H₂O₂-induced premature senescence via negative modulation of p53 by deacetylation of Lys-373, Lys-382 and Lys-320 in the human endothelium (61). Ota *et al* (62,63) demonstrated that the overexpression of




Figure 3. Compound C suppresses the beneficial effects of Rb1 on H₂O₂-induced antioxidant and anti-inflammatory functions. The HUVECs were pretreated with the indicated concentrations of compound C for 30 min and then pretreated with 20 μ M Rb1 for 30 min, followed by treatment with or without 60 μ M H₂O₂ for 1 h and then culture for 24 h. (A) Phosphorylation of AMPK in HUVECs pretreated with compound C (0, 2, 4, 6, 8 and 10 μ M) for 30 min detected by western blot analysis and semi-quantified by densitometry. Total AMPK and GAPDH were also determined. (B) The protein levels of PAI-1, eNOS, phosphorylated AMPK, AMPK and GAPDH were determined by western blot. (C) Quantitative analysis of PAI-1 expression ratio and PAI-1 expression levels. (D) Quantitative analysis of eNOS expression ratio and eNOS expression levels. (E) NO production measured by the Griess reaction and expressed as the percentage of the control. Results represent three independent experiments. *P<0.05 and **P<0.01 vs. control group; #P<0.05 vs. 60 μ M H₂O₂ group; &P<0.05 vs. 60 μ M H₂O₂+Rb1 group. Rb1, ginsenoside Rb1; H₂O₂, hydrogen peroxide; HUVEC, human umbilical vein endothelial cell; AMPK, AMP-activated protein kinase; PAI-1, plasminogen activator inhibitor-1; eNOS, endothelial nitric oxide synthase; p, phosphorylated; NO, nitric oxide; Comp C, compound C.

SIRT1 in the endothelium reverses H₂O₂-induced premature cellular senescence through an eNOS-dependent signaling pathway. The present study also confirmed that inhibition of SIRT1 by nicotinamide caused a stress-induced increase in PAI-1 with a concomitant decrease in eNOS expression and NO production, which are consistent with Cacicedo *et al* (64). Furthermore, the present study demonstrated that Rb1 induced AMPK phosphorylation, which was

abolished by nicotinamide. Previous studies have reported the functional connections between the two master regulators SIRT1 and AMPK: Gao *et al* (65) reported that activated SIRT1 functionally interacts with AMPK α and upregulates its phosphorylation in aortic endothelial and smooth muscle cells of Klotho knockout mice and other studies have shown that AMPK can also function as a SIRT1 activator by increasing the NAD⁺/NADH ratio (66,67). These studies




Figure 4. NAM blocks the protective effects of Rb1 against H₂O₂-induced senescence in HUVECs. (A) The protein levels of phosphorylated AMPK, AMPK, PAI-1, eNOS, SIRT1 and GAPDH were determined by western blot. (B) Quantitative analysis of phosphorylated AMPK expression ratio and phosphorylated AMPK levels. Total AMPK and GAPDH were also determined. (C) Quantitative analysis of PAI-1 expression ratio and PAI-1 expression levels. (D) Quantitative analysis of eNOS expression ratio and eNOS expression levels. (E) NO production measured by the Griess reaction and expressed as the percentage of the control. (F) The amount of ATP in the cells was measured using an ATP determination kit. Results represent three independent experiments. **P<0.01, vs. control group. #P<0.05 vs. 60 μM H₂O₂ group. &P<0.05, vs. 60 μM H₂O₂+Rb1 group. NAM, nicotinamide; Rb1, ginsenoside Rb1; H₂O₂, hydrogen peroxide; HUVEC, human umbilical vein endothelial cell; AMPK, AMP-activated protein kinase; PAI-1, plasminogen activator inhibitor-1; eNOS, endothelial nitric oxide synthase; SIRT1, Sirtuin-1; p, phosphorylated; NO, nitric oxide.

suggest that AMPK may crosstalk with SIRT1 to modulate downstream targets. The present study demonstrated that

inhibition of SIRT1 by nicotinamide functionally suppressed the phosphorylation of AMPK, which in turn abolished the

protective effect against Rb1 on H₂O₂-induced endothelial dysfunction. However, *in vivo* experiments are required to confirm the anti-aging and antioxidant effects of Rb1 demonstrated *in vitro*. Further research is also required to elucidate the complex molecular mechanisms underlying the involvement of AMPK and SIRT1 in anti-aging and antioxidant processes.

In conclusion, Rb1 improved H₂O₂-induced HUVEC senescence and dysfunction through the SIRT1/AMPK signaling pathway. The results of the present study suggested that the associated mechanisms may be related to decreased PAI-1 expression and upregulated eNOS expression and NO production. The present study provides evidence to support the novel role of AMPK in the beneficial effects of Rb1 on HUVEC senescence, which can be explored further in animal models and clinical studies.

Acknowledgements

The authors would like to thank Mr Mitchell Arico for editing the English text of a draft of this manuscript.

Funding

This study was supported by the grants from the National Natural Science Foundation of China (grant nos. 81370447; 81300707); Science and Technology Planning Project of Guangdong Province (grant no. 2016A050502014). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Availability of data and materials

The datasets used and/or analyzed during the present study are available from the corresponding author on reasonable request.

Authors' contributions

ZZ, JZ and XQ designed the study. ZZ, MW, CC, DL and LW conducted the research. ZZ and MW analyzed the data and wrote the manuscript. ZZ and MW revised the discussion section of the manuscript. All authors read and approved the final manuscript.

Ethics approval and consent to participate

The isolation procedure for HUVECs was approved by the Research Committee at the Third Affiliated Hospital of Sun Yat-sen University. The donors provided written informed consent to donate the umbilical cords.

Patient consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

1. Quyyumi AA: Endothelial function in health and disease: New insights into the genesis of cardiovascular disease. *Am J Med* 105: 32S-39S, 1998.
2. Minamino T, Miyauchi H, Yoshida T, Ishida Y, Yoshida H and Komuro I: Endothelial cell senescence in human atherosclerosis: Role of telomere in endothelial dysfunction. *Circulation* 105: 1541-1544, 2002.
3. Barton M: Obesity and aging: Determinants of endothelial cell dysfunction and atherosclerosis. *Pflugers Arch* 460: 825-837, 2010.
4. Ota H, Eto M, Kano MR, Kahyo T, Setou M, Ogawa S, Iijima K, Akishita M and Ouchi Y: Induction of endothelial nitric oxide synthase, SIRT1, and catalase by statins inhibits endothelial senescence through the Akt pathway. *Arterioscler Thromb Vasc Biol* 30: 2205-2211, 2010.
5. Ruan Y, Wu S, Zhang L, Chen G and Lai W: Retarding the senescence of human vascular endothelial cells induced by hydrogen peroxide: Effects of 17beta-estradiol (E2) mediated mitochondria protection. *Biogerontology* 15: 367-375, 2014.
6. Cai H: Hydrogen peroxide regulation of endothelial function: Origins, mechanisms, and consequences. *Cardiovasc Res* 68: 26-36, 2005.
7. Han X, Tai H, Wang X, Wang Z, Zhou J, Wei X, Ding Y, Gong H, Mo C, Zhang J, *et al*: AMPK activation protects cells from oxidative stress-induced senescence via autophagic flux restoration and intracellular NAD(+) elevation. *Aging Cell* 15: 416-427, 2016.
8. Conti V, Corbi G, Simeon V, Russomanno G, Manzo V, Ferrara N and Filippelli A: Aging-related changes in oxidative stress response of human endothelial cells. *Aging Clin Exp Res* 27: 547-553, 2015.
9. El Assar M, Angulo J and Rodriguez-Manas L: Oxidative stress and vascular inflammation in aging. *Free Radic Biol Med* 65: 380-401, 2013.
10. Sindler AL, Delp MD, Reyes R, Wu G and Muller-Delp JM: Effects of ageing and exercise training on eNOS uncoupling in skeletal muscle resistance arterioles. *J Physiol* 587: 3885-3897, 2009.
11. Kang LS, Reyes RA and Muller-Delp JM: Aging impairs flow-induced dilation in coronary arterioles: Role of NO and H(2)O(2). *Am J Physiol Heart Circ Physiol* 297: H1087-H1095, 2009.
12. Xiong Y, Shen L, Liu KJ, Tso P, Xiong Y, Wang G, Woods SC and Liu M: Antiobesity and antihyperglycemic effects of ginsenoside Rb1 in rats. *Diabetes* 59: 2505-2512, 2010.
13. Chen S, Li X, Wang Y, Mu P, Chen C, Huang P and Liu D: Ginsenoside Rb1 attenuates intestinal ischemia/reperfusion-induced inflammation and oxidative stress via activation of the PI3K/Akt/Nrf2 signaling pathway. *Mol Med Rep* 19: 3633-3641, 2019.
14. Sun Q, Meng QT, Jiang Y, Liu HM, Lei SQ, Su WT, Duan WN, Wu Y, Xia ZY and Xia ZY: Protective effect of ginsenoside Rb1 against intestinal ischemia-reperfusion induced acute renal injury in mice. *PLoS One* 8: e80859, 2013.
15. Wang M, Chen Y, Xiong Z, Yu S, Zhou B, Ling Y, Zheng Z, Shi G, Wu Y and Qian X: Ginsenoside Rb1 inhibits free fatty acids-induced oxidative stress and inflammation in 3T3-L1 adipocytes. *Mol Med Rep* 16: 9165-9172, 2017.
16. Liu DH, Chen YM, Liu Y, Hao BS, Zhou B, Wu L, Wang M, Chen L, Wu WK and Qian XX: Rb1 protects endothelial cells from hydrogen peroxide-induced cell senescence by modulating redox status. *Biol Pharm Bull* 34: 1072-1077, 2011.
17. Herzig S and Shaw RJ: AMPK: Guardian of metabolism and mitochondrial homeostasis. *Nat Rev Mol Cell Biol* 19: 121-135, 2018.
18. Zhang M, Wang S, Pan Z, Ou T, Ma J, Liu H, Li R, Yang P, Han W, Guan S, *et al*: AMPK/NF-KB signaling pathway regulated by ghrelin participates in the regulation of HUVEC and THP1 Inflammation. *Mol Cell Biochem* 437: 45-53, 2018.
19. Thornton CC, Al-Rashed F, Calay D, Birdsey GM, Bauer A, Mylroie H, Morley BJ, Randi AM, Haskard DO, Boyle JJ and Mason JC: Methotrexate-mediated activation of an AMPK-CREB-dependent pathway: A novel mechanism for vascular protection in chronic systemic inflammation. *Ann Rheum Dis* 75: 439-448, 2016.
20. Martinez-Martin N, Blas-Garcia A, Morales JM, Marti-Cabrera M, Monleon D and Apostolova N: Metabolomics of the effect of AMPK activation by AICAR on human umbilical vein endothelial cells. *Int J Mol Med* 29: 88-94, 2012.

21. Huang NL, Chiang SH, Hsueh CH, Liang YJ, Chen YJ and Lai LP: Metformin inhibits TNF- α -induced I κ B kinase phosphorylation, I κ B- α degradation and IL-6 production in endothelial cells through PI3K-dependent AMPK phosphorylation. *Int J Cardiol* 134: 169-175, 2009.
22. Nagata D, Mogi M and Walsh K: AMP-activated protein kinase (AMPK) signaling in endothelial cells is essential for angiogenesis in response to hypoxic stress. *J Biol Chem* 278: 31000-31006, 2003.
23. Ido Y, Carling D and Ruderman N: Hyperglycemia-induced apoptosis in human umbilical vein endothelial cells: Inhibition by the AMP-activated protein kinase activation. *Diabetes* 51: 159-167, 2002.
24. Zhou H, Fu B, Xu B, Mi X, Li G, Ma C, Xie J, Li J and Wang Z: Rosmarinic acid alleviates the endothelial dysfunction induced by hydrogen peroxide in rat aortic rings via activation of AMPK. *Oxid Med Cell Longev* 2017: 7091904, 2017.
25. Zaborska KE, Edwards G, Austin C and Wareing M: The role of O-GlcNAcylation in perivascular adipose tissue dysfunction of offspring of high-fat diet-fed rats. *J Vasc Res* 54: 79-91, 2017.
26. Baudin B, Bruneel A, Bosselut N and Vaubourdel M: A protocol for isolation and culture of human umbilical vein endothelial cells. *Nat Protoc* 2: 481-485, 2007.
27. Guo Y, Xing L, Qian C, Ding Z and Jin B: Involvement of flavonoids from the leaves of *Carya cathayensis* sarg. In sirtuin 1 expression in HUVEC senescence. *Evid Based Complement Alternat Med* 2018: 8246560, 2018.
28. Esfahanian N, Shakiba Y, Nikbin B, Soraya H, Maleki-Dizaji N, Ghazi-Khansari M and Garjani A: Effect of metformin on the proliferation, migration, and MMP-2 and -9 expression of human umbilical vein endothelial cells. *Mol Med Rep* 5: 1068-1074, 2012.
29. Zheng S, Li W, Xu M, Bai X, Zhou Z, Han J, Shyy JY and Wang X: Calcitonin gene-related peptide promotes angiogenesis via AMP-activated protein kinase. *Am J Physiol Cell Physiol* 299: C1485-C1492, 2010.
30. Green LC, Wagner DA, Glogowski J, Skipper PL, Wishnok JS and Tannenbaum SR: Analysis of nitrate, nitrite, and [15N] nitrate in biological fluids. *Anal Biochem* 126: 131-138, 1982.
31. Debaq-Chainiaux F, Erusalimsky JD, Campisi J and Toussaint O: Protocols to detect senescence-associated beta-galactosidase (SA- β gal) activity, a biomarker of senescent cells in culture and in vivo. *Nat Protoc* 4: 1798-1806, 2009.
32. Luo W, Wang Y, Yang H, Dai C, Hong H, Li J, Liu Z, Guo Z, Chen X, He P, *et al*: Heme oxygenase-1 ameliorates oxidative stress-induced endothelial senescence via regulating endothelial nitric oxide synthase activation and coupling. *Aging (Albany NY)* 10: 1722-1744, 2018.
33. Yan J, Wang J, Huang H, Huang Y, Mi T, Zhang C and Zhang L: Fibroblast growth factor 21 delayed endothelial replicative senescence and protected cells from H₂O₂-induced premature senescence through SIRT1. *Am J Transl Res* 9: 4492-4501, 2017.
34. Lin XL, Liu Y, Liu M, Hu H, Pan Y, Fan XJ, Hu XM and Zou WW: Inhibition of hydrogen peroxide-induced human umbilical vein endothelial cells aging by allicin depends on sirtuin1 activation. *Med Sci Monit* 23: 563-570, 2017.
35. Hahn HJ, Kim KB, An IS, Ahn KJ and Han HJ: Protective effects of rosmarinic acid against hydrogen peroxide-induced cellular senescence and the inflammatory response in normal human dermal fibroblasts. *Mol Med Rep* 16: 9763-9769, 2017.
36. Song Z, Liu Y, Hao B, Yu S, Zhang H, Liu D, Zhou B, Wu L, Wang M, Xiong Z, *et al*: Ginsenoside Rb1 prevents H₂O₂-induced HUVEC senescence by stimulating sirtuin-1 pathway. *PLoS One* 9: e112699, 2014.
37. Liu DH, Chen YM, Liu Y, Hao BS, Zhou B, Wu L, Wang M, Chen L, Wu WK and Qian XX: Ginsenoside Rb1 reverses H₂O₂-induced senescence in human umbilical endothelial cells: Involvement of eNOS pathway. *J Cardiovasc Pharmacol* 59: 222-230, 2012.
38. Erusalimsky JD and Kurz DJ: Cellular senescence in vivo: Its relevance in ageing and cardiovascular disease. *Exp Gerontol* 40: 634-642, 2005.
39. Shan HY, Bai XJ and Chen XM: Apoptosis is involved in the senescence of endothelial cells induced by angiotensin II. *Cell Biol Int* 32: 264-270, 2008.
40. Li CX, Gao JG, Wan XY, Chen Y, Xu CF, Feng ZM, Zeng H, Lin YM, Ma H, Xu P, *et al*: Allyl isothiocyanate ameliorates lipid accumulation and inflammation in nonalcoholic fatty liver disease via the Sirt1/AMPK and NF- κ B signaling pathways. *World J Gastroenterol* 25: 5120-5133, 2019.
41. Wang T, Gu J, Wu PF, Wang F, Xiong Z, Yang YJ, Wu WN, Dong LD and Chen JG: Protection by tetrahydroxystilbene glucoside against cerebral ischemia: Involvement of JNK, SIRT1, and NF- κ B pathways and inhibition of intracellular ROS/RNS generation. *Free Radic Biol Med* 47: 229-240, 2009.
42. Lee JH, Song MY, Song EK, Kim EK, Moon WS, Han MK, Park JW, Kwon KB and Park BH: Overexpression of SIRT1 protects pancreatic beta-cells against cytokine toxicity by suppressing the nuclear factor- κ B signaling pathway. *Diabetes* 58: 344-351, 2009.
43. Huang K, Li R and Wei W: Sirt1 activation prevents anti-Thy 1.1 mesangial proliferative glomerulonephritis in the rat through the Nrf2/ARE pathway. *Eur J Pharmacol* 832: 138-144, 2018.
44. Huang K, Chen C, Hao J, Huang J, Wang S, Liu P and Huang H: Polydatin promotes Nrf2-ARE anti-oxidative pathway through activating Sirt1 to resist AGEs-induced upregulation of fibronectin and transforming growth factor- β 1 in rat glomerular mesangial cells. *Mol Cell Endocrinol* 399: 178-189, 2015.
45. Chen AF, Chen DD, Daiber A, Faraci FM, Li H, Rembold CM and Laher I: Free radical biology of the cardiovascular system. *Clin Sci (Lond)* 123: 73-91, 2012.
46. Zhou P, Lu S, Luo Y, Wang S, Yang K, Zhai Y, Sun G and Sun X: Attenuation of TNF- α -induced inflammatory injury in endothelial cells by ginsenoside Rb1 via inhibiting NF- κ B, JNK and p38 signaling pathways. *Front Pharmacol* 8: 464, 2017.
47. Yuan Q, Jiang YW, Ma TT, Fang QH and Pan L: Attenuating effect of Ginsenoside Rb1 on LPS-induced lung injury in rats. *J Inflamm (Lond)* 11: 40, 2014.
48. Hwang HJ, Jung TW, Kim JW, Kim JA, Lee YB, Hong SH, Roh E, Choi KM, Baik SH and Yoo HJ: Protectin DX prevents H₂O₂-mediated oxidative stress in vascular endothelial cells via an AMPK-dependent mechanism. *Cell Signal* 53: 14-21, 2019.
49. Wang L, Han J, Shan P, You S, Chen X, Jin Y, Wang J, Huang W, Wang Y and Liang G: MD2 blockage protects obesity-induced vascular remodeling via activating AMPK/Nrf2. *Obesity (Silver Spring)* 25: 1532-1539, 2017.
50. Lu Q, Mundy M, Chambers E, Lange T, Newton J, Borgas D, Yao H, Choudhary G, Basak R, Oldham M and Rounds S: Alda-1 protects against acrolein-induced acute lung injury and endothelial barrier dysfunction. *Am J Respir Cell Mol Biol* 57: 662-673, 2017.
51. Kroller-Schon S, Jansen T, Hauptmann F, Schuler A, Heeren T, Hausding M, Oelze M, Viollet B, Keane JF Jr, Wenzel P, *et al*: α 1AMP-activated protein kinase mediates vascular protective effects of exercise. *Arterioscler Thromb Vasc Biol* 32: 1632-1641, 2012.
52. Morrow VA, Fougelle F, Connell JM, Petrie JR, Gould GW and Salt IP: Direct activation of AMP-activated protein kinase stimulates nitric-oxide synthesis in human aortic endothelial cells. *J Biol Chem* 278: 31629-31639, 2003.
53. Gaskin FS, Kamada K, Yusof M and Korhuis RJ: 5'-AMP-activated protein kinase activation prevents postischemic leukocyte-endothelial cell adhesive interactions. *Am J Physiol Heart Circ Physiol* 292: H326-H332, 2007.
54. Han F, Zhang S, Hou N, Wang D and Sun X: Irisin improves endothelial function in obese mice through the AMPK-eNOS pathway. *Am J Physiol Heart Circ Physiol* 309: H1501-H1508, 2015.
55. Winnik S, Auwerx J, Sinclair DA and Matter CM: Protective effects of sirtuins in cardiovascular diseases: From bench to bedside. *Eur Heart J* 36: 3404-3412, 2015.
56. Stein S and Matter CM: Protective roles of SIRT1 in atherosclerosis. *Cell Cycle* 10: 640-647, 2011.
57. Imai S, Armstrong CM, Kaerberlein M and Guarente L: Transcriptional silencing and longevity protein Sir2 is an NAD-dependent histone deacetylase. *Nature* 403: 795-800, 2000.
58. Yuan Q, Zhang D, Liu C, Zhang C and Yuan D: Chikusetsusaponin V inhibits LPS-activated inflammatory responses via SIRT1/NF- κ B signaling pathway in RAW264.7 cells. *Inflammation* 41: 2149-2159, 2018.
59. Qian Y, Chen C, Ma L, Wang Z, Wang LF, Zuo L, Yang Y, Huang X, Jiang M, Wang X, *et al*: CD38 deficiency promotes inflammatory response through activating Sirt1/NF- κ B-mediated inhibition of TLR2 expression in macrophages. *Mediators Inflamm* 2018: 8736949, 2018.
60. Li M, Hong W, Hao C, Li L, Wu D, Shen A, Lu J, Zheng Y, Li P and Xu Y: SIRT1 antagonizes liver fibrosis by blocking hepatic stellate cell activation in mice. *FASEB J* 32: 500-511, 2018.
61. Kao CL, Chen LK, Chang YL, Yung MC, Hsu CC, Chen YC, Lo WL, Chen SJ, Ku HH and Hwang SJ: Resveratrol protects human endothelium from H₂O₂-induced oxidative stress and senescence via Sirt1 activation. *J Atheroscler Thromb* 17: 970-979, 2010.

62. Ota H, Akishita M, Eto M, Iijima K, Kaneki M and Ouchi Y: Sirt1 modulates premature senescence-like phenotype in human endothelial cells. *J Mol Cell Cardiol* 43: 571-579, 2007.
63. Ota H, Eto M, Kano MR, Ogawa S, Iijima K, Akishita M and Ouchi Y: Cilostazol inhibits oxidative stress-induced premature senescence via upregulation of Sirt1 in human endothelial cells. *Arterioscler Thromb Vasc Biol* 28: 1634-1639, 2008.
64. Cacicedo JM, Gauthier MS, Lebrasseur NK, Jasuja R, Ruderman NB and Ido Y: Acute exercise activates AMPK and eNOS in the mouse aorta. *Am J Physiol Heart Circ Physiol* 301: H1255-H1265, 2011.
65. Gao D, Zuo Z, Tian J, Ali Q, Lin Y, Lei H and Sun Z: Activation of SIRT1 attenuates klotho deficiency-induced arterial stiffness and hypertension by enhancing AMP-activated protein kinase activity. *Hypertension* 68: 1191-1199, 2016.
66. Canto C, Gerhart-Hines Z, Feige JN, Lagouge M, Noriega L, Milne JC, Elliott PJ, Puigserver P and Auwerx J: AMPK regulates energy expenditure by modulating NAD⁺ metabolism and SIRT1 activity. *Nature* 458: 1056-1060, 2009.
67. Chau MD, Gao J, Yang Q, Wu Z and Gromada J: Fibroblast growth factor 21 regulates energy metabolism by activating the AMPK-SIRT1-PGC-1 α pathway. *Proc Natl Acad Sci USA* 107: 12553-12558, 2010.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.