

Neiguan (PC6)-based acupuncture pretreatment for myocardial ischemia reperfusion injury

A protocol for preclinical systematic review and meta-analysis

Qiqi Yang, MM^a , Huifang Mao, PhD^b, Xia Chen, MM^b, Yanji Zhang, MM^a, Xiaolei Zhang, MM^a, Zhenzhen Liu, MM^a, Gongdao Jiang, MM^{a,*}, Wei Huang, PhD^{b,*} 

Abstract

Background: Myocardial ischemia reperfusion injury (MIRI) is 1 of the leading causes of disability and mortality worldwide in the cardiovascular diseases. Acupuncture has been widely applied in the treatment and prevention of cardiovascular diseases in recent years. This systematic review protocol aims to provide the methods for evaluating the efficacy of Neiguan (PC6)-based acupuncture pretreatment in animal models of MIRI.

Methods and analysis: The electronic databases of PubMed, Embase, Cochrane Library, as well as the Chinese databases such as China National Knowledge Infrastructure, Chinese Science and Technology Periodical Database, China Biology Medicine Database and WanFang Database will be searched from inception to November 2019. The outcome measures were myocardial infarct size, the level of ST-segment elevation, left ventricular ejection fraction, shortening fraction, arrhythmia score, cardiac enzymes, and cardiac troponin. Study inclusion, data extraction and quality assessment will be performed independently by 2 reviewers. RevMan 5.3 software will be used for the data synthesis and the quality of each study will be assessed independently by using the Collaborative Approach To Meta-Analysis And Review Of Animal Data From Experimental Studies checklist with minor modification.

Results: This review will provide a high-quality synthesis of Neiguan (PC6)-based acupuncture pretreatment for MIRI in animal models

Conclusions: This systematic review will provide conclusive evidence for whether Neiguan (PC6)-based acupuncture pretreatment is an effective intervention in animal models of MIRI.

Trial registration number: PROSPERO CRD42020175144.

Abbreviations: MIRI = myocardial ischemia reperfusion injury, TCM = traditional Chinese medicine.

Keywords: acupuncture, myocardial ischemia reperfusion injury, Neiguan (PC6), pretreatment, protocol, systematic review

Ethical approval is not necessary since this protocol is only for systematic review and does not involve privacy data or conduct an animal experiment. This protocol will be disseminated by a peer-review journal or conference presentation.

Youth Fund Project of National Natural Science Foundation of China (no. 81704142), China Association of Science and Technology Youth Talents Invitation Project (no. 2017QNRC001), Wuhan young and middle-aged medical backbone talents (sixth batch) (no. 116, family planning tong [2018]), Hubei hospital of traditional Chinese medicine, the first Tanhualin famous doctor, student training project (Hubei traditional Chinese medicine yard: [2018] no. 72), and Hubei Provincial Second Medical Leading Talent Project Training Object and Hubei Famous Medical Studio Project (no. 47 [2019])

The funders had no role in the design, execution, or writing of the study.

The authors have no conflicts of interest to disclose.

Data sharing not applicable to this article as no datasets were generated or analyzed during the current study.

^a Hubei University of Chinese Medicine/The Co-innovation Center for Preventive Treatment of Disease of Acupuncture-moxibustion in Hubei Province, ^b Hubei Provincial Hospital of Traditional Chinese Medicine, Wuhan, China.

** Correspondence: Wei Huang, Hubei Provincial Hospital of TCM, No. 4 Garden Hill, Liangdao Street, Wuchang District, Wuhan, Hubei 430061, China (e-mail: huangwei@hbhctm.com), Gongdao Jiang, Hubei University of Chinese Medicine, No. 01 Tanhualin, Wuchang District, Wuhan, Hubei 430061, China (e-mail: 22108217@qq.com).*

Copyright © 2020 the Author(s). Published by Wolters Kluwer Health, Inc.

This is an open access article distributed under the Creative Commons Attribution License 4.0 (CCBY), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Yang Q, Mao H, Chen X, Zhang Y, Zhang X, Liu Z, Jiang G, Huang W. Neiguan (PC6)-based acupuncture pretreatment for myocardial ischemia reperfusion injury: A protocol for preclinical systematic review and meta-analysis. Medicine 2020;99:28(e20792).

Received: 11 May 2020 / Accepted: 21 May 2020

<http://dx.doi.org/10.1097/MD.00000000000020792>

1. Introduction

Acute myocardial infarction (AMI) is the myocardial necrosis caused by acute ischemia and hypoxia of coronary artery, which is 1 of the leading causes of disability and mortality worldwide.^[1,2] In the US, the overall prevalence for AMI is 3.0% in adults older than 20.^[3] Within China, the mortality rate of AMI in urban areas and rural areas is about 58.69/100,000 people and 74.72/100000 people respectively,^[4] and it is estimated that the total annual cost of hospitalization for AMI alone is up to RMB 19.85 billion.^[5] Although various timely and effective revascularization, such as percutaneous coronary intervention and coronary artery bypass grafting, have been widely used in clinic to rescue ischemic myocardium, and can save patients' lives to a certain extent.^[6] However, the process of reperfusion may further damage myocardial ultrastructure, metabolism and function, even lead to a further death of cardiomyocytes which contributes up to 50% of final myocardial damage.^[7,8] Moreover, it has been reported that nearly 900000 people have cardiovascular events caused by myocardial ischemia reperfusion injury (MIRI) such as perioperative myocardial infarction, heart failure and sudden cardiac death every year in the world, with a mortality rate of 10% to 15%.^[9] All in all, MIRI not only reduce the benefits of reperfusion therapies, lead to a significant increase in the economic burden of MIRI-related medical systems, but also seriously threatens patients' lives.^[7]

At present, prevention plays an important role in controlling MIRI. It has been proved that pretreatment before ischemia or reperfusion can make body generate moderate pressure, and active the endogenous adaptation mechanism to prevent or reduce the damage of subsequent diseases.^[10] The current means to activate the endogenous protection mechanism of body mainly include ischemic preconditioning,^[11] drug preconditioning,^[12] temperature preconditioning^[13] and exercise preconditioning,^[14] which can enhance the tolerance of the heart to the damage caused by myocardial I/R injury. However, these pretreatment measures are cumbersome to implement and inevitably cause trauma to the body, thereby greatly limiting their translation into clinical benefits. Therefore, seeking an effective, non-invasive and simple pretreatment method has become an urgent clinical need.

Acupuncture, as a key component of traditional Chinese medicine (TCM), is a therapy that uses a sterile needle to penetrate a specific acupoint in body to treat specific diseases.^[15] With minimal side effects and strong operability, it has been 1 of the most widely used complementary therapies in many countries.^[16] Previous studies have shown that acupuncture pretreatment can also activate endogenous protective mechanisms of body and improve immunity.^[17] In addition, the significance of acupoint selection in acupuncture therapy has also been affirmed. Neiguan (PC6) acupoint is the Luo-connecting point of pericardial meridian, which has the specific connection with the heart. Acupuncture at Neiguan (PC6) has dual-directional regulation effects on cardiovascular system, which has the characteristics of specific treatment of cardiovascular disease.^[18] A large number of studies have revealed that Neiguan (PC6)-based acupuncture pretreatment can exert cardio-protective effects in animal models of MIRI by improving coronary vasodilation and energy metabolism, inhibiting of oxidative stress, inflammatory response and apoptosis.^[19-22] Therefore, Neiguan (PC6)-based acupuncture pretreatment undoubtedly provides new inspiration for the clinical prevention and treatment of MIRI.

However, due to low quality, small samples of included studies and no study protocols reported in advance, 2 systematic reviews published in Chinese have failed to translate the efficacy of acupuncture into clinical benefits.^[23,24] And 1 review mainly focused on the add-on effect of electroacupuncture and ignored the critical effect of preconditioning in MIRI.^[24] Meanwhile, considerable new studies in this field have been published within past years. Thus, it is necessary to make a systematic review and meta-analysis based on the most comprehensive and up-to-date resources to provide a convincing conclusion whether Neiguan (PC6)-based acupuncture pretreatment is an effective cardio-protective intervention for MIRI.

2. Methods

2.1. Study registration

This systematic review protocol has been registered on PROSPERO with number CRD42020175144 (https://www.crd.york.ac.uk/prospero/display_record.php?RecordID=175144). Additionally, the current protocol report adheres to the Cochrane Handbook for Systematic Reviews and Meta-Analysis Protocol guidelines.^[25] Any change of the review will be described if needed.

2.2. Inclusion criteria

2.2.1. Types of studies. Only randomized controlled animal study published in English and Chinese about acupuncture for MIRI will be included.

2.2.2. Types of participants. Rat models of MIRI which were induced by ligation of the left anterior descending coronary artery (LAD) regardless of their age and sex.

2.2.3. Types of interventions. Only manual acupuncture or electroacupuncture (EA) at Neiguan (PC6) point alone or a combination of PC6 and other acupoints will be used in experimental group. The intervention time is limited to before reperfusion.

2.2.4. Types of comparators. Control intervention will be limited to no treatment or sham acupuncture.

2.2.5. Types of outcome measures. Myocardial infarct size, the level of ST-segment elvation, left ventricular ejection fraction (LVEF), shortening fraction (FS) or arrhythmia score will be evaluated as the primary outcomes. And the secondary outcomes were cardiac enzymes (creatine kinase, creatine kinase-MB, lactate dehydrogenase, etc) or cardiac troponin (cardiac troponin T, cardiac troponin I, cardiac troponin C).

2.3. Exclusion criteria

- (1) Case report or cross-over studies or review or meta-analysis or meeting abstract.
- (2) Non-rat animal model or in vitro animal model or in silico models.
- (3) MIRI animals with complications or induced by modeling methods other than ligation of LAD.
- (4) Acupuncture therapies combined with other forms of acupuncture (such as auricular acupuncture, acupoint injection, moxibustion and etc) or TCM.
- (5) Other acupuncture treatments were used in control group.
- (6) Duplicate publication or literature published by identical data.

- (7) Not published in peer-review journals.
- (8) No predetermined outcome index.

2.4. Search methods for identification of studies

2.4.1. Electronic searches. Experimental studies estimating the efficacy of Neiguan (PC6)-based acupuncture pretreatment in animal models of MIRI were systematically searched from EMBASE, PubMed, Cochrane Library, Wanfang database, China National Knowledge Infrastructure, Chinese Science and Technology Periodical Database, China Biology Medicine disc from inception to November 2019. No restrictions on language and publication were applied in the search strategy. Moreover, reference lists of potential studies were searched for relevant studies. And the searches will be re-run just before the final analyses to retrieve the most recent studies eligible for inclusion.

2.4.2. Searching strategy. The search strategy for PubMed is shown in Table 1, which includes all search terms. Other electronic databases will be used the same strategy.

2.5. Data collection and analysis

2.5.1. Selection of studies. All reviewers will receive professional training to understand the objective and process of the review before the selection of studies. All the retrieved studies will be managed with EndnoteX7, and the duplicated studies will be discarded. Two review authors (YQQ and ZYJ) screened independently titles and abstracts of studies to identify studies that potentially meet the inclusion criteria outlined above. Then the full text of these potentially eligible studies will be independently assessed for eligibility by 2 reviewers (YQQ and ZYJ). Any disagreement between them over the eligibility of studies will be discussed with the third reviewer (ZXL). The procedures of study selection will be performed in accordance with the Preferred Reporting Items for Systematic reviews and Meta-Analysis flow chart (see Fig. 1).

2.5.2. Data extraction and management. Two independent authors (ZXL and LZZ) extracted the following details from included studies:

- (1) name of first author, year of publication;
- (2) the detail information of animals for each study, including animal species, number, sex, and weight;
- (3) the use of anesthesia in the experiment and the methods to establish animal models;
- (4) information on treatment was obtained, including timing and duration for treatment, acupoint selection, types and method of acupuncture procedure;
- (5) all the outcome measures and its intergroup differences.

If the data for meta-analysis were missing or only expressed graphically, we attempted to contact the authors to obtain the detailed data. Any disagreement noticed in the process of data cross-checking will be discussed with the third reviewer.(ZYJ).

2.5.3. Assessment of risk of bias and reporting quality of included studies. We will evaluate the methodological quality of the included studies by using a checklist of the Collaborative Approach to Meta-Analysis and Review of Animal Data from Experimental Studies^[26] with minor modifications.^[27] Ten items are as follows:

- (1) publication in a peer-reviewed journal;
- (2) statement of temperature control;

Table 1	
Search strategy for PubMed database.	
Number	Search terms
1	Myocardial reperfusion injury [Mesh]
2	Myocardial reperfusion [Mesh]
3	Myocardial ischemia [Mesh]
4	Myocardial ischemic reperfusion injury [Title/Abstract]
5	Injuries, myocardial reperfusion [Title/Abstract]
6	Myocardial reperfusion injuries [Title/Abstract]
7	Reperfusion injuries, myocardial [Title/Abstract]
8	Reperfusion injury, myocardial [Title/Abstract]
9	Injury, myocardial reperfusion [Title/Abstract]
10	Reperfusion, myocardial [Title/Abstract]
11	Myocardial reperfusion [Title/Abstract]
12	Reperfusion, myocardial [Title/Abstract]
13	Coronary reperfusion [Title/Abstract]
14	Coronary reperfusion [Title/Abstract]
15	Reperfusion, coronary [Title/Abstract]
16	1 OR 2 OR 3 OR 4 OR 5 OR 6 OR 7 OR 8 OR 9 OR 10 OR 11 OR 12 OR 13 OR #14 OR 15
17	Acupuncture [Mesh]
18	Acupuncture Points [Mesh]
19	Acupuncture Therapy [Mesh]
20	Electroacupuncture [Mesh]
21	Manual acupuncture [Title/Abstract]
22	Silver needle [Title/Abstract]
23	Electro-acupuncture [Title/Abstract]
24	Meridian [Title/Abstract]
25	Needle pricking [Title/Abstract]
26	Pricking therapy [Title/Abstract]
27	Point application [Title/Abstract]
28	Acupotomy [Title/Abstract]
29	Trigger point [Title/Abstract]
30	De qi [Title/Abstract]
31	Acupressure [Title/Abstract]
32	17 OR 18 OR 19 OR 20 OR 21 OR 22 OR 23 OR 24 OR 25 OR 26 OR 27 OR 28 OR 29 OR 30 OR 31
33	Neiguan [All Fields]
34	Neiguan [All Fields]
35	PC 6 [All Fields]
36	33 OR 34 OR 35
37	32 AND 36
38	Animal [Mesh]
39	Animal Experimentation [All Fields]
40	animals [All Fields]
41	animal model [All Fields]
42	experimental model [All Fields]
43	38 OR 39 OR 40 OR 41 OR 42
44	16 AND 37 AND 43

mh = Mesh.

- (3) random allocation to groups;
- (4) allocation concealment;
- (5) blinded assessment of outcome;
- (6) use of anesthetic without significant intrinsic cardioprotective activity;
- (7) appropriate animal model (aged, diabetic, or hypertensive);
- (8) sample size calculation;
- (9) compliance with animal welfare regulations;
- (10) statement of potential conflict of interests.

The assessment of risk of bias will be carried out by 2 independent reviewers (LZZ and ZXL) Any disagreements will be arbitrated by a third reviewer (ZYJ).




Figure 1. Flow diagram of the study selection process.

2.5.4. Measurement of treatment effect. Efficacy data will be synthesized and statistically analyzed in RevMan 5.3 software. For dichotomous data, a risk ratio with 95% confidence intervals will be used for analysis. For continuous data, a mean difference or a standard mean difference with 95% confidence intervals will be used for analysis.

2.5.5. Unit of analysis issues. If outcomes are presented from the studies of animals at different time points, we extracted data from the last time point. For studies comparing different intensity or frequency of acupuncture treatment to a single control group, the data of treatment group with the maximum amount of treatment will be selected for analysis.

2.5.6. Dealing with missing data. The corresponding authors or relevant authors will be contacted to obtain insufficient data or missing data. If we received no response to our request, we will use GetData software to measure the data in the graph. If the complete data is still not available, then we will exclude the article of missing data from the data synthesis.

2.5.7. Assessment of heterogeneity. Heterogeneity between studies results will be investigated based on I^2 statistic. It will be considered significant heterogeneity while $I^2 > 50\%$. On the contrary, When $I^2 \leq 50\%$, study will be regarded as little or no heterogeneity.

2.5.8. Assessment of publication bias. A funnel plot analysis will be conducted to determine publication bias if 10 or more studies are included in the meta-analysis.

2.5.9. Data synthesis. All analyses will be performed with Revman 5.3 software by the Cochrane Collaboration. Meta-analysis will be performed for outcome measures reported in more than 2 studies. In the present meta-analysis, we will use the random effects model rather than the fixed effects model because heterogeneity between multi-studies has to be taken into account. If the I^2 test indicates unacceptable heterogeneity ($I^2 > 50\%$), sensitivity analysis and subgroup analysis will be performed to find out any possible reasons that may cause the heterogeneity. If meta-analysis is impossible, data will be reported by a descriptive

summary. Probability value $P < .05$ is considered statistically significant.

2.5.10. Subgroup analysis. Subgroup analyses on the following factors will be conducted to assess heterogeneity as well as possible: infarct time, reperfusion time, species of rat, type of acupuncture, acupoint prescription, the type of anesthetic, and the quality of study.

2.5.11. Sensitivity analysis. After conducting a quality assessment of the included studies, we will conduct a sensitivity analysis if there are studies of low quality. Sensitivity analysis will also be performed when heterogeneity testing suggests unacceptable heterogeneity between studies ($I^2 > 50\%$). Then we will obtain a stable consolidated result of our study.

3. Discussion

Acupuncture pretreatment, as a safe and reliable measure, is derived from the thought of preventive treatment in TCM. Before the occurrence of the disease, using acupuncture stimulate certain acupuncture points or meridians in advance can increase the pre-adaptability of the body to diseases, thereby reducing subsequent damage of diseases to organs.^[17] At present, this intervention is widely used in health care and provides broad application prospects in the prevention and treatment of cardiovascular diseases. Additionally, Neiguan (PC6) acupoint has always been the preferred acupoint for the acupuncture therapy of cardiovascular diseases, which is located 2 Cun proximal to the wrist, between the flexor carpi radialis and the palmaris longus tendons.^[28] Hence we chose Neiguan (PC6)-based acupuncture pretreatment as the intervention in this review. The therapeutic strategy not only combines the triple function of acupuncture, Neiguan acupoint and pretreatment, but also provide new treatment ideas in improving MIRI. Especially in clinical practice, this simple therapy is anticipated to protect heart against MIRI before revascularization.

Systematic reviews of pre-clinical animal data could clarify the underlying mechanism of human diseases,^[29] preclude unnecessary study replication and improve the likelihood of success of future clinical trials.^[30] Therefore, this systematic review will provide an in-depth summary and latest analysis of the latest evidence on the efficacy of Neiguan (PC6)-based acupuncture pretreatment in animal models of MIRI. We expect the findings of this study would provide reference basis for Chinese guidelines on the treatment of MIRI as well as promote acupuncture treatment and application of acupuncture points so as to benefit more patients in the future.^[31]

However, this systematic review also has potential limitations. Different types of acupuncture, needling techniques, and number of treatments may cause considerable heterogeneity in this review. In addition, limited to language ability, only studies in English and Chinese will be included, and reports in other languages may be ignored.

Author contributions

Qiqi Yang and, Huifang Mao and Xia Chen have contributed equally to this work. All authors have read and approved the publication of the protocol

Conceptualization: Qiqi Yang, Huifang Mao, Xia Chen.

Data curation: Xia Chen, Xiaolei Zhang, Zhenzhen Liu.

Formal analysis: Huifang Mao, Gongdao Jiang, Wei Huang.

Investigation: Qiqi Yang, Yanji Zhang.

Methodology: Xiaolei Zhang, Gongdao Jiang, Wei Huang.

Software: Qiqi Yang, Yanji Zhang, Zhenzhen Liu.

Supervision: Gongdao Jiang, Wei Huang.

Writing – original draft: Qiqi Yang, Huifang Mao, Xia Chen, Yanji Zhang.

Writing – review & editing: Xiaolei Zhang, Zhenzhen Liu, Gongdao Jiang, Wei Huang.

References

- [1] Thygesen K, Alpert JS, White HD, et al. Universal definition of myocardial infarction. *J Am Coll Cardiol* 2007;50:2173–95.
- [2] Santos-Gallego CG, Picatoste B, Badimón JJ. Pathophysiology of acute coronary syndrome. *Curr Atheroscler Rep* 2014;16:401.
- [3] Virani SS, Alonso A, Benjamin EJ, et al. Heart disease and stroke statistics-2020 update: a report from the American Heart Association. *Circulation* 2020;141:e139–596.
- [4] Ma LY, Wu YZ, Chen WW. Key points of the 2018 report on cardiovascular diseases in China [Chinese]. *Chin J Hypertension* 2019;27:712–6.
- [5] Hu ST, Yang YJ, Zheng Z, et al. Summary of the 2018 report on cardiovascular diseases in China [Chinese]. *Chin Circul J* 2019;34:209–20.
- [6] Heusch G, Gersh BJ. The pathophysiology of acute myocardial infarction and strategies of protection beyond reperfusion: a continual challenge. *Eur Heart J* 2017;38:774–84.
- [7] Yellon DM, Hausenloy DJ. Myocardial reperfusion injury. *N Engl J Med* 2007;357:1121–35.
- [8] Hausenloy DJ, Yellon DM. Myocardial ischemia-reperfusion injury: a neglected therapeutic target. *J Clin Invest* 2013;123:92–100.
- [9] Devereaux PJ, Goldman L, Cook DJ, et al. Perioperative cardiac events in patients undergoing noncardiac surgery: a review of the magnitude of the problem, the pathophysiology of the events and methods to estimate and communicate risk. *CMAJ* 2005;173:627–34.
- [10] Murry CE, Jennings RB, Reimer KA. Preconditioning with ischemia: a delay of lethal cell injury in ischemic myocardium. *Circulation* 1986;74:1124–36.
- [11] Przyklenk K, Bauer B, Ovize M, et al. Regional ischemic 'preconditioning' protects remote virgin myocardium from subsequent sustained coronary occlusion. *Circulation* 1993;87:893–9.
- [12] Szekeres L. Drug-induced delayed cardiac protection against the effects of myocardial ischemia. *Pharmacol Ther* 2005;108:269–80.
- [13] Donnelly TJ, Sievers RE, Vissern FL, et al. Heat shock protein induction in rat hearts. A role for improved myocardial salvage after ischemia and reperfusion? *Circulation* 1992;85:769–78.
- [14] Quindry JC, Hamilton KL, French JP, et al. Exercise-induced HSP-72 elevation and cardioprotection against infarct and apoptosis. *J Appl Physiol* 2007;103:1056–62.
- [15] Kaptchuk TJ. Methodological issues in trials of acupuncture. *JAMA* 2001;285:1015–6.
- [16] NIH Consensus Conference Acupuncture. *JAMA* 1998;280:1518–24.
- [17] Li XH. Effect and Mechanism of Preventive Moxibustion on Adjuvant Arthritis [Ph.D.]. Beijing, China: Beijing University of Chinese Medicine; 2004.
- [18] Yu J, Meng QL, Zhang YC, et al. Double-way regulation effect of electroacupuncture at neiguan point on cardiac arrhythmia rats and discussion on its central mechanism [Chinese]. *J Clin Acupunct Moxib* 2013;29:79–83.
- [19] Wang SK, Lu C, Wang YJ. Effects of electroacupuncture stimulation of "Neiguan" (PC 6) at different frequencies on plasma vasoactive substance levels in myocardial ischemic reperfusion rats. *Zhen Ci Yan Jiu* 2015;40:378–82.
- [20] Du T, Ren YL, He YG, et al. Action mechanism of acupuncture along meridians on improving energy metabolism in cardiomyocytes. *Zhongguo Zhen Jiu* 2017;37:1197–203.
- [21] Shao ML, Li Y, Cui HF, et al. Protective effect of acupuncture preconditioning on oxidative stress injury induced by myocardial ischemia-reperfusion injury in rats. *Zhongguo Zhen Jiu* 2017;37:285–90.
- [22] Lu SF, Huang Y, Wang N, et al. Cardioprotective effect of electroacupuncture pretreatment on myocardial ischemia/reperfusion injury via antiapoptotic signaling. *Evid Based Complement Alternat Med* 2016;2016:1–9.

- [23] Li H, Xiao N, Shao ML, et al. Systemic review on neiguan acupuncture pretreatment for myocardial ischemia reperfusion of rat experiments effects. *Chin J Basic Med Tradit Chin Med* 2006;22:1665–9.
- [24] Yang HS, Wu S, Zheng QY, et al. Meta-analysis on protective effect of electroacupuncture at “Nèigūān” (PC 6) in rats with reperfusion injury induced myocardial ischemia. *World J Acup-Mox* 2015;25:43–53.
- [25] Moher D, Shamseer L, Clarke M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Syst Rev* 2015;4:1.
- [26] Macleod MR, O’Collins T, Howells DW, et al. Pooling of animal experimental data reveals influence of study design and publication bias. *Stroke* 2004;35:1203–8.
- [27] Yu J, Zhang X, Zhang Y. Astragaloside attenuates myocardial injury in a rat model of acute myocardial infarction by upregulating hypoxia inducible factor-1 and Notch1/Jagged1 signaling. *Mol Med Rep* 2017; 15:4015–20.
- [28] Zotelli VL, Grillo CM, de Sousa Mda L. Nausea control by needling at acupuncture point Neiguan (PC6) during an intraoral impression-taking procedure. *J Acupunct Meridian Stud* 2014;7:318–23.
- [29] Sena ES, Currie GL, McCann SK, et al. Systematic reviews and meta-analysis of preclinical studies: why perform them and how to appraise them critically. *J Cereb Blood Flow Metab* 2014;34:737–42.
- [30] Murphy SP, Murphy AN. Pre-clinical systematic review. *J Neurochem* 2010;115:805.
- [31] Tian Y, Hu H, Zhang Y, et al. Zusanli (ST36) acupoint injection for acute diarrhea in children under 5 years old: a protocol of systematic review and meta-analysis of randomized clinical trials. *Medicine (Baltimore)* 2019;98:e16949.